Parental care results in a greater mutation load, for which it is also a phenotypic antidote

Sonia Pascoal[1], Hideyasu Shimadzu[2,3,4], Rahia Mashoodh[1, 5*] and Rebecca M. Kilner[1**]

[1]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, U.K.

[2]Department of Mathematical Sciences, Loughborough University, Leicestershire, LE11 3TU, UK

[3]Graduate School of Public Health, Teikyo University, Tokyo, Japan

[4]Department of Data Science, Kitasato University, Kanagawa, 252-0373, Japan

[5] Centre for Biodiversity and Environmental Research, Department of Genetics, Evolution and Environment, University College London, Gower Street, WC1E 6BT, London, UK

*co-corresponding author: r.mashoodh@ucl.ac.uk

** corresponding author: rmk1002@cam.ac.uk

Orcid IDs

Sonia Pascoal	0000-0002-8249-984X	Sonia.pascoal@bioresource.nihr.ac.uk
Hideyasu Shimadzu	0000-0003-0919-8829	H.Shimadzu@lboro.ac.uk
Rahia Mashoodh	0000-0003-3065-9044	
Rebecca Kilner	0000-0003-1159-0758	

Keywords: social evolution, mutation load, extinction rate

Abstract

Benevolent social behaviours, such as parental care, are thought to enable mildly deleterious

mutations to persist. We tested this prediction experimentally using the burying beetle

Nicrophorus vespilloides, an insect with biparental care. For 20 generations, we allowed

replicate experimental burying beetle populations to evolve either with post-hatching care

('Full Care' populations) or without it ('No Care' populations). We then established new

lineages, seeded from these experimental populations, which we inbred to assess their

mutation load. Outbred lineages served as controls. We also tested whether the deleterious

effects of a greater mutation load could be concealed by parental care by allowing half the

lineages to receive post-hatching care, while half did not. We found that inbred lineages from

the Full Care populations went extinct more quickly than inbred lineages from the No Care

populations – but only when offspring received no post-hatching care. We infer that Full Care

lineages carried a greater mutation load, but that the associated deleterious effects on fitness

could be overcome if larvae received parental care. We suggest that the increased mutation

load caused by parental care increases a population's dependence upon care. This could

explain why care is seldom lost once it has evolved.

Introduction

Classical population genetics models imagine that populations attain an equilibrium level of genetic variation known as mutation-selection balance [*e.g.* 1,2]. New genetic mutations arise spontaneously, through diverse mechanisms, and increase genetic variation in the population [*e.g.* 3,4]. However, since the majority of new mutations yield mildly deleterious phenotypes [*e.g.* 3,4] they are quickly purged by natural selection. Mutation-selection balance is theoretically achieved when the rate of input of new genetic variants through spontaneous mutation is perfectly balanced by the rate of their elimination by selection [*e.g.* 1,2].

Social behaviour can, in principle, play a key role in modulating mutation-selection balance [e.g. 2,5,6] and thence influences the extent of standing genetic variation within a population. This is particularly true for social activities that create a more benign environment by enhancing access to resources, or reducing exposure to pathogens, or yielding elaborate architecture that protects the inhabitants from the wider world. Actions like this are relatively commonplace in the many bird, mammal and insect species that provide parental care, or interact cooperatively in other ways [7].

There are three different mechanisms by which parental care, or indeed other types of cooperative interaction [5], might influence mutation-selection balance [6]. The environmental stress hypothesis focuses on the way in which the social environment potentially modulates the phenotypic expression of a mutation. It suggests that in a more benign social environment, any deleterious effect of the mutation on the phenotype might generally be less severe [8,9]. The net effect is that selection against the mutation is relaxed, and it persists. The compensation hypothesis differs slightly by proposing that the deleterious effect of the mutation is expressed, but then fully compensated by parental care so that it is undetectable phenotypically once offspring become independent of their parents. For example, parents may compensate for a mutation that causes offspring to exhibit low growth by increasing the rate at which they provision young [10,11]. Again, the net effect is that selection against the mutation is relaxed, and it persists. A third possibility is that a benign social environment does not directly influence the phenotype expressed but instead relaxes selection generally by buffering against the stressors in the environment that are otherwise a source of natural selection [12,13], such as attack by predators or pathogens or scarce

nutrition. In this more benign environment, more diverse genetic variants can persist [4,6].

Although there is empirical evidence that is consistent with each of these hypotheses within a

single generation [6,13], the cumulative effects over the generations are much less well

understood.

Furthermore, where the correlation between cooperative social behaviour and genetic

variation has been analysed before, usually in more complex animal societies, there exist

additional factors that can independently perturb the mutation-selection balance [5]. For

example, animals that breed cooperatively also tend to produce fewer, larger offspring. This

life history strategy is known to reduce genetic diversity [14] and could potentially oppose, or

even conceal, any increases in genetic variation that are due to cooperation buffering the

effects of natural selection. Cooperative animal societies are also commonly associated with a

high incidence of reproductive skew. Since only a few dominant individuals are typically able

to reproduce, the effective population size is greatly reduced [5]. This can lead to a reduction

in the efficiency of natural selection and a greater influence of genetic drift [4], potentially

confounding any increases in genetic variation that are due solely to relaxed selection.

Similarly, animal societies typically comprise related individuals that derive kin-selected

benefits from their cooperative social interactions. Theoretical analyses have shown that kin

selection acts more weakly than direct selection [15]. Consequently, loci under kin selection

are predicted to harbour more sequence variation than loci under direct selection [15].

To bypass some of these confounding difficulties, we investigated the effect of parental care

on genetic variation [12]. Our experiments focused on replicate laboratory populations of

burying beetles *Nicrophorus vespilloides*, which we evolved under sharply contrasting levels

of parental care, for several generations. We then compared the magnitude of the mutation

load between populations. Comparing populations within species allowed us to eliminate any

confounding effects of kin selection, and offspring size or number, on genetic diversity [14].

Focusing on parental care further eliminated confounding effects that could be due to

reproductive skew.

Burying beetles breed on the body of a small dead vertebrate [16], which the parents jointly

convert into a carrion nest by removing the fur or feathers, rolling the flesh into a ball,

covering it with anti-microbial anal exudates, and burying it. This is pre-hatching parental

care [17]. After hatching, parents also guard and feed larvae and maintain the carrion nest to

prevent putrefaction, though larvae can survive in the lab with no post-hatching care at all

[18]. In two of our evolving populations, larvae were able to receive both pre-hatching and

post-hatching parental care (these were called the 'Full Care' populations) while in two other

populations we prevented parents from supplying any post-hatching care by removing them

before the larvae hatched, after the carrion nest was complete (these were called the 'No

Care' populations). During the first 20 or so generations of experimental evolution, No Care

populations rapidly adapted to a life without parental care [19], through divergent phenotypic

change in both larval [*e.g.* 20]) and parental [*e.g.* 17] traits.

To determine whether parental care causes deleterious genetic variation to accumulate over

the generations, we inbred sub-populations, each derived from the replicate experimental

evolving populations, for 8 successive generations (we called this The Evolutionary History

Experiment). For these 8 generations, we measured the extent to which inbreeding reduced

measures of reproductive success in comparison with control outbred populations. To

determine whether parental care could temper the rate of extinction (as implied by [13]), in

half of all our treatments parents were allowed to provide care after their offspring hatched,

while in the remainder they were prevented from supplying post-hatching care. This

generated 8 different treatments in total (see Figure 1 for the design of the Evolutionary

History Experiment).

We used the data from the Evolutionary History Experiment, to test whether inbred lineages

derived from Full Care evolving populations had lower survival than equivalent inbred

lineages from the No Care evolving populations (reflecting a greater mutation load). We also

investigated whether a supply of post-hatching care modulated the survival of the inbred

lineages. The outbred populations acted as a control treatment for each test.

Methods

Nicrophorus vespilloides **natural history.**

The common burying beetle *N. vespilloides* breeds on a small dead vertebrate (like a songbird

or mouse). The larvae hatch from eggs laid nearby in the soil and crawl to their carrion nest,

which they can feed upon themselves [16]. Once at the carcass, larvae receive post-hatching

biparental care. Parents supply fluids to their offspring through oral trophallaxis, and defend

their brood and the carrion nest from attack by predators, microbes and rival beetles [16]. The

duration and extent of post-hatching care are highly variable, however. For example, when

wild beetles are brought into the lab to breed, roughly 5% of larvae receive no post-hatching

care at all, yet larvae can still survive to become reproductively competent adults [*e.g.* 18,21].

Within roughly a week of hatching, the larvae complete development and at this point (which

we refer to as 'dispersal'), they start to crawl away from the scant remains of the carcass to

pupate in the soil. The parents, meanwhile, fly off in search of a new carcass.

Experimental evolution

The experimental populations used in this work have been described in detail elsewhere [*e.g.*

17,20]. In brief, we established a large founding population of *N. vespilloides* by

interbreeding wild-caught individuals from four different woodlands. This was then divided

into four experimental populations. In two populations, larvae experienced 'Full Care' at each

generation, with both parents staying in the breeding box throughout the breeding bout had

the opportunity to provide post-hatching care as well as pre-hatching care. We have

previously shown that when parents are given the opportunity to provide post-hatching care,

more than 94% actually supply care [21]. In the other two 'No Care' populations, parents

engaged in pre-hatching care but at each generation they were removed from the breeding

box around 53 h after they were paired, so that they never interacted with their larvae. The

work reported here began when these populations had been exposed to 20 generations of

experimental evolution under these contrasting regimes of care.

Evolutionary History Experiment

Preparatory common garden generation

The experiment began by taking individuals from the four evolving populations (Full Care

replicated twice and No Care replicated twice) and exposing them, within each population, to

a common garden Full Care environment for one generation (N = 60 pairs for each No Care

population (to counter-balance the slightly lower breeding success caused by the No Care

environment) and N = 50 pairs for each Full Care population). In this way, we minimised any

potentially confounding transgenerational effects prior to starting the Evolutionary History

Experiment.

Overview (see Figure 1)

Broods from the common garden generation were used to seed new experimental lineages.

Half the lineages derived from the Full Care populations (FC_{POP}) while the other half derived

from the No Care populations (NC_{POP}). From Generation 1 onwards, half of the experimental

lineages were exposed to continuous inbreeding (full-sibling crosses) for up to 8 generations

(by which point all the inbred lineages had gone extinct) (N = c. 45 crosses per treatment at

Generation 1). The remaining experimental lineages were outbred in identical conditions to

provide a control baseline for comparison with the inbred lineages (N = c. 35-40 crosses per

treatment, per generation). Half of all inbred lineages, and half of the outbreeding lineages,

were allowed to provide post-hatching care for their young (Full Care environment). In

remaining lineages, parents were removed 53h after pairing and so were unable to provide

any post-hatching care (No Care environment). The experiment therefore had a 2 x 2 x 2

design, with 8 treatments in all (Full Care versus No Care evolving population of origin;

Inbred versus Outbred; Full Care environment versus No Care environment), with each

treatment replicated twice due to replicate Full Care and No Care populations (see Figure 1

for full overview of the design).

Detailed methods

Beetle maintenance was carried out following standard protocols [19]. Briefly, adult beetles

were kept individually in plastic boxes (12 x 8 x 6cm) filled with moist soil and fed twice a

week with raw beef mince. Adults were bred at 2-3 weeks post-eclosion in a breeding box

(17 x 12 x 6cm) with soil and a mouse carcass (11-13 g for all treatments except for the

individuals derived from the Full Care lines, that were outbred under Full Care conditions (8-

14 g)). To ease the considerable burden of work, data for broods in this treatment were

collected from the ongoing experimental evolution lines in the laboratory. Carcass size was

included, where appropriate, as a factor in the statistical analyses (see below).

For the inbreeding treatments, we paired full siblings (one pair per family) whereas for the

outbreeding treatments we paired males and females at random and did not pair siblings or

cousins. Each pair was given a breeding box with a dead mouse sitting on soil, and the

breeding boxes were placed in a dark cupboard to simulate natural underground conditions.

For broods assigned to a No Care environment, parents were removed around 53 h after

pairing. Eight days after pairing (which is when the larvae have completed their development

and start to disperse away from the carcass) we scored two standard measures of reproductive

success in burying beetles [17]: brood success (fail = no larvae produced; success = some

larvae produced) and brood size at dispersal. Larvae were then placed into cells (2 x 2 x 2cm)

in an eclosion box (10 x 10 x 2cm), with one eclosion box per brood, which was filled with

soil until larvae had developed into sexually immature adults (about 18 days after dispersal).

At this point, adults were transferred to individual boxes until they reached sexual maturity

roughly 2 weeks later. Both the eclosion boxes and the individual boxes were kept on shelves

in the laboratory at 21°C on a 16L:8D hour light cycle.

Statistical Analyses

All statistical tests were conducted in R version 3.5.1 [22] using the base 'statistics' and

'survminer' [23] R packages . Data handling and visualisation were carried out using the

'tidyverse' [24]. Model diagnostics were checked visually. All data and code presented in the

manuscript is available at this link [25].

Survival of inbred lineages derived from Full Care versus No Care populations, with and

without post-hatching care

To determine the effect of evolutionary history (*i.e.* derived from a No Care evolving

population or from a Full Care evolving population), and current care environment (*i.e.*

experienced No Care or Full Care) on the survival of the different lineages across

generations, we fit accelerated time hazard models with a log-logistic distribution using the

'survival' R package [24]. Carcass weight and block were included as covariates. A lineage

was considered to be extinct if it did not survive to reproduce in the subsequent generation.

We subsequently ran post-hoc analyses separately for the No Care and Full Care current care

environments to examine any interactions between evolutionary history and the current care

environment. We additionally used the non-parametric Kruskal Wallis test to determine if

median survival times of each inbred lineage differed, by comparing the effect of

evolutionary history in separate analyses, one for each current care environment.

The greatest decline in survival occurred in the first generation of inbreeding, so we

examined this generation in greater detail. Using binomial generalised linear models (GLM),

we tested the effect of evolutionary history, current care environment, and inbreeding

condition (*i.e.* inbred or outbred) on brood success. Models were fit with brood success as a

Bernoulli response with a complementary log-log link function. We defined brood success at

dispersal in the following way: broods that produced at least one larva that survived to breed

were defined as successful (score = 1) (following [17, 19]) whereas those that did not produce

any surviving young were classified as failures (score = 0). We subsequently ran analyses

separately for the inbreeding and outbreeding conditions to examine any interactions between

evolutionary history and the current environment. We included block and carcass weight as

covariates to ensure any effects we detected occurred over and above any variation in these

variables.

Results

Survival of inbred lineages derived from Full Care versus No Care populations, with and

without post-hatching care

Whilst all inbred lineages in our experiments eventually went extinct, outbred lineages were

still reproducing successfully at the point at which the experiment was terminated (Figure 2).

In general, a No Care current environment caused particularly rapid extinction of inbred

lineages (Figure 2A; Table 1). For the inbred lineages, there was an interaction between the

evolutionary history of a population and the extent of current post-hatching care received

(Table 1). To explore the source of this interaction we ran analyses of each current care

environment separately. This revealed that inbred lineages seeded from the Full Care

evolving populations went extinct more rapidly than inbred lineages seeded from the No Care

evolving populations when care was absent (Beta=0.20 [0.05-0.36], *p*<0.01; Figure 2A).

However, when parents supplied post-hatching care, this difference in survival between

inbred lineages was no longer apparent (Beta=-0.01 [-0.13-0.10], *p*=0.85; Figure 2A).

Indeed, lineages seeded from the Full Care evolving populations reached 50% extinction one

generation sooner under a No Care environment than inbred lineages seeded from the No

Care evolving populations (non-parametric Kruskal Wallis test: *H*(1)=4.59, *p*=0.03;

Supplementary Table 1).

Analysis of Generation 1

The greatest drop in lineage survival occurred in the first generation of inbreeding (Figure

2A), so next we compared lineages by focusing on this generation alone. In general, we found

that outbred populations had higher brood success than inbred populations in Generation 1

(Figure 3, Tables 2,3). Within the inbred lineages, brood success was markedly lower in a No

Care current environment but there was interaction with evolutionary history (Figure 3, Table

2). Post-hoc analyses indicated that within the No Care current environment, inbred lineages

derived from the Full Care evolving populations had lower breeding success than inbred

lineages derived from the No Care evolving populations (Figure 3, Table 2).

In post-hoc analyses, we split the data collected in Generation 1 by the current level of care

supplied, to be able to examine the effect of evolutionary history in more detail. In support of

our prediction, we found that inbred families derived from the Full Care populations had

lower brood survival than inbred families drawn from the No Care populations (Beta=1.12

[0.49-1.80], z=3.42, *p*<0.001) – though only when broods were raised in a No Care current

environment. No equivalent differences were observed in the Full Care current environment

(Beta=0.20 [-1.4,1.9], z=0.25, *p*=0.80). For the outbred families, the evolutionary history of

the lineage had no effect on breeding success, though broods were in general less successful

when they received no post-hatching care (Table 3).

Discussion

Burying beetles care for their offspring by making a nest for them to inhabit during

development, providing them with plentiful carrion to feed upon, feeding them via oral

trophallaxis and defending them from attack by rival microbes and animals [16]. Our

experiments suggest that the supply of post-hatching care is sufficient to perturb the

mutation-selection balance – as predicted generally by previous work [2,3,5,6,13].

We infer that when parents provided care then diverse genetic variants were able to persist,

just as previous work has shown [13]. Consequently, after 20 generations of experimental

evolution in these contrasting environments, the Full Care evolving populations carried a

greater mutation load than the No Care populations. This finding is independently supported

by genetic data presented in a companion paper [26] which uses SNPs to quantify the extent

of genetic variation in the two types of experimental population. We assume that the reduced

survival of inbred lineages derived from the Full Care populations is caused by their greater

level of genetic diversity, which presumably includes mildly deleterious mutations. It is

possible that epigenetic differences between the populations could have contributed to this

effect as well [27].

We infer that natural selection against mildly deleterious mutations is relaxed when parents

supply care, and that this contributed to a greater mutation load in the Full Care evolving

populations than in the No Care evolving populations. However, we cannot deduce from our

experimental design which of the hypotheses outlined in the Introduction is responsible for

relaxing selection. It could be that phenotypic expression of any mildly deleterious mutations

was modulated by parental care (as proposed by both the Environmental Stress hypothesis

and the Compensation hypothesis; [6]) or it could be that forces of natural selection from the

wider environment were buffered by parental care [12,13].

The difference in the survival of inbred lineages between those derived from No Care and

Full Care was especially pronounced during the first generation of inbreeding, and most

readily detectable when inbred individuals were prevented from supplying care. This suggests

that some of the additional mutations present in the Full Care populations were recessive and

/ or only mildly deleterious [3]. Given the relatively short timeframe of this experiment, we

presume that these mutations were present in the founding populations of wild-caught beetles

but were removed from the No Care populations by selection acting more strongly against

them. In this sense, our findings are similar to previous work on *Tribolium* which found that

deleterious genetic variation was purged when populations were exposed experimentally to

more intense sexual selection [28].

Although it is now well-understood why individuals evolve cooperative behaviour, the

mechanisms that cause cooperation to persist and diversify remain relatively unclear [29].

Recent theoretical work suggests that positive feedback cycles could play a key role in

entrenching cooperation, following its initial evolution [30]. Cooperative social interactions

facilitate the transfer of beneficial microbes, for example, upon which social partners might

then become dependent over evolutionary time, ensuring that cooperation must persist [e.g.

31-34]. Likewise, cooperative interactions can promote the division of labour between social

partners, causing a degree of interdependence that ensures cooperation must continue [35].

Our results, together with those obtained by Pilakouta et al. 2015 [13], suggest a third

mechanism through which cooperation can become entrenched, hinted at originally by Crow

1966 [2]. We have shown that parental care creates a problem (increased mutation load: our

results) for which it is also the solution (enhanced survival of all genetic variants: [13], our

results). By relaxing selection, parental care causes an increase mutation load which increases

the population's dependence upon care. Care ensures that the diverse genetic variants, whose

existence it has facilitated, are able survive until the end of development. This could explain

why parental care has evolved more frequently than it has been evolutionarily lost [12]. As

Crow 1966 [2] put it: 'there is no turning back…A return to the original conditions leads to

the immediate full impact of all the mutants that have accumulated during the period of

improved environment". In principle, this reasoning can be extended to any form of

cooperation that relaxes selection. Indeed, Crow 1966 [2] made the argument originally in the

context of environmental improvements in human societies and their effect on genetic

variation. Consistent with his predictions, recent comparative genomic analyses have

revealed a greater incidence of genetic pathologies in western industrialised populations than

in traditional, pre-industrial human societies which are more exposed to natural selection

[3,36-39].

Finally, we have focused on the immediate effects of parental care on genetic variation, but

the longer-term consequences are still unclear and need not match the effects seen in the

short-term. For example, although greater intensity of intrasexual selection is beneficial in the

short term, because it purges deleterious mutations from the population [28], in the longer run

more intense intrasexual selection can make lineages more prone to extinction [40]. This

might be due to a lack of beneficial genetic diversity. Likewise, although parental care

enables mildly deleterious mutations to persist in the short-term, perhaps in the longer-term it

builds up genetic diversity that could be beneficial and underpin rapid evolution, especially if

environmental conditions change suddenly, or if mutations promote novelty through

compensatory evolution [22]. In future work, it would be interesting to isolate the longer-

term effects of parental care on genetic diversity and the effects it might have on the

evolutionary resilience of wild populations in a changing world [41].

References

1. Wright, S. 1960. On the number of self-incompatibility alleles maintained in equilibrium by a given mutation rate in a population of a given size: a reexamination. *Biometrics* **16**, 61-85.

2. Crow JJ. 1966 The quality of people: Human evolutionary changes. *BioScience* **16,** 863-867.

3. Kondrashov A. 2017 *Crumbling Genome: The impact of deleterious mutations on human populations.* Wiley Online Library 2017.

4. Baer CF, Miyamoto MM, Denver DR. 2007 Mutation rate variation in multicellular eukaryotes: causes and consequences. *Nat. Rev. Gen.* **8,** 619-631. (doi:10.1038/nrg2158)

5. Rubinstein DR, Ågren JA, Carbone L, Elde NC, Hoekstra HE, Kapheim KM, Keller L, Moreau CS, Toth AL, Yeaman A et al. 2019 Coevolution of genomic architecture and social behavior. *Trends Ecol. Evol.* **34,** 844-855. (doi:10.1016/j.tree.2019.04.011)

6. Snell-Rood EC, Burger M, Hutton Q, Moczek AP. 2016 Effects of parental care on the accumulation and release of cryptic genetic variation: review of mechanisms and a case study of dung beetles. *Evol. Ecol.* **30**, 251-265. (doi: 10.1007/s10682-015-9813-4)

7. Davies NB, Krebs JR, West SA. 2012 *An Introduction to Behavioural Ecology* (Wiley-Blackwell, Oxford).

8. Kondrashov AS, Houle D. 1994. Genotype-environmental interactions and the estimation of the genomic mutation rate in *Drosophila melanogaster. Proc. R. Soc. B* **258,** 221-227. (doi: 10.1098/rspb.1994.0166)

9. Martin G, Lenormand T. 2006 The fitness effect of mutations across environments: a survey in light of fitness landscape models. *Evolution* **60,** 2413-2427. (doi:10.1111/j.0014-3820.2006.tb01878.x)

10. Lock JE, Smiseth PT, Moore PJ, Moore, AJ. 2007 Coadaptation of prenatal and postnatal maternal effects. *Am. Nat.***170,** 709-718. (doi:10.1111/j.0014-3820.2006.tb01878.x)

11. Monteith KM, Andrews C, Smiseth PT. 2012. Post-hatching parental care masks the effects of egg size on offspring fitness: a removal experiment on burying beetles. *J. Evol. Biol.* **25,** 1815-1822. (doi: 10.1002/ece3.1876)

12. Royle NJ, Smiseth PT, Kölliker M. 2012 *The Evolution of Parental Care* (Oxford
University Press, Oxford).

13. Pilakouta N, Jamieson S, Moorad JA, Smiseth PT. 2015. Parental care buffers against
inbreeding depression in burying beetles. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 8031-
8035. (doi:10.1073/pnas.1500658112)

14. Romiguier J, Gayrai P, Ballenghien M, Bernard A, Cahais V, Chenuil A, Chiari Y,
Dernat R, Faivre N, Loire E et al. 2014 Comparative population genomics in animals
uncovers the determinants of genetic diversity. *Nature* **515,** 261-263.
(doi:10.1038/nature13685)

15. Linksvayer TA, Wade MJ. 2016 Theoretical predictions for sociogenomic data: the
effects of kin selection and sex-limited expression on the evolution of social insect
genomes. *Front. Ecol. Evol.* **4,** 65. (doi: 10.3389/fevo.2016.00065)

16. Scott MP. 1998 The ecology and behavior of burying beetles. *Annu. Rev. Entomol.*
43, 595–618. (doi: 10.1146/annurev.ento.43.1.595)

17. Duarte A, Rebar D, Hallett AC, Jarrett BJM, Kilner RM. 2021 Evolutionary change in
the construction of the nursery environment when parents are prevented from caring
for their young directly. *Proc. Natl. Acad. Sci. U.S.A.* **48,** e2102450118. (doi:
10.1073/pnas.2102450118)

18. Eggert AK, Reinking M, Müller JK. 1998 Parental care improves offspring survival
and growth in burying beetles. *Anim. Behav.* **55**, 97–107. (doi:
10.1006/anbe.1997.0588)

19. Schrader M, Jarrett BJM, Kilner RM. 2015 Using experimental evolution to study
adaptations for life within the family. *Am. Nat.* **185**, 610-619. (doi: 10.1086/680500)

20. Rebar D, Bailey NW, Jarrett BJM, Kilner RM. 2020 An evolutionary switch from
sibling rivalry to sibling cooperation, caused by a sustained loss of parental care.
Proc. Natl. Acad. Sci. U.S.A. **117**, 2544-2550. (doi: 10.1073/pnas.1911677117)

21. Jarrett BJM, Evans E, Haynes HB, Leaf MR, Rebar D, Duarte A, Schrader M, Kilner
RM. 2018 A sustained change in the supply of parental care causes adaptive evolution
of offspring morphology. *Nature Comms* **9,** 1-8. (doi: 10.1038/s41467-018-06513-6)

22. Farkas Z, Kovács K, Sarkadi Z, Kalapis D, Fekete G, Birtyik F, Ayaydin F, Molnár C,
Horváth P, Pál C, Papp, B. 2022 Gene loss and compensatory evolution promotes the
emergence of morphological novelties in budding yeast. *Nature Ecol. Evol.* **6,** 763-
773. (doi: 10.1038/s41559-022-01730-1)

23. Kassambara A, Kosinski M, Biecek P. 2021 survminer: Drawing Survival Curves
using 'ggplot2'. R package version 0.4.9. https://CRAN.R-
project.org/package=survminer

24. Therneau T 2021. A Package for Survival Analysis in R. R package version 3.2-
13,<URL: https://CRAN.R-project.org/package=survival>.

25. doi:10.5061/dryad.b2rbnzsm5

26. Mashoodh R, Trowsdale AT, Manica A, Kilner RM. 2023 Parental care drives the
evolution of molecular genetic variation.
https://www.biorxiv.org/content/10.1101/2023.01.09.523216v1

27. Mashoodh R, Sarkies P, Westoby J, Kilner, RM. 2021. Evolved changes in DNA
methylation in response to the sustained loss of parental care in the burying beetle.
https://www.biorxiv.org/content/10.1101/2021.03.25.436923v2

28. Lumley AJ, Michalczyk L, Kitson JJN, Spurgin LG, Morrison CA, Godwin JL,
Dickinson ME, Martin OY, Emerson BC, Chapman T, Gage MJG 2015. Sexual
selection protects against extinction. *Nature* **522,** 470-473. (doi:
10.1038/nature14419)

29. West SA, Cooper GA, Ghoul MB, Griffin AS 2021. Ten recent insights for our
understanding of cooperation. *Nature Ecol. Evol.* **5,** 419-430. (doi: 10.1038/s41559-
020-01384-x)

30. Franz M, Schülke, Ostner J. 2013 Rapid evolution of cooperation in group-living
animals. *BMC Evolutionary Biology* **13,** 235 (doi:10.1186/1471-2148-13-235).

31. Nalepa CA, Bignell DE, Bandi C. 2001 Detritivory, coprophagy, and the evolution of
digestive mutualisms in Dictyoptera. *Insectes Sociaux* **48**, 194-201
(doi:10.1007/PL00001767).

32. Lombardo MP. 2008 Access to mutualistic endosymbiotic microbes: an
underappreciated benefit of group living. *Behav. Ecol. Sociobiol.* **62,** 479-497. (doi:
10.1007/s00265-007-0428-9)

33. Ezenwa VO, Archie EA, Craft ME, Hawley DM, Martin LB, Moore J, White L. 2016
Host behaviour – parasite feedback: an essential link between animal behaviour and
disease ecology. *Proc. R. Soc. B.* **283,** 20153078. (doi: 10.1098/rspb.2015.3078)

34. Dillard J, Benbow ME. 2020 From symbionts to societies: how wood resources have
shaped insect sociality. *Front Ecol. Evol*. **8,** 173. (doi: 10.3389/fevo.2020.00173)

35. Cooper GA, West SA. 2018 Division of labour and the evolution of extreme
specialisation. *Nature Ecol. Evol.* **2,** 1161-1167. (doi:10.1038/s41559-018-0564-9)

36. Lynch M. 2016. Mutation and human exceptionalism: Our future genetic load.
Genetics **202**, 869-875. (doi: 10.1534/genetics.115.180471)

37. You W, Henneberg H. 2017 Cancer incidence increasing globally: The role of relaxed
natural selection. *Evol. Appl.* **11,** 140-152. (doi: 10.1111/eva.12523).

38. Aris-Brosou S. 2019 Direct evidence of an increasing mutational load in humans.
Mol. Biol. Evol **36,** 2823-2829. (doi: 10.1093/molbev/msz192)

39. Saniotis A, Hennberg M, Mohammadi K. 2021 Genetic load and biological changes
to extant humans. *J. Biosoc. Sci.* **53,** 639-642. (doi: 10.1017/S0021932020000413)

40. Martins MJF, Puckett TM, Lockwood R, Swaddle JP, Hunt G. 2018 High male sexual
investment as a driver of extinction in fossil ostracods. *Nature* **556,** 366-369.
(doi:10.1038/s41586-018-0020-7).

41. Balmford A. 1996 Extinction filters and current resilience: the significane of past
selection pressure for conservation biology. *Trends Ecol. Evol.* **11,** 193-196. (doi:
10.1016/0169-5347(96)10026-4)

Acknowledgements

We thank Sue Aspinall and Chris Swannack for helping with beetle maintenance. We also thank Benjamin Jarrett and Darren Rebar for collecting data from the evolving populations. We are indebted to Allen Moore, Locke Rowe and anonymous reviewer for constructive and insightful comments on an earlier draft of this paper.

Funding statement: This project was supported by a Consolidator's Grant from the European Research Council (310785 Baldwinian_Beetles), by a Wolfson Merit Award from the Royal Society, The Leverhulme Trust (RPG-2018-232) and The Isaac Newton Trust (18.23(q)), each to RMK. RM was supported by a Biotechnology and Biological Sciences Research Council Future Leaders Fellowship (BB/R01115X/1). HS was partially supported by the Japan Society for the Promotion of Science (KAKENHI Grant Number: JP19K21569).

Author contributions: Conceived the idea and designed the Evolutionary History Experiment: RMK, SP; Performed the Evolutionary History Experiment: SP; Collected data from Experimentally evolving populations: RM; Analysed the data: RM (analyses presented here), HS (initial analyses of data); Wrote the manuscript: RMK, RM. All authors discussed the results and commented on the manuscript.

Competing interests: The authors declare no competing financial or nonfinancial interests.

Data and materials availability: Data and code needed to evaluate the conclusions in the paper are available from https://github.com/r-mashoodh/nves_MutationLoad.

 TABLES

Table 1. Summary of accelerated failure time hazard model estimates for inbred lineage

success in the Evolutionary History Experiment, for Inbred populations only. For each

analysis we tested whether brood success was predicted by the type of population in which

they Evolved (*i.e.* whether families were derived from the No Care or Full Care evolving

populations) and Current Environment (*i.e.* whether families experienced No Care or Full

Care in the current generation). Carcass weight and Block were included as covariates (see

Figure 1 for experimental design).

	Beta	95% CI[1]	p-value
Evolutionary History			
Full Care	—	—	
No Care	-0.01	-0.14, 0.12	0.9
Current Environment			
Full Care	**—**	**—**	
No Care	**-0.86**	**-1.0, -0.73**	**<0.001**
Block			
1	—	—	
2	-0.01	-0.11, 0.09	0.8
Carcass weight	0.06	-0.02, 0.14	0.2
Evolutionary History * Current Environment			
No Care * No Care	**0.22**	**0.03, 0.41**	**0.026**

[1]CI = Confidence Interval

Table 2. Summary of binomial generalised linear model estimates for brood success in all
treatments in Generation 1 of the Evolutionary History Experiment, predicted by the type of
population in which they Evolved (*i.e.* whether families were derived from the No Care or
Full Care evolving populations), Current Environment (*i.e.* whether families experienced No
Care or Full Care in the current generation) and Breeding Condition (*i.e.* whether families
were inbred or outbred). Carcass weight and Block were included as a covariate for inbred
lineages (see Figure 1 for experimental design).

Characteristic	Beta	95% CI[1]	p-value
Evolutionary History			
Full Care	—	—	
No Care	-0.05	-0.40, 0.30	0.8
Current Environment			
Full Care	—	—	
No Care	**-1.4**	**-1.7, -1.0**	**<0.001**
Breeding			
Inbred	—	—	
Outbred	**0.33**	**0.10, 0.56**	**0.006**
Block	-0.02	-0.24, 0.20	0.9
Carcass weight	0.12	-0.01, 0.25	0.10
Evolutionary History * Current Environment			
No Care * No Care	**0.48**	**0.02, 0.93**	**0.042**

[1]CI = Confidence Interval

Table 3 Summary of binomial generalised linear model estimates for brood success in

Generation 1 of the Evolutionary History Experiment (see Figure 1 for design of the

Evolutionary History Experiment). Models are shown for Inbred and Outbred lineages, which

were analysed separately. 'Evolutionary History' indicates whether lineages were derived

from the No Care or Full Care evolving populations. 'Current Environment' refers to whether

lineages experienced No Care or Full Care in the Evolutionary History Experiment.

'Breeding' indicates whether lineages were inbred or outbred). Carcass weight and Block

were included as a covariates.

Characteristic	Inbred			Outbred		
	Beta	95% CI[1]	p-value	Beta[1]	95% CI[1]	p-value
Evolutionary History						
Full Care	—	—		—	—	
No Care	0.08	-0.38, 0.54	0.7	-0.13	-0.71, 0.45	0.7
Current Environment						
Full Care	—	—		—	—	
No Care	**-1.7**	**-2.2, -1.3**	**<0.001**	**-0.92**	**-1.4, -0.43**	**<0.001**
Block	-0.09	-0.40, 0.22	0.6	0.11	-0.21, 0.45	0.5
Carcass weight	0.20	-0.05, 0.45	0.13	0.05	-0.13, 0.22	0.7
Evolutionary History * Current Environment						
No Care * No Care	**0.64**	**0.02, 1.3**	**0.043**	0.25	-0.46, 0.97	0.5

[1]CI = Confidence Interval

 FIGURE CAPTIONS

Figure 1. Overview of the Evolutionary History Experiment.

Beetles that had evolved in the Full Care (FC_{POP}) or No Care (NC_{POP}) populations for 20

generations were used to seed new experimental lines in the lab. Before inbreeding began, all

beetles from these lines experienced one generation of a Full Care common garden

environment to minimise potentially confounding transgenerational effects. Sequential

inbreeding or outbreeding was then applied for up to eight generations under both Full Care

and No Care environments. n: number of families per population in generation one of

inbreeding (*i.e.* the generation immediately after the common garden Full Care generation).

The design includes two replicate populations organised into Blocks (Block 1 and Block 2)

whose breeding was staggered by one week (to ease the workload of maintaining them). Grey

boxes: data for these two populations were collected from the evolving populations.

Figure 2. Survival of lineages across generations in the Evolutionary History Experiment.

A. Survival curves for **inbred** lineages and associated 95% confidence intervals. **B**. Survival curves

for **outbred** lineages and associated 95% confidence intervals**.** Lineages derived from Full Care

populations (FC_{POP}) are shown in red, lineages derived from No Care populations (NC_{POP}) are shown

in blue. A dashed line indicates the lineage was raised in its native environment ($FC_{POP}FC_{ENV}$ and

$NC_{POP}NC_{ENV}$), a solid line means it experienced the reciprocal current environment ($FC_{POP}NC_{ENV}$ and

$NC_{POP}FC_{ENV}$). Thus, red dashed line = FC derived lineage in NC environment; solid blue line = NC

derived lineage in NC environment; dashed blue line = NC derived lineage in FC environment; solid

red line = FC derived lineage in FC environment.

Figure 3. Breeding success of the eight different experimental lineages in Generation 1 of the

Evolutionary History Experiment.

Predicted brood survival probabilities ± S.E are shown, under both inbreeding and outbreeding.

Lineages derived from the Full Care evolving populations (FC_{POP}) are shown in red, those derived

from the No Care evolving populations (NC_{POP})are shown in blue. The Current Environment refers to

the opportunity for post-hatching care experienced in Generation 1 by each lineage: Full Care (FC) or

No Care (NC).

Figure 1.

Figure 2.

A.

B.

Figure 3.