**RanBP1 plays an essential role in directed migration of neural crest cells during**

**development.**

Elias H Bariga[1,2#], Delan Alassadi[1#], Chiara Mencarelli[3], Roberto Mayor[1]* and Franck

Pichaud[3]*

# Co-first authorship

[1]Cell and developmental Biology, University College London, London, WC1E 6BT, United

Kingdom.

[2]Mechanisms of Morphogenesis Lab, Instituto Gulbenkian de Ciencia, Oeiras, 2780-156,

Portugal.

[3]Laboratory for Molecular Cell Biology, University College London, London, WC1E 6BT,

United Kingdom.

* Corresponding authors: r.mayor@ucl.ac.uk and f.pichaud@ucl.ac.uk

Phone: (+44) 0 207 679 7817

Fax: (+44) 0 207 679 7805

**Running title: RanBP1 regulates neural crest cell migration**

*Keywords:* Nucleocytoplasmic transport, RanBP1, LKB1, Cell migration, Neural crest cells.

**ABSTRACT (180 words)**

Collective cell migration is essential for embryonic development, tissue regeneration and

repair, and has been implicated in pathological conditions such as cancer metastasis. It is, in

part, directed by external cues that promote front-to-rear polarity in individual cells.

However, our understanding of the pathways that underpin the directional movement of cells

in response to external cues remains incomplete. To examine this issue we made use of

neural crest cells (NC), which migrate as a collective during development to generate vital

structures including bones and cartilage. Using a candidate approach, we found an essential

role for Ran-binding protein 1 (RanBP1), a key effector of the nucleocytoplasmic transport

pathway, in enabling directed migration of these cells. Our results indicate that RanBP1 is

required for establishing front-to-rear polarity, so that NCs are able to chemotax. Moreover,

our work suggests that RanBP1 function in chemotaxis involves the polarity kinase

LKB1/PAR4. We envisage that regulated nuclear export of LKB1 through Ran/RanBP1 is a

key regulatory step required for establishing front-to-rear polarity and thus chemotaxis,

during NC collective migration.

**INTRODUCTION**

Polarity is a widespread feature in biology, and the function of many cell types is rooted in

their ability to become polarised. This is the case for neurons, with the soma-axon polarity axis,

for epithelial cells, with the apical (top) – basal (bottom) axis, and for migrating cells, which

can polarise along a front-to-rear axis. While progress continue to be made on our

understanding of the pathways that promote cell polarity, gaps remain, especially considering

how information in a cell's microenvironment can mobilise cellular machineries to generate or

remodel polarity. In previous work using a genetic approach to address this importat issue in

biology, we found that the Ran pathways, which controls nucleocytoplasmic exchange of

macromolecules, is required for radial migration and polarisation of cortical neurons (i.e. axon specification), during cortex development (Mencarelli et al., 2018). During nuclear export, RanBP1 promotes cargoes unloading from the Ran-export complex, which terminates cargo export and allows for the import of Ran, back to the nucleus (Bischoff and Gorlich, 1997). Remarkably, failure in differentiating an axon in cortical neurons deficient for RanBP1 could be rescued by expressing the polarity factor, serine/threonine kinase, LKB1/PAR4 (Mencarelli et al., 2018), a known Ran cargo (Baas et al., 2003; Dorfman and Macara, 2008; Matsuki et al., 2010). This led us to propose that regulated nuclear export of LKB1 is a key step in establishing polarity in cortical neurons, a notion well supported by other studies in these cells and also in epithelial cells (Baas et al., 2004; Barnes and Polleux, 2009; Matsuki et al., 2010; Shelly et al., 2007). Whether a similar RanBP1-LKB1 connection plays a role in directed migration has not been explore so far.

Next to cortical neurons, RanBP1 is also highly transcribed in the eye and in the cranial neural crest cells (NC) (Maynard et al., 2003). In addition, velo-cardio-facial/DiGeorge syndrome, a developmental disease caused by a micro-deletion (22q11.2) uncovering RanBP1 (Dunham et al., 1999; Merscher et al., 2001), includes developmental defects that can be attributed to the NC cells (McDonald-McGinn et al., 2015). Furthermore, NC migration requires LKB1 (Creuzet et al., 2016). These observations prompted us to examine whether the RanBP1-LKB1 pathway plays a role in NC migration . Early on in vertebrate development, the NCs migrate as a collective, and populate the embryo in a stereotyped manner to induce formation on many adult structures and organs, including bones, cartilage, and portions of the heart, for example (Etchevers et al., 2019; Szabo and Mayor, 2018). Collective cell migration can be defined as a cooperative mode of cellular motion whereby group of cells migrate in a highly coordinated manner. This mode of migration is vital for a range of biological processes during

embryogenesis, wound healing and cancer metastasis (Barriga and Mayor, 2019; Mayor and

Etienne-Manneville, 2016; Rorth, 2009, 2011). During collective cell migration, the direction

of migration is set though biochemical or biophysical cues, which are present in the

microenvironment (Barriga and Theveneau, 2020; Rorth, 2009; Shellard and Mayor, 2020).

These cues direct migration by activating signalling downstream of receptors, which in turn

can harness the activity of cell intrinsic pathways to promote directed cell motility.

Here, we examine the function of the RanBP1-LKB1 pathway in NC migration. We find that

RanBP1 is required for NC migration *in vivo*, and our results indicate that this is in part due to

a function in enabling chemotaxis. Furthermore, our results show that while LKB1 knockdown

affects NC migration, overexpression of this kinase can rescue the migration defect induced by

the knockdown of RanBP1. Altogether, our results suggest that RanBP1 requirment in NC

migration depends upon the polarity effector LKB1. Our work supports the idea that regulated

nucleocytoplasmic transport of this kinases plays an important role in cell across cell types.

**RESULTS**

**RanBP1 is required for neural crest cell migration *in vivo***

In order to test RanBP1 function in *Xenopus* NC cells *in vivo,* we microinjected a morpholino

(MO) specifically designed to inhibit RanBP1 expression (RanBP1-MO). The effect of this

morpholino on NC cells migration was analysed by monitoring NC position using *in situ*

hybridisation (ISH) against *snail2*, a marker for migrating NC. In these experiments, one side

was injected with the MO and the pattern of NC was compared to that of the contralateral

control side of the same embryo. We determined that injecting 6.5 ng of RanBP1-MO *per*

blastomere was an optimal concentration whereby we could achieve a strong inhibition of

RanBP1 as shown by western blot analysis (Figure 1A-B), without affecting embryo survival.

While NCs normally migrate by forming stereotypical streams from dorsal to ventral territories, we found that RanBP1-MO treated cephalic NC cells failed to migrate (Figure 1C-E). Instead, they remained as a group at the border of the neural folds, which is the region of the embryo where they are originally induced (Figure 1C and 1E). Rescue experiments using a human RanBP1 (hRanBP1) mRNA (Mencarelli et al., 2018), which cannot be targeted by the MO, confirmed that this phenotype was due to the specific loss of RanBP1 (Figure 1D-E). To ensure that the MO RanBP1 migration phenotype was not due to defects in sanil expression, we made use of other, complementary migratory NC makers, such as *FoxD3*, *Twist* and *Snail2*. These markers confirmed that NC migration was impaired when decreasing the expression of RanBP1 (Figure 2). In order to test wether RanBP1 depletion could have an effect on early NC specification or ectodermal development in general, we analyzed the expression of Pax3, Sox2 and Keratin. This analysis showed that RanBP1-MO does not affect the expression of early neural fold (Figure 3A,G) or early NC (Figure 3B-D,G) markers. Morevover, RanBP1-MO did not affect the expression of epidermal or neural plate markers (Figure 3E-G). To further characterize the RanBP1-MO phenotype in *Xenopus*, we also examined the effect of RanBP1 loss-of-function in craniofacial development, a well-known NC derivate tissue (Barriga et al., 2013). For this purpose, stage 46 larvae were stained with alcian blue to specifically visualise the cartilage. This revealed multiple defects in cranial cartilage morphology on the MO-injected side of the embryo when compared to the non-treated one (Figure 4). Altogether, these results are consistent with a requirement for RanBP1 in NC migration during development.

**RanBP1 is required for NC chemotaxis in response to SDF-1 exposure.**

Next, to examine the possible cause for the NC migration phenotype we observed in RanBP1-MO embryos, we turned to an *ex vivo* chemotaxis assay, making use of beads coated with Stromal Derived Factor 1 SDF-1(Theveneau et al., 2010). In this assay, the beads are fixed

into a fibronectin coated dish and NC migration can be recorder using time-lapse microscopy

(Figure 5A,B and Supplementary Movie S1). Wild type NC migrate towards the SDF-1 bead

as a collective showing high persistent directionality (Figure 5B). In contrast, RanBP1-MO

injected cells failed to chemotax towards SDF-1 (Figure 5B).

In order to gain further insight into this phenotype, we tracked the migrating NC to extract their

trajectories and then calculated how persistent their migration towards SDF-1 was, by

determining their forward migration index. A forward migration index close to 1 indicates

sustained directional cell migration. While control NC efficiently chemotax with fairly straight

trajectories and a forward migration index of ~ 1, we found that RanBP1-MO injected cells

displayed more randomised trajectories with an average forward migration index of ~0.4; a

typical value displayed by clusters that fail to show directionality during migration (Foxman et

al., 1999) (Figure 5C and Supplementary Movie S1). From this, we conclude that RanBP1 is

required for directed migration of NC, in response to SDF-1.

**RanBP1 is required for directed migration**

Failure in chemotaxis could be due to many reasons, including a failure for individual cells to

polarize along the front-to-rear axis. In order to examine this possibility, we tracked

individual NC to extract cell motility parameters including migratory speed, directionality

and circularity (Figure 6A-D and Supplementary Movie 2 and 3). Consistent, with our

previous observations, quantification of RanBP1-MO individual NCs showed that these cells

display slower migratory speed and less directionalities when compared to wild type cells

(Figure 6B-C; Supplementary Movie 2 and 3). In addition, many cells failed to extend

migratory protrusions and displayed a rounder morphology than that of  wild type cells,

which is indicative of poor front-to-rear polarity and defect in NC migration (Figure 5D and

Supplementary Movie 2 and 3). Altogether these results indicate that RanBP1 is required in

NC to support directed migration, in response to SDF-1.

**LKB1 mediates RanBP1 function during neural crest cell migration.**

In cortical neurons, RanBP1 funtion in polaraly is principaly mediated by the kinase LKB1

(Mencarelli et al., 2018). LKB1 is expressed and required in NC cells for their migration

(Creuzet et al., 2016), raising the issue that the RanBP1-LKB1 pathway is at play during this

process. To test this idea, we took advantage of a previously published *Xenopus* LKB1 kinase

dead dominant negative construct (Ossipova et al., 2003). We found that the injection of this

construct leads to a failure in NC migration that resembles that of RanBP1 injected animals

(Figure 6A-C). Then, to determine whether LKB1 could mediate part of RanBP1 function in

NC migration, we performed an epistatic experiment whereby RanBP1-MO was co-injected

with a wildtype version of LKB1. This co-injection of *LKB1* mRNA was sufficient to restore

the defects observed upon knocking down RanBP1 expression (Figure 7D-F). These results

are consistent with LKB1and RanBP1 functioning together to enable directed NC migration.

**DISCUSSION**

Nuclear export of LKB1 is regulated by Ran and its effector RanBP1 (Baas et al., 2003;

Dorfman and Macara, 2008), and is required for polarization of epithelial cells and rat cortical

neurons (Baas et al., 2004; Matsuki et al., 2010; Mencarelli et al., 2018; Shelly et al., 2007).

Here we show that RanBP1 and LKB1 are also required for directed migration of cephalic NCs.

In this context, our results indicate RanBP1 is required for NCs to polarize properly along the

front-to-rear axis, and thus for chemotaxis in response to a source of SDF-1. Altogether, our

work is compatible with a model whereby RanBP1 requirement in NC migration stems from

its function in promoting unloading of LKB1 from the nuclear export complex, essential for

releasing this kinase in the cytosol.

**RanBP1 is required for NC front-rear polarization**

RanBP1 is required in culture cortical neurons for axon formation and *in vivo*, for neuron

migration duing cortex development (Mencarelli et al., 2018). As it is the case in cortical

neurons, RanBP1 is essential for cell viability in NC cells (Data not shown). However, we

were able to calibrate our MO concentration to achieve an hypomorphic situation, whereby

the expression of this factor is not totally inhibited. In this hypomorphic situation, cell

viability is not affected, but we expect that RanBP1-dependent cargo release is less efficient

compared to wildtype cells. In this case, we find a clear requirement for RanBP1 in front-rear

polarity in individual migrating NC. RanBP1 has been associated with velo-cardio-

facial/DiGeorge syndrome {Dunham, 1999 #3338;Merscher, 2001 #3339, whch includes

phenotypes that can be linked to NC biology. Our results thefore raise the possibility that

RanBP1 deficiency contribute to this syndrome by affecting NC migration.

**The RanBP1-LKB1 axis in NC polarity**

Collective NC migration has been shown to be triggered by stiffening of their migratory

substrate {Barriga, 2018 #3327}. As they start migrating, these cells need to integrate several

extrinsic cues to orient their polarity and to move towards their target tissues (Barriga and

Theveneau, 2020; Shellard and Mayor, 2020). One of the most studied mechanisms of NCs

guidance is their chemotaxis in response the chemokine SDF-1, which is mediated by the

CCR4 receptor and the small GTPase Rac1 (Theveneau et al., 2010). However, how CCR4

activation translates into polarization of NC cell is not well understood. LKB1 has also been

proposed to promote NC migration through AMP-activated protein kinase and via

actomyosin regulation through the Rho-dependent kinase ROCK (Creuzet et al., 2016). Our

work confirms LKB1 is required for NC cell migration, and links this key polarity factor to

the nucleocytoplasmic transport pathway, through RanBP1.

LKB1 has been shown to regulate polarity in a range of cell types and systems. This kinase is

essential for enabling asymmetric division of the C. elegans zygote (Watts et al., 2000).  It

has also been shown to promote neuron polarity and dendrite specification by regulating

microtubule stability through the family of MARK-kinases and in particular through the

related  SAD-A/B kinases, and through Golgi regulation (Barnes et al., 2007; Huang et al.,

2014; Matsuki et al., 2010; Mencarelli et al., 2018; Shelly et al., 2007). Morevover, LKB1

regulates apical-basal polarity in epithelial cells (Amin et al., 2009; Baas et al., 2004; Martin

and St Johnston, 2003).  In neurons and human epithelial cells,  LKB1 function in polarity

starts with its nuclear export (Baas et al., 2004; Mencarelli et al., 2018). Our results show that

overexpression of LKB1 can rescue the RanBP1 loss-of-function phenotype.  While we

cannot rule out that LKB1 functions in paralelle to RanBP1 in NC migration, and that

overexpression of this kinase can compensate for the inhibition of RanBP1 expression, we

favor a model whereby RanBP1 function in NC migration is to control the LKB1 availabilty

in the cytosol of the migrating NC cells. In this model, response to SDF-1 stimulation

requires LKB1.

**FIGURE LEGENDS**

**Figure 1: RanBP1 is required for neural crest migration.** (**A**) Diagram depicting the

microinjection strategy. (**B**) Western blot of cephalic *Xenopus* regions enriched in NCs from

stage 21-22 animals, and gel densitometry showing RanBP1 levels in Control and RanBP1-

MO embryos ; histogram represents the media and bars the standard deviation of three

independent experiments, two-tailed t-test ***, P < 0.001 , N = 3 blots from 3 independent

experiments. (**C** and **D**) Dorsal and lateral views of embryos hybridized with a probe against

*snail2* to label migrating NCs. (**C**) a red asterisk marks the injected side of the embryo. Lateral

views show that while NCs from the control side of the embryo migrate normally by forming

dorso ventral streams, RanBP1-MO cells (red asterisk) stay in the dorsal part of the embryo.

(**D**) a red asterisk marks the injected side of the embryo and dorsal as well as lateral views

show that both Control and RanBP1-MO+RanBP1mRNA co-injected NCs migrate normally

by forming dorso ventral streams.  Scale bars represent 200 μm. (**E**) Normalized NC migration

and percentage of embryos displaying the representative phenotypes shown in (**C** and **D**).

Histograms represent the median and bars the standard deviation of three independent

experiments, N= 20 embryos for each condition. two-tailed t-test ***, P < 0.001 and ****, P

< 0.0001.

**Figure 2: Analysis of NC migration.** Embryos were injected as described in Figure 1, and NC

migration was analyzed by ISH against FoxD3 in stage 24 animals (**A**), Twist (**B**) and Slug

(**C**). Lateral views of the same embryo show the uninjected control side and injected side, as

indicated. Inhibition of NC migration is observed in stage 24 animals for the 3 markers upon

RanBP1 depletion. (**D**) Percentage of stage 24 embryos displaying normal NC migration.

Histograms represent the median and bars the standard deviation of three independent

experiments, N= 15 stage 24 embryos for each condition. two-tailed t-test ***, P < 0.001.

**Figure 3: RanBP1 depletion does not affect early NC or ectodermal specification.**  2-cell

stage embryos were injected with a Control MO or RanBP1 MO in one of the two blastomeres.

Embryos were fixed at stage 14 (**A**), 16 (**B-D**) and 18 (**E,F**), and the indicated genes were

analyzed by ISH. The injected side is indicated by an asterisk. No effect on the expression of

the markers was observed upon RanBP1 MO injection. (**G**) Quantification showing the % of

embryos with normal expression. At least 15 embryos were analyzed for each gene/condition.

**Figure 4. Decreasing RanBP1 expression leads to skeleton defects.** (**A-B**) Cartilage labelled

using Alcian Blue in control and RanBP1MO embryos. (**A**) is a wild type embryo and (**B**) one

treated using the RanBP1 MO. At least 10 embryos were analysed for each genotype, and the

picture presented in (**A-B**) are representative of the RanBP1 MO phenotype.

**Figure 5: RanBP1 is required for NC chemotaxis towards SDF-1.** (**A**) Diagram depicting

our chemotaxis assay. (**B**) Clusters of Control or RanBP1-MO injected NCs where their

trajectories have been colour coded as a function of time. The Scale bar represents 50 μm. (**C**)

Centred tracks showing the trajectories of multiple Control or RanBP1 treated NCs. (**D**) Box

plot showing the FMI and speed of migration of Control and RanBP1-MO cells, bars represent

median and wiskers the min and max values; two-tailed Mann–Whitney test \*\*\*, $P < 0.001$;

CI= 95%; $n_{control}$= 43, $n_{RanBP1-MO}$= 68 cells from three independent experiments.

**Figure 6: RanBP1 is required for NC polarity and motility.** (**A**) Drawing showing our single

NCs migration assay. (**B**) Representative images of migrating Control-MO or RanBP1-MO

NCs clusters, showing loss of rear-front polarity. (**C -D**) Parameters for individual cell motility

as obtained from Supplementary Movie 3. Box plots showing cell directionality and speed of

Control and RanBP1-MO cells in (**C**) and max protrusion area as well as circulatiry index in

(**D**). In (**C-D**) bars represent median and wiskers the min and max values. In (**C**), for single cell

directionality two-tailed Mann–Whitney test \*\*\*, $P < 0.0005$; $n_{control}$= 16, $n_{RanBP1-MO}$= 36 cells

from three independent experiments; and for single cell speed two-tailed t-test \*\*\*\*, $P <$

0.0001; CI= 95%; $n_{control}$= 20, $n_{RanBP1-MO}$= 34 cells from three independent experiments. In (**D**),

for max protrusion area two-tailed Mann–Whitney test ****, P < 0.0001; $n_{control}$= 185, $n_{RanBP1-}$

$_{MO}$= 191 protrusions from three independent experiments; and for circulatiry index two-tailed

t-test **, P = 0.0062; CI= 95%; $n_{control}$= 20, $n_{RanBP1-MO}$= 20 cells from three independent

experiments. Scale bar = 20 micron.

**Figure 7: LKB1 expression can rescue the RanBP1-MO migration phenotype.** (**A**)

Diagram showing our injection strategy and (**B**) histogram representing the percentage of

Control or LKB1-KO injected embryos that displayed NC migration defects. (**C-D**) Dorsal and

lateral views of embryos hybridized with a probe against *snail2* to label migrating NCs. (**C**) A

red asterisk marks the injected side of the embryo. Lateral views show that while NC form the

control side of the embryo migrate normally by forming dorso ventral streams, LKB1-KD cells

(red asterisk) remain in the dorsal part of the embryo. (**A**) A red asterisk denotes the injected

side of the embryo and dorsal as well as lateral views show that both Control and RanBP1-

MO+LKB1mRNA co-injected NCs normally migrate by forming dorso-ventral streams. Scale

bars represent 200 μm. (**E**) Normalized NC migration and percentage of embryos displaying

the representative phenotypes shown here. Histograms represent the media and bars the

standard deviation of three independent experiments, N= 20 embryos for each condition. two-

tailed t-test ***, P < 0.001 and ****, P < 0.0001.

**Supplementary Video 1. RanBP1 is required for NC chemotaxis towards SDF-1.**

*ex-vivo* time-lapse of control and RanBP1-MO NC migrating in an SDF-1 chemotaxis assays.

Time-lapse setting was set at 1 picture every 5 min; 35 frames are shown. Frame rate is 7

frames per second.

**Supplementary Video 2. RanBP1 is required for cluster cell spreading.**

*ex-vivo* time-lapse of control and RanBP1-MO NC migrating in a cluster spreading assay.

Time-lapse setting was  set at 1 picture every 5 min; 54 frames are shown.

Frame rate is 7 frames per second.

**Supplementary Video 3. RanBP1 is required for single cell motility.**

*ex-vivo* time-lapse of control and RanBP1-MO NC migrating in a single cell migration assay.

Time-lapse setting was set at 1 picture every 5 min; 31 frames are shown. Frame rate is 7

frames per second.

**EXPERIMENTAL PROCEDURES**

*Xenopus* **manipulation.**

Adult *X. laevis* were maintained in a range of temperatures 14-18 °C.  To obtain embryos we

followed established protocols (Barriga et al., 2013). Briefly, superovulation was induced in

adult female frogs by injecting chorionic gonadotrophin (Intervet), then obtained oocytes were

mixed with a sperm solution for *in vitro* fertilizations. Embryos were staged as previously

described (Hubrecht-Laboratorium (Embryologisch Institut), 1967) and maintained at 14ºC.

All animal experiments were approved by the Biological Service Unit at University College

London and complied with UK Home Office guidelines (Animal Act 1986); or by the Ethics

Committee and the Animal Welfare Body of the Instituto Gulbenkian de Ciencia and by the

Direção Geral de Alimentação e Veterinária (DGAV), Portugal.

***In situ* hybridisation and *in vitro* transcription.**

*In situ* hybridizations and templates for all *in vitro* transcription were generated as previously

described (Barriga et al., 2013; Barriga et al., 2019; Theveneau and Mayor, 2011).

Digoxigenin-labeled RNA probes against *snail2* (Mayor et al., 1995), *twist* (Hopwood et al.,

1989), *FoxD3* (Kelsh et al., 2000), *Sox2* (Kishi et al., 2000) , *Keratin* (Jonas et al., 1985),

*Pax3* (Alkobtawi et al., 2018) was used in this study. In brief,  digoxigenin-labeled probes

against target mRNA were transcribed by using an *in vitro* Transcription System (Promega

P1420) and all mRNAs were transcribed with a mMESSAGE mMACHINE SP6 and T7

Transcription Kits (Thermo-Fisher AM1340 and AM1344, respectively).

**Analysis of neural crest migration *in vivo*:** After imaging embryos subjected to *in situ*

hybridization, the dorso-ventral length of the NC was extracted with the built-in measurement

tool of ImageJ. This length was normalized against the total dorso-ventral length of the embryo

that was also extracted using ImageJ. Further analyses and comparisons were carried out as

described in the Statistical analysis section.

**Morpholino and mRNA injections**.

Injections were performed using glass needles calibrated to deliver 10nL. Briefly, at eight-

cell stage a ventral and dorsal animal blastomeres were injected with: 5ng of a morpholino

designed against *Xenopus RanBP1* (*RanBP1*-MO 5'-GCGGGTCTTCTGTTTCTACAATGTC

-3'), with the same amount of a standard control MO and/or with 1ng of mRNAs encoding for

Lkb1 or RanBP1. Similarly, to fluorescently tag NCs, 250 pg of membrane GFP (mGFP) and

or nuclear RFP (nRFP) were injected, as shown in each figure.

**Neural crests dissection**.

Dissections were performed after (Barriga et al., 2019). In brief, the vitellin membrane was mechanically removed, and embryos were immobilized in a dish containing modelling clay. Then, the epidermis was removed with a hair knife and the same tool was used to explant the NCs. NCs explants were maintained in Danilchik's medium (DFA; 1 mM MgSO4(7H2O), 5 mM Na2CO3, 4.5 mM KGluconate, 53 mM NaCl, 32 mM NaGluconate, 0.1% BSA and 1 mM CaCl2; after dissolving the components the pH was adjusted to 8.3 with Bicine). ENREF_1

**Single cell migration assay**.

To analyse motility at the single cell level, we used an assay in which explants were mechanically disaggregated to obtain single cells, and then platted onnto a dish coated with fibronectin (μ-Dish, 35 mm, Ibidi). NCs migration was recorded by time-lapse microscopy.

**Chemotaxis assay**.

NC response to SDF-1 was assesed after(Theveneau and Mayor, 2011). In brief, acrylic heparin beads (Sigma-Aldrich) were coated with 1 μg/ml Sdf-1 (Sigma-Aldrich) for 1 hour, washed with PBS. Silicon grease was used to fix these SDF-1-coated beads to a fibronectin-coated dish (Ibidi). NCs clusters (tagged with mRFP and or mGFP) were placed in front of the bead and their directional response was recorded by time-lapse.

**Time-lapse imaging**.

For chemotaxis and single cell migration assays, an emage was acquired every 5 min at 18 °C using an upright microscope (DMR XA2, Leica). This microscope was equipped with a motorized stage (Prior Scientific) to facilitate imaging of multiple samples and a high

resolution camera (Orca-5G Hamamatsu). Images acquisition was performed with a 20×

objective (HCX APO L 20×/0.50 W U-V-I FWD = 3.50 mm, Leica) and all microscopes

components were motorised and controlled through a dedicated software, SimplePCI

(Hamamatsu).

*In situ* **hybridization imaging**.

Embryo images were acquired by using 3.2× magnification using a Leica MZ FL III dissecting

microscope equipped with a Leica DFL420 camera, controlled through a Leica IM50 imaging

software.

**Cell tracking, chemotaxis, protrusion area and circularity index**.

The trajectories of individual cells or clusters were established using a standard Manual

tracking plugin for ImageJ and the Chemotaxis Tool plugin from the same software was used

to extract speeds of migration and directionality. Max protrusion area and circularity index

were extracted by using the built-in hand free selection and measurement tools of ImageJ. Data

were further analysed as described in Statistical analysis.

**Image processing**

The ImageJ software was used to generate *z*-stacks, maximum projections and to mount time-

lapse movies. ImageJ and/or Adobe Photoshop was used for linear adjustment of display map

levels, crop for re-sizing and application of pseudocolour. Adobe Illustrator was used to

assemble all figures.

**Statistical analysis**

Each experiment was repeated at least three times. To determine whether data set were

parametric or not we used  Kolmogorov–Smirnov, d'Agostino–Pearson or Shapiro–Wilk test

in Prism7 (GraphPad). When parametric, significance was compared by using unpaired $t$-test

(two-tailed, unequal variances) and in case of non-parametric distribution, significance

between ranks was determined with Mann–Whitney $U$-test (two-tailed). CI was set at 95% in

all cases.

**ACKNOWELDGEMENTS**

We are grateful to the Pichaud lab for discussions related to this study, and the LMCB, CDB

and IGC imaging, and the aquatic animal facility at UCL and IGC for their support. The

authors also thank Prof Jeremy Green for kindly providing the *Xenopus* LKB1 constructs.

This work was funded by an MRC (MC_UU_12018/3), BBSRC (BB/R000697) and Royal

Society grant (Award #181274) to F. Pichaud, and benefited from MRC core funding to the

LMCB, covering access to microscopy (MC_12266B).  E.H. Bariga was funded at UCL

through a BBSRC grant (Grant No: M008517) to R. Mayor. Work in R. Mayor lab is funded

by grants from the Medical Research Council (MR/S007792/1) and BBSRC (M008517).

Work in EHB lab is funded by the European Research Council (ERC) (grant agreement No.

950254), EMBO IG Project Number 4765 ; la Caixa Junior Leader Incoming (94978) and

support provided by Instituto Gulbenkian de Ciencia and Fundaçao Calouste Gulbenkian

(start-up fund I-411133.01).

**REFERENCES**

Alkobtawi, M., Ray, H., Barriga, E.H., Moreno, M., Kerney, R., Monsoro-Burq, A.H., Saint-
Jeannet, J.P., Mayor, R., 2018. Characterization of Pax3 and Sox10 transgenic Xenopus laevis
embryos as tools to study neural crest development. Dev Biol 444 Suppl 1, S202-S208.

Amin, N., Khan, A., St Johnston, D., Tomlinson, I., Martin, S., Brenman, J., McNeill, H., 2009. LKB1 regulates polarity remodeling and adherens junction formation in the Drosophila eye. Proc Natl Acad Sci U S A 106, 8941-8946.

Baas, A.F., Boudeau, J., Sapkota, G.P., Smit, L., Medema, R., Morrice, N.A., Alessi, D.R., Clevers, H.C., 2003. Activation of the tumour suppressor kinase LKB1 by the STE20-like pseudokinase STRAD. Embo J 22, 3062-3072.

Baas, A.F., Kuipers, J., van der Wel, N.N., Batlle, E., Koerten, H.K., Peters, P.J., Clevers, H.C., 2004. Complete polarization of single intestinal epithelial cells upon activation of LKB1 by STRAD. Cell 116, 457-466.

Barnes, A.P., Lilley, B.N., Pan, Y.A., Plummer, L.J., Powell, A.W., Raines, A.N., Sanes, J.R., Polleux, F., 2007. LKB1 and SAD kinases define a pathway required for the polarization of cortical neurons. Cell 129, 549-563.

Barnes, A.P., Polleux, F., 2009. Establishment of axon-dendrite polarity in developing neurons. Annual review of neuroscience 32, 347-381.

Barriga, E.H., Maxwell, P.H., Reyes, A.E., Mayor, R., 2013. The hypoxia factor Hif-1alpha controls neural crest chemotaxis and epithelial to mesenchymal transition. J Cell Biol 201, 759-776.

Barriga, E.H., Mayor, R., 2019. Adjustable viscoelasticity allows for efficient collective cell migration. Semin Cell Dev Biol 93, 55-68.

Barriga, E.H., Shellard, A., Mayor, R., 2019. In Vivo and In Vitro Quantitative Analysis of Neural Crest Cell Migration. Methods Mol Biol 1976, 135-152.

Barriga, E.H., Theveneau, E., 2020. In vivo Neural Crest Cell Migration Is Controlled by "Mixotaxis". Front Physiol 11, 586432.

Bischoff, F.R., Gorlich, D., 1997. RanBP1 is crucial for the release of RanGTP from importin beta-related nuclear transport factors. FEBS Lett 419, 249-254.

Creuzet, S.E., Viallet, J.P., Ghawitian, M., Torch, S., Thelu, J., Alrajeh, M., Radu, A.G., Bouvard, D., Costagliola, F., Borgne, M.L., Buchet-Poyau, K., Aznar, N., Buschlen, S., Hosoya, H., Thibert, C., Billaud, M., 2016. LKB1 signaling in cephalic neural crest cells is essential for vertebrate head development. Dev Biol 418, 283-296.

Dorfman, J., Macara, I.G., 2008. STRADalpha regulates LKB1 localization by blocking access to importin-alpha, and by association with Crm1 and exportin-7. Mol Biol Cell 19, 1614-1626.

Dunham, I., Shimizu, N., Roe, B.A., Chissoe, S., Hunt, A.R., Collins, J.E., Bruskiewich, R., Beare, D.M., Clamp, M., Smink, L.J., Ainscough, R., Almeida, J.P., Babbage, A., Bagguley, C., Bailey, J., Barlow, K., Bates, K.N., Beasley, O., Bird, C.P., Blakey, S., Bridgeman, A.M., Buck, D., Burgess, J., Burrill, W.D., O'Brien, K.P., et al., 1999. The DNA sequence of human chromosome 22. Nature 402, 489-495.

Etchevers, H.C., Dupin, E., Le Douarin, N.M., 2019. The diverse neural crest: from embryology to human pathology. Development 146.

Foxman, E.F., Kunkel, E.J., Butcher, E.C., 1999. Integrating conflicting chemotactic signals. The role of memory in leukocyte navigation. J Cell Biol 147, 577-588.

Hopwood, N.D., Pluck, A., Gurdon, J.B., 1989. A Xenopus mRNA related to Drosophila twist is expressed in response to induction in the mesoderm and the neural crest. Cell 59, 893-903.

Huang, W., She, L., Chang, X.Y., Yang, R.R., Wang, L., Ji, H.B., Jiao, J.W., Poo, M.M., 2014. Protein kinase LKB1 regulates polarized dendrite formation of adult hippocampal newborn neurons. Proc Natl Acad Sci U S A 111, 469-474.

Hubrecht-Laboratorium (Embryologisch Institut), N., P. D. & Faber, J. , 1967. Normal table of Xenopus laevis (Daudin). A systematical and chronological survey of the development from the fertilized egg till the end of metamorphosis. , in: Co., N.-H.P. (Ed.).

Jonas, E., Sargent, T.D., Dawid, I.B., 1985. Epidermal keratin gene expressed in embryos of Xenopus laevis. Proc Natl Acad Sci U S A 82, 5413-5417.

Kelsh, R.N., Dutton, K., Medlin, J., Eisen, J.S., 2000. Expression of zebrafish fkd6 in neural
crest-derived glia. Mech Dev 93, 161-164.
Kishi, M., Mizuseki, K., Sasai, N., Yamazaki, H., Shiota, K., Nakanishi, S., Sasai, Y., 2000.
Requirement of Sox2-mediated signaling for differentiation of early Xenopus neuroectoderm.
Development 127, 791-800.
Martin, S.G., St Johnston, D., 2003. A role for Drosophila LKB1 in anterior-posterior axis
formation and epithelial polarity. Nature 421, 379-384.
Matsuki, T., Matthews, R.T., Cooper, J.A., van der Brug, M.P., Cookson, M.R., Hardy, J.A.,
Olson, E.C., Howell, B.W., 2010. Reelin and stk25 have opposing roles in neuronal
polarization and dendritic Golgi deployment. Cell 143, 826-836.
Maynard, T.M., Haskell, G.T., Peters, A.Z., Sikich, L., Lieberman, J.A., LaMantia, A.S., 2003.
A comprehensive analysis of 22q11 gene expression in the developing and adult brain. Proc
Natl Acad Sci U S A 100, 14433-14438.
Mayor, R., Etienne-Manneville, S., 2016. The front and rear of collective cell migration. Nature
reviews. Molecular cell biology 17, 97-109.
Mayor, R., Morgan, R., Sargent, M.G., 1995. Induction of the prospective neural crest of
Xenopus. Development 121, 767-777.
McDonald-McGinn, D.M., Sullivan, K.E., Marino, B., Philip, N., Swillen, A., Vorstman, J.A.,
Zackai, E.H., Emanuel, B.S., Vermeesch, J.R., Morrow, B.E., Scambler, P.J., Bassett, A.S.,
2015. 22q11.2 deletion syndrome. Nat Rev Dis Primers 1, 15071.
Mencarelli, C., Nitarska, J., Kroecher, T., Ferraro, F., Massey, K., Riccio, A., Pichaud, F.,
2018. RanBP1 Couples Nuclear Export and Golgi Regulation through LKB1 to Promote
Cortical Neuron Polarity. Cell reports 24, 2529-2539 e2524.
Merscher, S., Funke, B., Epstein, J.A., Heyer, J., Puech, A., Lu, M.M., Xavier, R.J., Demay,
M.B., Russell, R.G., Factor, S., Tokooya, K., Jore, B.S., Lopez, M., Pandita, R.K., Lia, M.,
Carrion, D., Xu, H., Schorle, H., Kobler, J.B., Scambler, P., Wynshaw-Boris, A., Skoultchi,
A.I., Morrow, B.E., Kucherlapati, R., 2001. TBX1 is responsible for cardiovascular defects in
velo-cardio-facial/DiGeorge syndrome. Cell 104, 619-629.
Ossipova, O., Bardeesy, N., DePinho, R.A., Green, J.B., 2003. LKB1 (XEEK1) regulates Wnt
signalling in vertebrate development. Nat Cell Biol 5, 889-894.
Rorth, P., 2009. Collective cell migration. Annu Rev Cell Dev Biol 25, 407-429.
Rorth, P., 2011. Whence directionality: guidance mechanisms in solitary and collective cell
migration. Dev Cell 20, 9-18.
Shellard, A., Mayor, R., 2020. All Roads Lead to Directional Cell Migration. Trends Cell Biol
30, 852-868.
Shelly, M., Cancedda, L., Heilshorn, S., Sumbre, G., Poo, M.M., 2007. LKB1/STRAD
promotes axon initiation during neuronal polarization. Cell 129, 565-577.
Szabo, A., Mayor, R., 2018. Mechanisms of Neural Crest Migration. Annu Rev Genet 52, 43-
63.
Theveneau, E., Marchant, L., Kuriyama, S., Gull, M., Moepps, B., Parsons, M., Mayor, R.,
2010. Collective chemotaxis requires contact-dependent cell polarity. Dev Cell 19, 39-53.
Theveneau, E., Mayor, R., 2011. Beads on the run: beads as alternative tools for chemotaxis
assays. Methods Mol Biol 769, 449-460.
Watts, J.L., Morton, D.G., Bestman, J., Kemphues, K.J., 2000. The C. elegans par-4 gene
encodes a putative serine-threonine kinase required for establishing embryonic asymmetry.
Development 127, 1467-1475.
520

# Figure 1: RanBP1 is required for neural crest cell migration.

|  | Control Side | RanBP1-Mo Side |
|---|---|---|
| FoxD3 | | |
| Twist | | |
| Slug | | |

350mm

**A** FoxD3

**B** Twist

**C** Slug

**D**

NC Migration (%)

FoxD3 · Twist · Slug

*** *** ***

□ Control side
■ RanBP1-Mo side

# Figure 4: Decreasing RanBP1 expression leads to skeletal defects.

# Figure 5: RanBP1 is required for chemotaxis.

# Figure 6: RanBP1 is required for cell polarity and motility.

# Figure 7: LKB1 expression can rescue the RanBP1-MO migration phenotype