Optical mapping compendium of structural variants across global cattle breeds

Talenti A.[1], Powell J.[1], Wragg D.[1,2], Chepkwony M.[3,4], Fisch A.[5], Ferreira B.R.[5], Marcadante M.E.Z.[6], Santos I.M.[7], Ezeasor C.K.[8], Obishakin E.T.[9,10], Muhanguzi D.[11], Amanyire W.[11], Silwamba I.[12,13], Muma J.B.[12], Mainda G.[14], Kelly R.F.[1,2], Toye P.[3], Connelley T.[1,15]*, Prendergast J.[1,15]*

[1] The Roslin Institute, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG, United Kingdom
[2] Royal (Dick) School of Veterinary Studies, University of Edinburgh, Roslin, UK
[3] The International Livestock Research Institute, PO Box 30709, Nairobi, Kenya.
[4] Centre for Tropical Livestock Genetics and Health, ILRI Kenya, Nairobi, 30709-00100, Kenya
[5] Ribeirão Preto College of Nursing, University of Sao Paulo, Ribeirão Preto, SP, Brazil
[6] Institute of Animal Science, Agriculture Department of São Paulo Government, Sertãozinho, SP, 14.174-000, Brazil
[7] Ribeirão Preto School of Medicine, University of São Paulo, Ribeirão Preto, SP, 14049-900, Brazil
[8] Department of Veterinary Pathology and Microbiology, University of Nigeria, Nsukka. Enugu State, Nigeria
[9] Biotechnology Division, National Veterinary Research Institute, Vom, Plateau State, Nigeria.
[10] Biomedical Research Centre, Ghent University Global Campus, Songdo, Incheon, South Korea.
[11] School of Biosecurity, Biotechnology and Laboratory Sciences (SBLS), College of Veterinary Medicine, Animal Resources and Biosecurity, Makerere University, P.O Box 7062, Kampala, Uganda
[12] Department of Disease Control, School of Veterinary Medicine, University of Zambia, P.O BOX 32379, Lusaka, Zambia
[13] Department of Laboratory and Diagnostics, Livestock Services Cooperative Society, P.O. BOX 32025, Lusaka, Zambia
[14] Department of Veterinary Services, Ministry of Fisheries and Livestock, Central Veterinary Research Institute, P.O. Box 33980, Lusaka, Zambia,
[15] Centre for Tropical Livestock Genetics and Health, Easter Bush, Midlothian, EH25 9RG, UK

*: Contributed equally

Abstract

Structural variants (SV) have been linked to important bovine disease phenotypes, but due to the difficulty of their accurate detection with standard sequencing approaches, their role in shaping important traits across cattle breeds is largely unexplored. Optical mapping is an alternative approach for mapping SVs that has been shown to have higher sensitivity than DNA sequencing approaches. The aim of this project was to use optical mapping to develop a high-quality database of structural variation across cattle breeds from different geographical regions, to enable further study of SVs in cattle.

To do this we generated 100X Bionano optical mapping data for 18 cattle of nine different ancestries, three continents and both cattle sub-species. In total we identified 13,457 SVs, of which 1,200 putatively overlap coding regions. This resource provides a high-quality set of optical mapping-based SV calls that can be used across studies, from validating DNA sequencing-based SV calls to prioritising candidate functional variants in genetic association studies and expanding our understanding of the role of SVs in cattle evolution.

Background & Summary

Structural variants (SV) are a heterogeneous class of genetic variants involving large fragments of the genome (>50bp)[1]. These variants include genomic insertions and deletions (InDels), inversions, duplications, translocations and more complex rearrangements[2]. Single nucleotide polymorphisms (SNPs) have been the primary focus of studies trying to map genetic loci underlying important cattle phenotypes. However, there are multiple lines of evidence suggesting SVs likely underlie many important cattle traits. As many as 25-29% of all protein truncating events are thought to be caused by an SV in humans[1] and notably, despite being less well studied, SVs have already been tied to key livestock phenotypes. For example, a duplication of the CIITA class II major histocompatibility complex transactivator gene in cattle has been tied to resistance to intestinal nematodes[3] and a 12Kb copy number variant has been linked to mastitis in cattle[4]. Chromosomal translocations and duplications have been linked to skin pigmentation, a phenotype closely tied to environmental adaptation, and SVs across livestock species have been linked to phenotypes such as olfaction or resistance to adenocarcinoma-causing viruses[5]. Importantly SVs are responsible for approximately 5-10 times as many heritable nucleotide sequence differences between individuals than SNPs[6]. Unlike SNPs, that only effect a single basepair, and most often far from coding regions, SVs effect large regions and potentially multiple genes. Consequently, although smaller in number, any given novel SV event is more likely to have a phenotypic consequence.

The two most popular methods used to detect SVs are high-throughput sequencing (HTS) and array comparative genomic hybridisation (aCGH), both of which have been applied to European cattle[7–9], but with few studies performed in other cattle breeds[10–12]. Each technology has advantages and limitations. aCGH, for example, involves measuring binding to probes covering the reference genome, and therefore it can only detect relative copy number changes between sample pairs and cannot for example detect novel insertions. Resolution is also limited. A major advantage of HTS approaches is that theoretically they can detect SVs at base-pair resolution. However, accurate calling of SVs from HTS data has proven to be difficult for a number of reasons including poor reference assemblies, chimeric reads, aligners penalising reads that don't match the reference and the difficulties of sequencing and mapping to repetitive regions. This is exemplified by the generally poor agreement between SV callers even when run across the same samples[13,14]. Approaches using long reads

and *de novo* assembly can still have true positive rates as low as 77%, even when using
simulated data[15].

Optical mapping (OM), a light microscope-based method that labels and physically locates
specific motifs in the genome[16], offers an alternative protocol to accurately detect large SVs.
OM molecules can be consistently hundreds of Kb long, allowing for the detection of complex
rearrangements undetectable using HTS. Despite the limitation of not being able to detect
the actual sequence of the identified SVs, as well as missing smaller SVs, OM has a very high
sensitivity and specificity, allowing for the generation of high-quality catalogues of SVs in
individuals[17]. A study in humans successfully used OM reads to identify SVs in a total of 26
genomes revealing population-specific patterns of structural variation[18].
In this study, we generated the first catalogue of cattle OM data for 18 animals from 9
different global breeds, and three continents, to better characterise common SVs across the
cattle pan-genome. This data is a particularly valuable resource of SVs for the cattle species
to intersect with other datasets, for example, for the validation of SV calls from other
approaches.

Methods

Sample preparation

We selected a set of 18 cattle across 9 divergent European, African and Indian breeds
representative of Indicine, Sanga and Taurine ancestries (Table 1). Blood was collected by
jugular venipuncture into EDTA vacutainers. Somatic recombination in B cells and T cells
means the Ig and TCR loci in these cell types will be highly heterogenous, confounding
accurate reconstruction of germline SVs at these loci from whole blood samples.
Consequently, after the erthyrocyte lysis, monocytes were purified from the leukocytes using
a MACS positive selection protocol with an anti-bovine SIRPα mono-clonal antibody (ILA-
24[19]). Agarose plugs containing $5 \times 10^5 - 1 \times 10^6$ of isolated monocytes were prepared using the
Bionano Blood and cell culture DNA isolation kit (Bionano Genomics, San Diego, US) according
to the manufacturer's instructions and the extracted DNA used for analysis on the Bionano
Saphyr platform to generate ~100X optical mapping coverage of each genome.

Bionano Solve optical mapping processing

OM reads were filtered using the filter_SNR_dynamic.pl script with default parameters included with the Solve workflow, and then processed through the Bionano Solve[20] pipeline (v3.3_10252018) using two different releases of RefAligner to overcome bugs preventing the successful assembly of the reads (version 7915.7989rel and 10330.10436rel). We generated the reference CMAP for the ARS-UCD1.2 genome with the Y chromosome from the 1000 bulls genome project (https://sites.ualberta.ca/~stothard/1000_bull_genomes/) using fa2cmap_multi_color.pl (default options and specifying the DLE1 as enzyme). The resulting data were imported into the Bionano Access (v1.6) software, and single-sample SVs were filtered using the recommended thresholds for SVs generated using Bionano Solve prior to v1.6.0 with the sizes recommended to achieve 90% sensitivity (see https://bionanogenomics.com/wp-content/uploads/2018/04/30110-Bionano-Solve-Theory-of-Operation-Structural-Variant-Calling.pdf): minimum insertion size of 5Kb, minimum deletion size of 5Kb, minimum inversion size of 100Kb, and minimum duplication size of 150kb.

Filtered smap format files were converted to vcf format using smap_to_vcf_v2.py and sorted with bcftools (v1.10.2[21]). The resulting SVs were screened using bcftools and retained if 1) they had successfully been genotyped, 2) their size was >1Kb and 3) their quality was >= 20. The latter filtering largely removed all translocations, duplications, and complex events due to these having either very low (<1) or nil quality values.

We then defined the total amount of non-redundant reference sequence involved in a high-quality deletion. For each deletion, we calculated the central point in the genomic region affected by the SV:

$$Center = \frac{POS + abs(CIPOS) + END - abs(CIEND)}{2}$$

Where POS is the initial position, END is the end position, CIPOS is the confidence interval of POS and CIEND is the confidence intervals of END. Having defined the central point of the region, we defined the initial and final positions of the SV as:

$$BPI = Center - \frac{abs(SVLEN)}{2}; BPE = Center + \frac{abs(SVLEN)}{2}$$

Where BPI and BPE are the limits of the SV and SVLEN is the size of the SV.

We then concatenated the regions for all the individuals, sorted them and merged them using

bedtools sort and bedtools merge[22] to remove any redundancies among the regions.

Following filtering, we merged the resulting variants within samples using SURVIVOR (v1.0.7[2])

accounting for the SV type and collapsing those whose break points were within 1kb. We

represented the intersection of SVs across individuals by extracting the support vectors

generated by SURVIVOR[2] at merging time, and plotted them using the UpSet function from

the R[23] package ComplexHeatmap[24] (v2.8.0). We extracted the support value (i.e. how many

animals present a specific SV) and SV size for each variant in the combined VCF and tested

whether the SVs found in one individual only (support = 1) were significantly larger than those

shared among individuals (support > 1) by performing a Wilcoxon signed-rank test in a custom

R script.

Finally, we defined which of the final set of SVs were found to potentially affect a gene. We

ran VEP v105[25] to predict which SVs were likely to disrupt a gene's function, with the options

--sift b (both preditions score and term), --nearest symbol (report the gene symbol), and --

distance 200 (200 bp up and downstream consequence prediction). Those variants presenting

coordinates referring to the negative strand (end position smaller than initial position) were

manually fixed through an in-house script. We then investigated which SVs putatively overlap

a coding region annotated in the cow genome by intersecting merged SVs with coding

sequence intervals. Intersecting genes were investigated with FUMA[26] to identify enriched

gene ontologies and gene sets using all 35,142 gene elements with a unique Entrez gene ID

as the background list.

Data Records

The datasets presented here are stored at ENA under analysis IDs PRJEB47998 and PRJEB52551. The data are uploaded in Bionano BNX format compatible with downstream analyses. The output of the Solve workflows can be downloaded from Zenodo (DOIs: 10.5281/zenodo.6516993 and 10.5281/zenodo.6517172). The raw VCF files, converted using smap_2_vcf_v2.py, can be found on Zenodo with DOI: 10.5281/zenodo.6516933.

Technical Validation

Assembly statistics and SV calling

We aligned the Saphyr optical mapping reads to the ARS-UCD1.2 genome[27], expanded with the BTau5 Y chromosome generated by the 1000 bulls genome project, using Bionano Solve (v3.3 and 3.5) to assemble the genome maps and call SVs. The two NDama samples had previously been used to validate SV using graph genome approaches[28].

Workflow metrics are provided in Supplementary Data 1, summarising key metrics for the analysis of these samples in comparison to the recommended values from Bionano (see https://bionanogenomics.com/wp-content/uploads/2017/03/30255-Bionano-Access-Assembly-Report-Guidelines.pdf).

Unfiltered molecules had average read lengths of 131.9-219.8kb (recommended >150kb for both metrics) and molecule N50s ranged from 185.2-361.9kb across the samples (recommended >150Kbp). Following molecule filtering, all samples were within the recommended average length (245.5-383.1, recommended >230Kbp) and molecule N50 (245.0-426.5, recommended >230Kbp), and only 1 sample (Angoni 1) was slightly below the recommended label density (13.1-16.4, recommended 14-17). Importantly all samples passed the recommended values for the effective coverage of the reference (72.5-128.2, recommended >70) and of average confidence (30.1-43.2, recommended >20).

All samples also generated assemblies with high genome map N50s for both the diploid (71.7-85.0, recommended >50) and haploid (71.3-84.5, recommended >50) assemblies. Despite the low proportion of assembled contigs aligning to the reference genome (0.14-0.25, recommended >0.70), the high uniquely aligned length by reference length (0.835-0.906,

recommended >0.85) shows the presence of long assembled contigs. The contigs present a

high fraction of molecules aligned (0.77-0.94, recommended >0.80), effective coverage

assembly (37.7-66.7, recommended >40) and average confidence (38.5-51.3, recommended

>20).

Overall, 1 sample had 11 metrics within the recommended values, 6 had 12 metrics within

the recommended values, 9 had 13 metrics within the recommended values and 2 had 14

metrics within the recommended values.

The Bionano Solve workflow identified a number of SV in each sample, ranging from 4,944 to

11,184 for a Hereford and Nelore, respectively (Table 2). This mirrors the evolutionary

distance of each sample from the reference genome, with the European taurine possessing

fewer SVs (4,944-5,652) than the other samples, and an African taurine N'Dama possessing

the least among the non-European individuals (N = 6,254). Relative SV numbers consequently

broadly mirror prior expectations.

Variant statistics

SVs were filtered using Bionano Access, excluding SVs with unknown dosages, and retaining

those larger than 1Kb and with a quality >20. SVs for each individual were then combined

using SURVIVOR[2] if the breakpoints were within 1Kb, i.e. below the effective resolution of the

approach[2]. This process allowed us to select a catalogue of 13,457 SVs across the genome,

containing 8,262 insertions, 5,191 deletions and 4 inversions (see Supplementary Data 2 and

3 for the details on the type of SV identified). No duplications, inverted duplications and

translocations passed the quality filtering. Consistent with results from previous studies[28],

most of the post-filtering insertions and deletions identified fell into the smaller classes,

though 1,796 SVs (403 deletions, 1,389 insertions and 4 inversions) of over 50kb in length

were identified (Figure 1).

These SVs longer than 1Kb and of high quality involve a total of 2,656 unique regions, for an

estimated total of over 90Mb of non-redundant bases (Supplementary Data 4). This number

is comparable to what has been seen for novel sequences (i.e. insertions) using graph genome

approaches, where an extra 70Mb and 116Mb of novel sequence were reported on 5 and 4

cattle reference genomes, respectively[28,29]. After merging the filtered variants from all the

samples, most of the SVs were found to be private to an individual (Figure 2), consistent with

what has been observed in previous studies[1]. Individuals of indicine ancestry (Nelore and

Boran) carry almost twice as many SVs relative to the Hereford reference as taurine

individuals, further highlighting that the current reference poorly represents these breeds

(Figure 2).

Interestingly, we find that SVs only found in one animal (support = 1; n = 7,445, mean SV

length = 85,954.23bp) are generally larger (Wilcoxon rank-sum test P-value = $8.99 * 10^{-37}$)

than the SVs found in more than one animal (support > 1; n = 6,012, mean SV length =

27,747.17bp, Figure 1, Supplementary Figure 1). The list of all SVs, with their position, support

and size, are reported in Supplementary Data 5.

Finally, we investigated whether any of the high-quality SVs potentially impact annotated

genes. VEP successfully processed 12,999 out of 13,457 SVs (see HTML report on GitHub).

Some variants were too large to be successfully processed by VEP, and other were called as

incomplete by VEP. Of these, 6,946 were intergenic, and the remaining 5,934

overlapped5,780 genes and 17,386 transcripts, suggesting the potential for functional

variants among the SVs detected. A total of 1,200 SVs putatively overlap a coding sequence.

These coding sequences are included in a total of 884 unique gene elements in the cow

annotation (Ensembl v105), and of these 483 have an associated gene name (Supplementary

Data 6). A total of 292 out of 483 genes had an ID recognized by FUMA[26]. These 292 genes

belong to a number of gene sets such as the Hallmark bile acid metabolism and interferon γ

and α response sets (Supplementary Figure 2), as well as the olfactory receptor curated gene

set (Supplementary Figure 3). All gene set results from FUMA are reported in Supplementary

Data 7.

Usage Notes

Even with the ever-decreasing cost of long read sequencing making it increasingly tractable

to call SVs across sets of samples using HTS, validation of the SV calls remain challenging. This

compendium of SVs across global cattle breeds provides a validation set called using an

independent technology that can be used to assess the quality of cattle SV calls. In fact, optical

mapping data has previously been used to validate sequencing based SV calls[28], and we

believe this dataset provides the largest set of optical maps to date for a livestock species.

With many SVs shared across the two animals of each breed, the raw molecules in this dataset

can also be used to help scaffold and validate novel assemblies of cattle of breeds closely

related to the individuals represented here, potentially reducing the cost of future genome

assembly projects.

Unlike most cattle studies, this database is not focused just on European cattle breeds,

meaning this will be a valuable resource to researchers across the globe. Importantly, it will

allow for SVs to inform the interpretation of results from GWAS and population genetics

studies by providing candidate functional variants in relevant regions.

Ultimately, we expect the database to enable further insights into SVs, an understudied class

of genetic variation in cattle, giving access to a catalogue of thousands of variants present

across multiple breeds worldwide.

Code availability

262 The code used in this article were deposited at

263 https://github.com/evotools/CattleOManalyses.

References

1. Collins, R. L. *et al.* A structural variation reference for medical and population genetics.

Nature **581**, 444–451 (2020).

2. Jeffares, D. C. *et al.* Transient structural variations have strong effects on quantitative

traits and reproductive isolation in fission yeast. *Nat. Commun.* **8**, (2017).

3. Liu, G. E. *et al.* Initial analysis of copy number variations in cattle selected for resistance

or susceptibility to intestinal nematodes. *Mamm. Genome Off. J. Int. Mamm. Genome*

Soc. **22**, 111–121 (2011).

4. A 12 kb multi-allelic copy number variation encompassing a GC gene enhancer is

associated with mastitis resistance in dairy cattle.

https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1009331.

5.   Bickhart, D. M. & Liu, G. E. The challenges and importance of structural variation

detection in livestock. *Front. Genet.* **5**, (2014).

6.   Weischenfeldt, J., Symmons, O., Spitz, F. & Korbel, J. O. Phenotypic impact of genomic

structural variation: insights from and for human disease. *Nat. Rev. Genet.* **14**, 125–138

(2013).

7.   Chen, L., Chamberlain, A. J., Reich, C. M., Daetwyler, H. D. & Hayes, B. J. Detection and

validation of structural variations in bovine whole-genome sequence data. *Genet. Sel.*

Evol. **49**, 13 (2017).

8.   Couldrey, C. *et al.* Detection and assessment of copy number variation using PacBio

long-read and Illumina sequencing in New Zealand dairy cattle. *J. Dairy Sci.* **100**, 5472–

5478 (2017).

9.   Bickhart, D. M. *et al.* Diversity and population-genetic properties of copy number

variations and multicopy genes in cattle. *DNA Res.* **23**, 253–262 (2016).

10. Liu, G. E. *et al.* Analysis of copy number variations among diverse cattle breeds. *Genome*

Res. **20**, 693–703 (2010).

11. Mei, C. *et al.* Copy number variation detection in Chinese indigenous cattle by whole

genome sequencing. *Genomics* **112**, 831–836 (2020).

12. Upadhyay, M. *et al.* Introgression contributes to distribution of structural variations in

cattle. *Genomics* **113**, 3092–3102 (2021).

13. Alkan, C., Coe, B. P. & Eichler, E. E. Genome structural variation discovery and

genotyping. *Nat. Rev. Genet.* **12**, 363–376 (2011).

14. Pabinger, S. *et al.* A survey of tools for variant analysis of next-generation genome

sequencing data. *Brief. Bioinform.* **15**, 256–278 (2014).

15. Wala, J. A. *et al.* SvABA: genome-wide detection of structural variants and indels by local

assembly. *Genome Res.* **28**, 581–591 (2018).

16. Yuan, Y., Chung, C. Y.-L. & Chan, T.-F. Advances in optical mapping for genomic research.

Comput. Struct. Biotechnol. J. **18**, 2051–2062 (2020).

17. Lam, E. T. *et al.* Genome mapping on nanochannel arrays for structural variation analysis

and sequence assembly. *Nat. Biotechnol.* **30**, 771–776 (2012).

18. Levy-Sakin, M. *et al.* Genome maps across 26 human populations reveal population-

specific patterns of structural variation. *Nat. Commun.* **10**, (2019).

19. Ellis, J. A. *et al.* Differentiation antigens on bovine mononuclear phagocytes identified by

monoclonal antibodies. *Vet. Immunol. Immunopathol.* **19**, 325–340 (1988).

20. Chan, S. *et al.* Structural Variation Detection and Analysis Using Bionano Optical

Mapping. in *Copy Number Variants: Methods and Protocols* (ed. Bickhart, D. M.) 193–

203 (Springer, 2018). doi:10.1007/978-1-4939-8666-8_16.

21. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008

(2021).

22. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinforma. Oxf. Engl.* **26**, 841–2 (2010).

23. R core team. *R: a language and environment for statistical computing*. (R Foundation for

Statistical Computing, 2021).

24. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

25. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122–122

(2016).

26. Watanabe, K., Taskesen, E., Van Bochoven, A. & Posthuma, D. Functional mapping and

annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826–1826 (2017).

27. Rosen, B. D. *et al.* De novo assembly of the cattle reference genome with single-

molecule sequencing. *GigaScience* **9**, 1–9 (2020).

28. Talenti, A. *et al.* A cattle graph genome incorporating global breed diversity. *Nat.*

Commun. **13**, 910 (2022).

29. Crysnanto, D., Leonard, A. S., Fang, Z.-H. & Pausch, H. Novel functional sequences

uncovered through a bovine multiassembly graph. *Proc. Natl. Acad. Sci.* **118**,

e2101056118 (2021).

330
331

Acknowledgements

The study was funded by grant BB/R015155/1 to JGDP and Institute Strategic Programme Grant BBS/E/D/10002070 from the Biotechnology and Biological Sciences Research Council (BBSRC).

Author contribution

J.G.D.P. and T.C. conceived the study, A.T. and J.G.D.P. designed the analyses and A.T. performed them. J.G.D.P. and A.T. prepared the initial manuscript with all authors contributing to subsequent drafts. J.Po., M.C. and T.C. prepared the DNA and Bionano samples. D.W. and P.T. provided samples and expertise for the study. M.E.Z.M., I.M.S., A.F., B.R.F., C.E., E.T.O., D.M., W.A., I.S., J.B.M., G.M. and R.F.K provided samples for the study.

Figure captions

Figure 1: Histogram of the structural variant (SV) sizes. Histogram of the size of the identified SVs in bins of 5Kb.

Figure 2: Upset plot of the structural variants. Upset plot of the structural variants by individual for the 40 sets containing the most SVs.

Tables

Sampling Continent	Sampling Country	Group	Breed	ENA project ID	ENA sample ID
S. America	Brazil	Indicine	Nelore	PRJEB52551	ERS11891755
				PRJEB52551	ERS11891754
Africa	Kenya	Indicine	Boran	PRJEB52551	ERS11891767
				PRJEB52551	ERS11891766
Africa	Nigeria	Indicine	White Fulani	PRJEB52551	ERS11891768
				PRJEB52551	ERS11891769
Africa	Zambia	Indicine	Angoni	PRJEB52551	ERS11891764
				PRJEB52551	ERS11891765
Africa	Uganda	Sanga	Ankole	PRJEB52551	ERS11891756

				PRJEB52551	ERS11891757
Africa	Zambia	Taurine	Barotse	PRJEB52551	ERS11891762
				PRJEB52551	ERS11891763
Africa	Nigeria	Taurine	N'Dama	PRJEB47998	ERS8452869
				PRJEB47998	ERS8452868
Europe	United Kingdom	Taurine	Hereford	PRJEB52551	ERS11891760
				PRJEB52551	ERS11891761
Europe	United Kingdom	Taurine	Holstein-Friesian	PRJEB52551	ERS11891759
				PRJEB52551	ERS11891758

350   Table 1: Description of the samples. Table describing the breeds and ancestry of samples,

351   with the continent and country of origin. The identifiers, as well as the ENA accession codes,

352   for each of the two animals sampled per breed are also reported.

Sample	Dels	Insert.	Dups.	Inversion breakpoints	Interchr. translocation breakpoints	Intrachr. translocation breakpoints	Total
Angoni 1	4349	4505	45	91	13	4	9007
Angoni 2	4387	4673	64	100	11	7	9242
Ankole 1	4314	4324	67	111	10	8	8834
Ankole 2	3911	3984	66	101	8	5	8075
Barotse 1	3971	4044	42	52	11	6	8126
Barotse 2	4199	4159	67	106	7	9	8547
Boran 1	4935	5087	56	113	15	6	10212
Boran 2	4990	5007	68	138	6	14	10223
Hereford 1	2465	2380	43	41	11	4	4944
Hereford 2	2435	2437	77	88	9	7	5053
Holstein 1	2756	2759	48	52	15	18	5648
Holstein 2	2702	2801	59	76	9	5	5652
N'Dama 1	3411	3481	92	125	10	13	7132
N'Dama 2	3005	3082	67	86	6	8	6254
Nelore 2	5294	5508	58	113	11	5	10989
Nelore 1	5420	5499	96	136	15	18	11184
White Fulani 1	4467	4642	54	114	11	3	9291

White Fulani 2	4782	4805	41	45	17	14	9704

Table 2: Raw number of structural variants (SVs) and type detected in the different samples. This table summarises the number of raw SVs detected in each sample, and their classification (e.g. insertion, deletion, duplication, inversion and inter- and intra-chromosomal translocation).

Minimum variant size = 1Kb, QUAL = 20 , break points distance = 1000