Title:

Field margins as substitute habitat for the conservation of birds in agricultural wetlands

Authors:

Mallet Pierre [1,2,3,4]

Béchet Arnaud [2]

Sirami Clélia [4]

Mesléard François [2,3]

Blanchon Thomas [2]

Calatayud François [4]

Dagonet Thomas [2]

Gaget Elie [5]

Leray Carole [2]

Galewski Thomas [2]

1. ENGREF AgroParisTech, 75000, Paris, France

2. Tour du Valat Research Institute, Le Sambuc, 13200, Arles, France

3. Mediterranean Institute of marine and terrestrial Biodiversity and Ecology, Avignon

Université, UMR CNRS IRD Aix Marseille Université, IUT Site Agroparc, BP 61207, 84911

Avignon Cedex 09, France

4. UMR Dynafor, INRAE, Toulouse University, 31326 Castanet Tolosan, France

5. Department of Biology, University of Turku, Finland

Correspondence author:

Name: Pierre Mallet

Adresse: Institut de recherche de la Tour du Valat. Le Sambuc, 13200 Arles, France

E-mail: mallet@tourduvalat.org

ORCID: 0000-0002-0139-3745

Abstract

Breeding birds in agricultural landscapes have declined considerably since the 1950s and the beginning of agricultural intensification in Europe. Given the increasing pressure on agricultural land, it is necessary to identify conservation measures that consume little productive land. We tested the compensation hypothesis which states that field margins may represent substitute habitats for bird species in agricultural wetlands. We monitored bird species in 86 crop fields in rice paddy landscapes of Camargue (southern France), a wetland of international importance for birds. We investigated whether the area of three types of field margins (reed strips, grass strips and hedgerows) within a 500 meter buffer around the centroid of each crop fields had an effect on the abundance of bird species from three groups defined based on their primary habitat (reedbeds, grasslands, and forest edge species). We controlled for the area of each type of semi-natural habitat (wetlands, grasslands, and woodlands), crop diversity (rice, wheat, alfalfa, rape, and market gardening) and mean crop field size. Results show partial support of the compensation hypothesis with species-dependent responses to primary and substitute habitat area. Some species within the reedbed and grassland bird guilds are favored by the area of their primary habitat as well as by the area of field margins, in line with the compensation hypothesis. Eurasian reed warbler is favored by the area of both wetlands and reed strips. Corn bunting is favored by grassland and grass strip areas. We could not confirm the compensation hypothesis for other species. However, this may be due to the fact that most of these species did not respond to their primary habitat. These results therefore suggest that field margins may represent substitute habitats for some species but further studies, in contexts where species are strongly associated with their primary habitat, would be needed to confirm the generality of this hypothesis. Our results also suggest that species response to increasing the area of a field margin type may vary among guilds and even within guilds. Therefore, it may be difficult to favor all species within a given landscape and management actions may need to be tailored to whichever species are locally associated with the highest conservation priority. To tackle this challenge, it may be necessary to design landscape management actions at different spatial scales.

57   **Keywords**: bird conservation, biodiversity, landscape heterogeneity, grassland birds, forest

58   edge birds, reedbed birds, wetland, rice, habitat compensation, land sharing

1 Introduction

Farmland bird populations have experienced a massive decline worldwide in recent decades, primarily due to the loss of semi-natural habitats and intensification of agricultural practices (PECBMS, 2022; Stanton et al., 2018; Sundar and Subramanya, 2010). Agricultural areas represent 37 % of the European terrestrial area and host a large proportion of terrestrial biodiversity (DataBank, 2018; Herzog et al., 2013). It is therefore not practical to rely solely on the creation of protected areas to compensate for the declines in biodiversity observed in European agricultural environments (Meyer et al., 2013; Warren et al., 2021). Rather, conservation efforts should also focus on maintaining and increasing the capacity of agricultural landscapes to support biodiversity through the adoption of biodiversity-friendly agricultural practices and the protection of non-productive refuge areas, i.e. promote land sharing (Grass et al., 2021).

Patches of semi-natural habitats, such as woodlands, grasslands and wetlands, remaining within agricultural landscapes may provide permanent habitat for wildlife and host a large part of farmland biodiversity (Newton, 2017; Toffoli and Rughetti, 2017). However, these patches are scarce and under increasing pressure in Europe due to agricultural intensification which leads to their progressive conversion to arable land despite efforts from the European Union to slow down this trend through agri-environment schemes (Batáry et al., 2015). Hence, in some agricultural landscapes, field margins, i.e. linear elements covered by semi-natural vegetation along the edge of crops, are the only type of semi-natural habitat left (Marshall and Moonen, 2002). The habitat compensation hypothesis states that species may compensate for the loss of their primary habitat by using alternative habitats as a substitute (Norton et al., 2000). For instance, Montagu's harrier (*Circus pygargus*) primarily nests in shrublands and grasslands but, in some part of its distribution range, it now relies exclusively on crop fields for breeding and foraging (Norton et al., 2000). It has also been shown that aquatic invertebrates can use drainage ditches as substitute habitats for natural lakes and rivers (Dollinger et al., 2015). The habitat compensation hypothesis has been investigated in the context of farmland abandonment and in dry agricultural areas (e.g. Brotons et al., 2005; Saura et al., 2014; Vallecillo et al., 2008) but rarely in wetland agricultural areas (e.g. Decleer et al., 2015) despite their specific landscape characteristics and biodiversity.

One of the main crops cultivated in wetlands is rice, a flooded cereal which represents 22.8 % of the world cereal surface area (FAO, 2018; Singh et al., 2001). In such rice paddy landscapes, agricultural and semi-natural areas are generally intermingled with the presence of

large field margins along ditches. Among birds associated with these rice paddy landscapes,
there are both waterbirds (e.g. gulls, terns, herons, storks, ibises, waders…) and terrestrial bird
species. While the role of rice paddy landscapes as alternative habitat for waterbirds has been
largely studied, their role for terrestrial birds has received much less attention (Elphick, 2015).
Considering the long-term decline of terrestrial bird populations in agricultural landscapes
(Fraixedas et al., 2019), identifying conditions favoring them would be useful to improve
recommendations for agri-environmental management practices in rice paddy landscapes.
Terrestrial birds using rice paddy landscapes include different ecological guilds: reedbed birds,
which are primarily associated with freshwater marshes (Morganti et al., 2019); forest edge
species, which are originally associated with forest borders and clearings (Hinsley and Bellamy,
2019; Newton, 2017); and grassland species, which originally live in grassy or shrubby
vegetation with no tree cover (Di Giacomo et al., 2010). Field margins could provide resources
and nesting habitats for these species (Vickery et al., 2009), e.g. reed strips along ditches for
reedbed birds, hedgerows for forest edge species and grass strips for grassland species.
However, the role of field margins for terrestrial birds has rarely been considered in studies
conducted in rice paddy landscapes (King et al., 2010).

The Camargue (Rhône delta) is a biologically rich area listed in the Ramsar Convention
and classified as a Biosphere Reserve by UNESCO (Blondel et al., 2019). Natural areas cover
58,000 ha and agricultural areas 55,100 ha (Tamisier and Grillas, 1994). Rice represents 48 %
of the crop area and is mainly cultivated in rotation with wheat (19 %) and alfalfa (5 %). Within
this region, bird species associated with agricultural areas have experienced the greatest rate of
decline over the past 50 years compared to waterbirds (Fraixedas et al., 2019; Galewski and
Devictor, 2016). Hence, it is critical to assess whether field margins could constitute a lever for
bird conservation as their restoration and management may be readily changed by farmers.

In this paper, we tested the habitat compensation hypothesis in rice paddy landscapes of
Camargue by assessing whether field margins act as substitute habitats for reedbed birds, forest
edge birds and grassland birds. We conducted bird surveys in 86 crop fields in Camargue.
Specifically, we predicted that (i) forest edge species would be positively influenced by
woodlands and hedgerows; (ii) grassland birds would be positively influenced by grasslands
and grass strips and (iii) reedbed birds would be positively influenced by semi-natural wetland
areas and reed strips.

2   Material and methods

2.1   Study area

Our study was conducted in the Rhône River delta, a 180,000 ha polderized flood plain located in Southern France and known as "Camargue". Warm summers typical of the Mediterranean climate (average monthly temperature between May and October above 15°C; Blondel et al., 2019), as well as fresh water pumped from the Rhône River allows rice cultivation. This flooded crop is essential for washing out salt-rich soils and allows rotation with dry crops, mainly wheat and alfalfa. In Camargue, field margins are often wide (> 3 m) to be waterproof and keep the crop fields flooded during the rice cultivation period. Several types of vegetation can therefore co-occur within the same field margin, such as reed strips, hedgerows or grass strips. In Camargue, the area of semi-natural habitats decreased from 67 % to 39 % between 1942 and 1984 and since remained stable at around 58,000 ha (Mallet, 2022; Tamisier and Grillas, 1994). These semi-natural areas are spatially segregated in the delta; woodlands are mainly restricted to riparian areas along the Rhône River, wetlands occupy depressions and cover large areas in the center and south of the delta while grasslands (mostly constituted of meadows and salt steppes) surround the wetlands on slightly elevated areas (Appendix A).

2.2   Study design

We selected 86 crop fields belonging to 17 farms across the Camargue (Fig. 1). All fields were organic to limit confounding effects associated with variation in the intensity of agricultural practices. We selected crop fields covered by the crop types representative of the main agricultural production in Camargue (rice, wheat, alfalfa, rape, and market gardening). Crop fields were selected along two independent gradients of semi-natural cover and hedgerow cover using the methodology developed by Pasher et al., (2013). To do so, we measured semi-natural and hedgerow areas in a 500 meters square moving window with a step size of 100 meters around every agricultural land of Camargue thanks to land-use data from 2019 of the BD TOPO®, OSO Land Cover Map and the Regional Natural Park of the Camargue. No maps of grass strip or reed strip were available prior to crop field selection. Therefore, we checked for the distribution of sampled crop fields along gradients of explanatory variables once the selection and on-site mapping were completed. We also checked for correlation among the cover of different types of field margin and other landscape variables (see below).

156
Figure 1. Location of the 86 crop fields monitored for birds in Camargue, Rhône delta.
Triangles represent crop fields sampled in 2020 and stars represent crop fields sampled in
2021.

We calculated the area of each type of field margin and semi-natural habitat within a 500
meter buffer around the centroid of each crop field, following Chan et al. (2007). The maximum
size of the sampled crop fields was 14 ha, hence much smaller than this buffer. First, we
estimated the area of the three types of field margins: (1) hedgerows, tree lines and bushy areas;
(2) grass strips, grassy boundaries including grassy tracks or dirt roads used for the moving of
agricultural machinery; (3) reed strips that grow in and along irrigation or drainage earthen
ditches. Because we aimed at testing the hypothesis that field margins represent substitute
habitats whatever their shape, we calculated the area and not the length of field margins.
Second, we estimated the area of three categories of semi-natural areas: (1) woodlands (mainly
riparian forests dominated by white poplar (*Populus alba*), pinewoods (*Pinus pinaster*), and
tamarisk (*Tamarix gallica*) groves) and shrublands dominated by narrow-leaved mock privet
(*Phillyrea angustifolia*); (2) grasslands including dry grasslands extensively grazed by free-

range cattle, Mediterranean salt meadows and halophilous scrubs and fallow lands; (3) wetlands

including freshwater and brackish marshes, reedbeds and ponds. Landscape mapping was based

on field observations done after the bird monitoring in June 2020 and June 2021 (see below)

because fine scale assessment was not feasible based on remote sensing approaches only,

particularly for reed strips. Finally, to account for the possible confounding effect of crop field

heterogeneity, we also estimated within each 500 meter buffer the mean crop field size and the

Shannon diversity index of crop types ($Crop\_SHDI = -\sum_{i=1}^{n} pi \ln pi$, where pi corresponds to

the proportion of crop cover type i in the landscape), following the method implemented in

Sirami et al. (2019). As a result, we obtained values for eight landscape variables for each

sampled crop field.

2.3   Bird monitoring and traits

Birds were monitored over 5-minute point counts halfway along the longest field margin

of each crop field during the breeding period. Two visits were conducted at each site between

mid-April and mid-June with at least 4 weeks between the two visits, following the protocol

from the French common breeding bird census scheme (Jiguet, 2003). Flying birds were

removed from the analyses because they were not interacting directly with the landscape. Birds

landing outside the sampled crop field and its field margins were also removed to avoid

detection bias potentially generated by hedgerows preventing the observer to see birds beyond

trees. We used the maximum abundance per site between the two visits for each species for

further analyses.

We assigned each species to one of three guilds according to the primary habitat used for

breeding: reedbed, grassland and forest edge birds. Assignment was based on the EUNIS habitat

classification that describes species communities related to woodlands, wetlands, grasslands or

urban areas (Appendix B). Generalist birds, i.e. not linked to one habitat in particular, or birds

that use urban areas for breeding were discarded from the analyses. We modulated the EUNIS

data with information provided by a local expert (T.G.) to take into account ecological

particularities of the Camargue. To avoid extreme cases of zero-inflation, we only kept species

present in more than 15 % of the sampled crop fields (Marja and Herzon, 2012). Fourteen

species were retained for the analyses (Table 1).

Table 1. Species studied within the three guilds based on the EUNIS database combined with information provided by local experts to take into account ecological particularities of the Camargue (Appendix B).

Guilds	Species
Forest edge birds	European greenfinch (*Chloris chloris*)
	Carrion crow (*Corvus corone*)
	Melodious warbler (*Hippolais polyglotta*)
	Common nightingale (*Luscinia megarhynchos*)
	Great tit (*Parus major*)
	Eurasian magpie (*Pica pica*)
	European green woodpecker (*Picus viridis*)
	Eurasian blackcap (*Sylvia atricapilla*)
	Eurasian hoopoe (*Upupa epops*)
Grassland birds	Crested lark (*Galerida cristata*)
	Corn bunting (*Emberiza calandra*)
	Eurasian skylark (*Alauda arvensis*)
Reedbed birds	Eurasian reed warbler (*Acrocephalus scirpaceus*)
	Great reed warbler (*Acrocephalus arundinaceus*)

In order to check for the completeness of our data, we calculated the coverage of our
sampling, which is defined as the proportion of the total number of individuals in an assemblage
that belong to the species present in the sample (Chao and Jost, 2012). This index corresponds
to the probability of occurrence of the species observed in the sample. The coverage was
calculated by crop field for all 14 species considered within the present study. The overall
coverage of our sampling was 73.5 %, which reflects no undersampling issue (Mallet et al.,
2022). The sampling completeness per crop field was not correlated with any explanatory
variable (Pearson coefficient < 0.24, Appendix C), which suggests that the study design was
robust and not biased toward one or several landscape variables.

2.4   Data analysis

We ran one linear mixed-effect model with bird abundance as the response variable, while
fixed effects were species identity, the area of the three field margin types (hedgerows, grass
strips and reed strips), the area of the three semi-natural habitat types (woodlands, grasslands
and wetlands), crop diversity, mean crop field size and all two-way interactions between species
identity and the other explanatory variables. All explanatory variables were centered and scaled.
Crop type and site identity were added as random effects. We did not include variable 'year' in
our final models because this variable was never significant and was not relevant to our research

question. We accounted for spatial autocorrelation by using an exponential structure on crop
field coordinates, and checked for the absence of autocorrelation in the residuals. We used a
negative binomial error distribution (type 2: variance increases quadratically with the mean) to
deal with over-dispersion. We ran models with a log-link function. We conducted post hoc
comparisons of slopes using the emtrends function.
Statistical analyses were run using glmmTMB (Magnusson et al., 2020), entropart (Marcon
and Hérault, 2015) and emmeans in R 4.0.5 (R Core Team, 2017).
236

3   Results

The spatial variation in field margin area around the 86 organic crop fields was similar
across the three margin types; hedgerows (median = 3.67 ha; range: [0; 17.47]), reed strips
(median = 3.60 ha; range: [1.46; 8.72]) and grass strips (median = 3.29 ha; range: [0; 6.27]).
The dominant type of semi-natural habitat was grassland (median = 7.38 ha; range: [0; 45.23]),
followed by wetland (median = 1.37 ha; range: [0; 48.15]) and by woodland (median = 0.71 ha;
range: [0; 20.78]). Crop diversity was on average 0.93 ± 0.04 (median = 0.98; range: [0; 1.6]).
Crop mean field size was on average 2.32 ± 0.10 ha (median = 2.27 ha; range: [1.09; 5.85]).
There was no correlation among explanatory variables since all Pearson correlation coefficients
were under 0.45 (Appendix C).
247

3.1   Forest edge bird guild

Woodland area only had a positive effect on the abundance of one of the nine forest edge
species, great tit ($\beta = 0.10 \pm 0.03$, Table 2, Fig. 2), while the area of hedgerows had a positive
effect on the abundance of European greenfinch ($\beta = 0.15 \pm 0.07$, Table 2, Fig. 2).
Grassland area had a negative effect on the European green woodpecker ($\beta = -0.12 \pm 0.05$,
Table 2, Fig. 2).
The area of reed strips had a negative effect on the abundance of carrion crow, common
nightingale, European green woodpecker and Eurasian blackcap (respectively $\beta = -0.30 \pm 0.14$,
$\beta = -0.16 \pm 0.04$, $\beta = -0.55 \pm 0.28$, $\beta = -0.26 \pm 0.12$, Table 2, Fig. 2).
Crop diversity had a positive effect on the abundance of European greenfinch and great tit
(respectively $\beta = 2.09 \pm 0.92$, $\beta = 1.39 \pm 0.56$, Table 2), while crop mean field size had a
positive effect on the abundance of carrion crow ($\beta = 0.62 \pm 0.20$, Table 2).

There was no significant effect of wetland area or grass strip area on the abundance of
species of this guild (Table 2, Fig. 2).
262

3.2   Grassland bird guild
The abundance of corn bunting was positively related to both grassland area
($\beta = 0.12 \pm 0.03$, Table 2, Fig. 2) and the area of grass strips ($\beta = 0.46 \pm 0.18$, Table 2, Fig. 2).
Woodland area had a negative effect on the abundance of crested lark ($\beta = -0.33 \pm 0.15$,
Table 2, Fig. 2), while the area of hedgerows had a negative effect on the abundance of Eurasian
skylark and corn bunting (respectively $\beta = -0.29 \pm 0.10$, $\beta = -0.33 \pm 0.11$, Table 2, Fig. 2).
Wetland area had a positive effect on the abundance of Eurasian skylark and corn bunting
(respectively $\beta = 0.05 \pm 0.02$, $\beta = 0.06 \pm 0.02$, Table 2, Fig. 2), while the area of reed margins
had no effect on the abundance of grassland species (Table 2, Fig. 2).
Crop diversity had a positive effect on the abundance of corn bunting ($\beta = 2.33 \pm 0.87$,
Table 2), while crop mean field size had a negative effect on the abundance of Eurasian skylark
($\beta = -0.74 \pm 0.37$, Table 2).
275

3.3   Reedbed bird guild
The abundance of Eurasian reed warbler was positively related to both wetland area
($\beta = 0.04 \pm 0.01$, Table 2, Fig. 2) and the area of reed margins ($\beta = 0.26 \pm 0.09$, Table 2, Fig.
2).
The abundance of great reed warbler was negatively related to both grassland area ($\beta = -$
0.07 ± 0.02, Table 2, Fig. 2) and the area of grass strips ($\beta = -0.45 \pm 0.14$, Table 2, Fig. 2).
There was no significant effect of woodland area, hedgerow area, crop diversity or crop
mean field size on the abundance of species of this guild (Table 2, Fig. 2).
284
285

Figure 2. Estimates (± 95% confidence interval) of the effect of landscape variable for each

species studied. Each graph corresponds to a landscape variable; the habitat patches on the

left and the field margin of the right. The horizontal black line corresponds to 0.. If the 95%

confidence intervals does not overlap with zero, the effect of the landscape variable on the

abundance of the corresponding species is considered as significant.

Table 2. Averaged estimates of the effects of landscape variables for the three bird guilds

monitored in agricultural crop fields of the Camargue. The 95 % confidence intervals are in

brackets. Values in bold indicate significant effects.

Species name	Hedgerow	Grass strip	Reed strip	Woodland area	Grassland area	Wetland area	Crop diversity	Mean crop field size
European greenfinch (*Chloris chloris*)	**0.15 [0.01;0.29]**	0.12 [-0.19;0.43]	0.18 [-0.09;0.46]	0.03 [-0.11;0.16]	0.01 [-0.03;0.06]	0.01 [-0.04;0.06]	**2.09 [0.28;3.91]**	-0.21 [-0.92;0.49]
Carrion crow (*Corvus corone*)	0.05 [-0.07;0.18]	-0.24 [-0.56;0.08]	**-0.30 [-0.54;-0.03]**	-0.12 [-0.27;0.02]	-0.04 [-0.08;0.00]	-0.03 [-0.07;0.02]	-0.14 [-1.56;1.27]	**0.62 [0.21;1.03]**
Melodious warbler (*Hippolais polyglotta*)	0.04 [-0.05;0.12]	0.11 [-0.05;0.28]	-0.09 [-0.24;0.05]	0.01 [-0.05;0.07]	0.01 [-0.02;0.03]	-0.02 [-0.05;0.01]	0.19 [-0.66;1.03]	-0.08 [-0.41;0.24]
Common nightingale (*Luscinia megarhynchos*)	0.02 [-0.03;0.07]	-0.01 [-0.11;0.08]	**-0.16 [-0.24;-0.07]**	0.00 [-0.03;0.04]	0.00 [-0.01;0.02]	-0.01 [-0.03;0.00]	0.06 [-0.43;0.54]	0.17 [0.00;0.34]
Great tit (*Parus major*)	0.10 [-0.01;0.20]	-0.01 [-0.23;0.21]	-0.17 [-0.36;0.02]	**0.10 [0.03;0.16]**	0.02 [-0.01;0.05]	-0.01 [-0.04;0.03]	**1.39 [0.29;2.48]**	0.23 [-0.12;0.59]
Eurasian magpie (*Pica pica*)	-0.14 [-0.41;0.14]	0.02 [-0.34;0.39]	0.03 [-0.33;0.38]	-0.09 [-0.28;0.10]	-0.01 [-0.06;0.04]	-0.20 [-0.46;0.06]	-1.58 [-3.81;0.64]	0.23 [-0.44;0.90]
European green woodpecker (*Picus viridis*)	-0.17 [-0.52;0.18]	-0.50 [-1.04;0.05]	**-0.55 [-1.09;-0.01]**	-0.09 [-0.24;0.06]	**-0.12 [-0.21;-0.03]**	-0.40 [-0.98;0.04]	-1.32 [-3.39;0.74]	-0.17 [-1.10;0.76]
Eurasian blackcap (*Sylvia atricapilla*)	0.06 [-0.06;0.19]	-0.07 [-0.37;0.23]	**-0.26 [-0.50;-0.01]**	0.05 [-0.04;0.14]	-0.01 [-0.05;0.03]	0.00 [-0.03;0.04]	1.33 [-0.01;2.66]	0.36 [-0.09;0.81]
Eurasian hoopoe (*Upupa epops*)	0.02 [-0.16;0.20]	-0.09 [-0.47;0.28]	0.02 [-0.31;0.35]	-0.01 [-0.18;0.015]	0.02 [-0.03;0.07]	0.01 [-0.05;0.06]	1.09 [-0.83;3.01]	0.07 [-0.59;0.74]
Crested lark (*Galerida cristata*)	-0.12 [-0.26;0.03]	-0.07 [-0.32;0.17]	-0.07 [-0.29;0.14]	**-0.33 [-0.61;-0.05]**	0.01 [-0.02;0.04]	0.02 [-0.01;0.05]	0.22 [-1.02;1.45]	-0.07 [-0.57;0.42]
Corn bunting (*Emberiza calandra*)	**-0.33 [-0.55;-0.12]**	**0.46 [0.11;0.80]**	-0.26 [-0.54;0.02]	-0.14 [-0.41;0.14]	**0.12 [0.06;0.17]**	**0.06 [0.02;0.11]**	**2.33 [0.62;4.03]**	-0.13 [-0.77;0.50]
Eurasian skylark (*Alauda arvensis*)	**-0.29 [-0.48;-0.09]**	0.06 [-0.25;0.37]	-0.08 [-0.33;0.18]	-0.15 [-0.38;0.08]	0.03 [-0.01;0.07]	**0.05 [0.01;0.09]**	1.12 [-0.34;2.59]	**-0.74 [-1.46;-0.02]**
Eurasian reed warbler (*Acrocephalus scirpaceus*)	-0.08 [-0.18;0.03]	0.04 [-0.16;0.25]	**0.26 [0.09;0.43]**	0.01 [-0.08;0.11]	-0.01 [-0.04;0.02]	**0.04 [0.01;0.07]**	-0.40 [-1.57;0.78]	-0.07 [-0.51;0.36]

| Great reed warbler (*Acrocephalus arundinaceus*) | 0.00 [-0.10;0.10] | **-0.45** **[-0.74; -0.17]** | 0.08 [-0.14;0.31] | -0.17 [-0.35;0.01] | **-0.07** **[-0.12; -0.03]** | 0.02 [-0.00;0.05] | -1.42 [-2.86;0.02] | -0.49 [-1.19;0.21] |

4 Discussion

Our study shows that different types of field margins can provide alternative habitats to terrestrial birds in a rice paddy landscape, but species responses vary even within species guilds. We found that (i) grass strips represent a substitute habitat to grasslands for corn bunting and (ii) reed strips represent a substitute habitat to wetlands for the Eurasian reed warbler, in line with the habitat compensation hypothesis. For these two species, the positive effect of field margins on the abundance of species was even stronger than the effect of the corresponding semi-natural habitat patch. This result suggests that field margins are currently valuable habitat rather than substitute ones for these two species. It is consistent with the meta-analysis conducted by Riva and Fahrig (2022), which highlighted the higher value of small habitat patches for biodiversity conservation. In contrast, we could not confirm the compensation hypothesis for 12 out of 14 species. Such a lack of support to the compensation hypothesis could be explained by different methodological and ecological reasons. First, we observed a general lack of species responses to their primary habitat with only 3 species responding positively to the primary habitat surface area. This may result from the use of broad categories of habitat preferences, while species abundance may vary along ecological continuums. Also, semi-natural habitats have been grouped into three primary habitat categories, which may not be detailed enough to match species habitats preferences. For example, wetlands include reedbeds but also ponds without emergent vegetation which are likely not very attractive for reedbeds birds. A more detailed mapping of primary habitats or functional description of habitats, such as habitat quality, nesting opportunities or food resources would therefore be necessary to further test the habitat compensation hypothesis for several of the species considered. In addition, the observed species might potentially accommodate a diversity of habitats. Indeed, in the Camargue, some forest edge species like carrion crow, Eurasian magpie or common nightingale are known to be able to nest in very open landscape e.g. in isolated trees within a matrix of cultivated fields. Further studies aiming to test the habitat compensation hypothesis should therefore focus on species that are more strongly associated with their primary habitat.

Our results show that the compensation hypothesis cannot be generalized to all bird species

within the three guilds studied. Indeed, only some species benefited from the presence of field

margins as substitute habitat. Moreover, some species within these guilds were not even

recorded within sampled agricultural landscapes. For example, the bearded reedling (*Panurus*

biarmicus), a reedbed bird, the blue tit (*Cyanistes caeruleus*), a forest edge bird, or the tawny

pipit (*Anthus campestris*), a grassland bird, breed in Camargue but were not contacted at all

during our surveys.

The lack of effect of field margins on some species may be partly explained by both the

quality of field margins and the ecological preferences of these species. Indeed, in Camargue,

ditches are increasingly being lined with concrete or buried, like in Japanese rice paddy

landscapes for example (Yamada et al., 2011). Some studies have highlighted that earthen

ditches host much more aquatic fauna and flora than concrete ones (Katoh et al., 2009). It was

also shown that the density of intermediate egrets (*Egretta intermedia*) was twice as high in rice

fields with shallow earthen ditches than in rice fields with deep concrete-lined ditches

(Katayama et al., 2012). Here, we found a positive effect of reed field margins for the Eurasian

reed warbler but not for the great reed warbler, the latter requiring wetter and larger patches of

reedbeds than the Eurasian reed warbler (BirdLife International, 2022). The absence of the

bearded reedling is also consistent with the fact that this species requires larger areas of

reedbeds and is not encountered in reed strips along artificial ditches (P.M. pers. obs.).

Our results nearly support the hypothesis that hedgerows represent a substitute habitat for

great tit with a positive effect of woodland and a positive but no significant effect of hedgerow.

The European greenfinch is the only species significantly positively affected by hedgerow, a

result that may be useful to encourage farmers to maintain and restore hedgerows. Yet, the lack

of effect of hedgerows for the other species was surprising since hedgerows are known to

benefit a broad range of forest edge species (Batáry et al., 2010; Wilson et al., 2017). In

Camargue, the poor quality of hedgerows may explain the lack of response within a wider bird

community because several of them, i.e. coniferous or giant cane (*Arundo donax*) hedgerows,

are not suitable to forest edge birds as their volume and plant diversity are low (Graham et al.,

2018; Montgomery et al., 2020).

Our results highlight that grass strips have a stronger effect than grasslands for corn

bunting. The greater plant biomass of grass strips compared to Mediterranean salt meadows,

which constitute most of the grassland area habitat category, may explain this greater effect of

grass strips compared to other open habitats. The high density of seeds available in cultivated

fields where this species comes to feed (Madge and de Juana, 2020), can also be a confounding

effect. Unlike other types of field margins, grass strips are probably not used as a nesting habitat

due to disturbance from agricultural activity. In particular, these strips are frequently mowed

and used by farmers to move around the crop fields, which causes disturbances that might

prevent nesting (Vickery et al., 2009).

Further research should therefore assess the ecological value of field margins, for instance

by comparing the demographics of Eurasian reed warbler and corn bunting in the substitute

habitat and in natural habitat to ensure that field margins are not ecological traps (Horne, 1983).

This would also allow to develop recommendations on the most favorable field margin

management methods. It may also be relevant to study the role of different types of field

margins for generalist species. Indeed, a recent paper has highlighted the progressive

colonization of farmland habitats by generalist bird species over the last decades in Spain (Díaz

et al., 2022). Taking into account the response of generalist bird species may therefore help

avoiding the homogenization of bird communities in rice paddy landscapes. Finally, the value

of field margins may also depend on the availability of habitat patches within the landscape.

For instance, reedbeds may have a more positive effects when they are close to a large patch of

wetland. Testing such interactive effects would require an adequate study design with all

combinations of values for field margins and semi-natural patches, and a sample size large

enough to provide robust estimates of all parameters within associated statistical models.

Our study also highlights that increasing a type of field margins may have antagonistic

effects across different guilds. Indeed, four species within the forest edge bird guild were

negatively impacted by the area of reed strips. This result may be due to the fact that this type

of field margin provides too few resources in terms of food and nesting sites for forest edge

bird species (Shoffner et al., 2018). Similarly, the abundance of the great reed warbler is

negatively correlated to the area of grassland and grass strip as this species occur mainly in wet

habitats during the breeding season (Dyrcz, 2020). As expected, grassland birds were negatively

impacted by the area of hedgerows and woodland confirming previous studies that observed a

similar negative effect of wooded habitats on different species of grassland birds (e.g. Ellison

et al., 2013; Wilson et al., 2014). Woodland patches usually do not offer resources for grassland

birds and are avoided because they are a source of avian and mammalian predators (Burger et

al., 1994). Our study therefore confirms that it may not be possible to favor all bird species

within a single landscape and it may be necessary to focus on the type of field margins that

most favor species in need of conservation attention.

Our study confirms that increasing crop diversity and decreasing crop mean field size are

complementary levers to promote biodiversity in agricultural landscapes (Sirami et al., 2019).

Indeed, crop diversity benefited two of the nine forest edge species, European greenfinch and
great tit and one grassland species, corn bunting. Moreover, the decrease in crop field size had
a positive impact on Eurasian skylark. The results likely stem from the fact that higher
landscape heterogeneity provides readily available complementary resources (Batáry et al.,
2017; Sirami et al., 2019). On the other hand, we found a positive effect of the increase in crop
field size on the abundance of carrion crows. This effect is probably related to the fact that this
species feed in groups on the ground and may thus be favored by large open areas (Madge,
2020).

In conclusion, our results highlight that field margins are valuable landscape components
to improve biodiversity conservation but cannot be the only components to be promoted in rice
paddy landscapes. In Camargue, current conservation priorities concern the disappearance of
wetlands and grasslands as well as the degraded conservation status of species associated with
these habitats, whereas there is less concern for forest edge birds, which can be found in other
agricultural landscapes. Our study therefore suggests that conserving and restoring wetlands
and grasslands and the associates field margins, reed strips and grass strips, represent a
promising avenue to increase biodiversity in the agricultural landscapes of Camargue. On the
other hand, despite the negative impact of hedgerows on grassland birds and waterbirds
(Tourenq et al., 2001), they can host a diversity of auxiliary species as well as taxa of high
conservation importance in Camargue and other wetlands such as bats (Mas et al., 2021).
Hedgerows have also been shown to limit the presence of greater flamingos (*Phoenicopterus*
roseus), considered as a pest in rice fields (Ernoul et al., 2014). Taking into account the role of
hedgerows across taxa would be particularly relevant in the context of the current action plan
of replanting hedgerows carried out locally by the Regional Natural Park of the Camargue.
Land-use planning studies could be a good way to propose management actions to farmers and
stakeholders, maximizing both long-term agricultural benefits and biodiversity conservation.

Acknowledgements

We are grateful to all farmers and landowners who graciously permitted us to work in their

fields. We are particularly grateful to the company Biosud who helped with setting up the

network of organic farmers where we sampled avian biodiversity. Finally, particular thanks to

Fabien Laroche for his advices on statistical analysis.

Funding

This research received the support of the French Ministry of Agriculture [2019-2021], the

company "Alpina-Savoie" [2019-2021] and the Fondation de France [2021-2024].

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

Data, script and code availability

Supplementary information, dataset and statistical scripts are available here:

https://doi.org/10.5281/zenodo.7685771

Supplementary information

Appendix A: Map of habitat localization in Camargue

Appendix B: Table of species guilds

Appendix C: Correlation table between landscape explanatory variables and sampling

completeness (Cn).

References

Batáry, P., Dicks, L., Kleijn, D., Sutherland, W., 2015. The role of agri-environment schemes
in conservation and environmental management. Conserv. Biol. 29, 1006–1016.
https://doi.org/https://doi.org/10.1111/cobi.12536

Batáry, P., Gallé, R., Riesch, F., Fischer, C., Dormann, C.F., Mußhoff, O., Császár, P., Fusaro,
S., Gayer, C., Happe, A.K., Kurucz, K., Molnár, D., Rösch, V., Wietzke, A., Tscharntke,
T., 2017. The former Iron Curtain still drives biodiversity-profit trade-offs in German
agriculture. Nat. Ecol. Evol. 1, 1279–1284. https://doi.org/10.1038/s41559-017-0272-x

Batáry, P., Matthiesen, T., Tscharntke, T., 2010. Landscape-moderated importance of hedges
in conserving farmland bird diversity of organic vs. conventional croplands and
grasslands. Biol. Conserv. 143, 2020–2027. https://doi.org/10.1016/j.biocon.2010.05.005

BirdLife International, 2022. IUCN Red List for birds [WWW Document]. URL
http://datazone.birdlife.org/species/factsheet (accessed 2.25.22).

Blondel, J., Barruol, G., Vianet, R., 2019. L'Encyclopédie de la Camargue, 2nd edn. ed. Buchet-
Chastel, Paris.

Brotons, L., Wolff, A., Paulus, G., Martin, J.L., 2005. Effect of adjacent agricultural habitat on
the distribution of passerines in natural grasslands. Biol. Conserv. 124, 407–414.
https://doi.org/10.1016/j.biocon.2005.01.046

Burger, L.D., Burger, L.W., Faaborg, J., 1994. Effects of prairie fragmentation on predation on
artificial nests. J. Wildl. Manage. 58, 249. https://doi.org/10.2307/3809387

Chan, S.F., Severinghaus, L.L., Lee, C.K., 2007. The effect of rice field fragmentation on
wintering waterbirds at the landscape level. J. Ornithol. 148, 333–342.
https://doi.org/10.1007/s10336-007-0244-z

Chao, A., Jost, L., 2012. Coverage-based rarefaction and extrapolation: Standardizing samples
by completeness rather than size. Ecology 93, 2533–2547. https://doi.org/10.1890/11-
1952.1

DataBank, 2018. Terres agricoles (% du territoire) - European Union [WWW Document]. URL
https://donnees.banquemondiale.org (accessed 4.8.22).

Decleer, K., Maes, D., Van Calster, H., Jansen, I., Pollet, M., Dekoninck, W., Baert, L.,
Grootaert, P., Van Diggelen, R., Bonte, D., 2015. Importance of core and linear marsh
elements for wetland arthropod diversity in an agricultural landscape. Insect Conserv.
Divers. 8, 289–301. https://doi.org/10.1111/icad.12110

Di Giacomo, A.S., Vickery, P.D., Casaas, H., Spitznagel, O.A., Ostrosky, C., Krapovickas, S.,

Bosso, A.J., 2010. Landscape associations of globally threatened grassland birds in the
aguapey river important bird area, corrientes, Argentina. Bird Conserv. Int. 20, 62–73.
https://doi.org/10.1017/S0959270909990177

Díaz, M., Aycart, P., Ramos, A., Carricondo, A., Concepción, E.D., 2022. Site-based vs.
species-based analyses of long-term farmland bird datasets: Implications for conservation
policy evaluations. Ecol. Indic. 140, 109051.
https://doi.org/10.1016/j.ecolind.2022.109051

Dollinger, J., Dagès, C., Bailly, J.S., Lagacherie, P., Voltz, M., 2015. Managing ditches for
agroecological engineering of landscape. A review. Agron. Sustain. Dev. 35, 999–1020.
https://doi.org/10.1007/s13593-015-0301-6

Dyrcz, A., 2020. Great Reed Warbler (Acrocephalus arundinaceus), in: Birds of the World.
Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.grrwar1.01

Ellison, K.S., Ribic, C.A., Sample, D.W., Fawcett, M.J., Dadisman, J.D., 2013. Impacts of tree
rows on grassland birds and potential nest predators: a removal experiment. PLoS One 8,
e59151. https://doi.org/10.1371/journal.pone.0059151

Elphick, C.S., 2015. A history of ecological studies of birds in rice fields. J. Ornithol. 156, 239–
245. https://doi.org/10.1007/s10336-015-1194-5

Ernoul, L., Mesléard, F., Gaubert, P., Béchet, A., 2014. Limits to agri-environmental schemes
uptake to mitigate human-wildlife conflict: Lessons learned from Flamingos in the
Camargue, southern France. Int. J. Agric. Sustain. 12, 23–36.
https://doi.org/10.1080/14735903.2013.798897

FAO, 2018. FAO Statistical databases [WWW Document]. Food Agric. Organ. United Nations.
URL http://www.fao.org (accessed 4.3.19).

Fraixedas, S., Galewski, T., Ribeiro-Lopes, S., Loh, J., Blondel, J., Fontès, H., Grillas, P.,
Lambret, P., Nicolas, D., Olivier, A., Geijzendorffer, I.R., 2019. Estimating biodiversity
changes in the Camargue wetlands: An expert knowledge approach. PLoS One 14,
e0224235. https://doi.org/10.1371/journal.pone.0224235

Galewski, T., Devictor, V., 2016. When common birds became rare: historical records shed
light on long-term responses of bird communities to global change in the largest wetland
of France. PLoS One 11, e0165542. https://doi.org/10.1371/journal.pone.0165542

Graham, L., Gaulton, R., Gerard, F., Staley, J.T., 2018. The influence of hedgerow structural
condition on wildlife habitat provision in farmed landscapes. Biol. Conserv. 220, 122–131.
https://doi.org/10.1016/j.biocon.2018.02.017

Grass, I., Batáry, P., Tscharntke, T., 2021. Combining land-sparing and land-sharing in

European landscapes. Adv. Ecol. Res. 64, 251–303.
https://doi.org/10.1016/bs.aecr.2020.09.002

Herzog, F., Jeanneret, P., Ammari, Y., Angelova, S., Arndorfer, M., Al., E., 2013. Measuring
farmland biodiversity. Solutions 4, 52–58.

Hinsley, S.A., Bellamy, P.E., 2019. Birds of hedgerows and other field boundaries, in: The
Ecology of Hedgerows and Field Margins. pp. 210–232.
https://doi.org/10.4324/9781315121413-11

Horne, B. Van, 1983. Density as a misleading indicator of habitat quality. J. Wildl. Manage.
47, 893. https://doi.org/10.2307/3808148

Jiguet, F., 2003. Instructions pour le programme STOC-EPS. MNHN, pp. 1-18.

Katayama, N., Amano, T., Fujita, G., Higuchi, H., 2012. Spatial overlap between the
intermediate egret egretta intermedia and its aquatic prey at two spatiotemporal scales in
a rice paddy landscape. Zool. Stud. 51, 1105–1112.

Katoh, K., Sakai, S., Takahashi, T., 2009. Factors maintaining species diversity in satoyama, a
traditional agricultural landscape of Japan. Biol. Conserv. 142, 1930–1936.
https://doi.org/10.1016/j.biocon.2009.02.030

King, S., Elphick, C.S., Guadagnin, D., Taft, O., Amano, T., 2010. Effects of landscape features
on waterbird use of rice fields. Waterbirds 33, 151–159.
https://doi.org/10.1675/063.033.s111

Madge, S., 2020. Carrion Crow (Corvus corone), in: Birds of the World. Cornell Lab of
Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.carcro1.01

Madge, S., de Juana, E., 2020. Corn Bunting (Emberiza calandra), in: Birds of the World.
Cornell Lab of Ornithology, Ithaca, NY, USA.

Magnusson, A., Skaug, H., Nielsen, A., Berg, C., Kristensen, K., Maechler, M., Bentham, K.
van, Bolker, B., Sadat, N., Lüdecke, D., Lenth, R., O'Brien, J., Brooks, M., 2020.
Generalized linear mixed models using template model builder. Package glmmTMB.
Version 1.0.2.1 [WWW Document]. R Top. Doc. URL https://cran.r-
project.org/package=glmmTMB (accessed 10.19.21).

Mallet, P., 2022. Rôle des infrastructures et des pratiques agroécologiques pour la conservation
de la biodiversité dans les systèmes de grandes cultures en Camargue.
http://www.theses.fr. Avignon. https://doi.org/10.5281/zenodo.7652924

Mallet, P., Béchet, A., Galewski, T., Mesléard, F., Hilaire, S., Lefebvre, G., Poulin, B., Sirami,
C., 2022. Different components of landscape complexity are necessary to preserve
multiple taxonomic groups in intensively-managed rice paddy landscapes. Agric. Ecosyst.

Environ. 328, 107864. https://doi.org/10.1016/j.agee.2022.107864

Marcon, E., Hérault, B., 2015. Entropart: An R package to measure and partition diversity. J.

Stat. Softw. 67. https://doi.org/10.18637/jss.v067.i08

Marja, R., Herzon, I., 2012. The importance of drainage ditches for farmland birds in

agricultural landscapes in the Baltic countries: Does field type matter? Ornis Fenn. 89,

170–181.

Marshall, E.J.P., Moonen, A.C., 2002. Field margins in northern Europe: Their functions and

interactions with agriculture. Agric. Ecosyst. Environ. 89, 5–21.

https://doi.org/10.1016/S0167-8809(01)00315-2

Mas, M., Flaquer, C., Rebelo, H., López-Baucells, A., 2021. Bats and wetlands: synthesising

gaps in current knowledge and future opportunities for conservation. Mamm. Rev. 51,

369–384. https://doi.org/10.1111/mam.12243

Meyer, S., Wesche, K., Krause, B., Leuschner, C., 2013. Dramatic losses of specialist arable

plants in Central Germany since the 1950s/60s - a cross-regional analysis. Divers. Distrib.

19, 1175–1187. https://doi.org/10.1111/ddi.12102

Montgomery, I., Caruso, T., Reid, N., 2020. Hedgerows as ecosystems: Service delivery,

management, and restoration. Annu. Rev. Ecol. Evol. Syst. 51, 81–102.

https://doi.org/10.1146/annurev-ecolsys-012120-100346

Morganti, M., Manica, M., Bogliani, G., Gustin, M., Luoni, F., Trotti, P., Perin, V., Brambilla,

M., 2019. Multi-species habitat models highlight the key importance of flooded reedbeds

for inland wetland birds: Implications for management and conservation. Avian Res. 10,

1–13. https://doi.org/10.1186/s40657-019-0154-9

Newton, I., 2017. Farming and birds. HarperCollins UK, London.

Norton, M.R., Hannon, S.J., Schmiegelow, F.K.A., 2000. Fragments are not islands: Patch vs

landscape perspectives on songbird presence and abundance in a harvested boreal forest.

Ecography (Cop.). 23, 209–223. https://doi.org/10.1111/j.1600-0587.2000.tb00277.x

Pasher, J., Mitchell, S.W., King, D.J., Fahrig, L., Smith, A.C., Lindsay, K.E., 2013. Optimizing

landscape selection for estimating relative effects of landscape variables on ecological

responses. Landsc. Ecol. 28, 371–383. https://doi.org/10.1007/s10980-013-9852-6

PECBMS, 2022. PanEuropean Common Bird Monitoring Scheme [WWW Document]. Paneur.

Common Bird Monit. Scheme. URL https://pecbms.info/trends-and-indicators/indicators/

(accessed 1.20.22).

R Core Team, 2017. R: A Language and Environment for Statistical Computing.

https://doi.org/https://doi.org/10.1073/pnas.1906419116

Saura, S., Martín-Queller, E., Hunter, M.L., 2014. Forest landscape change and biodiversity
conservation, in: Forest Landscapes and Global Change: Challenges for Research and
Management. Springer New York, pp. 167–198. https://doi.org/10.1007/978-1-4939-
0953-7_7

Shoffner, A., Wilson, A.M., Tang, W., Gagné, S.A., 2018. The relative effects of forest amount,
forest configuration, and urban matrix quality on forest breeding birds. Sci. Rep. 8, 1–12.
https://doi.org/10.1038/s41598-018-35276-9

Singh, S., Sharma, S.N., Prasad, R., 2001. The effect of seeding and tillage methods on
productivity of rice-wheat cropping system. Soil Tillage Res. 61, 125–131.
https://doi.org/10.1016/S0167-1987(00)00188-4

Sirami, C., Gross, N., Baillod, A.B., Bertrand, C., Carrié, R., Hass, A., Henckel, L., Miguet, P.,
Vuillot, C., Alignier, A., Girard, J., Batáry, P., Clough, Y., Violle, C., Giralt, D., Bota, G.,
Badenhausser, I., Lefebvre, G., Gauffre, B., Vialatte, A., Calatayud, F., Gil-Tena, A.,
Tischendorf, L., Mitchell, S., Lindsay, K., Georges, R., Hilaire, S., Recasens, J., Solé-
Senan, X.O., Robleño, I., Bosch, J., Barrientos, J.A., Ricarte, A., Marcos-Garcia, M.Á.,
Miñano, J., Mathevet, R., Gibon, A., Baudry, J., Balent, G., Poulin, B., Burel, F.,
Tscharntke, T., Bretagnolle, V., Siriwardena, G., Ouin, A., Brotons, L., Martin, J.L.,
Fahrig, L., 2019. Increasing crop heterogeneity enhances multitrophic diversity across
agricultural regions. Proc. Natl. Acad. Sci. U. S. A. 116, 16442–16447.
https://doi.org/10.1073/pnas.1906419116

Stanton, R.L., Morrissey, C.A., Clark, R.G., 2018. Analysis of trends and agricultural drivers
of farmland bird declines in North America: A review. Agric. Ecosyst. Environ. 254, 244–
254. https://doi.org/10.1016/j.agee.2017.11.028

Sundar, K.S.G., Subramanya, S., 2010. Bird use of rice fields in the Indian subcontinent.
Waterbirds 33, 44–70. https://doi.org/10.1675/063.033.s104

Tamisier, A., Grillas, P., 1994. A review of habitat changes in the camargue: An assessment of
the effects of the loss of biological diversity on the wintering waterfowl community. Biol.
Conserv. 70, 39–47. https://doi.org/10.1016/0006-3207(94)90297-6

Toffoli, R., Rughetti, M., 2017. Bat activity in rice paddies: Organic and conventional farms
compared to unmanaged habitat. Agric. Ecosyst. Environ. 249, 123–129.
https://doi.org/10.1016/j.agee.2017.08.022

Tourenq, C., Aulagnier, S., Durieux, L., Lek, S., Mesléard, F., Johnson, A., Martin, J.L., 2001.
Identifying rice fields at risk from damage by the greater flamingo. J. Appl. Ecol. 38, 170–
179. https://doi.org/10.1046/j.1365-2664.2001.00581.x

Vallecillo, S., Brotons, L., Herrando, S., 2008. Assessing the response of open-habitat bird
species to landscape changes in Mediterranean mosaics. Biodivers. Conserv. 17, 103–119.
https://doi.org/10.1007/s10531-007-9233-z

Vickery, J.A., Feber, R.E., Fuller, R.J., 2009. Arable field margins managed for biodiversity
conservation: A review of food resource provision for farmland birds. Agric. Ecosyst.
Environ. 133, 1–13. https://doi.org/10.1016/j.agee.2009.05.012

Warren, M.S., Maes, D., van Swaay, C.A.M., Goffart, P., van Dyck, H., Bourn, N.A.D.,
Wynhoff, I., Hoare, D., Ellis, S., 2021. The decline of butterflies in Europe: Problems,
significance, and possible solutions. Proc. Natl. Acad. Sci. U. S. A. 118, e2002551117.
https://doi.org/10.1073/PNAS.2002551117

Wilson, J.D., Anderson, R., Bailey, S., Chetcuti, J., Cowie, N.R., Hancock, M.H., Quine, C.P.,
Russell, N., Stephen, L., Thompson, D.B.A., 2014. Modelling edge effects of mature forest
plantations on peatland waders informs landscape-scale conservation. J. Appl. Ecol. 51,
204–213. https://doi.org/10.1111/1365-2664.12173

Wilson, S., Mitchell, G.W., Pasher, J., McGovern, M., Hudson, M.A.R., Fahrig, L., 2017.
Influence of crop type, heterogeneity and woody structure on avian biodiversity in
agricultural landscapes. Ecol. Indic. 83, 218–226.
https://doi.org/10.1016/j.ecolind.2017.07.059

Yamada, S., Kusumoto, Y., Tokuoka, Y., Yamamoto, S., 2011. Landform type and land
improvement intensity affect floristic composition in rice paddy fields from central Japan.
Weed Res. 51, 51–62. https://doi.org/10.1111/j.1365-3180.2010.00815.x

630