Chromosomal position of ribosomal protein genes impacts long term evolution of

Vibrio cholerae.

Leticia Larotonda[1,2], Damien Mornico[3], Varun Khanna[3], Joaquín Bernal[4,5], Jean Marc

Ghigo[4], Marie-Eve Val[2], Diego Comerci[1] and Didier Mazel[2#], Alfonso Soler-Bistué[1,2#]

[1]Instituto de Investigaciones Biotecnológicas "Dr. Rodolfo A. Ugalde", IIB-IIBIO,

Universidad Nacional de San Martín-CONICET, San Martín, Buenos Aires, Argentina.

[2] Institut Pasteur, Université de Paris Cité, CNRS UMR 3525, Unité Plasticité du Génome

Bactérien, Paris F-75015, France.

[3] Institut Pasteur, Université de Paris Cité, CNRS UMR3756 Bioinformatics and

Biostatistics Hub, F-75015 Paris, France.

[4] Institut Pasteur, Université de Paris Cité, CNRS UMR 6047, Genetics of Biofilms

laboratory, Paris F-75015, France.

[5]Departamento de Genética, Facultad de Biología, Universidad de Sevilla, Apartado

1095, 41080 Sevilla, Spain.

#correspondence to: Alfonso Soler-Bistué: asoler@iib.unsam.edu.ar ; Didier Mazel

dmazel@pasteur.fr

Short title: Experimental evolution on ribosomal protein gene location

Abstract:

It is unclear how gene order within the chromosome influences bacterial evolution. The genomic location of genes encoding the flow of genetic information is biased towards the replication origin (*oriC*) in fast-growing bacteria. To study the role of chromosomal location on cell physiology we relocated the *S10-spec-α* locus (S10) , harboring half of ribosomal protein genes, to different chromosomal positions in the fast-growing pathogen *V. cholerae*. We found that growth rate, fitness and infectivity inversely correlated the distance between S10 and *oriC*. To gain insight into the evolutionary effect of RP genomic position, we evolved strains bearing S10 at its current *oriC*-proximal location or derivatives harboring the locus far from it (i.e. 1.5*M*bp). Populations deep sequencing on average 1 mutation fixed each 100 generations, mainly at genes linked to flagellum regulation, lipopolysaccharide synthesis, chemotaxis, biofilm and quorum sensing. Along the experiment, populations showed an increment in biofilm forming capacity. All populations increased their growth rate. However, growth rate advantage of populations bearing S10 at an *oriC*-proximal persisted along the experiment over those where the main ribosomal protein gene cluster locates at an *oriC*-distal position. This indicates that suppressor mutations cannot compensate S10 genomic location. We selected fast-growing clones displaying a ~10% growth rate increment finding that they harbored inactivating mutations at, among other sites, flagellum master regulators *flrAB* regardless S10 genomic location. The introduction of these mutations on naïve *V. cholerae* strains resulted in a ~10% increase in growth rate. Our study therefore demonstrates that the location of ribosomal protein genes conditions the evolutionary trajectory of growth rate in the long term. While genomic content is highly plastic in prokaryotes, gene order is an underestimated factor that conditions cellular physiology and lineage evolution. The lack of suppression enables artificial gene relocation for genetic circuit reprogramming.

Introduction:

In all known life forms, DNA stores the genetic information in a long polymer whose length surpasses the cell dimensions by thousands of times (1). As a consequence, the genetic material is highly packed by specific proteins that confer its structure and position within the cell (2-5). There is relatively little knowledge on the physiological role of spatial organization of the chromosome. The simplest model to tackle this issue is the bacterial cell due to its relatively small genome size and chromosome number. The bacterial chromosome is tightly organized within the confined space of the cytoplasm and along the cell cycle (6-13). This is physiologically relevant since DNA simultaneously serves as template for replication, transcription, segregation and the repair of the genetic material (1, 4, 14). Bacterial chromosomes have a single replication origin (*oriC*) from which DNA duplication begins bi-directionally until the terminal region (*ter*) where both replication forks meet. This configuration organizes the genome into two generally equally sized replichores along the *ori-ter* axis (6, 15, 16). Recent studies suggest that gene order contributes to link genome structure to cell physiology (14, 16-20). The genomic position of key gene groups is widely conserved along the *ori-ter* axis following the temporal pattern of gene expression (17, 18, 21, 22). Thus, coding sequences expressed in exponential phase are physically associated to the *oriC* region while factors transcribed in stress situations or in stationary phase cluster close to the *ter* region (17, 21, 23-26). Consistently, it was recently shown that some genes need a specific genomic location to achieve their function (16). In most cases, the genomic position of a gene conditions its genome-wide copy-number since those located close to *oriC* are overrepresented with respect to those close to the *ter* region during DNA replication (16, 27). This differential template abundance, called gene dosage effect, leads to increase the expression of genes located in the proximity of *oriC,* although this might be more

complex (28, 29). The *ori-ter* template abundance gradient is particularly important in

fast-growing bacteria during steady-state growth when cells display generation times

shorter required time to complete genome replication. In these conditions, successive

replication rounds overlap, a mechanism called Multi-Fork Replication (MFR) (16, 27).

In fact, many essential and highly expressed genes are clustered in this chromosomal

region (6, 24, 30-32). In parallel, some other genes require a precise location to function,

independently of their dosage (33, 34). Other coding sequences, particularly those highly

expressed, impact the chromosomal spatial structure (10, 11) suggesting an interplay

between gene order and chromosome structure(35).

Vibrio cholerae is a human pathogen of prime importance able to colonize the intestine

causing the cholera disease that kills around 100,000 persons yearly (36). This

microorganism is a common inhabitant of estuarine environments often associated to

plankton (37, 38). It is a highly motile bacterium thanks to a single, sheathed polar

flagellum. The flagellum biogenesis depends on a regulation cascade whose master

regulator is the FlrA protein (39). *V. cholerae* also alternates from a mobile to a sessile

lifestyle by forming structured biofilms, a key trait for both parts of *V. cholerae* life-cycle

(40-42). Importantly, flagellum-mediated swimming, chemotaxis, biofilm formation,

colonization, and virulence are intimately related processes connected through complex

regulatory cascades (40, 43). In particular, flagellum-mediated motility and biofilm

formation are inversely regulated by the concentration of the second messenger molecule

cyclic di-guanosine monophosphate (c-di-GMP)(43-45). This molecule is in turn

modulated by proteins carrying GGDEF and/or EAL domains displaying diguanylate

cyclase and phosphodiesterase activity respectively(43-46).

V. cholerae displays a short doubling time (~16 minutes) and high replication-associated

gene dosage effects (24). It is also a model of bacteria bearing multiple chromosomes

since it harbors a main chromosome (Chr1) of 2.96 Mbp and a 1.07-Mbp secondary

replicon (Chr2) (47). Replication of both chromosomes is coordinated. Chr1 first starts

its replication while Chr2 fires its replication origin (*ori2*) only after 2/3 of Chr1 are

duplicated and both chromosomes complete DNA synthesis synchronously (48).

Nevertheless replication-dependent gene-dosage effects exists in both replicons (30, 49,

50) while their relative timing modulates the gene dosage difference between both

replicons. In *Vibrionaceae,* transcription and translation genes map close to the *oriC* of

Chr1 (*ori1*) (31, 49).

We have previously studied the chromosomal position of ribosomal protein (RP) genes

to uncover the link between gene order and cell physiology (24, 51). For this we

manipulated the genomic location of *V. cholerae* major ribosomal protein locus *s10-spc-*

α (S10), a 13 Kbp array of essential genes highly conserved among prokaryotes and also

in eukaryotic organelles (52, 53). By relocating S10 locus along both chromosomes (49)

we generated a S10 *movant* strain set (i.e., isogenic strains where the genomic position of

the locus of interest is altered) whose growth rate, infectivity and fitness decreased as a

function of the distance between S10 and *ori1*. Relocation to the Chr2 showed a

detrimental effect accordingly. These phenotypes were caused by the reduction of S10

genome-wide copy number that were mostly consequence of macromolecular crowding

alterations rather than a deficit in the capacity to synthesize proteins (49, 50, 54). Further

increasing S10 copy-number was deleterious our neutral. Genomic position of S10

maximizes cell physiology and fitness along both parts of *V. cholerae* lifecycle.

Meanwhile, S10 is close to *ori1* among all *Vibronaceae.* Since this clade diverged from

enterobacteria some 500 million year ago (55), such genome positioning is likely to be

the result of selection in the long run rather than trait acquired through genetic drift. In

the last years, long term evolution experiment emerged as the experimental approach of

choice to directly assess evolutionary trajectories of (micro)organisms (56-59).

In this work, we analyzed how the genomic position of the major ribosomal protein locus

could condition *V. cholerae* evolution in the long term. For this, we experimentally

evolved strains in which the S10 locus is either placed at its original position or at the

terminal region of the chromosome. We showed that the growth rate reduction observed

in the latter case persisted over 1000 generations of evolution providing experimental

support to the biased genomic position of these genes in fast growing bacteria. Our

positional genetics approach also showed that increased biofilm formation and increased

growth rate at the expense of motility is a common adaptations of *V. cholerae* to in vitro

environment. Our study therefore contributes to improve our understanding of the role of

gene order on bacterial cell physiology and provides insights into the long-term

implication of gene order on genome evolution.

Results:

Vibrio cholerae **evolution experiments.** We conducted experimental evolution of single

clones of the parental or different *V. cholerae* movants for 1,000 generations. For this, we

started 3 populations from isolated colonies of each strain. Due to the fast growth of *V.*

cholerae, we passaged them twice a day in Lysogeny Broth (LB) supplemented with

streptomycin (Str) to avoid contamination (See Suppl. Material). On one hand, we

employed the parental strainS10 was not transposed (49), as reference (Populations P1,

P2 and P3, Figure 1A, Table S1). We also conducted the experimental evolution of

S10Tnp-35, a movant where the S10 locus was slightly relocated (P4 to P6, Figure 1A,

Table S1). This movant displays no fitness nor growth rate defect (Figure 1B), it is

isogenic with respect of to the rest of the movants and shows only 8 genes that are

differentially expressed with respect to the parental strain (54). These populations were

used as a control that any possible genetic change associated to S10 relocation *per se*

could impact the evolution of *V. cholerae*. In parallel, we evolved S10Tnp-1120 (P7 to

P9) and S10TnpC2+479 (P10 to P12) (Figure 1A, Table S1) where S10 was relocated to

the *ter* region of Chr1 and Chr2 respectively. These are the most phenotypically affected

movants displaying a ~15% loss of maximal growth rate and lower fitness than the

parental strain (Figure 1B) (50). The populations were frozen every 50 or 100 generations

(Supplementary Text). We also isolated single clones at 250 and 1000 generations for

later analysis.

Evolved *Vibrio cholerae* populations acquire increased biofilm formation capacity.

First, we noticed the emergence of strong biofilms on the Erlenmeyer flasks as early as

generation 81 (G81). By generation G100, most of the populations displayed strong

biofilms in the air-liquid interphase and/or dense cell aggregates in the medium (Figure

S1A). By G250 biofilms were present in all populations. Accompanying this

phenomenon, colonies of abnormal morphology appeared, particularly, rugose colonies

(Figure S1B). The subculture of isolated smooth (S) and rugose (R) colonies showed that

only the latter formed strong biofilms in the flasks (Figure S1C). To quantitatively address

biofilm-forming capacity of the populations along the experimental evolution, we

cultured them statically in 96-wells polystyrene plates, washed and then stained the wells

with crystal violet at G0, G100, G250, G500, G700 and G1000 (Figure 2). As positive

and negative controls we analyzed R1, a rugose clone isolated from P1 at G1000 (see

above) and the parental strain respectively. At the beginning of the experiment (G0), the

levels of crystal violet stain are minimal. Then biofilm-formation capacity increased

rapidly in all populations. In average, biofilm formation capacity reached a maximum at

~G250 (Figure 2) although some specific populations displayed an earlier peak near

~G100 (Figure S2). After G250, the biofilm formation capacity decreased in all evolved

populations without going back to the initial G0 baseline levels. The emergence of rugose

colonies paralleled evolution of biofilm forming capacity (Figure S1D). All populations

displayed a similar trend suggesting that it is selected along the experiment independently

of S10 genomic location (Figure 2 and Figure S2).

Evolved populations display growth rate increase. Previous studies showed that

bacterial populations increase their fitness along time compared to the ancestral clone (58,

60). Direct fitness determination from competition experiments results is difficult to

achieve in populations that form biofilms dynamically across generations. Therefore, we

employed maximum growth obtained rate (μ) as a proxy for fitness estimation. In

previous works, both parameters changed similarly as a function of S10 genomic location

(50). Therefore, the evolved populations were subjected to automated growth curves. For

this, at each time point, the twelve populations were diluted 1 in 1000 and cultured in LB

at 37°C measuring OD_{620nm} by triplicate at least twice. In each experiment, the 4 ancestral

strains were used as control. The value of μ was calculated from the growth curves and

relativized to the parental strains to remove variations between experiments. Then we

plotted the maximum growth rate as a function of elapsed generations (Figure 3A, Figure

S3A). During the first 150 generations, μ remained stable among populations. As

expected (49, 50), populations derived strains in which S10 is close to *ori1* (P1 to P6)

grew ~15% faster than those originated from the most affected movants (P7 to P12). After

this period, between G150 and G250, populations increase their μ. All populations

increased their growth rate along the experimental evolution but this rise occurred in a

stepwise manner, the most notorious occurring at G250. After G500, the alterations in

growth rate become noisier since populations originating from the same strain tend to

diverge (Figure S3A). Despite the general μ increment along generations, the differences

between populations evolving from faster strains (P1-P6) and slower ones (P7-P12)

persisted along the experiment (Figure 3B). Also, the evolutionary trajectory of this trait

was similar for all of them (Figure S3B). Importantly, we did not notice suppressor

mutants compensating the growth rate deficit in P7-P12. Hence, mutations able to

compensate the genomic position of S10 locus are very difficult to achieve, at least in this

time frame. This, strongly suggests that the S10 genomic location influences growth rate

along experimental evolution strongly indicating that the position of the main ribosomal

protein locus determines fitness in the long run.

Positive selection drives genome dynamics   in the first 250 generations of

experimental evolution. To better characterize mutations occurring along the

experiment, we deep sequenced each ancestral strain as reference and each population at

G250 and G1000. We speculated that mutations at G250 would correlate to the earlier

increase in growth rate and biofilm-formation capacity emergence.

Deep sequencing of the 12 populations at G250 revealed 128 mutations observed in more

than 25 % of the reads obtained from each culture (Dataset S1). Among them, we found

45 different mutations among all the populations. The number of fixed DNA changes

varied from 2 to 21 between each individual populations (Figure 4A). On average, each

characterized mutation occurred in 2-3 populations strongly indicating that they were

under positive selection. Most mutations (26 out of 45) corresponded to non-synonymous

mutations or frameshifts of coding sequences (Figure 4B). These changes are likely to

alter or interrupt their encoded functions. Meanwhile, 17 changes (42%) occurred at *a*

priori non-coding regions such as intergenic spaces and pseudogenes (Figure 4B). We

detected 2 (4.4%) synonymous mutations. When grouped by coding sequences (CDS) we

found that at least 17 genes suffered alterations (Dataset S1). Among them, 6 genes

presented more than one mutation across multiple populations: *aroG*. *ompT*, *flrA, flrB,*

flrC and *tagE2* (Table 1). First, *ompT* (61) is a highly expressed gene encoding one of the

main porins of *V. cholerae*. *tagE2* (VCA1043) was found altered in most of the

populations. There is little evidence of its encoded function although a recent study shows

it is an endopeptidase necessary for cell growth (62). The *aroG* gene is required for the

synthesis of aromatic amino acids although there is little work done on *V. cholerae*.

Finally, we found SNPs at *flr* genes, the main regulators of flagellum biosynthesis (39,

63, 64). FlrA is the master regulator of flagellum biosynthesis controlling the expression

of *flrB* and *flrC* which in turn, are downstream in the flagellar transcription hierarchy

(39). We reasoned that this multiplicity of independent mutational events within the

genetic pathway suggests a fitness benefit from motility inactivation. As a general

scenario, we found that at G250 mutations occur several times independently, suggesting

a positive selection of most of them rather than genetic drift. In particular, we observed

that a majority of the populations displayed mutations at the main flagellar regulators

(DataSet S1, Table 1).

Evolutionary forces change beyond G250: adaptive, non-adaptive mutations

and hypermutators after 1000 generations of experimental evolution. We speculated

that mutations observed at G1000 may reflect those that are fixed in the long run since

after ~G500 biofilm formation decreases and growth rate remains roughly constant. After

1000 generations of experimental evolution, we observed 512 fixed changes in at least

25% of the population at 396 sites of the genome of *V. cholerae* (Figure 4A, right bars).

Contrasting to G250 scenario, few mutational events occurred in more than one

population (1.3 events per populations on average). We observed a lower proportion of

mutated pseudogenes (4.5% vs 26.6%) and an increase in the proportion of synonymous

mutations (4.4% vs 19.4%) (Figure 4B). We also noticed mutational events at several

genes involved in DNA repair such as *mutT* (VC1342), *recBCD* (VC2319-VC2320),

mutS (VC0535), *recJ* (VC2417). Particularly, *mutL* (VC0345) was mutated in 8 out of 12

populations (1, 2, 3, 4, 5, 6, 7 and 11). Indeed, populations with intact DNA repair

mechanisms (i.e. P8, P10; P12) are those showing lowest number of mutations (Figure

4A). These data suggest that in most populations at G1000, many of the identified

mutations may result from an hypermutator phenotype rather than selection. All, fast-

growing populations (P1 to P6) showed one or more mutations in DNA repair systems.

In contrast half of the slower-growing populations (P7 to P12) displayed an intact DNA

repair pathway (Figure 4A). Nevertheless, we also detected many DNA changes in the

same gene or operon occurring many times independently or emerging in parallel in more

than one population likely indicating positive fitness selection along the evolution

experiment (Table 2). On top of *aroG, ompT, flrABC* and *tagE2* already acquired at G250,

we found many mutations in interlinked cellular processes (38, 43, 44) such as rugosity,

biofilm structure and regulation (*vpsA, vpsB, vpsP, vpsR, capK and rbmB*) (65-68),

quorum-sensing (*luxP, luxO* and *luxU*)(69), flagellum biosynthesis (*flgA, flgI, flgK, fliG,*

fliO, flag, flhA)(39) and chemotaxis (*cheY, cheR2*) (39, 70). We also identified several

altered functions such as extracellular macromolecular structures linked to virulence

(toxin coregulated pilus (*tcp*), mannose-sensitive hemagglutinin pilus (MSHA), type 6

secretion system (T6SS)), iron incorporation and metabolism (particularly vibriobactin

pathway genes and *iscR*), lipopolysaccharide biosynthesis (*rfbA, rfbB, rfbV, pgi*) and

phosphate sensing and incorporation (*phoR, phoU, phoH*). Interestingly, the latter have

been linked to biofilm downregulation (71). Most of them were non-synonymous or led

to frameshifts likely to be detrimental for their biological function (Data Set S2). We

interpret these mutations in the light of a change from a host-environmental life cycle to

the relatively constant culture condition of the evolution experiment. Functions such as

virulence, swimming, chemotaxis, pili production, T6SS and micronutrient scavenging

are energetically costly structures under strong selection in *V. cholerae* natural

environment that may become a burden during *in vitro* evolution.

Mutations linked to rugose phenotype. Along the experimental evolution we observed

the emergence of biofilms and aggregates linked to a rugose phenotype described earlier

(Figures 2, S1 and S2). To uncover genomic changes associated to biofilm emergence we

fully sequenced the genome of a rugose and a smooth clone from a slow growing (P1)

and a fast-growing population (P12). After 1000 generations, clones originated from P1

and P12 fixed 9 and 6 mutations respectively (Figure 4C) independently of their smooth

(S) or rugose (R) phenotype (1 mutation fixed every 133±30 generations). Although some

DNA changes were common among R and S colonies (Data Set S3), we found some

differential mutations potentially involved in biofilm and rugose phenotype emergence

(Table 3). In particular, mutations in *vpsR, vpsB, cdgB* and VCA1031 were previously

linked to rugose phenotype by Yildiz and co-workers (46, 72, 73). The *vpsR* is a positive

regulator of the *vps cluster* that suffered a non-synonymous mutation (D40Y) in the S

colony derived from P1. The *vpsB* gene codes for an enzyme participating in the synthesis

of nucleotide sugar precursor necessary for polysaccharide biosynthesis. We detected a

premature stop codon at the S colony collected from P12 (P12S). The P12R displayed a

non-synonymous mutation (D101G) at the *cdgB,* a GGDEF protein upregulated in rugose

clones (73, 74). It also showed a non-synonymous mutation in the *iscR* gene which codes

for a transcription factor regulating genes related to iron starvation, oxidative stress and

oxygen limitation in *Escherichia coli* and linked to virulence in other bacteria (75).

Meanwhile, the P1S colony displayed a mutation at VCA1031, encoding a putative

methyl-accepting chemotaxis proteins previously reported to be overexpressed in rugose

strains having a genetic *ΔcdgC* background(46). In parallel, P1R showed an intergenic

mutation also found at G250, G1000 and on fast-growing clones (see above). Finally,

P12S displayed a synonymous mutation in the *rpoA* gene, a main component of the RNA

polymerase. We found two other differential mutations unlikely to be causal rugose

phenotype but that might have been co-selected. In summary, the S and R clones showed

alterations in genes previously linked to biofilm generation and c-di-GMP regulation. We

detected, however, other changes in R clones such as in *iscR* and at the intergenic region

between VC0448-VC0449.

Increase growth rate is often associated to motility loss.

To uncover mutations associated to fast growth, we isolated 4 smooth colonies and a

rugose clone (S1 to S4 and R respectively) from P1 to P12 at G250 the moment along

experimental evolution where we noticed a sudden growth rate increase in all populations.

Next, we performed automated growth curves of these clones using as reference the

parental strain Par-1120 (Table S1). We calculated μ for each strain (Figure 5A) and we

selected the fastest growing clones from P1 to P6 and from P7 to P12. We called these

clones F̲ast G̲rowing clones followed by population of origin and its smooth or rugose

status. Thus, we isolated the clones FG-2-S2, FG-3-S2, FG-7-S1, FG-10-S1 and FG-11-

S1. Interestingly, the 3 former clones grew 7 to 10% faster than the parental strain.

Meanwhile, FG-G250-7-S1, FG-G250-10-S1 and FG-G250-11-S1 improved their growth

rate compared to their ancestors, they still displayed a μ reductions of ~7% with respect

to the parental strain (Figure 5B). To identify the mutations responsible of μ increase, we

fully sequenced FG-G250-2-S2, FG-G250-3-S2, FG-G250-7-S1, FG-G250-10-S1 and

FG-G250-11-S1 genomes and compared them to the ancestral strains. Overall, strains

displayed 1 to 3 changes in the DNA sequence (1 mutation fixed every ~140 generations)

(Figure 4C). In the 5 strains we found 8 different mutations (Table 4). All the isolated

strains harbored alterations either at *flrA* (4 out of 5) or at *flrB,* (1 strain) the main

flagellum regulators which were previously observed at the population level at G250

(Table 1). Three other mutations accompanied the latter: a non-silent mutation at a

magnesium transporter (*mgtE*), a point mutation at an intergenic base between genes

VC0448 and VC0449 and a frameshift mutation at *ompT*. Except for the latter, we found

the exact same mutations at G250 and most of them at G1000 (Dataset S4).

To demonstrate the role of these mutations on growth rate enhancement we performed

Multiplex Genome Editing by Natural Transformation (MuGENT) (76, 77) on naïve *V.*

cholerae N16961 to introduce point mutations, small indels or short insertions. *flrA* is

higher in the hierarchy of the flagellum regulation (39) and it is more prevalent on FG

strains. FlrA is a 488 aminoacid-long protein bearing a REC domain, a central ATPase

Associated with diverse cellular Activities (AAA+) and a C-terminal Helix-Turn-Helix

domain responsible for DNA binding (78). The detected *flrA* frameshifts mutations on

populations or on FG-G250 clones occurred exclusively at the end of this latter domain,

on positions 418-488 of the protein. The non-synonymous mutations occurred at the

AAA+ domain (positions 122 to 417 of FlrA). Using a *V. cholerae* wild type background,

we generated a frameshift at the same in *flrA* position (Fig S4) obtaining several

independent mutant clones that we tested on automated growth curves (Figure 6A). This

mutation increased growth rate with respect to wild type strain. Similarly, the introduction

of a frameshift in *flrB* resulted in a growth rate improvement. Also, the introduction of a

non-synonymous mutation at *mgtE* significantly improved growth rate (Fig 6A). To

quantify the effect of individual mutations we determined the percentage of generation

time variation (%GT). Mutants bearing the *flrA* frameshift displayed the strongest

reduction with a 13.41±4.7 % reduction of their generation time (95% CI [14.9-11.9],

p<0.0001). Meanwhile, *flrB* and *mgtE* mutants displayed a 6% of generation time

reduction (95% CI of [6.9-5.4] and [7.5-4.6] respectively, p<0.0001). These results

coincide with the observed increase in growth rate of FG clones. Simultaneously,

frameshifts in *flrA* and *flrB* caused a reduction of motility as shown by swimming assays

(Fig. 6B) demonstrating that growth rate increase is associated to the loss of *flrA* and *flrB*

function.

In summary, these results demonstrate that the mutations observed in the FG strains

increased growth rate in *V. cholerae*. They also suggest some of the mutations detected

at G250 and maybe at G1000 also contributed to the growth optimization of this

microorganism.

Discussion:

Increasing evidence suggests that gene order within the bacterial chromosome contributes

to cellular homeostasis by coordinating key entangled processes such as chromosome

structuration, cell cycle, replication and the expression of genetic information. While

genome reshuffling experiments have not been yet performed in bacterial systems (79)

approaches such as comparative genomics (17, 18, 21, 23, 31, 80), systems biology (20),

large DNA inversions (81-83) and relocation of individual gene sets (16, 19, 84-86) have

provided support to this notion. Genes encoding for the genetic information flow are

interesting models to test the role of genomic position on cellular physiology. The

chromosomal position of rRNA, some tRNAs, RP and RNA polymerase genes is biased

towards the *oriC*, particularly in fast-growing bacteria (17, 22, 24, 51). Relocation of S10,

the main RP locus harboring half of its encoding genes, far from *oriC* constrains

bacterial growth (49). Competition experiments of S10 movants against wild type strains

suggested that RP genomic position implied a strong fitness cost that must had

evolutionary consequences (50). In this work, we took this "positional genetics" approach

(in which genes of interest are systematically relocated within the genome) a step further

by observing how movants bearing S10 at different genomic locations evolve *in vitro* for

1000 generations in fast growing conditions, those in which phenotypic effects were

stronger (Figure 1).

Comparison with previous experimental evolution approaches. Pioneer work by

Richard Lenski and co-workers on long term evolution experiment was done for many

more generations than ours (i.e. more than 75,000 (and counting!) for *E. coli*) however

the strongest fitness improvement occurred during the first 2,000 generations (58, 60).

Therefore, although we performed a significantly shorter experiment, we believe that we

observed a representative part of the evolutionary trajectory. Meanwhile we developed a

different approach: we used rich media and a model organism that grows almost twice as

fast as *E. coli*. On the other hand, our system used 1 log larger bottlenecks that previous

studies improving the chance of mutant selection and lowering the effects of genetic drift

(i.e. we transferred 10^9 instead of 10^8 CFU). Also we evolved populations with same

genotype but bearing the key locus S10 at different genomic locations.

Biofilm dynamics along *V. cholerae* evolution. Along this evolution experiment we

found the emergence of abundant biofilms and aggregates in every population. As in prior

studies (67, 68, 72, 73), this phenotype was linked to the appearance of colonies of rugose

morphology. Usually there are different types of rugose phenotypes in *V. cholerae* (65),

but exploring their emergence along the experimental evolution was beyond the

objectives of our study. Other experimental evolution approaches also led to biofilms

and/or aggregates. In *Acinetobacter baylyii* ADP1 (87) it was the consequence of IS

action that led to evolve aggregating clones. The studies on *Burkholderia cenocepacia*

(88) and *Bacillus subtilis* (89) actively selected for biofilm emergence with a wide variety

of clones with clear fitness advantages such as labor division and heterogenicity

emergence within the structure. In our system, these traits emerged rapidly in all

populations. Then, biofilm formation was reduced along the generations (Figure 2). Such

dynamic behavior may reflect its selection by association to other traits such as flagellum

loss, followed by decline (see above), the differentiation of more specialized sub-

populations that are more efficient to specific microenvironments (such as air-liquid

interphase), the frequent but stochastic emergence of this trait or the beneficial nature of

this characteristic early along *in vitro* evolution. These possibilities are not exclusive.

Meanwhile, we also characterized some mutations that could explain the emergence of

rugose clones associated to biofilms. A limitation of our study is that we sequenced rugose

clones at G1000 instead of G250 when most of biofilms are observed. At this timepoint,

many other unrelated mutations could be co-selected with the DNA changes causing

rugose phenotypes. This is reinforced by the fact that, similarly to previous work, we

found mutations in several DNA repair systems. Whereas, understanding biofilm

development was not the primary goal of our work, we described its emergence along the

experimental evolution and we observed some mutations co-occurring with this

phenotype that are in agreement with previous seminal papers by Yilditz and co-workers

(38, 66-68, 72, 74).

Growth rate evolution along generations. Another clear phenotype that arose along the

experiment was growth improvement of all *V. cholerae* populations (Figure 3). This trait

was not initially noticed in the long term evolution experiment performed in *E. coli* by

Lenski and co-workers. Meanwhile, fitness and cell size increment was observed since

the initial steps of the experiment (60, 90, 91). Only recently, an increase in growth rate

was also measured (92) which is in concordance with previous studies that show a close

correlation between growth rate, cell size and fitness (93). In *V. cholerae* experimental

evolution, the reduction of populations doubling times was clearer than in *E. coli* since

growth rate could be detected using automated high-throughput growth measurements.

This behavior could be the result of employing a richer broth, the evolution strategy

(wider bottlenecks), and/or the faster growth nature of *V. cholerae*. Meanwhile, growth

rate is an acceptable proxy for fitness particularly in our system as we observed in

previous work (50, 94, 95). Therefore we propose that the increment in growth rate

observed along *V. cholerae* experimental evolution reflects, at least in part, the expected

fitness improvement. The stabilization of growth after G500 can be the result of high

divergence between populations or the consequence of terminating the experiment

relatively early, since further generations accumulation probably should lead to new

increases in growth rate in a stepwise manner.

Two phases in the evolution dynamics of *V. cholerae* genome. Deep sequencing of the

ancestral clones and the populations at G250 and at the end of the experiment suggested

different selection forces at each time point. At G250 each mutation occurred in parallel

in ~3 populations strongly suggesting a positive selection. Since these populations

displayed a sudden gain in growth rate and biofilm forming capacity, we could associate

these mutations (Table 1 and DataSet1) to the observed phenotypes (Figs. 2 and 3). This

scenario changes at G1000 since we find that most mutations are represented in single

populations suggesting that most of them arise by chance not being under strong selection.

This is reinforced by the fact that many populations show inactivation of DNA repair

systems. As expected, populations displaying mutations in repair systems resulted on

average on more DNA changes after 1000 generations of evolution (Figure 4a).

Interestingly, populations evolved from faster strains (P1-6) displayed more alteration in

repair systems and concurrently more mutations at G1000. We speculate that due to their

faster growth rate P1-P6 spent more time at stationary phase along the experimental

evolution. These repeated stress situations might have increased the chances of mutator

phenotype emergence.

Some DNA changes arose in more than one flask in particular inactivating genes encoding

functions that might become unnecessarily costly during in vitro continuous culture (i.e.

virulence, motility and nutrient scavenging). Interestingly, those acquired at G250 are not

lost suggesting their continuous selection along the experiment. Moreover, in the case of

flagellum biosynthetic pathway we found mutations in transcriptional regulators of lower

hierarchy such as *flhA* or components of the machinery such as *flgA*, *flgI* or *flgK*. We also

noticed several mutations at quorum-sensing and chemotaxis-related genes at G1000 (see

above). The former seems to be a common feature of laboratory domesticated *V. cholerae*

strains as previously described by Blockesh Laboratory (96).

At both timepoints, we detected several mutations occurring at *a priori* non-coding

regions of the chromosome such as intergenic regions and pseudogenes. We cannot

discard a biological effects through interfering promoter or regulatory regions, potential

uncharacterized RNA genes or pseudogenes expressing truncated but functional proteins.

This might be the case of those occurring in several populations independently such as

the point mutations occurring at VCA0168, a pseudogene located in the secondary

chromosome, mutated in 9 out of 12 populations (Dataset S1-S3). Interestingly,

VCA0168 is actively transcribed from an upstream transcription start site. It forms an

operon with VCA0169 and VCA0170 that is conserved among other *V. cholerae* strains.

Mutations further increasing *V. cholerae* growth. Importantly, we were able to

distinguish mutations that improved bacterial growth from those increasing biofilm-

forming capacity. Notably, the master regulator for flagellum biosynthesis, *flrA,* and *flrB*

downstream in the same regulation cascade, displayed mutations at G250 and on the FG

smooth clones. These genes were intact in rugose clones isolated at G1000 indicating that

we could discriminate between DNA changes improving growth rate from those involved

on biofilm formation (Fig. 5 and Table 4). In the case of *flrA*, we detected 3 non

synonymous mutations and two frameshifts occurring only in the last 20-30 amino acids

of this 488 codon ORF at the HTH domain. Mutations at *flrA* emerged in 4 and 11 out of

12 populations at G250 and G1000 respectively. In most cases, these DNA alterations led

to the described frameshifts (Dataset S1, S2 and S4). The introduction of a similar

frameshifts into *flrA* and *flrB* in a wild type background increased *V. cholerae* growth

demonstrating its effect on growth and strongly suggesting that the rest of the detected

mutations contributed to the enhanced growth of the populations (Fig 6A). Both

mutations also caused a reduction in motility (Fig 6B) indicating that these mutations alter

the DNA binding capacity of FlrA and FlrB. Recent studies from Sourjik and colleagues

provide an ecological and physiological interpretation of the genes that mutated along our

experimental evolution (97-99). Their work pointed out a clear trade-off between motility

and growth rate. In changing environmental conditions, cells deal several simultaneous

energetically costly processes such as growth, motility chemotaxis and metabolism.

Energy allocation to flagellar genes production, rotation of flagellum and, to a lesser

extent chemotaxis, significantly reduces growth in *E. coli* (98, 99). It is advantageous to

dedicate energy to motility and chemotaxis in poor and heterogenous environments. The

FG-G250 clones displayed mutations on the main regulators of flagella (*flrAB*) which

permit the allocation of more resources for growth outcompeting wild type strains.

Consistently, at G1000 we also noticed many mutations related to other swimming and

chemotaxis genes (e.g. *fliO, flrC, cheY, cheR2, flgA*). In summary, many of the mutations

detected along our experiment alter the motility/growth interplay in all the studied

populations. However many other mutations that contribute to improve the fitness remain

to be better characterized. For instance, mutations on *mgtE*, which is not clearly related

to motility also improved growth rate. These opens the possibility of finding other

mutations on genes not related to motility capable of improving the growth of an already

very fast growth bacterium.

Conclusions. In this study, we showed that the genomic location of ribosomal proteins

conditions cell physiology in the long term. Growth impairment caused by ectopic

location of the S10 locus was not suppressed by mutations elsewhere in the genome. After

1000 generations of experimental evolution, although all populations shortened their

generation time, those harboring S10 locus close to *ori1* remained the fastest. This

evidences the great potential of positional genetics approach (49) as a Synthetic Biology

tool to condition cell physiology in a permanent avoiding the escape of suppressor

variants. The latter conditions the design of stable genetic circuits. For instance, recent

work by Izard *et al*. (100) created a growth switch in *E. coli* by controlling RNA

polymerase expression from an inducible promoter than required 3 copies of the main

repressor to avoid regulation escape. Despite the powerful nature of this synthetic circuit

some escape could still be detected. Positional genetics offer a new way to cope with this

problem to generate robust genetic circuits immune to suppressor mutations.

By applying the positional genetics approach on ribosomal protein genes, we contributed

to understand the role of gene order on bacterial cell physiology. The present work

incorporates the evolutionary aspect to studies on positional genetics expanding our

knowledge on the long-term implication of gene order on genome evolution. The role of

the genomic position of other key genes remain to be explored such as ribosomal

RNAs(101), tRNA[UBI](22, 24), RNA polymerase(51), ATP synthase(17), among others.

This is central to understand how genome's primary structure impacts cell physiology, in

particular growth rate and genome evolution. Such knowledge will enable bacterial

growth repurposing while allowing to understand the behavior of more complex

biological systems thus promising a deep impact in genome design, bioengineering and

biotechnology.

Materials and methods:

Bacterial strains and plasmids and culture conditions. All strains derive from *V.*

cholerae serotype O1 biotype El Tor strain N16961 with is Str^R(102). Bacterial growth

was done in Lysogeny Broth (LB) at 37ºC agitating at 200 rpm. Strains and plasmids

used in this study are listed in Table S1.

General procedures. Genomic DNA was extracted using the GeneJET Genomic DNA

Purification Kit while plasmid DNA was extracted using the GeneJET Plasmid Miniprep

Kit (Thermo Scientific) or ADN Puriprep B and P (InbioHighway, Argentina). PCR

assays were performed using DreamTaq, Phusion High-Fidelity PCR Master Mix

(Thermo Scientific) or MINT 2X (InbioHighway, Argentina). Oligonucleotides,

described in Table S2 were purchased from Macrogen, Ko.

Experimental evolution setting: Starting from individual colonies, 3 independent

cultures of each strain were propagated in 250 mL (7 generations) or 500 mL (8

generations) Erlenmeyer flasks. For each passage 1 mL of the previous flask is transferred

to the next step. Populations were regularly checked for contamination by plating in LB

Str. Colonies with unusual morphologies were checked phenotypically and by PCR

amplifying the *V. cholerae* specific gene *rctB (3398-3407*, Table S2*)* or a specific region

of the secondary chromosome (Tgt4_1-Tgt4_4, Table S2). Every 50 or 100 generations,

the populations were frozen to build the "Frozen Fossil Record" by centrifuging 10 mL

of each culture at 5000 rpm and resuspended them on 1 mL of LB supplemented with

10% of dimethyl sulfoxide (Merck). Then cultures were frozen at -80°C until later use.

(For detailed description see Supplements)

Automated growth curve measurements: ON cultures of the indicated microorganism

were diluted 1/1000 in LB. Bacterial preparations were distributed by triplicate, for

evolved populations, and sextuplicates, for the ancestral clones, in 96-well microplates.

Growth-curve experiments were performed using a TECAN Infinite Sunrise microplate

reader, with absorbance (620nm) taken at 5-minutes intervals for 5 hs on agitation at

37°C. Experiments were done at least twice independently. Slopes during exponential

phase were directly obtained using a home-made Python script coupled to growthrates

program (73). To filter noise due to growth variations among experiments, the obtained

µ was relativized to the ancestral clones and the relativized values were multiplied by the

average of the ancestral clones along all the experiments. This normalization did not

affect the results since raw data provide a similar picture as depicted in Figure S3. The %

GT was calculated as follows: $\%GT = (GT/GT^{WT}-1)*100$

Biofilm formation assays: PVC 9-well microtiter plates (BD Falcon) were used to

monitor biofilm formation as described previously (103). Briefly, LB was inoculated with

a 1/100 dilution directly obtained from the liquid overnight cultures of each population

in rich medium. After inoculation, microtiter plates were incubated at 37 °C for 24 h,

rinsed and 150 µL of a 0.1% solution of crystal violet was added to each well. The plates

were incubated at room temperature for 30 min and rinsed, and biofilm formation was

tested as follows: crystal violet was solubilized by addition of 150 µl of ethanol-acetone

(80:20), and the OD_{570} was determined. Results were presented as the mean of four

replicate wells in three independent experiments.

Deep sequencing analysis: The quality of Illumina reads was visualized by FastQC

v0.10.1   Brabraham   Bioinformatics

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). All reads of 32 samples

were aligned against the two chromosomes of *Vibrio cholera* (Genbank NC_002505 and

NC_002506, 17-DEC-2014), using paired end mapping mode of BWA 'mem'

v0.7.4(104) with the option '-M'. The reference chromosome I was modified since the

reference genome sequence (AE003852) showed an inversion around ori1 flanked by two

ribosomal RNA operons (*rrnB* and *rrnG-H*) (48). Output SAM files were converted to

BAM files using SAMtools v0.1.19 (105). The 'Add Read Groups' step was made by

Picard v1.131 (http://picard.sourceforge.net/). The aligned reads in BAM files were

realigned with the command 'IndelRealigner' implemented in GATK v2.2.16 (106).

'MarkDuplicates' step from Picard flagged duplicates reads. We only kept uniquely

mapped reads, using SAMtools (option 'view –bq 20'). Then, mpileup files were

generated using SAMtools without BAQ adjustments (option 'mpileup –B'). SNPs and

INDELs were called by the option 'pileup2cns' of Varscan2 v2.3.2 (107) with a minimum

depth of 10 reads, a threshold of 20 for minimum quality, a threshold of 0.25 and 0.8 for

minimum variant allele frequency for heterogeneous population strains and subclone

isolated strains respectively. The coverage around the mutation was checked with a

python script with a minimum depth of 7 reads. The annotation of identified variants was

processed by SnpEff v4.1 (108). Common mutations between all sequenced strains were

ignored, indicating differences with the reference sequence. All data generated and

analyzed during this study is included in this published article, its supplementary

information files, and publicly available repositories. The Illumina Deep Sequencing

datasets are deposited at the GenBank SRA as BioProject PRJNA816505.

Multiplex Genome Editing by Natural Transformation (MuGENT): To introduce the

mutations observed along the evolution experiment we employed MuGENT (76, 77).

Briefly, the wild type strain was grown overnight from a single colony at 30°C in LB with

agitation. This ON culture was diluted 10^{-3} into fresh medium, and the strain was grown

to an OD_{600nm} ~0.5. Cells were washed by centrifugation and resuspended at the original

volume in 1× Instant Ocean (IO) Sea Salts (7 g/L) (Blacksburg) and autoclaved chitin

powder (Sigma Aldrich) and incubated overnight without agitation   at   30°C   to

induce competence. Next, DNA fragments containing the desired chromosomal alteration

and a spectinomycin resistance cassette directed to a neutral locus at the intergenic space

between VC1903 and VC1902 region were co-transformed (109). After ON incubation

at 30°C without shaking sample was vortexed was plated in LB supplemented with

spectinomycin. Correct genome editing was then confirmed by MASC-PCR (77) and

Sanger sequencing (Macrogen, Ko) of the targeted loci.

Acknowledgements: We are grateful to Eduardo P. Rocha, José Antonio Escudero,

Alexandra Nivina, Céline Loot, Soledad Guidolín and Inés Marchesini for useful

discussions. We thank the technical assistance of Laurence Ma and Christiane Bouchier

from the Institut Pasteur Genomics Platform for DNA sequencing. We thank Pierre Faure

and Julie Lambert for their technical help. This study was supported by the Institut

Pasteur, the Centre National de la Recherche Scientifique (UMR3525), the French

National Research Agency grants ANR-10-BLAN593 131301 (BMC) and ANR-14-

CE10-0007 (MAGISBAC), the French government's Investissement d'Avenir Program,

Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" (Grant

n°ANR-10-LABX-62-IBEID, to JMG and DM), the ECOS-SUD France-Argentina

Program (A18ST06 to ASB and DM), the Agencia Nacional de Promoción de la

Investigación, el Desarrollo Tecnológico y la Innovación of Argentina (PICT-2017-0424

& PICT-2018-0476 to ASB), and International Center for Genetic Engineering and

Biotechnology (CRP/ARG18-06_EC to ASB). ASB and DJC are Career Members of

CONICET. The funders had no role in study design, data collection and analysis, decision

to publish, or preparation of the manuscript.

Table 1: Recurrent mutations occurring on evolved *V. cholerae* populations after 250
generations. We included genes mutated more than once in multiple populations.
Dataset S1 includes the full list and details of all mutations.
622

Gene	Function	Population	Type of mutation[1]
ompT (VC1854)	Main porin	9,10 ,11 and 12	FS, S'
flrA (VC2137)	Flagellar master regulator	1, 6	FS, NS
flrB (VC2136)	Flagellar regulator (Class II)	3, 4	FS, NS
flrC (VC2135)	Flagellar regulator (Class II)	2, 9, 10	FS, NS
aroG (VCA1036)	Aromatic amioacid synthesis	1, 6, 8, 9, 12	NS
tagE2 (VCA1043)	Endopeptidase	1,4,5,7,9,11,12	NS

623 [1]S: Synonymous mutation; NS: non-synonymous mutation; FS: Frame-shift mutation.
624

Table 2: Mutations occurring in multiple populations of evolved *V. cholerae* populations after 1000 generations. All mutations recovered and the details are described in Dataset S2.

Gene/ORF[1]	Function	Populations[2]	Type of mutation[3]
rfbB (VC0242)	Phospho-mannose mutase	2, 3	NS
rfbV (VC0259)	Lipopolysaccharide biosynthesis	2, 5	NS
manA-1 (VC0269)	Lipopolysaccharide biosynthesis Mannose-6-phosphate isomerase	1, 3, 4, 6, 7, 8 , 9, 11	FS
intergenic (VC0341(orn),tRNAGly)	Unknown	1, 6	-
mutL (VC0345)	Mismatch repair	1, 3, 4, 5, 6, 7, 11	NS, FS
mshH(VC0398)	MSHA biogenesis	1, 9	NS
intergenic tRNA33-VC0449	Unknown	1, 5 ,8	-
intergenic VC0511-12	Unknown	4, 7, 8 ,9 ,12	-
mutS (VC0535)	Missmatch repair	2, 4	FS
VC0590	Permease	7, 11	NS
VC0603	Unknown (PNPLA domain-containing protein)	1 ,5 ,6 ,7 ,11	FS
iscR (VC0747)	Fe-S cluster sensor status and biogenesis	1, 3, 4, 5, 6, 9, 10, 11	NS,S,FS
tcpC (VC0831)	Toxin corregulated pilus	1, 3, 6 ,7 ,9	FS
gcvA (VC0896)	Trascriptional activation of the glycine cleavage system.	4, 9	NS
vpsA (VC0917)	Biofilm formation	2 ,6	NS
luxO (VC1021)	Quorum sensing regulation, Biofilm formation	1, 2, 3, 5 ,6 ,10	NS
VC1724	Hypothetical protein	5, 6	FS
VC1761	Hypotetical protein (VPI-2)	4, 9	NS
ompT (VC1854)	Main porin	11, 12	NS,S,FS
flhA(VC2069)	Flagellar biosynthesis protein	1, 6	FS, NS
VC2105	Putative Thioesterase	1, 4, 5, 7, 9, 12	NS
flrA	Flagellar master regulator	1, 2, 3 ,4, 5, 6, 7, 9, 10, 11, 12	FS, NS

Intergenic (VC2286-VC2887)	Unknown	2, 3, 9	-
Intergenic (VC2387-2388)	Unknown	1, 6	-
rimO (VC2620)	Ribosome biogenesis	1, 6, 7	FS, NS
VC2684	Bifunctional aspartokinase/homoserine dehydrogenase	7, 10	NS
VC2707	Unknown	1, 9, 12	NS
VCA0044 (Pseudogene)	Diguanylate cyclase	8, 12	NS
VCA0088 (Pseudogene)	Proton/glutamate symporter	6, 7, 12	NS
VCA0168 (pseudogene)	BatD family protein	1,2, 4 6, 7, 8 ,9 ,11, 12	-
VCA0354 (Pseudogene)	DedA family protein	6, 9, 12	-
Intergenic (VCA0591-VCA0592)	Unknown	6, 9	-
luxP (VCA0737)	Quorum Sensing	3, 7, 9	NS
VCA1031 (Pseudogene)	Chemotaxis Like protein (10.1016/j.femsle.2004.08.039)	2, 4, 5, 6, 7, 8, 9, 11, 12	-
tagE2 (VCA1043)	Endopeptidase	1, 4, 5, 7, 8, 9, 12	NS,FS
intergenic (VCA1052-3)	Unknown	1, 3, 4, 5, 7	-
aroG (VCA1036)	Aromatic amioacid synthesis	5, 7	NS
VCA1080	HlyD family secretion protein	1, 2, 6, 8	NS

[1]We included intergenic regions specifying the flanking genes [2]S: Synonymous mutation; NS: non-synonymous mutation; FS: Frame-shift mutation.

632 **Table 3:** Mutations linked to rugose phenotype on *V. cholerae* populations evolved for
633 1000 generations. Dataset 3 includes the full list of mutations and their details.

Gene	Function	Population	Type of mutation[1]
vpsR (VC0665)	Positive Biofilm regulator	1S	NS
vpsB (VC0918)	Biofilm	12S	NS
cdgB (VC1029)	GGDEF protein	12R	NS
VCA1031 (Pseudogene)	Chemotaxis Like protein	1S	NS
iscR (VC0747)	Fe-S cluster sensor status and biogenesis	12R	NS
rpoA (VC2571)	α subunit of RNA polymerase	12S	S
Intergenic (VC0341(*orn*), tRNA-Gly	Unknown	1R	-

634 [1] S: Synonymous mutation; NS: non-synonymous mutation; FS: Frame-shift mutation.
635

636 **Table 4:** Mutations observed in fast-growing clones isolated from G250. Details are
637 described in Dataset S4.

Strain	Gene	Reference	Mutation	Result
FG-G250-2	*flrA* (VC2137)	A	G	I303T
FG-G250-2	Intergenic space (tRNA33-VC0449)	G	T	ND
FG-G250-3	*flrB* (VC2136)	TGTGAGCG AGTGGTGA A	T	FS(294)
FG-G250-7	*flrA* (VC2137)	AC	A	FS(461)
FG-G250-10	*flrA* (VC2137)	A	AC	FS(461)
FG-G250-10	*mgtE* (VCA0818)	G	C	G430R
FG-G250-11	*flrA* (VC2137)	A	C	L356R

Legends to Figures:

Figure 1: Genotype and associated phenotype of the evolved strains at the start of the experiment. A) Genome structure of the parental, the movants strains subjected to long term evolution for 1000 generations. Chromosome I and II are depicted as ovals- The orange arrow represents S10 at its genomic position with respect to *ori1* (green dot) or *ori2* (yellow dot). Colors are strain specific and we indicate the population numbering assigned. **B)** The variation of maximum growth rate (49)(black squares, right axis) and the relative fitness (50) (white squares, left axis) with respect to the parental strain is plotted for each strain.

Figure 2: Evolution of biofilm formation capacity along the evolution experiment. Biofilm formation phenotype of twelve evolved populations from the parental strains and 3 movants was evaluated on polystyrene microtiter plates as described in methods section. **A)** Representative wells where cells were stained with crystal violet are shown for P1-P12 at different time points. **B)** The crystal violet stain was quantified for all populations along generations. The parental (Violet, P1-P3) and S10Tnp-35 (Cyan, P4-P6), S10Tnp-1120 (Red, P7-P9) and S10TnpC2+479 (Blue, P10-P12) movants populations values were averaged for simplicity (Figure S2 shows the disaggregated data). Data averaged the means from three independent experiments done in quadruplicates.

Figure 3: S10 genomic location impacts growth rate evolution in the long term. A) ***V. cholerae* increases its growth rate along evolution.** At every time point, the populations from the frozen "fossil record" were subjected to automated growth curve and relativized to their ancestral strains. Parentals (P1-3), S10Tnp-35 (P4-6), S10Tnp-1120 (P7-9) and S10TnpC2+479(P10-12) are shown in tones of violet, cyan, red and blue

respectively. The graph shows the average at least two experiments done by triplicate of

μ ± SD as a function of the generations elapsed. The average μ from the Parental (Violet)

strain and S10Tnp-35 (Cyan), S10Tnp-1120 (Red) and S10TnpC2+479 (Blue) is shown

as dotted lines for reference. **B) Growth rate defect of movants persists along 1000**

generations of evolution. The obtained μ for each movant population (P4-P12) was

relativized to growth rate of the Parental populations (P1-P3) for each time point. Then

the mean with SD of percentage of μ variation was plotted as a function of generations.

The dotted lines represent the average deviation from growth of the parental for each

ancestral strain (0.91% for S10Tnp-35 and -13% for S10Tnp-1120 and S10TnpC2+479).

Figure 4: Evolutionary trajectories of the evolved populations: A) The graph shows

the number mutations fixed to at least of 25% of the population. For each population the

left and right bars indicate the mutations detected at G250 and G1000 respectively. The

colors indicate the ancestral strain for the population (violet and nuances for parental;

Cyan and nuances for S10Tnp-35; red and nuances for S10Tnp-1120; and blue and

nuances for S10TnpC2+479). The filled circles indicate populations showing mutations

at the DNA repair genes: *mutL* (green); *mutS* (yellow); others (*e.g. recBCD, recJ,* orange).

B) Each column represent the total of the observed mutations at G250 and G1000 . The

colors represent the proportion of each type of mutation observed. The numbers within

the color areas indicate the corresponding percentage. **C)** Total mutations fixed in the

single clones sequenced. The white columns represent the fast-growing clones (FG)

isolated at G250. The dark columns represent the S and R clones isolated at G1000.

Figure 5: Specific mutations associated with fast growth rate. A) Screening fast-

growing clones: 5 isolated clones per population (four S and one R) were tested per

triplicate. The graph shows the average µ of each clones compared to Parental- strain

(black). Clones indicated in red were selected for further study. **B)** The clones selected in

A) were subjected to growth curves and the % variation of µ with respect to the parental

strain is shown. The indicated clones were subjected to Deep sequencing. **C)** The

mutations identified in clones from B were introduced in a parental naïve genetic

background.

Figure 6: Fast-growth associated mutations increase growth rate in a *V. cholerae*

naïve context. A) Mutations associated with fast growth (Table 4) were introduced wild

type *V. cholerae* using MuGENT. The growth rate of these mutants was determined in

automated growth curves performed by triplicate (n=11). Left panel shows a

representative growth curve. Right panel represents the median ± CI of the % generation

time (GT) variation of the indicated mutants with respect to the wild type (WT).

Statistical significance was analyzed by one-way ANOVA. Then Dunnet´s test was

done for multiple comparisons. **B)** Mutations in *flrA* and *flrB* increasing growth rate

reduce *V. cholerae* swimming capacity. Left panel show a representative swimming

assay of the WT, the parental strain bearing the selection cassette (Parental) and the

clone bearing the indicated mutation (*flrA** and *flrB**). The right panel shows the

analyzed data for all the assays performed per triplicate (n=9). Statistical significance

was analyzed by one-way ANOVA. Then Tukey´s test was done to compare the mean

values. ** $p<0.01$ and **** $p<0.0001$.

References:

1.   Badrinarayanan A, Le TB, Laub MT. Bacterial chromosome organization and
segregation. Annual review of cell and developmental biology. 2015;31:171-99.
2.   Brickner J. Genetic and epigenetic control of the spatial organization of the
genome. Molecular biology of the cell. 2017 Feb 1;28(3):364-9.

3.   Misteli T. The Self-Organizing Genome: Principles of Genome Architecture and
Function. Cell. 2020 Oct 1;183(1):28-45.
4.   Surovtsev IV, Jacobs-Wagner C. Subcellular Organization: A Critical Feature of
Bacterial Cell Replication. Cell. 2018 Mar 8;172(6):1271-93.
5.   Lioy VS, Junier I, Boccard F. Multiscale Dynamic Structuring of Bacterial
Chromosomes. Annual review of microbiology. 2021 Oct 8;75:541-61.
6.   Touchon M, Rocha EP. Coevolution of the Organization and Structure of
Prokaryotic Genomes. Cold Spring Harbor perspectives in biology. 2016 Jan
4;8(1):a018168.
7.   Rocha EP. The organization of the bacterial genome. Annual review of genetics.
2008;42:211-33.
8.   Le TB, Laub MT. New approaches to understanding the spatial organization of
bacterial genomes. Current opinion in microbiology. 2014 Dec;22:15-21.
9.   Dorman CJ. Genome architecture and global gene regulation in bacteria: making
progress towards a unified model? Nature reviews Microbiology. 2013 May;11(5):349-
55.
10.   Le TB, Imakaev MV, Mirny LA, et al. High-resolution mapping of the spatial
organization of a bacterial chromosome. Science. 2013 Nov 08;342(6159):731-4.
11.   Le TB, Laub MT. Transcription rate and transcript length drive formation of
chromosomal interaction domain boundaries. The EMBO journal. 2016 Jul
15;35(14):1582-95.
12.   Lioy VS, Junier I, Lagage V, et al. Distinct Activities of Bacterial Condensins
for Chromosome Management in Pseudomonas aeruginosa. Cell reports. 2020 Nov
3;33(5):108344.
13.   Lioy VS, Cournac A, Marbouty M, et al. Multiscale Structuring of the E. coli
Chromosome by Nucleoid-Associated and Condensin Proteins. Cell. 2018 Feb
8;172(4):771-83 e18.
14.   Meyer S, Reverchon S, Nasser W, et al. Chromosomal organization of
transcription: in a nutshell. Current genetics. 2018 Jun;64(3):555-65.
15.   Rocha EP. The replication-related organization of bacterial genomes.
Microbiology. 2004 Jun;150(Pt 6):1609-27.
16.   Slager J, Veening JW. Hard-Wired Control of Bacterial Processes by
Chromosomal Gene Location. Trends in microbiology. 2016 Oct;24(10):788-800.
17.   Sobetzko P, Travers A, Muskhelishvili G. Gene order and chromosome
dynamics coordinate spatiotemporal gene expression during the bacterial growth cycle.
Proceedings of the National Academy of Sciences of the United States of America.
2012 Jan 10;109(2):E42-50.
18.   Kosmidis K, Jablonski KP, Muskhelishvili G, et al. Chromosomal origin of
replication coordinates logically distinct types of bacterial genetic regulation. NPJ
systems biology and applications. 2020 Feb 17;6(1):5.
19.   Gerganova V, Maurer S, Stoliar L, et al. Upstream binding of idling RNA
polymerase modulates transcription initiation from a nearby promoter. The Journal of
biological chemistry. 2015 Mar 27;290(13):8095-109.
20.   Yubero P, Poyatos JF. The Impact of Global Transcriptional Regulation on
Bacterial Gene Order. iScience. 2020 Apr 24;23(4):101029.
21.   Muskhelishvili G, Travers A. Order from the order: how a spatiotemporal
genetic program is encoded in a 2-D genetic map of the bacterial chromosome. Journal
of molecular microbiology and biotechnology. 2014;24(5-6):332-43.

22.   Hu XP, Lercher MJ. An optimal growth law for RNA composition and its partial
implementation through ribosomal and tRNA gene locations in bacterial genomes.
PLoS genetics. 2021 Nov;17(11):e1009939.
23.   Sobetzko P, Glinkowska M, Travers A, et al. DNA thermodynamic stability and
supercoil dynamics determine the gene expression program during the bacterial growth
cycle. Molecular bioSystems. 2013 Jul;9(7):1643-51.
24.   Couturier E, Rocha EP. Replication-associated gene dosage effects shape the
genomes of fast-growing bacteria but only for transcription and translation genes.
Molecular microbiology. 2006 Mar;59(5):1506-18.
25.   Japaridze A, Yang W, Dekker C, et al. DNA sequence-directed cooperation
between nucleoid-associated proteins. iScience. 2021 May 21;24(5):102408.
26.   Muskhelishvili G, Sobetzko P, Mehandziska S, et al. Composition of
Transcription Machinery and Its Crosstalk with Nucleoid-Associated Proteins and
Global Transcription Factors. Biomolecules. 2021 Jun 22;11(7).
27.   Block DH, Hussein R, Liang LW, et al. Regulatory consequences of gene
translocation in bacteria. Nucleic acids research. 2012 Oct;40(18):8979-92.
28.   Scholz SA, Diao R, Wolfe MB, et al. High-Resolution Mapping of the
Escherichia coli Chromosome Reveals Positions of High and Low Transcription. Cell
systems. 2019 Mar 27;8(3):212-25 e9.
29.   Bryant JA, Sellars LE, Busby SJ, et al. Chromosome position effects on gene
expression in Escherichia coli K-12. Nucleic acids research. 2014 Oct;42(18):11383-92.
30.   Dryselius R, Izutsu K, Honda T, et al. Differential replication dynamics for large
and small Vibrio chromosomes affect gene dosage, expression and location. BMC
genomics. 2008 Nov 26;9:559.
31.   Sonnenberg CB, Kahlke T, Haugen P. Vibrionaceae core, shell and cloud genes
are non-randomly distributed on Chr 1: An hypothesis that links the genomic location of
genes with their intracellular placement. BMC genomics. 2020 Oct 6;21(1):695.
32.   van Eijk E, Boekhoud IM, Kuijper EJ, et al. Genome Location Dictates the
Transcriptional Response to PolC Inhibition in Clostridium difficile. Antimicrobial
agents and chemotherapy. 2019 Feb;63(2).
33.   Gerganova V, Berger M, Zaldastanishvili E, et al. Chromosomal position shift of
a regulatory gene alters the bacterial phenotype. Nucleic acids research. 2015 Sep
30;43(17):8215-26.
34.   Fitzgerald S, Dillon SC, Chao TC, et al. Re-engineering cellular physiology by
rewiring high-level global regulatory genes. Scientific reports. 2015 Dec 03;5:17653.
35.   Moffitt JR, Pandey S, Boettiger AN, et al. Spatial organization shapes the
turnover of a bacterial transcriptome. eLife. 2016 May 20;5.
36.   Charles RC, Ryan ET. Cholera in the 21st century. Current opinion in infectious
diseases. 2011 Oct;24(5):472-7.
37.   Reen FJ, Almagro-Moreno S, Ussery D, et al. The genomic code: inferring
Vibrionaceae niche specialization. Nature reviews Microbiology. 2006 Sep;4(9):697-
704.
38.   Teschler JK, Zamorano-Sanchez D, Utada AS, et al. Living in the matrix:
assembly and control of Vibrio cholerae biofilms. Nature reviews Microbiology. 2015
May;13(5):255-68.
39.   Echazarreta MA, Klose KE. Vibrio Flagellar Synthesis. Frontiers in cellular and
infection microbiology. 2019;9:131.
40.   Silva AJ, Benitez JA. Vibrio cholerae Biofilms and Cholera Pathogenesis. PLoS
neglected tropical diseases. 2016 Feb;10(2):e0004330.

41.   Alam M, Sultana M, Nair GB, et al. Viable but nonculturable Vibrio cholerae
O1 in biofilms in the aquatic environment and their role in cholera transmission.
Proceedings of the National Academy of Sciences of the United States of America.
2007 Nov 6;104(45):17801-6.
42.   Faruque SM, Biswas K, Udden SM, et al. Transmissibility of cholera: in vivo-
formed biofilms and their relationship to infectivity and persistence in the environment.
Proceedings of the National Academy of Sciences of the United States of America.
2006 Apr 18;103(16):6350-5.
43.   Conner JG, Zamorano-Sanchez D, Park JH, et al. The ins and outs of cyclic di-
GMP signaling in Vibrio cholerae. Current opinion in microbiology. 2017 Apr;36:20-9.
44.   Wu DC, Zamorano-Sanchez D, Pagliai FA, et al. Reciprocal c-di-GMP
signaling: Incomplete flagellum biogenesis triggers c-di-GMP signaling pathways that
promote biofilm formation. PLoS genetics. 2020 Mar;16(3):e1008703.
45.   Floyd KA, Lee CK, Xian W, et al. c-di-GMP modulates type IV MSHA pilus
retraction and surface attachment in Vibrio cholerae. Nature communications. 2020 Mar
25;11(1):1549.
46.   Lim B, Beyhan S, Yildiz FH. Regulation of Vibrio polysaccharide synthesis and
virulence factor production by CdgC, a GGDEF-EAL domain protein, in Vibrio
cholerae. Journal of bacteriology. 2007 Feb;189(3):717-29.
47.   Val ME, Soler-Bistue A, Bland MJ, et al. Management of multipartite genomes:
the Vibrio cholerae model. Current opinion in microbiology. 2014 Dec;22:120-6.
48.   Val ME, Marbouty M, de Lemos Martins F, et al. A checkpoint control
orchestrates the replication of the two chromosomes of Vibrio cholerae. Science
advances. 2016 Apr;2(4):e1501914.
49.   Soler-Bistue A, Mondotte JA, Bland MJ, et al. Genomic location of the major
ribosomal protein gene locus determines Vibrio cholerae global growth and infectivity.
PLoS genetics. 2015 Apr;11(4):e1005156.
50.   Soler-Bistue A, Timmermans M, Mazel D. The Proximity of Ribosomal Protein
Genes to oriC Enhances Vibrio cholerae Fitness in the Absence of Multifork
Replication. mBio. 2017 Feb 28;8(1).
51.   Vieira-Silva S, Rocha EP. The systemic imprint of growth and its uses in
ecological (meta)genomics. PLoS genetics. 2010 Jan 15;6(1):e1000808.
52.   Coenye T, Vandamme P. Organisation of the S10, spc and alpha ribosomal
protein gene clusters in prokaryotic genomes. FEMS microbiology letters. 2005 Jan
1;242(1):117-26.
53.   Stoebe B, Kowallik KV. Gene-cluster analysis in chloroplast genomics. Trends
in genetics : TIG. 1999 Sep;15(9):344-7.
54.   Soler-Bistue A, Aguilar-Pierle S, Garcia-Garcera M, et al. Macromolecular
crowding links ribosomal protein gene dosage to growth rate in Vibrio cholerae. BMC
biology. 2020 Apr 29;18(1):43.
55.   Ochman H, Wilson AC. Evolution in bacteria: evidence for a universal
substitution rate in cellular genomes. Journal of molecular evolution. 1987;26(1-2):74-
86.
56.   Barrick JE, Lenski RE. Genome dynamics during experimental evolution.
Nature reviews Genetics. 2013 Dec;14(12):827-39.
57.   Kawecki TJ, Lenski RE, Ebert D, et al. Experimental evolution. Trends in
ecology & evolution. 2012 Oct;27(10):547-60.
58.   Lenski RE. Experimental evolution and the dynamics of adaptation and genome
evolution in microbial populations. The ISME journal. 2017 Oct;11(10):2181-94.

59.   Remigi P, Masson-Boivin C, Rocha EPC. Experimental Evolution as a Tool to
Investigate Natural Processes and Molecular Functions. Trends in microbiology. 2019
Jul;27(7):623-34.
60.   Lenski RE, Travisano M. Dynamics of adaptation and diversification: a 10,000-
generation experiment with bacterial populations. Proceedings of the National Academy
of Sciences of the United States of America. 1994 Jul 19;91(15):6808-14.
61.   Wibbenmeyer JA, Provenzano D, Landry CF, et al. Vibrio cholerae OmpU and
OmpT porins are differentially affected by bile. Infection and immunity. 2002
Jan;70(1):121-6.
62.   Murphy SG, Murtha AN, Zhao Z, et al. Class A Penicillin-Binding Protein-
Mediated Cell Wall Synthesis Promotes Structural Integrity during Peptidoglycan
Endopeptidase Insufficiency in Vibrio cholerae. mBio. 2021 Apr 6;12(2).
63.   Klose KE, Mekalanos JJ. Distinct roles of an alternative sigma factor during
both free-swimming and colonizing phases of the Vibrio cholerae pathogenic cycle.
Molecular microbiology. 1998 May;28(3):501-20.
64.   Buck M, Gallegos MT, Studholme DJ, et al. The bacterial enhancer-dependent
sigma(54) (sigma(N)) transcription factor. Journal of bacteriology. 2000
Aug;182(15):4129-36.
65.   Fong JCN, Syed KA, Klose KE, et al. Role of Vibrio polysaccharide (vps) genes
in VPS production, biofilm formation and Vibrio cholerae pathogenesis. Microbiology
(Reading). 2010 Sep;156(Pt 9):2757-69.
66.   Fong JC, Yildiz FH. The rbmBCDEF gene cluster modulates development of
rugose colony morphology and biofilm formation in Vibrio cholerae. Journal of
bacteriology. 2007 Mar;189(6):2319-30.
67.   Yildiz FH, Liu XS, Heydorn A, et al. Molecular analysis of rugosity in a Vibrio
cholerae O1 El Tor phase variant. Molecular microbiology. 2004 Jul;53(2):497-515.
68.   Yildiz FH, Schoolnik GK. Vibrio cholerae O1 El Tor: identification of a gene
cluster required for the rugose colony type, exopolysaccharide production, chlorine
resistance, and biofilm formation. Proceedings of the National Academy of Sciences of
the United States of America. 1999 Mar 30;96(7):4028-33.
69.   Ball AS, Chaparian RR, van Kessel JC. Quorum Sensing Gene Regulation by
LuxR/HapR Master Regulators in Vibrios. Journal of bacteriology. 2017 Oct 1;199(19).
70.   Ortega DR, Kjaer A, Briegel A. The chemosensory systems of Vibrio cholerae.
Molecular microbiology. 2020 Sep;114(3):367-76.
71.   Pratt JT, McDonough E, Camilli A. PhoB regulates motility, biofilms, and cyclic
di-GMP in Vibrio cholerae. Journal of bacteriology. 2009 Nov;191(21):6632-42.
72.   Beyhan S, Bilecen K, Salama SR, et al. Regulation of rugosity and biofilm
formation in Vibrio cholerae: comparison of VpsT and VpsR regulons and epistasis
analysis of vpsT, vpsR, and hapR. Journal of bacteriology. 2007 Jan;189(2):388-402.
73.   Lim B, Beyhan S, Meir J, et al. Cyclic-diGMP signal transduction systems in
Vibrio cholerae: modulation of rugosity and biofilm formation. Molecular
microbiology. 2006 Apr;60(2):331-48.
74.   Beyhan S, Tischler AD, Camilli A, et al. Transcriptome and phenotypic
responses of Vibrio cholerae to increased cyclic di-GMP level. Journal of bacteriology.
2006 May;188(10):3600-13.
75.   Miller HK, Kwuan L, Schwiesow L, et al. IscR is essential for yersinia
pseudotuberculosis type III secretion and virulence. PLoS pathogens. 2014
Jun;10(6):e1004194.

76.   Dalia AB, McDonough E, Camilli A. Multiplex genome editing by natural
transformation. Proceedings of the National Academy of Sciences of the United States
of America. 2014 Jun 17;111(24):8937-42.
77.   Dalia AB. Natural Cotransformation and Multiplex Genome Editing by Natural
Transformation (MuGENT) of Vibrio cholerae. Methods in molecular biology.
2018;1839:53-64.
78.   Srivastava D, Hsieh ML, Khataokar A, et al. Cyclic di-GMP inhibits Vibrio
cholerae motility by repressing induction of transcription and inducing extracellular
polysaccharide production. Molecular microbiology. 2013 Dec;90(6):1262-76.
79.   Dymond J, Boeke J. The Saccharomyces cerevisiae SCRaMbLE system and
genome minimization. Bioengineered bugs. 2012 May-Jun;3(3):168-71.
80.   Hu X-P, Lercher MJ. An optimal growth law for RNA composition and its
partial implementation through ribosomal and tRNA gene locations in bacterial
genomes. bioRxiv. 2021:2021.02.05.429890.
81.   Esnault E, Valens M, Espeli O, et al. Chromosome structuring limits genome
plasticity in Escherichia coli. PLoS genetics. 2007 Dec;3(12):e226.
82.   Savic DJ, Nguyen SV, McCullor K, et al. Biological Impact of a Large-Scale
Genomic Inversion That Grossly Disrupts the Relative Positions of the Origin and
Terminus Loci of the Streptococcus pyogenes Chromosome. Journal of bacteriology.
2019 Sep 1;201(17).
83.   Campo N, Dias MJ, Daveran-Mingot ML, et al. Chromosomal constraints in
Gram-positive bacteria revealed by artificial inversions. Molecular microbiology. 2004
Jan;51(2):511-22.
84.   Slager J, Kjos M, Attaiech L, et al. Antibiotic-induced replication stress triggers
bacterial competence by increasing gene dosage near the origin. Cell. 2014 Apr
10;157(2):395-406.
85.   Bogue MM, Mogre A, Beckett MC, et al. Network Rewiring: Physiological
Consequences of Reciprocally Exchanging the Physical Locations and Growth-Phase-
Dependent Expression Patterns of the Salmonella fis and dps Genes. mBio. 2020 Sep
8;11(5).
86.   Narula J, Kuchina A, Lee DD, et al. Chromosomal Arrangement of
Phosphorelay Genes Couples Sporulation and DNA Replication. Cell. 2015 Jul
16;162(2):328-37.
87.   Renda BA, Dasgupta A, Leon D, et al. Genome instability mediates the loss of
key traits by Acinetobacter baylyi ADP1 during laboratory evolution. Journal of
bacteriology. 2015 Mar;197(5):872-81.
88.   Poltak SR, Cooper VS. Ecological succession in long-term experimentally
evolved biofilms produces synergistic communities. The ISME journal. 2011
Mar;5(3):369-78.
89.   Kovacs AT, Dragos A. Evolved Biofilm: Review on the Experimental Evolution
Studies of Bacillus subtilis Pellicles. Journal of molecular biology. 2019 Nov
22;431(23):4749-59.
90.   Lenski RE, Rose MR, Simpson SC, et al. Long-Term Experimental Evolution in
Escherichia coli. I. Adaptation and Divergence During 2,000 Generations. The
American Naturalist. 1991;138(6):1315-41.
91.   Grant NA, Abdel Magid A, Franklin J, et al. Changes in Cell Size and Shape
during 50,000 Generations of Experimental Evolution with Escherichia coli. Journal of
bacteriology. 2021 Apr 21;203(10).

92. Atolia E, Cesar S, Arjes HA, et al. Environmental and Physiological Factors
Affecting High-Throughput Measurements of Bacterial Growth. mBio. 2020 Oct
20;11(5).
93. Jun S, Si F, Pugatch R, et al. Fundamental principles in bacterial physiology-
history, recent progress, and the future with focus on cell size control: a review. Reports
on progress in physics Physical Society. 2018 May;81(5):056601.
94. Wiser MJ, Lenski RE. A Comparison of Methods to Measure Fitness in
Escherichia coli. PloS one. 2015;10(5):e0126210.
95. Concepcion-Acevedo J, Weiss HN, Chaudhry WN, et al. Malthusian Parameters
as Estimators of the Fitness of Microbes: A Cautionary Tale about the Low Side of
High Throughput. PloS one. 2015 Jun 26;10(6).
96. Stutzmann S, Blokesch M. Comparison of chitin-induced natural transformation
in pandemic Vibrio cholerae O1 El Tor strains. Environmental microbiology. 2020
Oct;22(10):4149-66.
97. Colin R, Ni B, Laganenka L, et al. Multiple functions of flagellar motility and
chemotaxis in bacterial physiology. FEMS microbiology reviews. 2021 Nov 23;45(6).
98. Ni B, Ghosh B, Paldy FS, et al. Evolutionary Remodeling of Bacterial Motility
Checkpoint Control. Cell reports. 2017 Jan 24;18(4):866-77.
99. Ni B, Colin R, Link H, et al. Growth-rate dependent resource investment in
bacterial motile behavior quantitatively follows potential benefit of chemotaxis.
Proceedings of the National Academy of Sciences of the United States of America.
2020 Jan 7;117(1):595-601.
100. Izard J, Gomez Balderas CD, Ropers D, et al. A synthetic growth switch based
on controlled expression of RNA polymerase. Molecular systems biology. 2015 Nov
23;11(11):840.
101. Gyorfy Z, Draskovits G, Vernyik V, et al. Engineered ribosomal RNA operon
copy-number variants of E. coli reveal the evolutionary trade-offs shaping rRNA operon
number. Nucleic acids research. 2015 Feb 18;43(3):1783-94.
102. Heidelberg JF, Eisen JA, Nelson WC, et al. DNA sequence of both
chromosomes of the cholera pathogen Vibrio cholerae. Nature. 2000 Aug
3;406(6795):477-83.
103. Martinez-Gil M, Goh KGK, Rackaityte E, et al. YeeJ is an inverse
autotransporter from Escherichia coli that binds to peptidoglycan and promotes biofilm
formation. Scientific reports. 2017 2017/09/12;7(1):11326.
104. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics. 2009 Jul 15;25(14):1754-60.
105. Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and
SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078-9.
106. DePristo MA, Banks E, Poplin R, et al. A framework for variation discovery and
genotyping using next-generation DNA sequencing data. Nature genetics. 2011
May;43(5):491-8.
107. Koboldt DC, Zhang Q, Larson DE, et al. VarScan 2: somatic mutation and copy
number alteration discovery in cancer by exome sequencing. Genome research. 2012
Mar;22(3):568-76.
108. Cingolani P, Platts A, Wang le L, et al. A program for annotating and predicting
the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of
Drosophila melanogaster strain w1118; iso-2; iso-3. Fly. 2012 Apr-Jun;6(2):80-92.
109. de Lemos Martins F, Fournes F, Mazzuoli MV, et al. Vibrio cholerae
chromosome 2 copy number is controlled by the methylation-independent binding of its

monomeric initiator to the chromosome 1 crtS site. Nucleic acids research. 2018 Nov
2;46(19):10145-56.

1009
1010

A

S10 movants

ori1
S10

ori2

Parental
(Pop. 1-3)

ori1

ori2

S10Tnp-35
(Pop. 4-6)

ori1

ori2

S10Tnp-1120
(Pop. 7-9)

ori1

ori2

S10TnpC2+479
(Pop. 10-12)

B

□ Fitness
■ Growth

W (rel)

1.1

1.0

0.9

0.8

0.7

Par   -35   -1120   C2+479

A

	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12
G0												
G100												
G250												
G500												
G700												
G1000												

Par

R1

B

Abs 570 nm

Generation

- Parental
- -35
- -1120
- C2+479