Enterobacterales plasmid sharing amongst human bloodstream infections, livestock, wastewater, and waterway niches in Oxfordshire, UK

William Matlock*[,1], Samuel Lipworth*[,1,2], Kevin K. Chau[1], Manal Abu Oun[3], Leanne Barker[1], James Kavanagh[1], Monique Andersson[2], Sarah Oakley[2], Marcus Morgan[2], Derrick W. Crook[1,2,4,5], Daniel S. Read[6], Muna Anjum[3], Liam P. Shaw[#,7]/Nicole Stoesser[#,1,2,4,5], REHAB Consortium

* Contributed equally as first authors

# Contributed equally as senior authors

[1] Nuffield Department of Medicine, University of Oxford, Oxford, UK

[2] Oxford University Hospitals NHS Foundation Trust, Oxford, UK

[3] Animal and Plant Health Agency, Addlestone, UK

[4] NIHR Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance at University of Oxford in partnership with Public Health England, Oxford, UK

[5] NIHR Biomedical Research Centre, Oxford, UK

[6] Centre for Ecology and Hydrology, Wallingford, UK

[7] Department of Biology, University of Oxford, Oxford, UK

Correspondence: William Matlock (william.matlock@ndm.ox.ac.uk) and Nicole Stoesser (nicole.stoesser@ndm.ox.ac.uk)

Keywords: plasmid, *Enterobacterales*, antimicrobial resistance (AMR), bloodstream

infections (BSI), One Health, genomic epidemiology, *Escherichia coli*

Abstract

Plasmids enable the dissemination of antimicrobial resistance (AMR) in common

Enterobacterales pathogens, representing a major public health challenge. However, the

extent of plasmid sharing and evolution between *Enterobacterales* causing human infections

and other niches remains unclear, including the emergence of resistance plasmids. Dense,

unselected sampling is highly relevant to developing our understanding of plasmid

epidemiology and designing appropriate interventions to limit the emergence and

dissemination of plasmid-associated AMR. We established a geographically and temporally

restricted collection of human bloodstream infection (BSI)-associated, livestock-associated

(cattle, pig, poultry, and sheep faeces, farm soils) and wastewater treatment work (WwTW)-

associated (influent, effluent, waterways upstream/downstream of effluent outlets)

Enterobacterales. Isolates were collected between 2008-2020 from sites <60km apart in

Oxfordshire, UK. Pangenome analysis of plasmid clusters revealed shared "backbones", with

phylogenies suggesting an intertwined ecology where well-conserved plasmid backbones

carry diverse accessory functions, including AMR genes. Many plasmid "backbones" were

seen across species and niches, raising the possibility that plasmid movement between these

followed by rapid accessory gene change could be relatively common. Overall, the signature

of identical plasmid sharing is likely to be a highly transient one, implying that plasmid

movement might be occurring at greater rates than previously estimated, raising a challenge

for future genomic One Health studies.

Funding

This study was funded by the Antimicrobial Resistance Cross-council Initiative supported by

the seven research councils and the NIHR, UK.

Introduction

Enterobacterales are found both in human niches (e.g., hospital patients(1,2) and

wastewater(3)) and non-human niches (e.g., livestock-associated(4,5) and waterways(6)). In

recent decades, widespread carriage of antimicrobial resistance (AMR) genes has

complicated the treatment of *Enterobacterales* infections(7,8). The dissemination of AMR

genes between *Enterobacterales* occurs in a 'Russian-doll'-style hierarchy of nested,

mobilisable genetic structures(9): genes not only move between bacterial hosts on

mobilisable or conjugative plasmids but can also be transferred within and between plasmids

and chromosomes by smaller mobile genetic elements (MGEs) such as insertion

sequences(10,11). Despite gene gain/loss events, many plasmids have been shown to have a

persistent structure encoding replication and transfer machinery(12,13).

Many plasmids can transfer between species and are seen across different niches(14) but the

extent to which they are shared between human and non-human niches remains poorly

understood. Previous studies investigating this topic have often been limited in size given the

genetic diversity in these niches(15,16), and/or restricted to single species(17) or drug-

resistant isolates(18), or are systematic studies, pooling geographically/temporally disparate

samples (19,20). Further, fragmented genome assemblies in many cases make recovering

complete plasmids, and other MGEs, impossible(21).

Instances of cross-niche transfer of plasmids are well-described, but the frequency of such

events is poorly characterised. There are multiple instances where AMR genes have emerged

from non-human niches and subsequently become major clinical problems in human

Enterobacterales infections, highlighting the relevance of inter-niche transfer in AMR gene

dissemination (e.g., bla_{CTX-M}, *mcr-1*(22) and bla_{NDM-1}(23)). In general, environmental bacteria

are believed to be the original source of AMR genes that eventually become prevalent in

clinical settings after transfer into clinical pathogens. However, we know little about natural

rates of inter-niche transfer beyond these high-profile examples. It remains unclear how

plasmids evolve within natural populations, meaning we understand little about the wider

context in which AMR genes emerge and disseminate.

To explore *Enterobacterales* plasmid diversity and sharing across niches in a geographically

and temporally restricted context, we studied hybrid assemblies (i.e., using both long and

short reads) of large *Enterobacterales* isolate collections in Oxfordshire, UK, from (i) human

bloodstream infections (BSI; 2008-2018), (ii) livestock-associated sources (faeces from

cattle, pigs, poultry, sheep; surrounding environmental soils; [all 2017 except poultry 2019-

2020], and (iii) wastewater treatment work (WwTW)-associated sources (influent, effluent,

waterways upstream/downstream of effluent outlets; Oxfordshire, 2017).

Results

Our dataset of $n=3,697$ plasmids from $n=1,458$ isolates (Fig. 1a, Table 1) contained bacteria

from human bloodstream infections (BSI; $n=1,880$ plasmids from $n=738$ isolates), livestock-

associated sources (cattle, pig, poultry, and sheep faeces, soils surrounding livestock farms;

$n=1,155$ plasmids from $n=512$ isolates), and from wastewater treatment works (WwTW)-

associated sources (influent, effluent, waterways upstream/downstream of effluent outlets;

$n=662$ plasmids from $n=208$ isolates). All sampling sites were <60km apart (Fig. 1b) and

timeframes overlapped (2008-2020; Fig. 1c). Isolates had a median 2 plasmids (IQR=1-4,

range=0-16). Major *Enterobacterales* genera represented included: $n=1,044$ *Escherichia*,

$n=211$ *Klebsiella*, $n=125$ *Citrobacter*, and $n=63$ *Enterobacter*.

Table 1. Isolate niche breakdown.

Niche	Sample type(s)	No. isolates	No. plasmids
Bloodstream infections (BSI)	Community, nosocomial, other healthcare associated infections	738	1,880
Livestock-associated	Cattle faeces	133	215
	Sheep faeces	113	286
	Pig faeces	104	352
	Poultry faeces	34	112
	Soil surrounding livestock farms	128	190
Wastewater treatment work (WwTW)-associated	Influent	88	313
	Upstream waterways	25	60
	Effluent/downstream waterways	95	289
Total		**1,458**	**3,697**

Table 2. Isolate genus breakdown.

Niche	Isolate genus					Total
	Citrobacter	*Enterobacter*	*Escherichia*	*Klebsiella*	Other	
Bloodstream infections (BSI)	6	11	547	161	13	**738**
Livestock-associated	54	10	433	14	1	**512**

Wastewater treatment work (WwTW)-associated	65	42	64	37	0	**208**
Total	**125**	**63**	**1,044**	**212**	**14**	**1,458**

Sampling niche was strongly associated with isolate genus (Fisher's test, *p*-value<0.001;

Table 2). *Klebsiella* isolates were disproportionately derived from BSI versus other niches

(76% [161/212] versus 51% [738/1,458]). *Citrobacter* and *Enterobacter* were

disproportionately derived from WwTW-associated versus other niches (52% [65/125] and

67% [42/63] versus 14% [208/1,458]). Chromosomal Mash trees (see Supplementary

Methods) for the two most common species in the dataset, *E. coli* (72% [1,044/1,458]; see

Fig. S1) and *K. pneumoniae* (11% [163/1,458]; Fig. S2) demonstrated intermixing of human

and non-human isolates within clades, consistent with species-lineages not being structured

by niche.

We contextualised our plasmids within known plasmid diversity using 'plasmid taxonomic

units' (PTUs; using COPLA, see Supplementary Methods), designed to be equivalent to a

plasmid 'species'. We found 32% (1,193/3,697) of plasmids were unclassified, highlighting

the substantial plasmid diversity within this geographically restricted dataset. In total, we

found *n*=67 known PTUs, containing a median 9 plasmids (IQR=4-30, range=1-556), with

the largest PTU-F_E (556/2,504), corresponding to F-type *Escherichia* plasmids.

Near-identical plasmid sharing observed between human and livestock-associated

Enterobacterales

We screened for near-identical plasmids shared across isolates by grouping those with a low

Mash distance (*d*<0.0001) and highly similar lengths (longest plasmid ≤1% longer than

shorter plasmids; see Supplementary Methods). We found n=225 near-identical groups of ≥2

members, recruiting 19% (712/3,697) plasmids. Bootstrapping accumulation curves for near-

identical plasmid groups and singletons per the number of isolates (ACs; see Supplementary

Methods), we revealed a highly 'open' accumulation (Heap's parameter γ=0.97, Fig. S3)

suggesting further isolate sampling would detect more unique plasmids approximately

linearly. Restricted to BSI/livestock-associated isolates alone, we found similar curves for

both niches (BSI γ=0.98, livestock-associated γ=0.94), suggesting they had similar levels of

plasmid diversity.

Near-identical pairs of plasmids were most common, representing 71% (159/225) of groups

(group size IQR=2-3, range=2-32). Plasmid members of near-identical groups represented

multiple bacterial host STs (25% [56/225]), species (4% [9/225]), and genera (4% [9/225]),

consistent with plasmids capable of inter-lineage/species/genus transfer. Further, 8% (17/225)

of near-identical groups contained plasmids found across human BSIs and ≥1 other sampling

niche (livestock-associated/WwTW-associated), suggesting inter-niche transfer (i.e., 'cross-

niche groups'; Fig. 2a). Within cross-niche groups, n=3/17 contained plasmids from multiple

bacterial species (Fig. 2b), and most consisted of conjugative plasmids (n=5/17 conjugative,

n=9/17 mobilisable, n=3/17 non-mobilisable; Fig. 2c). AMR genes were carried by plasmids

in n=6/17 cross-niche groups (Fig. 2d), with n=5/6 of these groups containing ≥1 beta-

lactamase protein encoding gene.

Sharing between BSI and livestock-associated isolates was supported by 8/17 cross-niche

groups (n=45 plasmids). Of these, n=2/8 contained non-mobilisable Col-type plasmids (one

group contained BSI/pig/poultry/influent plasmids, and one group contained BSI/poultry

plasmids); n=4/8 contained mobilisable Col-type plasmids (two groups contained BSI/pig

plasmids, one group contained BSI/sheep plasmids, and one group contained

BSI/cattle/pig/poultry/sheep/influent plasmids), of which one group contained BSI/pig

plasmids carrying the AMR genes *aph(3ʺ)-Ib*, *aph(6)-Id*, *dfrA14*, and *sul2* (see

Supplementary Methods). The remaining 2/8 groups contained conjugative FIB-type

BSI/sheep plasmids. One group contained plasmids, carrying the AMR genes *aph(3ʺ)-Ib*,

aph(6)-Id, *bla$_{TEM-1}$*, *dfrA5*, *sul2*, and the other group contained plasmids carrying the MDR

efflux pump protein *robA*.

The beta-lactamase *bla*$_{TEM-1}$ was the most common AMR gene detected (8% of total AMR

gene annotations [424/5402]; see Supplementary Methods). In terms of sequence length (bp),

plasmids made up 3.1% of the overall dataset but 13.8% of the *bla*$_{TEM-1}$ -carrying proportion.

Of the plasmid clusters, 16% (39/247) carried *bla*$_{TEM-1}$, and of these 9 clusters were seen in

human BSI and at least one other niche. Plasmid clusters either variably or always carrying

bla$_{TEM-1}$ were strongly associated with BSI (p<0.01, Chi-squared test X^2=8.19, 33/161 of BSI

clusters containing *bla*$_{TEM-1}$ vs. 5/86 for non-BSI clusters) and carried a higher number of

other AMR genes (p<0.01, Wilcoxon text of *bla*$_{TEM-1}$-plasmid clusters vs. others; see Fig.

S4).

Plasmid clustering reveals a diverse but intertwined population structure across niches

Near-identical plasmids shared across niches are a likely signature of recent transfer events,

but we also wanted to examine the wider plasmid population structure. We therefore

agnostically clustered all plasmids based on alignment-free sequence similarity (clusters were

groups of *n*≥3 plasmids; see Supplementary Methods and Figs. S5-6). We defined *n*=247

plasmid clusters with median 5 members (IQR=3-10, range=3-123) recruiting 71%

(2,627/3,697) of the plasmids. The remainder were either singletons (i.e., single, unconnected

plasmids; 19% [718/3,697]) or doubletons (i.e., pairs of connected plasmids; 10%

[352/3,697]). By bootstrapping b=1,000 ACs for plasmid clusters, doubletons, and singletons

found against number of isolates sampled (Fig. S7; see Supplementary Methods), we

estimated that the rarefaction curve had a Heap's parameter γ=0.75, suggesting further isolate

sampling would likely detect more plasmid diversity and clusters.

Of the plasmid clusters, n=69/247 (28%) plasmid clusters had ≥10 members, representing

50% (1,832/3,697) of all plasmids (Fig. 3a). 122/247 (49%) clusters contained BSI plasmids

and plasmids from ≥1 other niche. This included 73/247 (30%) of clusters with both BSI and

livestock-associated plasmids, representing n=38 unique plasmid replicon haplotypes (i.e.,

combinations of replication proteins) of which only 24% (9/38) were Col-type plasmids,

which are often well-conserved and carry few genes(24). 72/247 (29%) of clusters contained

both BSI, and influent/effluent/downstream plasmids, reflecting a route of *Enterobacterales*

dissemination into waterways. In contrast, only 18/247 (7%) of clusters contained both BSI

and upstream waterway plasmids, of which most (13/18 [72%]) also contained

influent/effluent/downstream plasmids.

Overall, plasmid clusters scored high homogeneity (h) but low completeness (c) with respect

to biological and ecological characteristics (non-putative PTUs [h=0.99, c=0.66]; replicon

haplotype [h=0.92, c=0.69]; bacterial host ST [h=0.84, c=0.14] in Fig. 3b; predicted mobility

[h=0.93, c=0.20] in Fig. 3c). This indicated that clustered plasmids often had similar

characteristics, but the same characteristics were often observed in multiple clusters. The

imperfect homogeneity is to be anticipated as replicon haplotypes and mobilities can vary

within plasmid families, and plasmid families can have diverse host ranges(14).

Plasmids carrying AMR genes were found in 21% (52/247) of the plasmid clusters (i.e.,

'antimicrobial resistance gene (ARG)-carrying clusters'), representing n=550 plasmids (Fig.

3d). Of the ARG-carrying clusters, 92% (48/52) contained at least one beta-lactamase-

carrying plasmid (n=437 plasmids in total). AMR genes were present in a median proportion

67% of ARG-carrying cluster members (IQR=28-100%, range=3-100%). This highlights that

AMR genes are not necessarily widespread on genetically similar plasmids and can be

potentially acquired multiple different times through the activity of smaller MGEs (e.g.,

transposons) or recombination. For example, cluster 12 was a group of n=42 conjugative,

PTU-F$_E$ plasmids found in BSI, wastewater, and waterways. Of these, 31% (13/42) carried

the AMR gene bla_{TEM-1}, and in a range of genetic contexts: n=9/13 bla_{TEM-1} genes were found

within Tn3 and n=4/13 were carried without a transposase, of which n=2/4 were found with

the additional AMR genes $aph(6)$-Id, $aph(3'')$-Ib, and $sul2$. AMR genes were

disproportionately carried by F-type plasmids (61% [337/550] ARG-carrying cluster

plasmids versus 34% [891/2627] of the total clustered plasmids), underlining the known role

of F-type plasmids in AMR gene dissemination(13).

An intertwined ecology of plasmids across human and livestock-associated niches

Plasmids can change their genetic content, particularly when subject to new selective

pressures(25,26). Many plasmids have a structure with a 'backbone' of conserved core genes

and a 'cargo' of variable accessory genes(12,13,27). We wanted to explore evidence for

cross-niche plasmids with minimal mutational evolution in a shared backbone (compatible

with ~years of evolutionary separation) but variable accessory gene repertoires.

We first conducted a pangenome-style analysis (see Supplementary Methods) on the

n=69/247 plasmid clusters with ≥10 members. For each cluster, we determined "core" (genes

found in ≥95% of plasmids) and "accessory" gene repertoires (found in <95% of plasmids).

Within clusters, we found median 9 core genes (IQR=4-53, range=0-219), and median 9

accessory genes (IQR=3-145, range=0-801) (Fig. 3e). Core genes comprised a median

proportion 42.2% of the total pangenome sizes (IQR=20.9-66.7%). At an individual plasmid

level, core genes shared by a cluster comprised a median proportion 62.5% of each plasmid's

gene repertoire (IQR=37.4-83.3%; Fig. 3e). Putatively conjugative plasmids carried a

significantly higher proportion of accessory genes in their repertoires than mobilisable/non-

mobilisable plasmids (Kruskal-Wallis test [$H(2)=193.01$, p-value<0.001] followed by

Dunn's test).

Using multiple sequence alignments of the core genes within each cluster, we produced

maximum likelihood phylogenies (see Supp. File 1 and Supplementary Methods). For this

step, we only considered the n=62/69 clusters where each plasmid had ≥1 core gene. With the

n=27/62 clusters that contained both BSI and livestock-associated plasmids, we measured the

phylogenetic signal for plasmid sampling niche using Fritz and Purvis' D (see Table S1 and

Supplementary Methods). The analysis indicated that the evolutionary history of plasmid

clusters is neither strictly segregated by sampling niche nor completely intermixed, but

something intermediate.

Alongside the core gene phylogenies, we generated gene repertoire heatmaps (example

cluster 2 in Fig. 4a-b; all clusters and heatmaps in Supp. File 1). By visualising the genes in a

consensus synteny order (see Supplementary Methods), the putative backbone within each

plasmid cluster is shown alongside its accessory gene and transposase repertoire. This

highlights how plasmids might gain/lose accessory functions within a persistent backbone.

Log-transformed linear regression revealed a significant relationship between Jaccard

distance of accessory genes presence against core gene cophenetic distance

(y=0.080log(x)+0.978, R^2=0.47, F(1,52988)=4.75e4, p-value <0.001; see Fig. S8 and

Supplementary Methods).

Plasmid evolution between human and livestock-associated niches is not structured by

bacterial host

Alongside vertical inheritance, conjugative and mobilisable plasmids are capable of inter-host

transfer, crossing between bacterial lineages, species, up to phyla(14). Phylogenetic analysis

can determine whether plasmid evolution between BSI and livestock-associated niches is

driven by host clonal expansion or other means, as well as allow us to explore the early

emergence of AMR gene carrying plasmids.

As a detailed example, we evaluated the largest plasmid cluster containing both human and

livestock-associated plasmids (cluster 2, n=100 members). All plasmids carried at least one

F-type replicon and were all putatively conjugative, with 75% (75/100) and 25% (25/100)

assigned PTU-F_E and a putative PTU, respectively. Further, 48% (48/100) plasmids carried

bla_{TEM-1}, and 51% (51/100) carried >1 AMR gene. All host chromosomes were $E.$ $coli$ except

OX-BSI-481_2 ($S.$ $enterica$ ST 2998; hereon omitted from the analysis). The n=99 $E.$ $coli$

isolates represented six phylogroups: A (5/99), B1 (18/99), B2 (52/99), C (14/99), D (7/99),

and G (3/99; see Supplementary Methods).

Figure 4b-c shows the plasmid core gene phylogeny ($T_{plasmid}$) and the $E.$ $coli$ host core gene

phylogeny ($T_{chromosome}$). The $E.$ $coli$ phylogeny was structured by six clades corresponding to

the six phylogroups (see Supplementary Methods). We found low congruence between the

plasmid core-gene phylogeny and the chromosomal core-gene phylogeny as seen in the

central 'tanglegram' (i.e., lines connecting pairs of plasmid and chromosome tips from the

same isolate). Additionally, we calculated a Robinson-Foulds distance $RF(T_{\text{plasmid}}$,

$T_{\text{chromosome}})=162$, reflecting a high number of structural differences between the phylogenies

(see Supplementary Methods). There was some evidence of plasmid structuring by niche

(Fritz and Purvis' $D=0.24$; see Supplementary Methods).

Within the plasmid phylogeny, there was a clade of $n=44$ plasmids (support 100%; circled in

grey in Figure 4b) containing both BSI and livestock-associated plasmids, which were within

median 4 core gene SNPs of each other (IQR=2-8, range=0-59). Estimating plasmid

evolution at an approximate rate of one SNP per year (see Supplementary Methods) would

give a median time to most recent common ancestor of the backbone at approximately 4

years prior to sampling, consistent with recent movement between human and livestock-

associated niches. This plasmid clade was mainly present in phylogroup B2 (20/44), but also

A (3/44), B1 (9/44), C (8/44), and D (4/44), suggesting plasmid movement. Further, 77%

(34/44) of plasmids within the clade carried bla_{TEM-1} (BSI: 25/34, Livestock-associated: 8/34,

WwTW-associated: 1/34), and 82% (36/44) carried ≥1 AMR gene, highlighting the role of

plasmids in cross-niche dissemination of AMR.

To examine the evolution of entire plasmid sequences within the clade, we represented all

$n=44$ plasmids as a 'pangraph' (Figure 4d; see Supplementary Methods). Briefly, pangraph

converts input sequences into a consensus graph, where each sequence is a path along a set of

homologous sequence alignments i.e., 'blocks', which in series form 'pancontigs'. Filtering

for 'core blocks' (i.e., those found in ≥95% plasmids), we found 4 pancontigs (40 blocks

total), with the longest 98,269bp (total length 125,369bp), indicating a putative plasmid

backbone (Fig. 4e). Then, filtering for 'accessory blocks' (i.e., those found in <95%

plasmids), we found 18 pancontigs (39 blocks total), with median length 2,380bp (total length

63,753bp), forming the accessory gene repertoire (Fig. 4f). This points to a persistent plasmid

backbone structure with loss/gain events at particular 'hotspots' as well as rearrangements.

Discussion

Sharing of plasmids between different niches is normally focused on those carrying AMR

genes that are of particular current clinical concern, such as extended-spectrum beta-

lactamase (ESBL) or carbapenemase genes, meaning we lack information on the vast

'denominator' of background plasmid sharing, and on the dissemination of other AMR genes

which are now widespread in clinical isolates and from which important insights might be

gained. By analysing a dataset of n=3,697 systematically collected *Enterobacterales* plasmids

sampled from human BSI, livestock- and WwTW-associated sources in a geographically and

temporally restricted context, we found evidence supporting significant plasmid

dissemination across niches, putting those which carry AMR genes of current major clinical

concern into context. We found 225 instances of shared, near-identical plasmid groups, 25%

of which were found across multiple bacterial STs, 4% across multiple bacterial species, and

8% in both human BSI and ≥1 non-BSI niche. Beyond this near-identical sharing, we

analysed 'clusters' of plasmids and found that 73/247 clusters contained plasmids seen in

both human BSIs and other contexts. Approximately a fifth (52/247) of plasmid clusters

contained plasmids carrying AMR genes (n=550 plasmids). Our results suggest the need for

broad, unselected, and detailed sampling frames to fully understand plasmid diversity and

evolution, and to evaluate the "One Health" risk of AMR associated with plasmid-sharing

across niches.

Whilst many plasmid clusters were strongly structured by host phylogeny and isolate source,

some plasmids from human BSIs were highly genetically related to those in other niches,

including livestock. However, not all of these carried AMR genes. Our results highlight the

potential routes for transfer that exist through similar plasmids. However, recovering these

instances of putative sharing is a sampling challenge. Accumulation curve analyses suggested

increasing the size of our dataset would have led to further near-identical matches at an

approximately linear rate, meaning even a dataset of this size captures only a small fraction of

the true extent of plasmid sharing between human clinical and other non-human/clinical

niches. This presents a challenge for designing appropriately powered studies. Had we only

sampled n=100 livestock-associated isolates (i.e., around 20% of our actual sample), there

was only a 39% chance that we would have detected ≥5 matches with BSI plasmids (Fig. S9).

Understanding the evolutionary history, distribution, and epidemiology of well-known genes

in environmental plasmids may offer insights into the future trajectories of more recently

emerged genes. For example, the first plasmid-encoded beta-lactamase to be described was

bla_{TEM-1}, identified in 1965 in an *E. coli* isolate in Greece(28) and now widely prevalent in

Enterobacterales(29). bla_{TEM-1} has a narrow spectrum of activity and is now less clinically

concerning than newer genes which mediate broad-spectrum resistance, but in our dataset

bla_{TEM-1} was strongly associated with plasmid clusters seen in BSI and with the carriage of

other AMR genes. bla_{TEM-1} may continue to play an important role in the spread of AMR-

carrying plasmids which can transfer recently emerged genes, and similarities in its

association with plasmids and other smaller transposable mobile genetic elements may reflect

the future trajectory of other AMR genes of more recent clinical concern such as ESBLs and

carbapenemases.

Given that plasmids observed in BSI isolates represent small proportion of human

Enterobacterales diversity, many more sharing events may occur in the human gut(30) which

we only sampled incompletely using wastewater influent as a proxy. The human colon

contains around 10^{14} bacteria(31), with large ranges of *Enterobacteriaceae* abundance.

Further, even small numbers of across-niche sharing events, such as transfer events of

important AMR genes from species-to-species or niche-to-niche, may have significant

clinical implications, as has been seen with several important AMR genes globally. Future

studies need to carefully consider the limitations of sampling frames in detecting any genetic

overlap, given both substantial diversity and the effects of niches and geography(11,16).

By examining plasmid relatedness compared to bacterial host relatedness, we demonstrated

that cross-niche plasmid spread is not driven by clonal lineages. Using a pangenome-style

analysis, we showed that plasmids can share sets of near-identical core genes alongside

diverse accessory gene repertoires. While plasmids with more distantly related core genes

tended to have dissimilar accessory gene content, plasmids with more closely related core

genes shared a wide range of accessory gene content. This would be consistent with a

hypothesis of persistent 'backbone' structures gaining and losing accessory functions as they

move between hosts and niches. We suggest that this mode of transfer might be worth

considering. Evolutionary models for plasmids which can accommodate well-conserved

backbone evolution alongside accessory structural changes and gain/loss events are urgently

needed. Estimating plasmid evolutionary rates remains a challenge, with little known about

appropriate values for mutation rates in plasmids, and even less for non-mutational processes

such as gene gain/loss.

Our study had several limitations. Our non-BSI isolates were not as temporally varied as the

BSI isolates, meaning we could not fully explore temporal evolution. Isolate-based

methodologies are limited in evaluating the true diversity of the niches sampled; composite

approaches including metagenomics might shed additional insight in future studies. Further,

the exact source of an isolate is poorly defined for wastewater/waterway isolates as they act

as a confluence of multiple sources, although they represent important niches in their own

right. We only analysed plasmids from complete genomes i.e., where the chromosome and all

plasmids were circularised, meaning we disregarded ~23% and ~33% of BSI and non-BSI

assemblies, respectively. The exclusive use of complete assemblies was to ensure full

plasmid sequences could be examined in their full genomic context. We only focused on

plasmids as horizontally transmissible elements here; detailed study of other smaller mobile

genetic elements across-niches would represent interesting future work. We have also

investigated a limited subset of *Enterobacterales*: plasmid sharing likely extends to other

bacterial hosts not investigated here. Lastly, our isolate culture methods for livestock-

associated samples may not have been as sensitive for the identification of *Klebsiella* spp. as

for other *Enterobacterales* such as *Escherichia*, as we did not use enrichment and selective

culture on Simmons citrate agar with inositol(32).

In conclusion, this study presents to our knowledge the largest evaluation of systematically

collected *Enterobacterales* plasmids across human and non-human niches within a

geographically and temporally restricted context. Plasmids can clearly disseminate between

niches, although this dynamic likely varies by cluster; the overall number of near-identical

plasmid groups identified across niches consistent with recent transfer events was 8%

(17/225) and influenced by sample size. We demonstrate a likely intertwined ecology of

plasmids across human and non-human niches, where different plasmid clusters are variably

but incompletely structured and putative 'backbone' plasmid structures can rapidly gain and

lose accessory genes following cross-niche spread. Future "One Health" studies require dense

and unselected sampling, and complete/near-complete plasmid reconstruction, to

appropriately understand plasmid epidemiology across niches.

Materials and Methods

Livestock-associated isolates

n=247 *Enterobacterales* isolates from farm-proximate soils and poultry faeces (n=19 farms;

n=5 cattle, n=4 pig, n=5 poultry, n=5 sheep) were collected and sequenced for this study in

2017-2020. DNA extraction and sequencing was performed as in Shaw *et al.*, 2021(11).

Genomes were hybrid assemblies reconstructed using Unicycler(33) (v. 0.4.4; default hybrid

assembly parameters except min_component_size 500 and --min_dead_end_size 500). Only

complete assemblies (plasmids and chromosomes) were considered (n=162/247).

BSI isolates

Sequenced Human BSI Enterobacterales isolates from patients presenting to n=4 hospitals

within Oxfordshire, UK, September 2008-December 2018, as described in Lipworth *et al.*,

2021(34) were also included. Although all patients were sampled in Oxfordshire, a total of

n=505/738 patients resided in Oxfordshire, n=133/738 in surrounding counties, and

n=100/738 had location information omitted. Only complete assemblies (n=738/953 total

assembled) were considered.

Other livestock-associated and WwTW-associated isolates

Enterobacterales isolates from faeces from the n=14 non-poultry farms and wastewater

influent, effluent, and waterways upstream/downstream of effluent outlets surrounding n=5

WwTWs, across 3 seasonal timepoints in 2017 (as in (11)) were included. Only complete

assemblies (n=558/827 total assembled) were considered.

Statistical analysis and bioinformatics

Chromosome sequence types (STs) were determined with mlst (v. 2.19.0; see Supplementary

Methods). To generate accumulation curves (ACs), new plasmid diversity was recorded for

each isolate sampled randomly, without replacement. A bootstrapped average of $b=1,000$

ACs was used to estimate a Heap's parameter (γ) by fitting a linear regression to log-log

transformed data see (Supplementary Methods). We adopted three approaches to plasmid

classification, using COPLA to classify plasmids into broad plasmid taxonomic units (PTUs),

and also grouping and clustering plasmids into smaller clusters using alignment-free

distances (see Supplementary Methods). Within plasmid clusters, we identified core genes

with Panaroo (v. 1.2.9), aligned them with mafft (v7.407) and produced trees with IQ-tree (v.

2.0.6). Plots were primally produced using the R library ggplot2, with additional graphics in

BioRender. More information can be found in the Supplementary Methods.

Data availability

Study metadata is provided in Table S2. Accessions for poultry and environmental soil isolate

reads are given in Table S3, and assemblies will shortly be made available on NCBI.

Accessions for existing BSI and REHAB reads and assemblies can be found in Lipworth *et*

al., 2021(34) (BioProject PRJNA604975) and Shaw *et al.*, 2021(11) (BioProject

PRJNA605147) respectively.

Code availability

Analysis scripts can be found in the GitHub repository

https://github.com/wtmatlock/oxfordshire-overlap.

REHAB Consortium.

Manal AbuOun[2], Muna F. Anjum[2], Mark J. Bailey[3], Brett H[8], Mike J. Bowes[3], Kevin K.

Chau[1], Derrick W. Crook[1,6,7], Nicola de Maio[1], Nicholas Duggett[2], Daniel J. Wilson[1,9],

Daniel Gilson[2], H. Soon Gweon[3,4], Alasdair Hubbard[10], Sarah J. Hoosdally[1], William

Matlock[1], James Kavanagh[1], Hannah Jones[2], Timothy E. A. Peto[1,6,7], Daniel S. Read[3], Robert

Sebra[5], Liam P. Shaw[1], Anna E. Sheppard[1,6], Richard P. Smith[2], Emma Stubberfield[2], Nicole

Stoesser[1,6,7], Jeremy Swann[1], A. Sarah Walker[1,6,7], Neil Woodford[11]

[1] Nuffield Department of Medicine, University of Oxford, Oxford, UK

[2] Animal and Plant Health Agency, Weybridge, Addlestone, UK

[3] UK Centre for Ecology & Hydrology, Wallingford, UK

[4] University of Reading, Reading, UK

[5] Icahn Institute of Data Science and Genomic Technology, Mt Sinai, NY, USA

[6] NIHR HPRU in healthcare-associated infection and antimicrobial resistance, University of

Oxford, Oxford, UK

[7] NIHR Oxford Biomedical Research Centre, University of Oxford, Oxford, UK

[8] Thames Water Utilities, Clearwater Court, Vastern Road, Reading, UK

[9] Wellcome Trust Centre for Human Genetics, University of Oxford, Roosevelt Drive,

Oxford, UK

[10] Department of Tropical Disease Biology, Liverpool School of Tropical Medicine,

Liverpool, UK

[11] Antimicrobial Resistance and Healthcare Associated Infections (AMRHAI) Reference Unit,

National Infection Service, Public Health England, London, United Kingdom

Declarations of interests. The authors declare no interests.

Role of the funding source.

This work was funded by the Antimicrobial Resistance Cross-council Initiative supported by

the seven research councils [grant NE/N019989/1]. The UKCEH component of the REHAB

consortium was supported by the Natural Environment Research Council (NERC)

[grant NE/N019660/1]. DWC, SG, TEAP and NS are supported by the National Institute for

Health Research Health Protection Research Unit (NIHR HPRU) in Healthcare-Associated

Infections and Antimicrobial Resistance at the University of Oxford in partnership with

Public Health England (PHE) [grant HPRU-2012–10041 and NIHR200915]. DWC and

TEAP are also supported by the NIHR Oxford Biomedical Research Centre. The

computational aspects of this research were funded from the NIHR Oxford BRC with

additional support from a Wellcome Trust Core Award Grant [grant 203141/Z/16/Z]. The

views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the

Department of Health or Public Health England. WM and KKC are supported by a

scholarship from the Medical Research Foundation National PhD Training Programme in

Antimicrobial Resistance Research (MRF-145-0004-TPG-AVISO). NS is an Oxford Martin

Fellow and a Senior NIHR BRC Oxford Fellow. LPS is a Sir Henry Wellcome Postdoctoral

Fellow funded by Wellcome (grant 220422/Z/20/Z).

Bibliography

1. Linh TD, Thu NH, Shibayama K, Suzuki M, Yoshida LM, Thai PD, et al. Expansion of
KPC-producing Enterobacterales in four large hospitals in Hanoi, Vietnam. J Glob
Antimicrob Resist. 2021;27.
2. Kraftova L, Finianos M, Studentova V, Chudejova K, Jakubu V, Zemlickova H, et al.
Evidence of an epidemic spread of KPC-producing Enterobacterales in Czech
hospitals. Sci Rep. 2021;11(1).
3. Cahill N, O'Connor L, Mahon B, Varley Á, McGrath E, Ryan P, et al. Hospital effluent: A
reservoir for carbapenemase-producing Enterobacterales? Science of the Total
Environment. 2019;672.
4. Subramanya SH, Bairy I, Metok Y, Baral BP, Gautam D, Nayak N. Detection and
characterization of ESBL-producing Enterobacteriaceae from the gut of subsistence
farmers, their livestock, and the surrounding environment in rural Nepal. Sci Rep.
2021;11(1).
5. Abuoun M, Jones H, Stubberfield E, Gilson D, Shaw LP, Hubbard ATM, et al. A genomic
epidemiological study shows that prevalence of antimicrobial resistance in
enterobacterales is associated with the livestock host, as well as antimicrobial usage.
Microb Genom. 2021;7(10).
6. Díaz-Gavidia C, Barría C, Rivas L, García P, Alvarez FP, González-Rocha G, et al.
Isolation of Ciprofloxacin and Ceftazidime-Resistant Enterobacterales From
Vegetables and River Water Is Strongly Associated With the Season and the Sample
Type. Front Microbiol. 2021;12.
7. Buchy P, Ascioglu S, Buisson Y, Datta S, Nissen M, Tambyah PA, et al. Impact of
vaccines on antimicrobial resistance. Vol. 90, International Journal of Infectious
Diseases. 2020.
8. Ruppé E, Cherkaoui A, Charretier Y, Girard M, Schicklin S, Lazarevic V, et al. From
genotype to antibiotic susceptibility phenotype in the order Enterobacterales: a
clinical perspective. Clinical Microbiology and Infection. 2020;26(5).
9. Sheppard AE, Stoesser N, Wilson DJ, Sebra R, Kasarskis A, Anson LW, et al. Nested
Russian doll-like genetic mobility drives rapid dissemination of the carbapenem
resistance gene blakpc. Antimicrob Agents Chemother. 2016;60(6).
10. Che Y, Yang Y, Xu X, Brinda K, Polz MF, Hanage WP, et al. Conjugative plasmids
interact with insertion sequences to shape the horizontal transfer of antimicrobial
resistance genes. Proc Natl Acad Sci U S A. 2021;118(6).
11. Shaw LP, Chau KK, Kavanagh J, AbuOun M, Stubberfield E, Gweon HS, et al. Niche and
local geography shape the pangenome of wastewater-and livestock-associated
Enterobacteriaceae. Sci Adv. 2021;7(15).
12. Orlek A, Stoesser N, Anjum MF, Doumith M, Ellington MJ, Peto T, et al. Plasmid
classification in an era of whole-genome sequencing: Application in studies of
antibiotic resistance epidemiology. Vol. 8, Frontiers in Microbiology. 2017.
13. Matlock W, Chau KK, AbuOun M, Stubberfield E, Barker L, Kavanagh J, et al. Genomic
network analysis of environmental and livestock F-type plasmid populations. ISME
Journal. 2021;15(8).
14. Redondo-Salvo S, Fernández-López R, Ruiz R, Vielva L, de Toro M, Rocha EPC, et al.
Pathways for horizontal gene transfer in bacteria revealed by a global map of their
plasmids. Nat Commun. 2020;11(1).

15.   Mounsey O, Schubert H, Findlay J, Morley K, Puddy EF, Gould VC, et al. Limited
phylogenetic overlap between fluoroquinolone-resistant Escherichia coli isolated on
dairy farms and those causing bacteriuria in humans living in the same geographical
region. Journal of Antimicrobial Chemotherapy. 2021;76(12).
16.   Hanage WP. Two health or not two health? That is the question. mBio.
2019;10(2):e00550-19.
17.   Ludden C, Raven KE, Jamrozy D, Gouliouris T, Blane B, Coll F, et al. One health
genomic surveillance of escherichia coli demonstrates distinct lineages and mobile
genetic elements in isolates from humans versus livestock. mBio. 2019;10(1).
18.   Shen C, Zhong LL, Yang Y, Doi Y, Paterson DL, Stoesser N, et al. Dynamics of mcr-1
prevalence and mcr-1-positive Escherichia coli after the cessation of colistin use as a
feed additive for animals in China: a prospective cross-sectional and whole genome
sequencing-based molecular epidemiological study. Lancet Microbe. 2020;1(1):e34–
43.
19.   Cherak Z, Loucif L, Moussi A, Rolain JM. Epidemiology of mobile colistin resistance
(mcr) genes in aquatic environments. Vol. 27, Journal of Global Antimicrobial
Resistance. 2021.
20.   Carlos Bastidas-Caldes, Jacobus H. de Waard, María Soledad Salgado, María José
Villacís, Marco Coral-Almeida, Yoshimasa Yamamoto, et al. Worldwide Prevalence of
mcr-mediated Colistin-Resistance Escherichia coli in Isolates of Clinical Samples,
Healthy Humans, and Livestock—A Systematic Review and Meta-Analysis. Pathogens.
2022 Jun;11(6).
21.   Hilpert C, Bricheux G, Debroas D. Reconstruction of plasmids by shotgun sequencing
from environmental DNA: Which bioinformatic workflow? Brief Bioinform.
2021;22(3).
22.   Wang R, van Dorp L, Shaw LP, Bradley P, Wang Q, Wang X, et al. The global
distribution and spread of the mobilized colistin resistance gene mcr-1. Nat Commun.
2018;9(1):1–9.
23.   Sekizuka T, Matsui M, Yamane K, Takeuchi F, Ohnishi M, Hishinuma A, et al. Complete
sequencing of the bla NDM-1-positive IncA/C plasmid from Escherichia coli ST38
isolate suggests a possible origin from plant pathogens. PLoS One. 2011;6(9).
24.   Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, Guerra B,
et al. Plasmids carrying antimicrobial resistance genes in Enterobacteriaceae. Journal
of Antimicrobial Chemotherapy. 2018;73(5).
25.   Rodríguez-Beltrán J, DelaFuente J, León-Sampedro R, MacLean RC, San Millán Á.
Beyond horizontal gene transfer: the role of plasmids in bacterial evolution. Vol. 19,
Nature Reviews Microbiology. 2021.
26.   Pesesky MW, Tilley R, Beck DAC. Mosaic plasmids are abundant and unevenly
distributed across prokaryotic taxa. Plasmid. 2019;102.
27.   Coluzzi C, Garcillán-Barcia MP, de La Cruz F, Rocha EPC. Evolution of Plasmid Mobility:
Origin and Fate of Conjugative and Nonconjugative Plasmids. Mol Biol Evol.
2022;39(6).
28.   Datta N, Kontomichalou P. Penicillinase synthesis controlled by infectious R factors in
enterobacteriaceae. Nature. 1965;208(5007).
29.   Bush K, Bradford PA. Epidemiology of β-lactamase-producing pathogens. Vol. 33,
Clinical Microbiology Reviews. 2020.

30.   Forster SC, Kumar N, Anonye BO, Almeida A, Viciani E, Stares MD, et al. A human gut
bacterial genome and culture collection for improved metagenomic analyses. Nat
Biotechnol. 2019;37(2).

31.   Sender R, Fuchs S, Milo R. Revised Estimates for the Number of Human and Bacteria
Cells in the Body. PLoS Biol. 2016;14(8).

32.   Rodrigues C, Hauser K, Cahill N, Ligowska-Marzęta M, Centorotola G, Cornacchia A, et
al. High Prevalence of Klebsiella pneumoniae in European Food Products: a
Multicentric Study Comparing Culture and Molecular Detection Methods. Microbiol
Spectr. 2022;10(1).

33.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. PLoS Comput Biol. 2017;13(6).

34.   Lipworth S, Vihta KD, Chau K, Barker L, George S, Kavanagh J, et al. Ten-year
longitudinal molecular epidemiology study of Escherichia coli and Klebsiella species
bloodstream infections in Oxfordshire, UK. Genome Med. 2021;13(1).

 Fig. 1. A diverse sample of geographically and temporally restricted *Enterobacterales*

 (a) Number of chromosomes and plasmids by niche, stratified by isolate genus. **(b)** Map of

 approximate, relative distances between sampling sites, coloured by niche (human

 bloodstream infection [BSI], livestock-associated (cattle, pig, poultry, and sheep faeces, soils

 nearby livestock sites), and wastewater treatment work (WwTW)-associated sources

 (influent, effluent, waterways upstream/downstream of effluent outlets). Number in circles

 indicates how many of the *n*=1,458 isolates are from that location. **(c)** Sampling timeframe

 for BSI and REHAB (non-BSI) isolates.

601 **Fig. 2 Cross-niche, near-identical plasmids.**

602 **(a)** Size of cross-niche, near-identical plasmid groups, coloured by niche (total *n*=84

603 plasmids). Median length (bp) of plasmids within groups increases from left to right. **(b)**

604 Proportion of plasmid host species by group. **(c)** Predicted mobility of plasmid. **(d)** AMR

605 gene carriage in plasmid.

 Fig. 3. Genetically similar plasmids share between niches

 (a) Size of plasmid clusters with at least 10 members, coloured by niche. Size of clusters

 decreases from left to right. **(b)** Proportion of plasmid host species by cluster. **(c)** Plasmid

 mobility class and size: Left hand axis shows proportion of plasmids with a predicted

 mobility class by cluster. Right hand axis shows plasmid length boxplots by cluster. **(d)**

 Proportions of AMR gene carriage by cluster. **(e)** Plasmid core and accessory genomes: Left

 hand axis shows the count of core and accessory coding sequences (CDS) by cluster. Right

 hand axis shows plasmid core gene proportion (i.e., plasmid core CDS/total plasmid CDS)

614	boxplots by cluster.

 Fig. 4. Cluster 2 plasmid and host evolution

 (a) Consensus gene ordering for plasmid cluster 2, coloured by gene type (total *n*=99

 plasmids; *n*=1 *S. enterica* isolate omitted). Genes are coloured by core, accessory, or

 transposase. **(b)** Plasmid core gene phylogeny with tips coloured by sampling niche. The grey

 circle highlights the clade of *n*=44 plasmids which were further analysed. **(c)** Plasmid host

 chromosome core gene phylogeny with tips coloured by sampling niche. Plasmid and host

 phylogeny tips are connected in a 'tanglegram' which connects pairs of plasmids and

 chromosomes from the same isolate. **(d)** Visualisation of the pangraph for *n*=44 plasmids in

 the grey-circled clade in (b). Blocks are coloured by presence in plasmids. **(e)** Core blocks

 (found in at least 95% of the *n*=44 plasmids). **(f)** Accessory blocks (found in less than 95% of

 the *n*=44 plasmids).

a Plasmid consensus gene synteny   **b** Plasmid core gene phylogeny   **c** Chromosome core gene phylogeny

Gene type
- Absent
- Core
- Accessory
- Transposase

Niche
- BSI
- Livestock–associated
- WwTW–associated

Phylogroup
- A
- B1
- B2
- C
- D
- G

No. nucleotide substitutions per site

0   0.01   0.02   0.04   0.02   0

d Pangraph   **e** Core blocks   **f** Accessory blocks

Block type
- Core
- Accessory