Alone in a crowd: Effect of a nonfunctional lateral line on expression of the social hormone

parathyroid hormone 2.

Alexandra Venuto[1], Cameron P. Smith[1,2], Marybelle Cameron-Pack[3], and Timothy Erickson[3]

1. East Carolina University, Greenville, NC, USA

2. East Carolina University School of Dental Medicine, Greenville, NC, USA

3. University of New Brunswick, Fredericton, NB, CAN

Corresponding author: Timothy Erickson, timothy.erickson@unb.ca

SUMMARY STATEMENT

Expression of the pro-social neuropeptide *pth2* is downregulated in larval zebrafish with a congenital loss of lateral line function. Thus, even in social environments, fish with compromised lateral lines may experience chronic loneliness.

ABSTRACT

Parathyroid hormone 2 (Pth2) is a vertebrate-specific neuropeptide whose thalamic expression is upregulated by social contact with conspecifics. However, social interactions fail to stimulate *pth2* expression in isolated zebrafish whose lateral line hair cells have been chemically ablated. These results suggest that modulation of *pth2* by social context is acutely dependent on mechanosensory information from the lateral line. However, it is unclear how a congenital loss of lateral line function influences the ability of zebrafish to interpret their social environment. In this study, we measure *pth2* levels in zebrafish mutants lacking hair cell function in either the lateral line only, or in both the inner ear and lateral line. Socially-raised lateral line mutants express lower levels of *pth2* relative to wild type siblings, but there is no further reduction when all sensory hair cells are nonfunctional. However, social isolation of hair cell mutants causes a further reduction in *pth2* expression, pointing to additional unidentified sensory cues that influence *pth2* production. Lastly, we report that social context modulates fluorescent transgenes driven by the *pth2* promoter. Altogether, these data suggest that lateral line mutants experience a chronic sense of isolation, even when raised in a social environment.

INTRODUCTION

The complex relationship between the brain and social behavior has been studied across animal species for decades (Brothers, 1990; Chen and Hong, 2018; Kravitz and Huber, 2003; Robinson et al., 2008). Different social contexts can evoke a wide range of emotions, from joy and positivity (Burgdorf and Panksepp, 2006; Panksepp and Burgdorf, 2003) to fear and aggression (Amaral, 2002; Neumann et al., 2010) with a high level of plasticity (Chen and Hong, 2018; Kolb and Whishaw, 2003). In situations of social isolation, many animals exhibit anxious behaviors in addition to neurochemical changes in the brain (Amaral, 2002; Mumtaz et al., 2018; Shams et al., 2017; Weiss et al., 2004).

Social behavior is modulated by a suite of neuropeptides (Adkins-Regan, 2009; O'Connell
and Hofmann, 2011; Oldfield and Hofmann, 2011; Panksepp et al., 2007; Schoofs et al., 2017).
Parathyroid hormone 2 (Pth2, formerly known as tuberoinfundibular peptide of 39 residues
or TIP39) is a neuropeptide expressed in the vertebrate thalamus, which is a sensory integration
and relay center (Anneser et al., 2020; Bhattacharya et al., 2011; Dobolyi et al., 2012; Mueller,
2012). In rodents, the PTH2 receptor is expressed in several nuclei throughout the central nervous
system (Dobolyi et al., 2012). Zebrafish have two *pth2r* paralogs, *pth2ra* and *pth2rb,* each with
distinct expression patterns in the fish brain (Seitz, 2019). Activation of PTH2R increases
intracellular cAMP levels and can modulate neuroendocrine activity in the hypothalamic–
pituitary–adrenal (HPA) axis (Dimitrov and Usdin, 2010). PTH2-PTH2R signaling influences
several behaviors, including maternal care (Cservenák et al., 2013), pain sensitivity (Dimitrov et
al., 2010; Dimitrov et al., 2013; LaBuda and Usdin, 2004), and anxiety-like responses (Anneser et
al., 2022; Coutellier and Usdin, 2011; LaBuda et al., 2004).

Recent work suggests that Pth2 signaling promotes social interactions between
conspecifics. In zebrafish, *pth2* mutants exhibit decreased preference for conspecifics (Anneser et
al., 2022), while chemogenetic activation of the Pth2-positive region of the rodent brain promotes
social contact (Keller et al., 2022). Consistent with this, expression of *pth2* itself positively scales
with the quantity of social interactions with conspecifics (Anneser et al., 2020; Anneser et al.,
2022; Keller et al., 2022). In rodents, physical touch from conspecifics activates *Pth2* expression
(Cservenák et al., 2013; Keller et al., 2022). In addition to tactile touch sensation, fish also have
another form of "distant touch" through the mechanosensory lateral line (Dijkgraaf, 1963). The
mechanosensory lateral line is a hair cell-based system in aquatic vertebrates that senses
hydrodynamic information, such as that created by conspecifics (Butler and Maruska, 2015; Butler
and Maruska, 2016a; Butler and Maruska, 2016b; Groneberg et al., 2020; Montgomery et al.,
2013). In zebrafish, chemical ablation of the mechanosensory lateral line prevents the rescue of
pth2 expression by a social environment following isolation (Anneser et al., 2020). Therefore,
social "touch" (direct contact in rodents and hydrodynamic communication in zebrafish) seems to
be a primary sensory activator of *pth2* expression.

Despite this knowledge, there remain several unanswered questions regarding the social
regulation of *pth2* expression. First, how do zebrafish interpret social context when they are born
without lateral line function? Secondly, do mechanosensory hair cells in the inner ear contribute

to the ability of zebrafish larvae to respond to social context? Thirdly, is *pth2* expression regulated
solely via mechanosensation, or can fish use additional sensory inputs to sense the presence of
conspecifics? Here, we report that the congenital loss of mechanosensory hair cell function leads
to chronically depressed levels of *pth2* expression in socially-housed larval zebrafish.
Interestingly, social isolation of hair cell mutants further decreases *pth2* levels, suggesting that
larval fish use additional sensory information to sense conspecifics and modulate *pth2* expression.
Lastly, we utilize fluorescent reporters driven by the *pth2* regulatory region to evaluate the
morphology of *pth2*-expressing cells and visualize *pth2* dynamics *in vivo*. Overall, this study offers
additional insight into the sensory basis for social interactions in zebrafish and introduces novel
tools to track *pth2* expression in living fish.

RESULTS
Decreased *pth2* expression in larval zebrafish with a genetically disabled lateral line.
pth2 expression decreases when fish are raised in isolation but increases following a brief
period of social interaction with conspecifics. However, when the mechanosensory lateral line is
ablated in isolated fish prior to social exposure, *pth2* expression levels remain low (Anneser et al.,
2020). This implicates the lateral line sensory system in recognizing conspecifics to stimulate *pth2*
expression. To further evaluate the role of the lateral line in social recognition, we raised lateral
line-deficient mutants (*lhfpl5b*[-/-]) and their siblings in separate social groups and detected *pth2*
expression by mRNA *in situ* hybridization (ISH). Socially-raised lateral line mutants exhibit a
decrease in *pth2* expression compared to wild type (WT) siblings (Fig. 1B, C), both in terms of a
qualitative decrease in the intensity of staining as well as a significant decrease in the number of
detectable *pth2*-positive cells (One-tailed Welch's t-test, WT Soc: M = 29.9, SD = 1.7; MUT Soc:
M = 9.7, SD = 4.3; $p < 0.0001$) (Fig. 1D). We also ablated lateral line hair cells in socially raised
WT larvae using repeated neomycin treatments (Venuto and Erickson, 2021) (Figure 1E). Even
though these larvae had social experience prior to losing lateral line input, they exhibited low levels
of *pth2* expression similar to lateral line mutants. These results confirm that the loss of lateral line
function impairs the ability of *pth2* cells to respond to social context.

Mechanosensory input from inner ear hair cells do not contribute to *pth2* expression.

A low level of *pth2* expression remains in socially-raised lateral line mutants (Fig. 1B, C).

To determine if mechanosensory input from the inner ear was responsible for this remaining *pth2*

expression, we used three different types of mutations that disable hair cells of both the lateral line

and inner ear to compare to the lateral line mutant (*lhfpl5b$^{-/-}$*) and wild type zebrafish. Two of these

mutations (*pcdh15a$^{-/-}$* and *tomt$^{-/-}$*) render the mechano-electrical transduction channel

nonfunctional in all hair cells, similar to how the *lhfpl5b* mutation specifically disables lateral line

hair cells (Erickson et al., 2017; Erickson et al., 2020; Maeda et al., 2017). The third mutant

disables the voltage-gated calcium channel in hair cells (*cacna1da$^{-/-}$*), thereby blocking stimulus-

dependent glutamate release (Sheets et al., 2012; Sidi et al., 2004). Again, we measured *pth2*

expression via mRNA ISH (Fig. 2B-F) as well as quantitative PCR (qPCR) (Fig. 2G). All hair cell

mutants exhibit similarly low levels of *pth2* expression when raised socially, with no additional

decrease seen in zebrafish with genetically disabled inner ear and lateral line sensory function

(One-way ANOVA with Tukey post-test, $\alpha = 0.05$). From these experiments, we find that inner

ear mechanosensory hair cells do not contribute to the regulation of *pth2* expression.

Social isolation further reduces *pth2* expression in sensory hair cell mutants.

Chemosensory and visual perception of conspecifics does not significantly rescue *pth2*

expression in larval zebrafish raised in isolation (Anneser, et al., 2020). However, since disabling

both the lateral line and the inner ear does not completely eliminate *pth2* expression, other non-

hair cell-based sensory systems may contribute to the modulation of *pth2* by social context. To test

this, we compared *pth2* expression in social and isolated *lhfpl5b* and *tomt* mutants by both mRNA

ISH (Fig. 3A-C) and qPCR (Fig. 3D). If *pth2* responds to conspecific signals solely via the lateral

line, we expect to see equivalently low expression in mutants raised in social and isolated

environments. However, we find this not to be the case. Instead, isolated *lhfpl5b* and *tomt* mutants

express less *pth2* than social mutant siblings (One-way ANOVA with Tukey post-test, $\alpha = 0.05$).

pth2 levels in isolated mutants and wild types are indistinguishable. Because social isolation

further reduces *pth2* expression in lateral line mutants, we conclude that *pth2* responds to

multisensory input during social experience.

Morphology of *pth2*-expressing cells using fluorescent reporters.

Fluorescent reporters provide information about the morphology and function of cells, along with the ability to quantify gene expression in living organisms (Choe et al., 2021; Soboleski et al., 2005). Therefore, we created stable lines of transgenic zebrafish expressing GFP or RFP under control of the presumptive *pth2* promoter. These transgenes allow us to visualize the morphology of the *pth2*-expressing cells in living larvae (Fig. 4 and S1). Imaging of transgenic larvae reveals the two bilateral cell clusters with complex projection patterns (Fig. 4B). The anterior ipsilateral projections converge into a dense neuropil in the forebrain, while other fibers project ventrally and posteriorly toward the midbrain. Additionally, we find ventral projections that cross the midline between the two cell clusters. This morphology is consistent in both the GFP and RFP transgenes.

The *pth2* transgenes respond to lateral line input and social context.

Next, we tested if the transgene responds to social context (Fig. 5). Consistent with our mRNA ISH and qPCR results, transgenic *lhfpl5b* mutants (Fig. 5A-C) and isolates (Fig. 5D-F) both exhibit a significant decrease in the number of fluorescent cells relative to control siblings (Fig. 5C: One-tailed Welch's t-test, WT Soc: M = 43.2, SD = 3.9; MUT Soc: M = 34.2, SD = 2.8; $p < 0.0001$) (Fig. 5F: One-tailed Welch's t-test, WT Soc: M = 37.3, SD = 2.6; WT Iso: M = 28.3, SD = 2.8; $p < 0.0001$). To gauge whether isolation specifically diminishes expression from the *pth2*-driven transgenes, we compared the effect of social isolation on transgene expression in *Tg(pth2:TagRFP)unb3* and *Tg(slc6a3:EGFP)ot80* larvae. The *slc6a3:*EGFP transgene expresses GFP under control of the dopamine transporter promoter and marks a population of dopaminergic cells in the pretectum, which lies dorsally to the *pth2*-expressing cells. The number of dopaminergic cells in the pretectum do not change in response to social context (One-tailed Welch's t-test, WT Soc: M = 44.0, SD = 4.3; WT Iso: M = 45.1, SD = 4.7; $p = 0.3$) (Fig. 5D-F). These experiments demonstrate that the *pth2* transgenes respond to social context and allows for a wide variety of future experiments to better understand how *pth2* is regulated.

DISCUSSION

Previous work revealed that larval zebrafish use lateral line information to sense the presence of conspecifics, leading to upregulation of the social hormone Pth2. Here, we show that

pth2 levels are chronically depressed in socially-raised larvae born without lateral line function.

However, social isolation further reduces *pth2* levels in lateral line mutants, suggesting that other

socially-derived sensory information can modulate *pth2* expression.

The mechanosensory lateral line mediates *pth2* expression.

Anneser, et al. (2020) implicated hydrodynamic stimuli from conspecifics as a key social

cue that upregulates production of the neuropeptide Pth2. Isolated larvae with chemically ablated

lateral lines fail to upregulate *pth2* when returned to a social environment. In the present study, we

use mutant zebrafish with a congenital loss of lateral line function to confirm that hydrodynamic

information modulates *pth2* expression. Furthermore, we find that repeated chemical ablation of

the lateral line also impairs the ability of social larvae to express *pth2*, even when exposed to

conspecifics prior to loss of lateral line function. Our approach complements that of Anneser et al.

by providing genetic evidence that *pth2* is modulated by lateral line sensory information (Fig. 1).

We conclude that, in terms of Pth2 production, zebrafish born without a lateral line effectively feel

alone even when raised in a social context.

Auditory and vestibular stimuli do not modulate *pth2* expression in larval zebrafish.

Inner ear hair cells are functionally and structurally similar to lateral line hair cells, and

fish use auditory and vestibular sensory information in a variety of behaviors (Bagnall and

Schoppik, 2018; Kohashi et al., 2012; Nicolson et al., 1998; Privat et al., 2019). Given that some

pth2 expression remains in lateral line mutants, we examined whether sensory information from

the inner ear contributes to social modulation of *pth2*. Using zebrafish with mutations that disable

both inner ear and the lateral line hair cells, we find that there is no additional loss of *pth2*

expression compared to the lateral line-only mutant (Fig. 2). In addition, we asked if different types

of mutations to hair cells impact *pth2* expression. Three of the mutant lines used in this study,

lhfpl5b, tomt and *pcdh15a,* render the mechanotransduction (MET) channel non-functional

(Erickson et al., 2017; Erickson et al., 2020; Maeda et al., 2017), and one mutant line, *cacna1da /*

cav1.3a, affects the voltage-gated calcium channel (Sheets et al., 2012; Sidi et al., 2004). While

both components are necessary for proper hair cell signal transduction, there are key differences

between the types of mutations. For mutants affecting the MET channel, potassium and calcium

ions are unable to enter the cell in response to stimuli, but the potential for spontaneous calcium

influx and neurotransmitter release remains (Trapani and Nicolson, 2011). Whereas for the

calcium channel mutation, ions enter through the MET channel but there is no evoked or

spontaneous neurotransmitter release. Regardless, all mutations that disrupt hair cell function

result in the same decrease of *pth2* expression in socially raised fish compared to their wild type

siblings (Fig. 2). Therefore, we conclude that regulation of *pth2* by social context is primarily via

the mechanosensory lateral line in larval zebrafish and that auditory and vestibular information

does not contribute.

Additional socially-derived sensory cues modulate *pth2* expression.

Anneser et al. (2020) demonstrated that neither chemical nor visual cues from conspecifics

significantly restored *pth2* expression in isolated larvae. In our study, we found that isolation

further reduced *pth2* expression in lateral line mutants to the same level as isolated wild types (Fig.

3). This indicates that additional sensory modalities contribute to the regulation of *pth2* expression.

It will be informative to test whether the photo- or chemosensory systems can modulate *pth2*

expression under sensitized circumstances in which hydrosensory information is compromised.

Additionally, it is possible that tactile stimuli are contributing to *pth2* regulation in zebrafish,

consistent with the effects of social touch in rodents. Future work is necessary to test this theory.

Visualizing *pth2* expression *in vivo*.

We used *pth2* promoter-driven fluorescent transgenes to visualize *pth2*-expressing cells

and their projections. Consistent with the Anneser, et al. (2020) whole-mount immunostains, we

see Pth2-positive cells located in bilateral clusters in the thalamus that feature lateral neurites,

anterior ipsilateral projections terminating as a loosely organized neuropil in the telencephalon, as

well as posterior ipsilateral projections. The transgene reveals additional fibers that project

ventrally from the cell bodies and cross the midline, potentially contacting the contralateral *pth2*-

expressing cluster (Fig. 4 and Supplemental Movie 1). Investigating the morphology of these cells

will help us understand the cellular targets of Pth2 signaling and how activity of the two clusters

is coordinated.

We also tested the *pth2:EGFP* and *pth2:TagRFP* transgenic lines as a proxy for *pth2*

expression *in vivo* and found that both lateral line mutants and isolated wild type larvae exhibit

fewer GFP/RFP-positive cells than socially raised siblings (Fig. 5). In contrast, the number of

dopaminergic cells located dorsally to the *pth2* clusters exhibited no change to social context.

Dopamine has been implicated in sensation and social interaction (Haehnel-Taguchi et al., 2018;

Saif et al., 2013; Shams et al., 2018), and our results indicate that the social modulation of *pth2* is

distinct from the dopaminergic system in this brain region in larval zebrafish. While our results

with the *pth2* transgenes agree with our *in situ* hybridization and qPCR results (Fig. 1, 3), we note

that the effects of isolation and lateral line deficiency on transgene expression are not as

pronounced as when assaying for endogenous *pth2* mRNA and protein. It could be that the post-

transcriptional and / or post-translation regulation of fluorescent proteins do not reflect regulation

of endogenous *pth2* transcription and protein stability. It is also possible that the *pth2* promoter

region we use to drive the transgene is missing some critical regulatory elements. Future work will

strive to distinguish between these possibilities and to optimize the transgene for capturing *pth2*

properties in vivo.

Overall, our results confirm that *pth2* expression is modulated by the mechanosensory

lateral line (Fig. 1). We also show that the lateral line is the only hair-cell based sensory organ

regulating *pth2* expression (Fig. 2). However, due to expression levels being slightly higher in

social lateral line mutants compared to the same mutants or wild type larvae raised in isolation, we

conclude that there must be unidentified sensory system(s) that contribute to the modulation of

pth2 (Fig. 3)*.* Finally, we created transgenic lines to visualize the morphology of *pth2+* cells *in*

vivo (Fig. 4), and report that the transgenes respond to social context (Fig. 5). These transgenic

lines can be used in future studies to elucidate the role of *pth2* in zebrafish. Taken together, this

work suggests that zebrafish without a functioning lateral line experience a chronic sense of

isolation, even in a social context.

MATERIALS AND METHODS

Ethics statement

Animal research complied with guidelines stipulated by the Institutional Animal Care and

Use Committees at East Carolina University (Greenville, NC, United States) and at University of

New Brunswick (Fredericton, New Brunswick, Canada). Zebrafish (*Danio rerio*) were maintained

and bred using standard procedures (Westerfield, 2000). All experiments used larvae at 1–5 days

post-fertilization (dpf), which are of indeterminate sex at this stage.

Mutant and transgenic fish lines

The following previously described zebrafish mutant and transgenic lines were used in this study: *lhfpl5b*[vo35], *pcdh15a*[th263b], *tomt*[tk256c], *cacna1da*[tc323d], and *Tg(slc6a3:EGFP)ot80* (Erickson et al., 2017; Erickson et al., 2020; Maeda et al., 2017; Sidi et al., 2004; Xi et al., 2011). To make the *Tg(pth2:EGFP)unb2* and *Tg(pth2:TagRFP)unb3* transgenic lines, 2067 base pairs immediately upstream of the *pth2* coding region (including the 5' UTR and first intron) were PCR amplified with primers containing attB4-B1 recombination sites (*pth2*-specific sequence underlined: Fwd: GGGGACAACTTTGTATAGAAAAGTTG<u>GTAAAGGACACTCTTTGAGATCCT</u>, Rvs: GGGGACTGCTTTTTTGTACAAACTTGC<u>CTGCAGATGAATAAGTTGAATAATACA</u>. The PCR product was cloned into the Gateway pDONR-P4-P1R entry vector by a standard BP clonase reaction and an Multisite LR reaction was performed to create *pth2:EGFP-pA, cryaa:EGFP* and *pth2:tagRFP-pA, cryaa:mCherry* plasmids. pDestTol2pACryGFP and pDESTtol2pACrymCherry plasmids were gifts from Joachim Berger & Peter Currie (Addgene plasmids # 64022, 64023). The expression constructs were co-injected with *tol2* transposase mRNA into 1-cell zebrafish embryos (Kwan et al., 2007). F0 founders were outcrossed to produce stable F1 transgenics. At a minimum, progeny from F2 transgenics were used in this study.

Social isolation experiments for *in situ* hybridization, qPCR, and transgenic lines

Zebrafish larvae were sorted for *lhfpl5b*[vo35], *pcdh15a*[th263b], *tomt*[tk256c] homozygotes at 2 dpf with FM1-43 dye labelling. *cacna1da*[tc323d] homozygotes were identified behaviorally at 4 dpf. *Tg(pth2:EGFP)* and *Tg(pth2:TagRFP)* transgenics were identified by their GFP or mCherry lens markers, respectively. *Tg(slc6a3:EGFP)*-positive larvae were identified by GFP expression in the brain. Identified larvae were divided into social groups of thirty larvae, while another thirty larvae were placed in isolation. Both social and isolated larvae were housed in 6-well tissue culture plates containing 5 ml of E3 embryo media. Strips of paper were placed between each well to prevent visual access between wells. Larvae intended for mRNA *in situ* hybridization (ISH) were grown in E3 with 200 μM N-Phenylthiourea (PTU) (Alfa Aesar) to prevent pigment formation. Each group was incubated at 28.5°C with a 14L:10D photoperiod from 2 to 4 dpf. At 4 dpf, larvae were either fixed for *in situ* hybridization, processed for qPCR, or imaged to visualize the transgene.

Repeated neomycin treatments

The repeated neomycin treatments were performed essentially as previously described
(Venuto & Erickson, 2021). At 24 hpf, WT embryos were divided into two groups of 30 and
grown continuously in media supplemented with 200 µM PTU to prevent pigment formation.
Larvae were manually dechorionated at 2 dpf. Starting at 9:00 AM, 3 dpf larvae were exposed to
either 0 or 50 µM neomycin sulfate hydrate (Alfa Aesar) for 30 minutes at 28ºC every 12 hours.
At noon on 4 dpf, treated and untreated groups of larvae were anesthetized with 0.016% MS-222
and fixed in 4% PFA for mRNA ISH. The experiment was repeated twice using 30 larvae per
treatment group.

mRNA in situ hybridization

The *pth2* in situ probe sequence was amplified using the Invitrogen SuperScript IV One-
Step RT-PCR kit (ThermoFisher) using the following forward and reverse primers: F:
CATTGCATGGACGATTTACG; R: TGCCATGTCATTCAAAATCC. The RT-PCR product
was purified by gel extraction and propagated in the pCR4-TOPO vector. The pCR4-*pth2* plasmid
was linearized with NotI and the antisense DIG-labeled ISH probe was made using the T3 RNA
polymerase promoter. Probe synthesis and in situ hybridization were performed essentially as
described (Erickson et al., 2010). Larvae were imaged at 20X magnification using an Olympus
BH-2 microscope outfitted with a Jenoptik Gryphax Arktur camera.

RNA isolation and real time quantitative PCR (qPCR)

For RNA isolation, 10 whole larval heads were used per replicate. Samples were collected
on ice, placed in 1.7 mL RNase-free microcentrifuge tubes containing 200 µl RNAzol (Molecular
Research Center, Inc., OH. Catalog: RN 190) and homogenized immediately. Total RNA was
extracted from homogenized solutions according to the manufacturer's protocol. For each sample,
cDNAs was synthesized using 1 µg total RNA and a high-capacity cDNA Reverse Transcription
kit (Thermo Fisher Scientific, Waltham, Massachusetts, USA, Catalog#4368814) following the
manufacturer's instructions.

The levels of *pth2* transcripts were determined by quantitative real-time PCR (qPCR) using
SYBR green dye (Invitrogen) and a CFX Connect real-time thermal cycler (Bio-Rad Laboratories,
Hercules, California, USA). The qPCR reaction was conducted with initial denaturation at 95 °C

for 2 min, followed by 30 cycles of 30 s denaturation at 95 °C, 30 s annealing at 57 °C, and 30 s extension at 72 °C using the specific primers (Forward: CCACGCAACACACAGTCAAG, Reverse: GCAAGTTACTTTGCAGAGGTC) and GoTaq G2 DNA polymerase (Promega, Madison, Wisconsin, USA). Each PCR mixture (15 μl) consisted of 7.795 μl DNase free water, 3 μl 5XGoTaq buffer, 1.5 μl 25 mM MgCl2, 0.3 μl 10 mM dNTP mix, 0.15 ml 10μM forward or reverse primer, 2 μl cDNA, 0.03 μl 100X SYBR green dye (final concentration 0.2X), and 0.075 μl Taq. The absolute values (copies/μg total RNA) were determined using Ct values of samples and a standard curve generated from serial known concentrations of plasmid containing the target region of *pth2*. The relative expression was computed by calculating the fold change compared to the mean value of the mutant isolated condition. The efficiency of the PCR and authentic PCR products was further confirmed by analyses of melting curve and gel electrophoresis.

Transgenic imaging and quantification

Live zebrafish larvae were anesthetized with E3 plus 0.016% MS-222 and mounted dorsally on a depression slide in 1.2% low melting point agarose in E3. EGFP was imaged using a Zeiss LSM800 laser-scanning confocal with a 40x water objective and Zeiss Zen software. To quantify results, cell counts were performed using Fiji/ImageJ image processing software (Schindelin et al., 2012). Depth encoded 2D and 3D projections were created in Fiji/ImageJ using the *Z-stack Depth Color Code 0.0.2* plugin. Figures were assembled in Adobe Photoshop and equally adjusted for brightness and contrast among corresponding treatment groups.

Statistical tests

All graphs and statistical tests were done using R (R Core Team, 2021). For *in* situ hybridization cell counts, a one-tailed Welch's t-test was used (two replicates, $n = 7$-9 larvae per condition per replicate). For the qPCR experiment, a one-way ANOVA with a Tukey post-hoc test was conducted (three replicates, $n = 10$ per condition per replicate). For the transgenic cell counts, a one-tailed Welch's t-test was used (three replicates, $n = 5$ per condition per replicate). A p-value of less than 0.05 was considered significant. One-way ANOVAs were used to compare the whole experimental condition (genotype plus social context) as single independent variables. One-tailed t-tests were used due to only having an expectation for a decrease in *pth2* expression following isolation or loss of lateral line sensory information. Sample sizes were determined by the number

of fish used in the experiment. Replicates indicate how many times the experiment was conducted.

No samples were excluded from the analysis.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Yong Zhu for providing his reagents and assistance

with qPCR experiments, and Dr. Bryan Crawford for helpful comments on the manuscript. We

also thank members of the Erickson and Issa labs for their assistance with zebrafish husbandry.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

This work was funded by East Carolina University's Division of Research, Economic

Development and Engagement (REDE) and The Natural Science and Engineering Research

Council (NSERC) of Canada Discovery Grant RGPIN-2021-03166 to T.E.

DATA AVAILABILITY

The datasets generated for this study are available on request to the corresponding author.

REFERENCES

Adkins-Regan, E. (2009). Neuroendocrinology of social behavior. *ILAR Journal* **50**, 5–14.

Amaral, D. G. (2002). The primate amygdala and the neurobiology of social behavior:

implications for understanding social anxiety. *Biological Psychiatry* **51**, 11–17.

Anneser, L., Alcantara, I. C., Gemmer, A., Mirkes, K., Ryu, S. and Schuman, E. M. (2020).

The neuropeptide Pth2 dynamically senses others via mechanosensation. *Nature 2020*

588:7839 **588**, 653–657.

Anneser, L., Gemmer, A., Eilers, T., Alcantara, I. C., Loos, A.-Y., Ryu, S. and Schuman, E.

M. (2022). The neuropeptide Pth2 modulates social behavior and anxiety in zebrafish.

iScience **25**, 103868.

Bagnall, M. W. and Schoppik, D. (2018). Development of vestibular behaviors in zebrafish.

Curr Opin Neurobiol **53**, 83.

Bhattacharya, P., Yan, Y. L., Postlethwait, J. and Rubin, D. A. (2011). Evolution of the
vertebrate pth2 (TIP39) gene family and the regulation of PTH type 2 receptor (pth2r) and
its endogenous ligand pth2 by hedgehog signaling in zebrafish development. *Journal of*
Endocrinology **211**, 187–200.

Brothers, L. (1990). The neural basis of primate social communication. *Motivation and Emotion*
1990 14:2 **14**, 81–91.

Burgdorf, J. and Panksepp, J. (2006). The neurobiology of positive emotions. *Neuroscience &*
Biobehavioral Reviews **30**, 173–187.

Butler, J. M. and Maruska, K. P. (2015). The mechanosensory lateral line is used to assess
opponents and mediate aggressive behaviors during territorial interactions in an African
cichlid fish. *J Exp Biol* **218**, 3284–94.

Butler, J. M. and Maruska, K. P. (2016a). The Mechanosensory Lateral Line System Mediates
Activation of Socially-Relevant Brain Regions during Territorial Interactions. *Frontiers in*
Behavioral Neuroscience **10**, 93.

Butler, J. M. and Maruska, K. P. (2016b). Mechanosensory signaling as a potential mode of
communication during social interactions in fishes. *The Journal of Experimental Biology*
219, 2781–2789.

Chen, P. and Hong, W. (2018). Neural Circuit Mechanisms of Social Behavior. *Neuron* **98**, 16–
30.

Choe, C. P., Choi, S.-Y., Kee, Y., Kim, M. J., Kim, S.-H., Lee, Y., Park, H.-C. and Ro, H.
(2021). Transgenic fluorescent zebrafish lines that have revolutionized biomedical research.
Laboratory Animal Research 2021 37:1 **37**, 1–29.

Coutellier, L. and Usdin, T. B. (2011). Enhanced long-term fear memory and increased anxiety
and depression-like behavior after exposure to an aversive event in mice lacking TIP39
signaling. *Behavioural Brain Research* **222**, 265–269.

Cservenák, M., Szabó, É. R., Bodnár, I., Lékó, A., Palkovits, M., Nagy, G. M., Usdin, T. B.
and Dobolyi, A. (2013). Thalamic neuropeptide mediating the effects of nursing on
lactation and maternal motivation. *Psychoneuroendocrinology* **38**, 3070–3084.

Dijkgraaf, S. (1963). The functioning and significance of the lateral-line organs. *Biological*
Reviews **38**, 51–105.

Dimitrov, E. and Usdin, T. B. (2010). Tuberoinfundibular peptide of 39 residues modulates the mouse hypothalamic-pituitary-adrenal axis via paraventricular glutamatergic neurons. *J Comp Neurol* **518**, 4375–4394.

Dimitrov, E. L., Petrus, E. and Usdin, T. B. (2010). Tuberoinfundibular peptide of 39 residues (TIP39) signaling modulates acute and tonic nociception. *Experimental Neurology* **226**, 68–83.

Dimitrov, E. L., Kuo, J., Kohno, K. and Usdin, T. B. (2013). Neuropathic and inflammatory pain are modulated by tuberoinfundibular peptide of 39 residues. *Proceedings of the National Academy of Sciences* **110**, 13156–13161.

Dobolyi, A., Dimitrov, E., Palkovits, M. and Usdin, T. B. (2012). The Neuroendocrine Functions of the Parathyroid Hormone 2 Receptor. *Frontiers in Endocrinology* **3**, 121.

Erickson, T., French, C. R. and Waskiewicz, A. J. (2010). Meis1 specifies positional information in the retina and tectum to organize the zebrafish visual system. *Neural Development* **5**, 22.

Erickson, T., Morgan, C. P., Olt, J., Hardy, K., Busch-Nentwich, E., Maeda, R., Clemens, R., Krey, J. F., Nechiporuk, A., Barr-Gillespie, P. G., et al. (2017). Integration of Tmc1/2 into the mechanotransduction complex in zebrafish hair cells is regulated by Transmembrane O-methyltransferase (Tomt). *Elife* **6**,.

Erickson, T., Pacentine, I. v., Venuto, A., Clemens, R. and Nicolson, T. (2020). The lhfpl5 Ohnologs lhfpl5a and lhfpl5b Are Required for Mechanotransduction in Distinct Populations of Sensory Hair Cells in Zebrafish. *Frontiers in Molecular Neuroscience* **12**, 320.

Groneberg, A. H., Marques, J. C., Martins, A. L., Diez del Corral, R., de Polavieja, G. G. and Orger, M. B. (2020). Early-Life Social Experience Shapes Social Avoidance Reactions in Larval Zebrafish. *Current Biology* **30**, 4009-4021.e4.

Haehnel-Taguchi, M., Fernandes, A. M., Böhler, M., Schmitt, I., Tittel, L. and Driever, W. (2018). Projections of the diencephalospinal dopaminergic system to peripheral sense organs in larval zebrafish (Danio rerio). *Frontiers in Neuroanatomy* **12**, 20.

Keller, D., Láng, T., Cservenák, M., Puska, G., Barna, J., Csillag, V., Farkas, I., Zelena, D., Dóra, F., Barteczko, L., et al. (2022). A thalamo-preoptic pathway promoting social touch. *bioRxiv* 2022.01.11.475648.

Kohashi, T., Nakata, N. and Oda, Y. (2012). Effective Sensory Modality Activating an Escape
Triggering Neuron Switches during Early Development in Zebrafish. *The Journal of*
Neuroscience **32**, 5810.

Kolb, B. and Whishaw, I. Q. (2003). Brain Plasticity and Behavior.
http://dx.doi.org/10.1146/annurev.psych.49.1.43 **49**, 43–64.

Kravitz, E. A. and Huber, R. (2003). Aggression in invertebrates. *Current Opinion in*
Neurobiology **13**, 736–743.

Kwan, K. M., Fujimoto, E., Grabher, C., Mangum, B. D., Hardy, M. E., Campbell, D. S.,
Parant, J. M., Yost, H. J., Kanki, J. P. and Chien, C.-B. (2007). The Tol2kit: a multisite
gateway-based construction kit for Tol2 transposon transgenesis constructs. *Dev Dyn* **236**,
3088–99.

LaBuda, C. J. and Usdin, T. B. (2004). Tuberoinfundibular peptide of 39 residues decreases
pain-related affective behavior. *NeuroReport* **15**, 1779–1782.

LaBuda, C. J., Dobolyi, A. and Usdin, T. B. (2004). Tuberoinfundibular peptide of 39 residues
produces anxiolytic and antidepressant actions. *Neuroreport* **15**, 881–885.

Maeda, R., Pacentine, I. v., Erickson, T. and Nicolson, T. (2017). Functional Analysis of the
Transmembrane and Cytoplasmic Domains of Pcdh15a in Zebrafish Hair Cells. *The Journal*
of Neuroscience **37**, 3231–3245.

Montgomery, J., Bleckmann, H. and Coombs, S. (2013). Sensory Ecology and Neuroethology
of the Lateral Line. 121–150.

Mueller, T. (2012). What is the Thalamus in Zebrafish? *Frontiers in Neuroscience* **6**, 64.

Mumtaz, F., Khan, M. I., Zubair, M. and Dehpour, A. R. (2018). Neurobiology and
consequences of social isolation stress in animal model—A comprehensive review.
Biomedicine & Pharmacotherapy **105**, 1205–1222.

Neumann, I. D., Veenema, A. H. and Beiderbeck, D. I. (2010). Aggression and anxiety: Social
context and neurobiological links. *Frontiers in Behavioral Neuroscience* **4**, 12.

Nicolson, T., Rüsch, A., Friedrich, R. W., Granato, M., Ruppersberg, J. P. and Nüsslein-
Volhard, C. (1998). Genetic Analysis of Vertebrate Sensory Hair Cell Mechanosensation:
the Zebrafish Circler Mutants. *Neuron* **20**, 271–283.

O'Connell, L. A. and Hofmann, H. A. (2011). Genes, hormones, and circuits: An integrative

approach to study the evolution of social behavior. *Frontiers in Neuroendocrinology* **32**,

320–335.

Oldfield, R. G. and Hofmann, H. A. (2011). Neuropeptide regulation of social behavior in a

monogamous cichlid fish. *Physiology & Behavior* **102**, 296–303.

Panksepp, J. and Burgdorf, J. (2003). "Laughing" rats and the evolutionary antecedents of

human joy? *Physiology & Behavior* **79**, 533–547.

Panksepp, J., Burgdorf, J., Beinfeld, M. C., Kroes, R. A. and Moskal, J. R. (2007). Brain

Regional Neuropeptide Changes Resulting From Social Defeat. *Behavioral Neuroscience*

121, 1364–1371.

Privat, M., Romano, S. A., Pietri, T., Jouary, A., Boulanger-Weill, J., Elbaz, N., Duchemin,

A., Soares, D. and Sumbre, G. (2019). Sensorimotor Transformations in the Zebrafish

Auditory System. *Current Biology* **29**, 4010-4023.e4.

R Core Team (2021). R: A Language and Environment for Statistical Computing.

Robinson, G. E., Fernald, R. D. and Clayton, D. F. (2008). Genes and social behavior. *Science*

322, 896–900.

Saif, M., Chatterjee, D., Buske, C. and Gerlai, R. (2013). Sight of conspecific images induces

changes in neurochemistry in zebrafish. *Behavioural Brain Research* **243**, 294–299.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source

platform for biological-image analysis. *Nature Methods 2012 9:7* **9**, 676–682.

Schoofs, L., de Loof, A. and van Hiel, M. B. (2017). Neuropeptides as Regulators of Behavior

in Insects. *http://dx.doi.org/10.1146/annurev-ento-031616-035500* **62**, 35–52.

Seitz, N. (2019). Characterization Of Pth Type-2 Receptor (pth2rb) And Its Endogenous Ligand

Pth2 in Zebrafish Development. *Theses and Dissertations* 1131.

https://ir.library.illinoisstate.edu/etd/1131.

Shams, S., Seguin, D., Facciol, A., Chatterjee, D. and Gerlai, R. (2017). Effect of social

isolation on anxiety-related behaviors, cortisol, and monoamines in adult zebrafish.

Behavioral Neuroscience **131**, 492–504.

Shams, S., Amlani, S., Buske, C., Chatterjee, D. and Gerlai, R. (2018). Developmental social

isolation affects adult behavior, social interaction, and dopamine metabolite levels in

zebrafish. *Developmental Psychobiology* **60**, 43–56.

Sheets, L., Kindt, K. S. and Nicolson, T. (2012). Presynaptic CaV1.3 Channels Regulate

Synaptic Ribbon Size and Are Required for Synaptic Maintenance in Sensory Hair Cells.

The Journal of Neuroscience **32**, 17273.

Sidi, S., Busch-Nentwich, E., Friedrich, R., Schoenberger, U. and Nicolson, T. (2004).

gemini Encodes a Zebrafish L-Type Calcium Channel That Localizes at Sensory Hair Cell

Ribbon Synapses. *The Journal of Neuroscience* **24**, 4213.

Soboleski, M. R., Oaks, J. and Halford, W. P. (2005). Green fluorescent protein is a

quantitative reporter of gene expression in individual eukaryotic cells. *The FASEB journal :*

official publication of the Federation of American Societies for Experimental Biology **19**,

440.

Trapani, J. G. and Nicolson, T. (2011). Mechanism of Spontaneous Activity in Afferent

Neurons of the Zebrafish Lateral-Line Organ. *The Journal of Neuroscience* **31**, 1614.

Venuto, A. and Erickson, T. (2021). Evaluating the Death and Recovery of Lateral Line Hair

Cells Following Repeated Neomycin Treatments. *Life* **11**, 1180.

Weiss, I. C., Pryce, C. R., Jongen-Rêlo, A. L., Nanz-Bahr, N. I. and Feldon, J. (2004). Effect

of social isolation on stress-related behavioural and neuroendocrine state in the rat.

Behavioural Brain Research **152**, 279–295.

Xi, Y., Yu, M., Godoy, R., Hatch, G., Poitras, L. and Ekker, M. (2011). Transgenic zebrafish

expressing green fluorescent protein in dopaminergic neurons of the ventral diencephalon.

Developmental Dynamics **240**, 2539–2547.

 FIGURES AND LEGENDS:

Figure 1: Decreased *pth2* expression in lateral line-deficient larval zebrafish. A: Location of *pth2*-expressing cells in the thalamus. B-C: Representative images of mRNA *in situ* hybridizations (ISH) showing *pth2*-positive cells in socially-reared B) wild type and C) lateral line mutant (*lhfpl5b^{vo35}*) zebrafish at 4 dpf. D) Boxplots of *pth2*-positive cell counts from ISH images (WT = 14, MUT = 18 larvae from two replicates). E) Representative mRNA ISH image showing reduced *pth2* expression in a socially-reared larva at 4 dpf whose lateral line was ablated with repeated neomycin treatments. Scale bar is 50 μm and applies to all images. A one-tailed Welch's t-test was used to assess significance, **** = p < 0.0001.

Figure 2: Mechanosensory input from inner ear hair cells does not contribute to *pth2*
expression. A: Location of *pth2*-expressing cells in the thalamus. B-F: Representative images of
mRNA *in situ* hybridizations (ISH) showing *pth2*-positive cells in socially-reared B) wild type, C)
lhfpl5b^{vo35}, D) *tomt^{tk256c}*, E) *pcdh15a^{th263b}*, and F) *cacna1da^{tc323d}* larvae at 4 dpf. Scale bar = 50 μm,
applies to all images. G: Boxplots of qPCR results showing *pth2* expression in socially-reared
mutants and siblings. Three replicates per condition, n = 10 larvae per condition per replicate. One
way ANOVA with Tukey post-test to assess significance, *** = p < 0.001, ** = p < 0.01.

Figure 3: Social isolation further reduces *pth2* expression in sensory hair cell mutants. A-C:

mRNA *in situ* hybridization showing representative examples of *pth2*-positive cells in A) socially-

reared *lhfpl5b^vo35^* mutants, B) isolated wild type larvae, and C) isolated *lhfpl5b^vo35^* mutants. Scale

bar = 50 μm, applies to all images. D: Boxplots of qPCR results comparing *pth2* expression in

socially-reared *lhfpl5b^vo35^* and *tomt^tk256c^* mutants relative to isolated mutant and wild type siblings.

Three replicates per condition, n = 10 larvae per condition per replicate. One way ANOVA with

Tukey post-test to assess significance, *** = p < 0.001, * = p < 0.05, ns = not significant.

Figure 4: Morphology of *pth2*-expressing cells. A: Representative depth-encoded image of the

Tg(pth2:TagRFP)unb3 transgenic line at 4 dpf. The anterior-posterior (A-P) axis and dorsal-

ventral (D-V) depth color scale are indicated at the bottom. B: Black and white version of the

image in A. From the top: the dashed box indicates the anterior neuropil, the starred bracket

highlights the ventral neurites that cross the midline, and the bilateral open arrows point to

posterior neurites. Scale bar in B is 50 µm, applies to both images. See Supplemental Movie 1 for

a 3D rendering of the depth-encoded *pth2* transgene.

Figure 5: Effect of lateral line loss of function and social isolation on *pth2* and *slc6a3* expression using fluorescent reporters. A-B: Representative images of *Tg(pth2:EGFP)unb2* transgenic expression in socially-reared A) wild type and B) lateral line mutant (*lhfpl5b^{vo35}*) zebrafish. C) Boxplots of GFP-positive cell counts in wild type (WT) and *lhfpl5b* mutant (MUT) siblings. D-E: Representative images of *Tg(pth2:TagRFP)unb3* (false-colored green, top) and *Tg(slc6a3:EGFP)ot80* (false-colored magenta, middle) transgenes in D) socially-reared and E) isolated larvae. Merges of the two channels are shown at the bottom. F) Boxplots of *pth2* and *slc6a3* cell counts. Scale bar is 50 μm. A one-tailed Welch's t-test was used to assess significance, **** = p < 0.0001, ns = not significant.