1 **Vertical gradients in photosynthetic physiology diverge at the latitudinal range extremes of**
2 **white spruce**
3

Stephanie C. Schmiege[1,2*], Kevin L. Griffin[1,3,4], Natalie T. Boelman[4], Lee A. Vierling[5,6], Sarah

G. Bruner[1], Elizabeth Min[3], Andrew J. Maguire[5,6], Johanna Jensen[1], Jan U. H. Eitel[5,6]

[1]Department of Ecology, Evolution and Environmental Biology, Columbia University, New

York, NY 10027, USA

[2] New York Botanical Garden, Bronx, NY 10458

[3]Department of Earth and Environmental Sciences, Columbia University, Palisades, NY 10964,

USA

[4]Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA

[5]Department of Natural Resources and Society, College of Natural Resources, University of

Idaho, Moscow, ID 83843, USA

[6]McCall Outdoor Science School, College of Natural Resources, University of Idaho, McCall, ID

83638, USA

*Current affiliations: Plant Resilience Institute, Michigan State University, East Lansing, MI

48824 USA & Department of Biology, Western University, London, Ontario, Canada N6A 5B7

CORRESPONDING AUTHOR: Stephanie C. Schmiege

Email address: schmie18@msu.edu

Telephone: 646-234-7728

ABSTRACT (200 words)

Light availability drives vertical canopy gradients in photosynthetic functioning and carbon (C)

balance, yet patterns of variability in these gradients remain unclear. We measured light

availability, photosynthetic CO_2 and light response curves, foliar C, nitrogen (N) and pigment

concentrations, and the photochemical reflectance index (PRI) on upper and lower canopy

needles of white spruce trees (*Picea glauca*) at the species' northern and southern range

extremes. We combined our photosynthetic data with previously published respiratory data to

compare and contrast canopy C balance between latitudinal extremes. We found steep canopy

gradients in irradiance, photosynthesis, and leaf traits at the southern range limit, but a lack of

variation across canopy positions at the northern range limit. Thus, unlike many tree species

from tropical to mid-latitude forests, high latitude trees may not require vertical gradients of

metabolic activity to optimize photosynthetic C gain. Consequently, accounting for self-shading

is less critical for predicting gross primary productivity at northern relative to southern latitudes.

Northern trees also had a significantly smaller net positive leaf C balance than southern trees

suggesting that, regardless of canopy position, low photosynthetic rates coupled with high

respiratory costs may ultimately constrain the northern range limit of this widely distributed

boreal species.

SHORT TITLE: Variation in photosynthetic canopy gradients of white spruce at the species'

range limits.

KEY WORDS: *Picea glauca*, canopy gradients, carbon balance, photosynthesis, photochemical

reflectance index, Arctic treeline

SUMMARY STATEMENT

Canopy gradients in photosynthetic capacity of white spruce diminish at high compared to low

latitudes. Low carbon balance in high latitude trees may determine the extent of northern

treeline.

INTRODUCTION

Illuminating the factors driving species' distributional ranges, i.e., the boundaries within which a species can grow and reproduce successfully, remains a fundamental ecological, evolutionary, and biogeographic challenge. Extensive research has uncovered a number of factors that play important roles. These include environmental tolerance, niche breadth, metapopulation dynamics, genetic diversity, phenotypic plasticity and dispersal ability (Brown, Stevens & Kaufmnan 1996; Gaston 1996; Lowry & Lester 2006). The diversity of possible factors highlights that no universal cause of species' distributions has yet emerged, but rather that a combination of historical, physiological, and biotic filters may define a species' distributional range (Lambers & Oliveira 2019). Nevertheless, the ability of a plant to tolerate the local growth environment remains paramount. This ability is intimately linked to a species' physiological performance. The physiological processes of photosynthesis and respiration, resulting in the exchange of carbon, are heavily affected by environmental conditions. Together, these two processes determine the carbon balance of a plant which must remain positive if a plant is to grow, survive and reproduce in a given environment. Thus, photosynthesis and respiration fundamentally act as a physiological filter determining a species' distributional range (Lambers & Oliveira 2019). They are also critical to quantifying carbon exchange as well as the ecological effects of climate change on future carbon sequestration.

The evolution from modeling vegetation as a single "big leaf" to "two big leaves" including distinct parameters for sun and shade leaves, to multi-layer canopies including gradients of physiological function (Field 1983; Sands 1995) has improved estimates of vegetative carbon gain in many ecosystems (Harley & Baldocchi 1995; de Pury & Farquhar 1997; Bonan, Oleson, Fisher, Lasslop & Reichstein 2012; Rogers *et al.* 2017). These multi-layer canopy models are parameterized using physiological and biochemical traits that vary predictably with vertical gradients in environmental conditions such as light, temperature, humidity and vapor pressure deficit throughout the tree crown (Sellers, Berry, Collatz, Field & Hall 1992; Bond, Farnsworth, Coulombe & Winner 1999). Of particular importance to modeling photosynthesis are gradients in light availability. For example, traits such as maximum photosynthesis (A_{max}) and leaf nitrogen content, as well as leaf mass per area (LMA) have been found to correlate positively with changes in irradiance (Field 1983; Hirose & Werger 1987; Ellsworth & Reich 1993; Niinemets, Kull & Tenhunen 1998; Niinemets 2007) leading to the

common finding that high canopy, high light foliage has higher photosynthetic capacity.
Pigment pools have similarly been found to vary with canopy position and irradiance with high
light foliage frequently containing larger pools of photoprotective pigments that can dissipate
excess light energy and protect chlorophyll-filled photosystems (Demmig-Adams & Adams III
1992). Even the ratio of Chl$a$:Chl$b$ frequently reflects canopy gradients of light availability
because shade-adapted leaves contain more light harvesting antenna complexes (LHCII)
associated with photosystem II that are high in Chl$b$ (Dale & Causton 1992; Hikosaka &
Terashima 1995; Kitajima & Hogan 2003; Ruban 2015; Magney *et al.* 2016; Scartazza, Di
Baccio, Bertolotto, Gavrichkova & Matteucci 2016). Thus, the incorporation of canopy
gradients into carbon exchange modeling comes from a strong foundation of work that
demonstrates how differences in photosynthetic functioning and resource allocation depend on
canopy position and light availability.
Other environmental factors may also play a role in determining canopy gradients in
photosynthesis. For example, photosynthesis is not only dependent on light availability, but also
on water transport to leaves since high photosynthetic rates tend to require high rates of stomatal
conductance and transpiration (Wong, Cowan & Farquhar 1978; Schulze, Kelliher, Körner,
Lloyd & Leuning 1994). Thus, limitations in water availability may constrain vertical canopy
gradients even if light is available in abundance, resulting in shallower canopy gradients in
photosynthesis and leaf nitrogen (Peltoniemi, Duursma & Medlyn 2012). Therefore, we cannot
assume that light availability is the only driver of canopy gradients without quantifying its effect
on photosynthesis, especially in species whose ranges span large latitudinal differences over
which environmental conditions can vary dramatically. In this study we examine the strength of
the relationships between light availability and canopy gradients of physiological traits across the
range extremes of an important boreal species.
White spruce (*Picea glauca* (Moench) Voss) has a massive, transcontinental range
distribution stretching from the Forest Tundra Ecotone (FTE) in northern Alaska where it is one
of the dominant species, to mixed forests on the eastern seaboard of Canada and New England
(Viereck, Van Cleve & Dyrness 1986; Little Jr. 1999; Eitel *et al.* 2019). At the northern edge of
the distribution, white spruce are exposed to unique environmental challenges for plant growth
and survival including low winter temperatures, shallow soil depth underlain by permafrost, and
brief growing seasons. During the growing season, white spruce canopies are exposed to a

unique light environment caused by high solar zenith angles and continuous photoperiod (i.e. 24

hours of light for several weeks). Furthermore, tree crown structure at the FTE exhibits an

almost ubiquitous tall, vertical, narrow and untapering crown shape (Kuuluvainen 1992) which

creates an open canopy forest structure with widely spaced trees. By contrast, white spruce at

the southern range extreme are exposed to much lower solar zenith angles and a shorter

photoperiod during the growing season. The southern tree crowns are also generally much

denser, wider and more tapered, and typically grow with a higher stand density than in the north.

These contrasting light environments may impact canopy gradients of light interception, which

may, in turn, drive canopy gradients of photosynthetic physiology, biochemistry and pigment

concentrations.

In addition to photosynthetic, biochemical and pigment traits, we also examine canopy

differences in the photochemical reflectance index (PRI). This index is a more successful

photosynthetic proxy than the commonly used normalized difference vegetation index (NDVI) in

conifer species because it is sensitive to changes in pigment pools such as the ratio of

chlorophylls to carotenoids (Gamon *et al.* 2016). PRI has been successfully employed to track

seasonal changes in photosynthetic phenology in coniferous forests (Wong & Gamon 2015a, b;

Gamon *et al.* 2016) and at the FTE (Eitel *et al.* 2019, 2020), and a broadband analog Chlorophyll

Carotenoid Index (CCI), can be derived from global Moderate Resolution Imaging

Spectroradiometer (MODIS) imagery (Gamon *et al.* 2016). Thus, PRI presents a promising

prospective avenue for further understanding carbon dynamics across broadly distributed conifer

forests.

In this study we focus specifically on the photosynthetic traits of white spruce and their

relationship to light environment. However, to gain a full picture of the physiological limits at

latitudinally distant range extremes, a carbon balance approach should be employed. In fact,

recent work on the physiology of white spruce has focused specifically on the respiratory cost

associated with growth and survival at the FTE. In a previous study, we found that white spruce

in this environment have extremely high respiration compared to white spruce growing at the

southernmost range extreme as well as higher respiration than the average respiration of the

boreal biome and needle-leaved evergreen trees (Griffin *et al.* 2021). Griffin et al. (2021) posit

that this high respiratory cost may constrain the northern limit of white spruce and the Arctic

treeline in general. With the addition of photosynthetic traits measured on the same trees, we are

now in a unique position to examine the degree to which this respiratory cost is matched by

photosynthetic carbon gains and whether a lower carbon balance may contribute to the northern

limits of white spruce's latitudinal range.

Thus, the overarching aims of this research are twofold: first, to provide a comprehensive

assessment of the photosynthetic physiology of white spruce and its relationship to canopy and

latitudinal light environment at its range extremes; and second, to examine whether the leaf

carbon balance of this species constrains its northernmost range limit. First, we hypothesize that

foliar light availability, physiological traits, biochemical traits and PRI will converge across high

and low vertical canopy positions in white spruce growing at the FTE. In contrast, steep canopy

gradients of light availability and all foliar traits will be apparent at the southernmost range

extreme. Second, we hypothesize that the carbon balance of white spruce at the northern

location will be much lower than the southern location reflecting the challenging growth

environment and providing a physiological mechanism for the northernmost range extreme of

this important boreal species.

MATERIALS AND METHODS

Site Descriptions: This study was conducted in two locations at the northern and southern range

extremes of white spruce (*Picea glauca* (Moench) Voss). In the north, data were collected from

six sites chosen to represent the FTE (Eitel *et al.* 2019). These sites are located along a 5.5 km

long stretch of the Dalton Highway, in the Brooks Range in northern Alaska (67°59'40.92" N

latitude, 149°45'15.84" W longitude). At the FTE, white spruce is the dominant tree species

with deciduous shrubs and sedges present in the understory (Eitel *et al.* 2019). The forest

structure is relatively open with approximately 407 stems ha^{-1} (Jensen pers. comm.). Three white

spruce study trees (with DBH greater than 10cm) were chosen at each site for a total of 18 study

trees at the FTE (tree characteristics are described in Table 1). Mean annual precipitation was

485.4 mm and mean annual temperature was -8.1°C as determined from a SNOTEL site at the

nearby Atigun Pass (https://wcc.sc.egov.usda.gov/nwcc/site?sitenum=957) between 2007 and

2016. During the Alaska measurement campaign in July 2017, photoperiod ranged from 22 to

24 hours. All measurements were taken at a high and low south-facing canopy position on each

of 18 trees for a total of 36 measurements. The low canopy position was at approximately 1.37

m (diameter at breast height), and the high canopy position was approximately 1m below the
apical meristem.

At the southern edge of the species' range, data were collected from six trees located in
Black Rock Forest, New York (BRF; 41°24'03.91" N latitude, 74°01'28.49" W longitude; Table
1). Black Rock Forest is an oak-dominated forest with approximately 760 stems ha^{-1} (Whitehead
et al. 2004). Mean annual precipitation from 1982 – 2010 was 1285 mm and mean annual
temperature was 10.9°C (Arguez *et al.* 2012). During our BRF measurement campaign in June
2017, photoperiod was approximately 15 hours. BRF is an oak-dominated northern temperate
deciduous forest (Schuster *et al.* 2008; Patterson *et al.* 2018). Measurements were also taken at
high and low south-facing canopy positions on each tree. Due to the lower count of available
trees, three measurements were taken at the low and high canopy positions on each of the six
trees for a total of 36 measurements. As at the FTE, the low canopy position was at
approximately 1.37 m and the high canopy position was approximately 1m below the apical
meristem.

Environmental Measurements: Hemispherical photographs were taken with a digital camera
(CoolPix 4500, Nikon Corporation, Tokyo Japan) and an attached fisheye lens to assess canopy
openness at the high and low canopy positions chosen for each tree (described above). The
camera was positioned immediately adjacent to the branch chosen for measurements, with the
camera pointing north. Once the camera was level, the hemispherical canopy photo was taken.
Using these photographs, light environment was assessed and modeled using the free R code and
documentation of ter Steege (2018). These scripts are based on the widely used Hemiphot and
Winphot programs (ter Steege 1993, 1997). Modeling of the light environment using this code
takes into account GPS location as well as the measurement dates in order to correct for
differences in solar angle (see ter Steege (2018) for a fuller description). Average Photosynthetic
Photon Flux Density (PPFD ; mol m^{-2} day^{-1}) at the high and low canopy positions was
calculated by averaging the diurnal light courses over the measurement campaign days (June 21
– June 27 at BRF, and July 4 – July 20 at the FTE). At the FTE, air temperature (°C) was
collected using a meteorological sensor (VP-4, METER, Pullman, WA) and local solar open sky
radiation readings were collected using a PYR Solar Radiation Sensor (METER, Pullman, WA).
Solar radiation (W m^{-2}) was converted to PPFD (μmol m^{-2} s^{-1}) by multiplying by 2.02 (Mavi &

Tupper 2004). At BRF, air temperature and local open sky PPFD were collected from a nearby

BRF-run weather station (https://blackrock.ccnmtl.columbia.edu/portal/weather/) equipped with

a temperature/humidity sensor (HC2S3, Rotronic, Hauppauge, NY) and a quantum sensor (LI-

190SB Quantum Sensor, LICOR, Lincoln, NE).

Gas Exchange Measurements: Gas exchange was measured at high and low canopy positions on

each tree at the two locations. At both locations, measurements were taken on a branch tip

inserted into the cuvette. At BRF, all measurements were taken on the tree with high canopy

reached using an articulating boom lift (600AJ, JLG, McConnellsburg, PA). At the FTE, low

canopy branches were measured on the tree, but high canopy branches had to be detached using

a pole-clippers (Fiskars fiberglass pole pruner, Fiskars Brands Inc, Middleton, WI) as there was

no way to reach samples otherwise. Detaching foliage for physiological measurements is

common, with no observable differences usually found between attached and detached samples

(Akalusi, Meng & Bourque 2021). For detached samples, cut branches at least 30 cm in length

were placed immediately in water, and recut underwater. All measurements on cut branches

were taken within 12 hours. CO_2 (A-C_i) and light response (A-Q) curves were collected using

two portable photosynthesis systems (LI-6800, LiCor, Lincoln, NE). Needles were sealed in a

cuvette in which the temperature and relative humidity were set to 20°C and 50% humidity at the

FTE, and 25°C, and 60% humidity at BRF in order to mirror ambient conditions in each location.

For the A-C_i curves, irradiance was kept constant at 1000 PPFD and the CO_2 concentrations were

stepped through the following concentrations (ppm): 1200, 1000, 800, 600, 400, 300, 200, 100,

75, 50, 25, 0. Values of V_{cmax} (the maximum rate of RuBisCO carboxylation) and J_{max} (the

maximum rate of electron transport) were determined according to the photosynthesis model of

Farquhar, von Caemmerer and Berry (Farquhar, von Caemmerer & Berry 1980) and adjusted to

25°C using the plantecophys package in R (Duursma 2015).

For the A-Q curves, reference CO_2 concentration was set to 400 ppm and photosynthesis

was measured at the light intensities: 1800, 1500, 1200, 1000, 800, 400, 200, 150, 100, 90, 80,

75, 70, 65, 60, 55, 50, 45, 40, 35, 30, 25, 20, 15, 10, 5, 0 μmol m^{-2} s^{-1}. After 15 minutes at zero

irradiance, respiration in the dark (R_D) was recorded. Maximum likelihood estimation (executed

in the bblme package for R (Bolker & R Development Core Team 2020)) was used to fit the

following model (describing a non-rectangular hyperbola) to the data (Herrmann, Schwartz &

Johnson 2020):

$$A = \frac{\Phi * I + A_{max} - \sqrt{(\Phi * I + A_{max})^2 - 4 * \theta * \Phi * I * A_{max}}}{2 * \theta} - R_D$$

Where A is the measured rate of photosynthesis, θ is the curvature factor and I is the irradiance.

Parameters extracted and calculated from the fit of the model to each light curve included the

maximum rate of CO_2 fixation (A_{max}), photosynthesis at 1500 $\mu mol\ m^{-2}\ s^{-1}$ (A_{1500}), the apparent

quantum yield (Φ), the light compensation point (LCP – the light level required for zero net

carbon flux) and the light saturation point (LSP – the light level required to obtain 75% of A_{1500}).

Respiration in the Light: Respiration in the light (R_L) was obtained from the A-Q curves

according to the methods of Kok (1948); Sharp, Matthews & Boyer (1984); and Heskel, Atkin,

Turnbull & Griffin (2013). These curves contain a subtle change in slope near the light

compensation point (the light intensity where photosynthesis and respiration equal each other).

The break point where this change in slope occurs divides the photosynthetic response to light

into two linear sections. Extrapolating from the upper linear portion of the curve to the y-axis

highlights an alternate respiration value which was interpreted by Bessel Kok to be the amount

of respiration occurring in the light (Kok 1948; Heskel *et al.* 2013; Tcherkez *et al.* 2017).

An important assumption of the Kok method is that carbon assimilation is only

influenced by light. Consequently, corrections were made to account for any changes in internal

CO_2 concentration (C_i) following the methods of Kirschbaum & Farquhar (1987) (see Ayub *et*

al. (2011) for a full description). It is possible that R_L measured with the Kok method may be

influenced by factors such as mesophyll conductance (Farquhar & Busch, 2017; *n.b.* Buckley *et*

al., 2017); however, such investigations are beyond the scope of this study.

Leaf Trait Measurements: After collecting A-C_i and A-Q curves, needles were detached from the

branch, photographed to determine projected surface area (cm^2) using ImageJ (Schnieder,

Rasband & Eliceiri 2012), placed into coin envelopes and oven dried at 60°C for 48 hours.

Leaves were reweighed post-drying for leaf dry mass (g). Specific leaf area (SLA; cm^2g^{-1}) was

calculated. Leaf samples were then ground using a ball mill (SPEX 8000 Mixer/Mill, Metuchen,

NJ, USA) and prepared for analysis of % carbon and % nitrogen using a carbon-nitrogen flash

analyzer (CE Elantech, Lakewood, NJ, USA). Using these percentages, foliar nitrogen per area

(N_{area}; mg cm^{-2}) and the carbon to nitrogen ratio (C:N) were calculated.

Pigment measurements: Needles growing on branches adjacent to those selected for gas

exchange at both high and low canopy positions were selected for pigment analysis. Branches

were wrapped in wet paper towels and foil, and transported back to the laboratory in a cooler

with ice. Foliar pigment extraction began no later than three hours after returning from the field,

and any samples not processed in this time were frozen until pigment extraction could take place.

Between 5 and 10 needles were removed from each branch and their wet weight and

needle area measured. Needles were ground in 100% acetone until no identifiable green needle

tissue remained. A small amount of sand, $MgCO_3$ and ascorbic acid were added to help with

grinding and to prevent acidification and pigment degradation. The extracts were then

centrifuged and diluted. Using a spectrophotometer (Go Direct SpectroVis Plus

Spectrophotometer, Vernier, OR, USA), the absorbance of the supernatant was measured at 470,

645, 662 and 710 nm. Pigment pools were calculated on a leaf area basis including chlorophyll *a*

($Chla_{area}$), chlorophyll *b* ($Chlb_{area}$), and bulk carotenoids (Car_{area}; ug/cm^2 for all pools) were

calculated according to Lichtenthaler (1987). The total chlorophyll pool size ($Chl_{area} = Chla_{area} +$

$Chlb_{area}$) and ratios $Chla{:}Chlb_{area}$ and $Chl{:}Car_{area}$ were also calculated.

Spectral reflectance: PRI: Needle spectral reflectance was measured with a spectroradiometer

(UniSpec SC, PP Systems, Haverhill MA, USA) with an attached fiber optic probe (UNI400, PP

Systems, Haverhill MA, USA) and specialized needle clip (UNI501, PP Systems, Haverhill MA,

USA). The spectroradiometer allowed sampling of the spectrum between 310 and 1100 nm with

a 3.3 nm sampling interval which was afterwards reduced to 1 nm using linear interpolation.

After collecting dark and white standard (Spectralon, LabSphere, North Sutton, NY, USA)

reference measurements, three spectral measurements were made on each branch to capture

spatial heterogeneity. Reflectance values were calculated from the spectral measurements by

dividing each foliage measurement by the measurement from the white standard. PRI was

calculated according to the equation: $(R531 - R570)/(R531 + R570)$ (Gamon, Penuelas &

Field 1992; Penuelas, Filella & Gamon 1995) where R531 and R570 are the reflectances at 531

nm and 570 nm, respectively.

Statistical analyses: Relationships between measured foliar parameters (physiology, pigments

and leaf traits), and integrated daily PPFD and location (BRF or the FTE) were assessed using

linear mixed effects models. The interaction between PPFD and location was included so as to

examine the variations in slope between locations. Individual tree (FTE n = 18, BRF n = 6) was

included as a random effect to account for repeat measures on trees (Parameter ~ PPFD *

location + (1|tree)). If necessary, parameters were log-transformed to meet assumptions of

normality of the residuals. Parameter estimates were considered significantly different from zero

if $P<0.05$. Linear mixed effects analysis was conducted using the lme4 package (Bates,

Maechler, Bolker & Walker 2015) with the lmerTest package loaded to provide p-values

(Kuznetsova, Brockhoff & Christensen 2017). All analyses took place in R v. 4.1.0 (R Core

Team 2022).

The effects of canopy position and location on foliar traits were also assessed using

mixed effects models implemented in the lme4 package (Bates *et al.* 2015). Again, individual

tree was included as a random effect to account for repeat measures on trees. Pairwise

comparisons between canopy position (high or low) and location (FTE or BRF) were assessed

using the emmeans package in R and found to be significant if $P<0.05$ (Lenth 2020). All data

reported below are means ± SE. Finally, the relationships between physiological parameters and

either N_{area} or total chlorophyll (Chl+Car$_{area}$) were also examined using linear mixed effects

models. A description of the full models can be found in the Supplemental Information Methods

S1.

RESULTS

Tree Characteristics and Growth Environment: Tree characteristics for the trees measured in this

study have been previously reported in Griffin *et al.* (2022) but have been summarized in Table

1. Briefly, trees from BRF tended to have a larger diameter at breast height (DBH; measured at

1.37 m from the ground) with BRF trees having an average diameter of 23.1±1.99 cm and FTE

trees having an average diameter of 16.7±0.94 cm (Table 1). Trees from the two locations were

also extremely different in their average dripline areas (27.34±2.00 m^2 at BRF vs. 3.52±0.37 m^2

at the FTE; Table 1). Nevertheless, despite these differences trees from the two locations had

very similar average heights (9.91±0.73 m at BRF and 9.32±0.57 m at the FTE; Table 1).

Irradiance at the range extremes of white spruce, from the northernmost location at the

FTE in Alaska, to the southernmost location at BRF in New York, showed pronounced

differences in both intensity and day length (Fig 1a, b, d). Average ambient PPFD over the

measurement campaigns (Fig 1a; orange) estimated from daily diurnal light courses modeled

from canopy photos (e.g., July 4[th] 2017; Fig 1b) was 36% higher at BRF than at the FTE

(54.4±0.03 compared to 39.9±0.06 mol m^{-2} day^{-1}). However, day length was much longer at the

FTE than at BRF (Fig 1b). At the canopy positions where physiological and biochemical

measurements were made, average daily PPFD was significantly greater at high canopy positions

than low canopy positions at both BRF and the FTE (34.97±0.16 vs. 9.20±0.13 mol m^{-2} day^{-1} at

BRF, and 29.24±0.16 vs. 21.14±0.16 mol m^{-2} day^{-1} at the FTE; Fig 1a). Even so, the difference

in the daily mean PPFD experienced from the upper to the lower canopies was much smaller at

the FTE than at BRF (only 8 mol m^{-2} day^{-1} at the FTE vs. 25 mol m^{-2} day^{-1} at BRF). This

narrowing of the irradiance gradient across vertical canopy positions at the FTE was decoupled

from canopy height, given that trees at both locations were of similar heights.

In addition to differences in the light environment, there were also pronounced

differences in the daytime air temperatures between the two locations (Fig 1c). During the

respective study campaigns (as marked by the green (BRF) and blue (FTE) shaded areas (Fig

1c), BRF mean daytime temperature was 23.4°C, with a minimum temperature of 12.8°C and a

maximum temperature of 29.1°C, whereas the FTE mean temperature was 17°C, with a

minimum temperature of 6.5°C and a maximum temperature of 28°C. Thus, the FTE

experienced colder temperatures and greater temperature variability.

Relationships between light environment and gas exchange, biochemistry, pigments and PRI:

In general, significant relationships were found between average PPFD and physiological traits,

biochemical traits, pigment concentrations and PRI at BRF. In contrast, few significant

relationships were found between average PPFD and foliar traits at the FTE (Table 2).

Photosynthesis and Respiration: A significant positive relationship was observed between

average PPFD and A_{1500} at BRF with high canopy foliage having a higher A_{1500} than low canopy

foliage. No significant relationship was observed between A_{1500} and average PPFD at the FTE,

nor were there any significant differences between the canopy positions (Fig 2a, Tables 2, S1 &

S2). Across all samples at each location, mean A_{1500} was found to be 40% higher at BRF than at

the FTE (10.8±0.5 at BRF vs. 7.7±0.4 µmol m^{-2} s^{-1} at the FTE; Table S2). The differences

between BRF and the FTE in the relationship of A_{1500} to PPFD correspond to differences in

several other parameters commonly extracted from light response curves. Namely the following

were observed at BRF: a significant negative relationship between average PPFD and Φ (Fig 2b,

Table 2), with significantly lower Φ in the high than the low canopy (Fig 2b, Tables S1 & S2);

and significant positive relationships between average PPFD and LSP, LCP, R_D, and R_L with

higher values in the high than the low canopies (Figs 2c, 2d, 2e & 2f, Tables 2, S1 & S2). At the

FTE, as in the case of A_{1500}, no significant relationship was found between average PPFD and Φ,

LCP, R_D or R_L (Figs 2b, 2d, 2e & 2f, Table 2); however, a significant positive relationship was

found between average PPFD and LSP with a correspondingly higher LSP in the high canopy

than the low canopy foliage at both locations (Fig 2c, Tables 2, S1 & S2). Across all samples at

each location, Φ was significantly lower at the FTE than at BRF (0.03±0.001 at the FTE vs.

0.05±0.002 at BRF; Tables S1 & S2), and LSP, LCP, R_D and R_L were significantly higher at the

FTE than at BRF (LSP = 644±14 at the FTE vs. 522±18.6 µmol m^{-2} s^{-1} at BRF; LCP = 90.7±4.5

at the FTE vs. 36.8±2.1 µmol m^{-2} s^{-1} at BRF (LCP means are backtransformed); R_D = 2.46±0.11

at the FTE vs. 1.98±0.14 µmol m^{-2} s^{-1} at BRF; R_L = 2.22±0.12 at the FTE vs. 1.58±0.16 µmol m^{-2}

s^{-1} at BRF; Tables S1 & S2). In particular, the differences in mean R_L between the two locations

were dramatic, with the FTE having 41% higher R_L than at BRF.

The parameters extracted from the A-C_i curves (V_{cmax} and J_{max} normalized to 25°C) also

followed the positive relationship between PPFD and A_{1500} at BRF. Both V_{cmax} and J_{max} had

positive relationships with PPFD and significantly higher values in the high than the low canopy

at BRF (Fig 3, Tables 2, S1 & S2). However, the expected lack of a relationship with PPFD at

the FTE was not observed. Instead, FTE V_{cmax} and J_{max} both had significant negative

relationships with PPFD, with significantly lower values in the high than the low canopy.

Consequently, when values from canopy positions were pooled to compare the two locations, no

significant differences were observed between the FTE and BRF (V_{cmax} = 48.5±1.7 at the FTE

and 50.3±2.2 µmol m^{-2} s^{-1} at BRF; J_{max} = 80.8±2.7 at the FTE and 85.9±3.4 µmol m^{-2} s^{-1} at BRF;

Tables S1 & S2).

<u>Leaf traits</u>: A significant negative relationship was observed between SLA and PPFD at BRF

while no relationship was observed between SLA and PPFD at the FTE (Fig 4a, Table 2).

Across canopy positions, high canopy foliage had significantly lower SLA at both locations (Fig

4a, Tables S1 & S2). The FTE had significantly lower SLA than BRF (32.0±1.1 at the FTE vs.

50.4±1.2 cm^2g^{-1} at BRF, Tables S1 & S2). %N showed no significant relationships with PPFD,

and no significant differences between canopy positions in either location; although the FTE did

have a small but significantly lower %N than BRF (0.98±0.03 at the FTE vs. 1.1±0.04 at BRF;

Fig 4c, Tables 2, S1 & S2). Similarly, C:N showed no significant relationships with PPFD and

no significant differences between canopy positions in either location; however, C:N was

significantly lower at BRF compared to the FTE (Tables 2, S1 & S2). Due to the strong

relationship between SLA and PPFD, N_{area} also had a significant positive relationship with PPFD

at BRF and no relationship at the FTE (Fig 4e, Table 2).

<u>Pigments and Spectral Indices</u>: With regard to pigment concentrations, positive relationships

were seen both at BRF and the FTE in the ratio of chlorophyll *a* and chlorophyll *b*

(Chl*a*:Chl*b*$_{area}$) and in the Carotenoids (Car$_{area}$) with higher Chl*a*:Chl*b*$_{area}$ and higher Car$_{area}$ in the

high than the low canopies (Figs 4b & 4d, Tables 2, S1 & S2). The expected decrease in the

ratio of chlorophyll to carotenoids (Chl:Car$_{area}$) with increasing PPFD was observed at BRF, but

no relationship was found at the FTE (Fig 2f, Table 2). However, the FTE did have 46% more

carotenoids than BRF (14.5±0.5 at the FTE vs. 9.9±0.8 µg cm^{-2} at BRF; Tables S1 & S2) and a

significantly lower Chl:Car$_{area}$ (6.7±0.2 at the FTE vs. 9.0±0.3 at BRF; Tables S1 & S2). Finally,

PRI had the expected positive relationship with Chl:Car$_{area}$ at BRF (Fig 5, Tables 3, S3) and it

decreased significantly with increasing PPFD at BRF (Fig 5, Table 2). However, no

relationships between PRI and Chl:Car$_{area}$ or PPFD were found at the FTE (Fig 5, Tables 2, 3,

S3).

<u>Relationships between gas exchange traits and nitrogen</u>: As with PPFD, positive relationships

were found between N_{area} and A_{1500}, and N_{area} and V_{cmax} at BRF while no relationships were

observed at the FTE (Fig 6a & 6b, Tables 3, S3). In contrast, both BRF and the FTE had positive

relationships between N_{area} and R_D (Fig 6c, Tables 3, S3), between N_{area} and R_L (Tables 3, S3)

and between N_{area} and total pigments (chl$a$ + chl$b$ + carotenoids; Fig 6d, Tables 3, S3).

Carbon balance at the range extremes of white spruce: The collection of gas exchange

parameters such as V_{cmax} and J_{max}, which are commonly used to predict photosynthesis, together

with values of respiration at 25°C (R_{25}) previously reported by Griffin *et al.* (2022), provide a

unique opportunity to examine the carbon balance across the range extremes of white spruce.

We find that, when all parameters are normalized to 25°C, the ratios of V_{cmax} and J_{max} to R_{25} are

228% higher at BRF than at the FTE (Fig 7, Tables S1 & S2; $V_{cmax}/R_{25} = 21.4\pm1.39$ at the FTE

vs. 70.2 ± 7.47 at BRF; $J_{max}/R_{25} = 36.5\pm2.45$ at the FTE vs. 119.9 ± 13.21 at BRF). No significant

differences were observed in the ratios between canopy positions at each location, although there

was a trend for larger ratios in the low canopy than the high canopy at BRF (Fig 7, Tables S1 &

S2).

DISCUSSION

Local environmental conditions drive differences in tree structure between white spruce range

extremes

Vertical canopy gradients of foliar physiological and biochemical traits differ between white

spruce growing at its northern- and southern-most range limits. Whereas steep canopy gradients

in physiological functioning are found at the southern range limit in Black Rock Forest (BRF), a

clear lack of variation in foliar physiological capacity is observed between the high and low

canopies of white spruce at its northernmost range extreme in the Forest-Tundra Ecotone (FTE).

As our two study locations delimit the distributional range of white spruce and are separated by a

distance of more than 5000 km, these differences are likely in response to local environmental

conditions. Solar zenith angles, day length, air and soil temperatures, soil moisture and vapor

pressure deficit vary substantially between these two locations. Of these environmental

conditions, the impacts of local light environment are of particular interest for the following

reasons: 1) light energy is a critical driver of photosynthetic carbon gain (Hirose & Werger

1987); 2) abundant evidence shows that canopy gradients in photosynthetic traits are strongly

correlated to gradients in irradiance throughout tree crowns (Field 1983; Hirose & Werger 1987;

Ellsworth & Reich 1993; Bond *et al.* 1999; Niinemets, Keenan & Hallik 2015), but little work

assesses the impacts of latitudinal location on these gradients within a single species; and 3)

these gradients are key parameters in ecosystem models used to calculate gross primary

productivity across latitudinal space (Bonan *et al.* 2012; Rogers *et al.* 2017). Day length and

solar zenith angles increase and light intensity decreases as one moves towards the northernmost

latitudes of the globe (Nilsen 1983; Slaughter & Viereck 1986). These local light environments

are further modified by differences in forest and tree crown structure between the two locations.

Trees at the southern location exist within a complex forest canopy comprised of deciduous

hardwood species where they must compete for light energy and other above-ground resources.

Consequently white spruce tree crowns are distinctly conical in shape with a wide dripline area

in the southern location, presumably to minimize self-shading and maximize light energy

capture. In contrast, trees at the northern location are widely spaced with narrow and untapering

tree crowns. The open forest canopy minimizes competition for resources such as light.

Furthermore, the cylindrical shape promotes more equal exposure of foliage to irradiance in a

region of the globe with extremely high solar zenith angles, ultimately increasing the efficiency

of light capture for photosynthesis (Kuuluvainen 1992). Overall, these solar and structural

differences lead to a narrowing of the intra-canopy light gradients as one moves from the

southern (where light availability is ~35 and ~9 mol m^{-2} day^{-1} at the high and low canopies,

respectively) to the northern (where light availability is ~29 and ~21 mol m^{-2} day^{-1} at the high

and low canopies, respectively) latitudinal limits of this species.

Vertical gradients in physiological function are coordinated to latitudinal differences in light

environment

Inherent to the structural differences of trees in these two locations is the need for tree crowns to

organize their physiological and biochemical traits in order to optimize photosynthetic carbon

gain (Hirose & Werger 1987; Niinemets *et al.* 1998). Therefore, at BRF, where there are strong

intra-canopy light gradients, we find strongly positive relationships between PPFD and A_{1500},

J_{max} and V_{cmax} indicating the upregulation of the light reactions (in which RuBP is produced) and

the carbon reactions (in which rubisco acts on RuBP to fix carbon) of photosynthesis in the high

canopy (Givnish 1988; Valladares & Niinemets 2008). The elevated metabolic activity in the

high canopy results in higher respiratory activity in order to support the costs of growth and

protein maintenance (Reich *et al.* 1998; Lewis, McKane, Tingey & Beedlow P. A. 2000; Griffin,

Tissue, Turnbull, Schuster & Whitehead 2001). Biochemical traits also follow the inherent light

gradients. High light needles have lower SLA and a greater N_{area} reflecting the greater leaf

thickness of high light leaves from added palisade layers and the presence of nitrogen in both the

enzyme rubisco and in chlorophyll (Evans 1989; Smith, Vogelmann, DeLucia, Bell & Shepherd

1997; Evans & Poorter 2001). Finally, high light leaves have greater Car_{area} and a smaller

Chl:Car_{area} ratio reflecting the need for greater photoprotection from excess light energy

(Demmig-Adams & Adams III 1992). In all cases, biochemical and physiological traits of low

latitude white spruce conform to the established literature showing strong correlations with

canopy gradients of irradiance. At the high latitude location where intra-canopy light gradients

become less pronounced, we see the absence of canopy gradients in many physiological and

biochemical traits. What is most striking is the lack of relationships between irradiance and each

of photosynthesis (A_{1500}), light respiration, and dark respiration. Together with the lack of

relationships between irradiance and the apparent quantum yield, the light compensation point,

specific leaf area and N_{area}, these findings suggest that the uneven distribution of metabolic

activity throughout the canopy required to optimize photosynthetic carbon gain in

correspondence with canopy irradiance gradients at low latitudes does not occur at high latitudes

where irradiance varies minimally throughout the canopy profile.

The parameters discussed above suggest that northern latitude trees have few

distinguishable canopy gradients in photosynthetic physiology. However, positive relationships

between irradiance and the ratio of Chl*a* to Chl*b*, the light saturation point and carotenoid

concentrations suggest that even though the magnitude of the light gradient narrows at northern

latitudes, there is still a subtle intra-canopy light gradient to which pigments are responding.

Thus, in the higher light upper canopy we see greater investments in Chl*a* than Chl*b* to maximize

light capture and a greater investment in carotenoids to dissipate excess light energy and protect

photosynthetic machinery (Dale & Causton 1992; Hikosaka & Terashima 1995; Kitajima &

Hogan 2003; Ruban 2015; Magney *et al.* 2016; Scartazza *et al.* 2016). It is possible that these

subtle canopy gradients in pigments may be a result of the 24 hours of light exposure during the

growing season at the FTE.

In contrast to the lack of relationships found between irradiance and the majority of traits,

and the subtle positive relationships found between irradiance and pigments at treeline, we find a

surprising downregulation of J_{max} and V_{cmax} in the high canopy foliage at the FTE. It is unclear

why we find these negative relationships between irradiance and the component processes of

photosynthesis at the northern locations; however, there are several possible explanations and

hypotheses that could be further explored. One possibility is that the combination of diminutive

vertical canopy gradients in light, SLA and nitrogen, the latter of which is critical to the

performance of J_{max} and V_{cmax} because of the presence of nitrogen in chlorophyll and the enzyme

rubisco (Evans 1989; Smith *et al.* 1997; Evans & Poorter 2001), has led to the observed negative

relationships. However, strong relationships do exist between nitrogen and total pigment

concentrations even though there is no apparent relationship between nitrogen and V_{cmax}.

Furthermore, we still find weak positive relationships between irradiance and both carotenoid

concentrations and Chl*a*:Chl*b* even with a narrow light gradient. Consequently, we conclude

that a more mechanistic explanation must exist for the puzzling downregulation of V_{cmax} and

J_{max}. While in this study we focused on the latitudinal relationships between canopy light

environments and foliar physiological traits, it is possible that intra-canopy gradients in other

environmental conditions may also impact V_{cmax} and J_{max} measurements (Martin, Hinckley,

Meinzer & Sprugel 1999; Zweifel, Böhm & Häsler 2002; Bauerle, Bowden, Wang & Shahba

2009; Bachofen, D'Odorico & Buchmann 2020; Buckley 2021). Perhaps of particular relevance

with respect to rubisco carboxylation is the reliance of this process on the enzyme rubisco, whose

activity is well-known to be temperature dependent (Yamori & von Caemmerer 2009; Benomar

et al. 2018). Work at alpine treelines has clearly documented that as trees grow in stature, their

foliage becomes increasingly coupled to atmospheric conditions such as temperature and wind

stress (Körner 2012). These stresses may be detrimental to high canopy photosynthetic

performance (lowering V_{cmax} and J_{max}). In contrast, low canopy foliage is less coupled to

atmospheric conditions and may therefore be more protected from temperature and wind

extremes (Martin *et al.* 1999; Germino & Smith 1999; Johnson, Germino & Smith 2004;

Maguire *et al.* 2019).

Temperature and wind stress are not the only environmental conditions likely to cause

reduced gradients in physiological functioning. Hydraulic conductance has also been found to

vary in response to water availability, impacting canopy gradients in nitrogen and photosynthesis

(Peltoniemi *et al.* 2012). It is equally possible that winter temperature extremes may impact

hydraulic function in exposed foliage. Consequently, throughout the canopy of Alaskan trees

there may be a constant balancing act with regard to the resource allocation necessary to improve

whole canopy photosynthetic carbon gain. Two factors compete with each other: namely the

need to devote resources to light capture in the high canopy where irradiance is slightly more

available; and the need to devote resources either to foliage protection in the high canopy or

increased carbon gain in the low canopy when harsh conditions prevail. The result, ultimately, is

what we find across the majority of physiological and biochemical traits, i.e., a lack of variation

in photosynthetic carbon gain across the high and low canopy in high latitude trees. Without

detailed information on temperature, wind, and hydraulic profiles throughout the canopy, this

hypothesis cannot be fully tested. Yet, it seems clear from the linkages between irradiance,

photosynthesis and tree structure across the latitudinal extremes that the environment is an

integral driver of species' geographic range. We suggest that expanding to examine additional

environmental conditions such as temperature, wind and water stress may be a compelling future

direction, thereby adding to our overall understanding of the mechanisms controlling canopy

carbon exchange across the massive forest tundra ecotone and more broadly across latitudinal

gradients.

The accurate portrayal of canopy gradients is important to the precise prediction of

carbon exchange and gross primary productivity (GPP) (Sands 1995; Bonan *et al.* 2012).

However, the lack of variation in photosynthesis at the FTE suggests that complex canopy

models used to account for canopy self-shading at mid-latitudes may not be necessary to model

GPP in trees from high latitudes. The lack of variation in the photochemical reflectance index

(PRI) across vertical canopy positions at the FTE similarly suggests that accounting for canopy

self-shading may be less critical at northern latitudes as opposed to mid-latitudes. PRI has

previously been demonstrated to be an accurate predictor of whole canopy gross primary

productivity in broad-leaved species when used in concert with terrestrial Light Detection and

Ranging (LiDAR) to account for canopy self-shading (Hilker *et al.* 2008; Coops, Hermosilla,

Hilker & Black 2017). Although our own study does not employ LiDAR, the presence of

canopy differences in PRI at the southernmost location suggests that such a technique could

work equally well in mid-latitude coniferous forests. Furthermore, the fact that PRI mirrors A_{1500}

regardless of latitude suggests that this remotely sensed index may be a valuable future tool for

accurately assessing gross primary productivity of other species with different structural

properties and with large latitudinal distributions.

Physiological and biochemical traits may contribute to the range limits of white spruce
The detailed data on the physiology and biochemistry of white spruce not only allow for the
examination of canopy differences across latitudinal range extremes, but also for the examination
of locational differences in traits that may further illuminate possible drivers contributing to the
range limits of this important species. Here, we observed important differences in physical and
biochemical traits such as SLA, % N, C:N and carotenoid concentration between the southern
and northern range extremes. Most relevant to the above discussion of light environment and its
impacts on photosynthetic physiology is the finding of significantly higher carotenoid
concentrations in needles of northern white spruce compared to southern white spruce. This
indicates a greater investment in photoprotection at northern latitudes during the growing season.
Elevated carotenoids at the northern range extreme may be in response to the 24 hour
photoperiod at the Arctic tundra boreal forest ecotone. Past studies have found that very high
levels of light protecting xanthophyll pigments can be a feature of Arctic plants subjected to
continuous light (Magney *et al.* 2019). Lower overall SLA in Alaskan needles compared to BRF
needles indicates that foliage at the northern edge of the distribution is thicker or denser.
Although decreasing SLA is commonly found in response to increasing light availability due to
the added density of additional photosynthetic machinery (as seen throughout the BRF canopy),
SLA is also commonly positively correlated to ambient growth temperatures (Atkin, Loveys,
Atkinson & Pons 2006; Poorter, Niinemets, Poorter, Wright & Villar 2009). Thus, a low SLA at
the FTE may indicate a greater investment in needle structure rather than photosynthetic
machinery, a hypothesis that is corroborated by our finding of higher carbon to nitrogen ratios in
Alaskan needles (González-Zurdo, Escudero, Babiano, García-Ciudad & Mediavilla 2016).
Investment in structure is potentially adaptive in the harsh forest tundra ecotone where needles
must survive multiple years of winter temperatures, snow or ice accumulation and windblown ice
abrasion. Leaf nitrogen was also found to be slightly but significantly lower in needles from the
northern range limit. This result confirms the findings of Griffin *et al.* (2022) and agrees with
past studies that show negative correlations between nitrogen and both temperature and latitude
(Körner 1989; Yin 1993; Reich & Oleksyn 2004). Nitrogen biogeochemistry is intimately linked
to environmental and ecological factors. For example, harsh temperatures and unique hydrologic
and permafrost dynamics inherent to the Arctic can slow decomposition rates and depress
nitrogen fixation (Schimel & Stuart Chapin 1996; Schimel, Bilbrough & Welker 2004). The

result is the limited availability of nitrogen in the harsh northern latitudes of the globe. The

differences in biochemical and structural traits between our two study locations hint at

environmental drivers, in addition to light availability, that should be considered in future studies

examining latitudinal differences in canopy gradients.

Finally, in addition to biochemical differences between locations, metabolic differences

are key in understanding the mechanisms controlling the range limits of this species. Previous

work in white spruce shows this species to have extremely high rates of dark respiration at its

northern compared to southern range limit, a finding that has led to the conclusion that

respiratory carbon loss may be a crucial factor constraining the northern range limit of this

species (Griffin *et al.* 2022). We find similar patterns in the current study, with significantly

higher rates of dark respiration (as estimated from the light response curves) in white spruce in

Alaska than in New York. Furthermore, we find that these differences are apparent not only in

dark respiration, but also in light respiration. Given that the northern range extreme of this

species experiences 24 hours of sunlight during the majority of the growing season, the greater

rates of light respiration in FTE compared to BRF trees provides important corroboration of the

constancy of the extreme respiratory cost at this northern range limit and the role it may play in

determining the location of northern treeline. It is likely that these high respiratory fluxes to the

atmosphere are related to high maintenance costs given the harsh environmental conditions of

the region.

Northern white spruce are not only characterized by high carbon losses. The additional

information on photosynthetic processes provided here further suggests that these carbon costs

are not matched by similarly high carbon gains. We do not directly compare the ratios of

photosynthetic carbon gain to respiratory carbon loss due to the use of different measurement

temperatures chosen to reflect ambient conditions in each location, but the ability to normalize

both V_{cmax} and J_{max} to 25°C allows us to assess the carbon balance of white spruce at its

latitudinal range extremes. The results of this comparison are dramatic and demonstrate that

northern trees function within a much narrower margin to maintain a positive carbon balance. It

seems that it is not only high respiratory costs that result in slow growth rates of white spruce

(Jensen et al., *pers. com.*) and constrain the latitudinal range limit of northern treeline. Instead,

our study suggests that it is the combined effect of high respiration and low V_{cmax} and J_{max} that

ultimately limit the northern range of this widely distributed and important boreal species.

Conclusions

White spruce spans a massive distribution, the extremes of which are characterized by vastly different environmental conditions. Here we demonstrate that differences in local light environment correlate to differences in vertical canopy physiology, resource allocation and crown structure. Thus, at high latitudes where light intensity is low, solar zenith angles are high and day length is long, there is a clear lack of variation between high and low canopy white spruce traits. In contrast, white spruce at lower latitudes exist in a complex forest canopy of deciduous hardwood species and experience greater differences in local and canopy light environments leading to steep gradients in physiological and biochemical traits between high and low canopy positions. Consequently, accounting for self-shading may be less critical for predicting gross primary productivity at northern relative to southern latitudes. Our findings point to light environment as an important driver in latitudinal variation in physiology within a single species and emphasize the importance of the connection between environment and physiology in determining a species' distributional range. Furthermore, our findings highlight that additional environmental drivers such as temperature, and water availability may also contribute both to canopy and latitudinal differences in physiological traits.

These data not only allowed us to examine the role of physiology in determining a species' distributional range, but they also permitted us to question the role of a species' carbon balance in determining the extreme limit of tree growth form. Previous work has highlighted that extreme respiratory costs may be a critical component leading to the latitudinal location of northern treeline. Here we demonstrate that it is not only extreme respiratory costs but also minimal photosynthetic carbon gains that together may determine the latitudinal location of the world's largest ecotone and the transition from boreal forest to treeless tundra.

ACKNOWLEDGEMENTS

This work was supported by NASA ABoVE grant NNX15AT86A and the Arctic LTER (NSF Grant No. 1637459 & 2220863). We thank Sarah Sacket from the NASA ABoVE support team in Fairbanks, AK, for her energetic and prompt logistical support during Alaska field campaigns. At Black Rock Forest, we also wish to acknowledge the unflagging support of BRF staff

including Ben Brady and Matthew Munson as well as Dr. William Schuster, the Executive

Director of Black Rock.

AUTHOR CONTRIBUTION

SCS, KLG, NTB, LAV and JUHE designed the research. They were assisted in data collection

by SGB, LM, and JJ. SCS analyzed the data and wrote the first draft of the manuscript. All

authors edited and approved the final version for submission.

DATA AVAILABILITY

Data will be archived with ORNL DAAC upon acceptance of the manuscript.

REFERENCES

Akalusi M.E., Meng F.R. & Bourque C.P.-A. (2021) Photosynthetic parameters and stomatal

conductance in attached and detached balsam fir foliage. *Plant-Environment Interactions* **2**,

206–215.

Arguez A.I., Durre I., Applequist S., Squires M.F., Vose R.S., Yin X., … Owen T. (2012)

NOAA's 1981-2010 U.S. Climate Normals: An Overview. *Bulletin of the American*

Meteorological Society **93**, 1687–1697.

Atkin O.K., Loveys B.R., Atkinson L.J. & Pons T.L. (2006) Phenotypic plasticity and growth

temperature: understanding interspecific variability. *Journal of Experimental Botany* **57**,

267–281.

Ayub G., Smith R.A., Tissue D.T. & Atkin O.K. (2011) Impacts of drought on leaf respiration in

darkness and light in *Eucalyptus saligna* exposed to industrial-age atmospheric CO_2 and

growth temperature. *New Phytologist* **190**, 1003–1018.

Bachofen C., D'Odorico P. & Buchmann N. (2020) Light and VPD gradients drive foliar

nitrogen partitioning and photosynthesis in the canopy of European beech and silver fir.

Oecologia **192**, 323–339.

Bates D., Maechler M., Bolker B. & Walker S. (2015) Fitting Linear Mixed-Effects Models

using lme4. *Journal of Statistical Software* **67**, 1–48.

Bauerle W.L., Bowden J.D., Wang G.G. & Shahba M.A. (2009) Exploring the importance of

within-canopy spatial temperature variation on transpiration predictions. *Journal of*

Experimental Botany **60**, 3665–3676.

Benomar L., Lamhamedi M.S., Pepin S., Rainville A., Lambert M.-C., Margolis H.A., …

Beaulieu J. (2018) Thermal acclimation of photosynthesis and respiration of southern and

northern white spruce seed sources tested along a regional climatic gradient indicates

limited potential to cope with temperature warming. *Annals of Botany* **121**, 443–457.

Bolker B. & R Development Core Team (2020) bbmle: Tools for General Maximum Likelihood

Estimation. R package version 1.0.23.1. https://CRAN.R-project.org/package=bbmle.

Bonan G.B., Oleson K.W., Fisher R.A., Lasslop G. & Reichstein M. (2012) Reconciling leaf

physiological traits and canopy flux data: Use of the TRY and FLUXNET databases in the

Community Land Model version 4. *Journal of Geophysical Research: Biogeosciences* **117**,

1–19.

Bond B.J., Farnsworth B.T., Coulombe R.A. & Winner W.E. (1999) Foliage physiology and

biochemistry in response to light gradients in conifers with varying shade tolerance.

Oecologia **120**, 183–192.

Brown J.H., Stevens G.C. & Kaufmnan D.M. (1996) The Geographic Range: Size, Shape,

Boundaries, and Internal Structure. *Annual Review of Ecology and Systematics* **27**, 597–

623.

Buckley T.N. (2021) Optimal carbon partitioning helps reconcile the apparent divergence

between optimal and observed canopy profiles of photosynthetic capacity. *New Phytologist*

230, 2246–2260.

Buckley T.N., Vice H. & Adams M.A. (2017) The Kok effect in *Vicia faba* cannot be explained

solely by changes in chloroplastic CO_2 concentration. *New Phytologist* **216**, 1064–1071.

Coops N.C., Hermosilla T., Hilker T. & Black T.A. (2017) Linking stand architecture with

canopy reflectance to estimate vertical patterns of light-use efficiency. *Remote Sensing of*

Environment **194**, 322–330.

Dale M.P. & Causton D.R. (1992) Use of the Chlorophyll a/b Ratio as a Bioassay for the Light

Environment of a Plant. *Functional Ecology* **6**, 190–196.

Demmig-Adams B. & Adams III W.W. (1992) Photoprotection and other responses of plants to

high light stress. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **43**, 599–626.

Duursma R.A. (2015) Plantecophys--An R Package for Analysing and Modelling Leaf Gas

Exchange Data. *PloS One* **10**, e0143346.

Eitel J.U.H., Griffin K.L., Boelman N.T., Maguire A.J., Meddens A.J.H., Jensen J., …
Jennewein J.S. (2020) Foliar remote sensing tracks radial tree growth dynamics. *Global*
Change Biology **26**, 4068–4078.

Eitel J.U.H., Maguire A.J., Boelman N., Vierling L.A., Griffin K.L., Jensen J., … Sonnentag O.
(2019) Proximal remote sensing of tree physiology at northern treeline: Do late-season
changes in the photochemical reflectance index (PRI) respond to climate or photoperiod?
Remote Sensing of Environment **221**, 340–350.

Ellsworth D.S. & Reich P.B. (1993) Canopy Structure and Vertical Patterns of Photosynthesis
and Related Leaf Traits in a Deciduous Forest. *Oecologia* **96**, 169–178.

Evans J.R. (1989) Photosynthesis and nitrogen relationships in leaves of C_3 plants. *Oecologia* **78**,
9–19.

Evans J.R. & Poorter H. (2001) Photosynthetic acclimation of plants to growth irradiance: the
relative importance of specific leaf area and nitrogen partitioning in maximizing carbon
gain. *Plant, Cell and Environment* **24**, 755–767.

Farquhar G.D. & Busch F.A. (2017) Changes in the chloroplastic CO_2 concentration explain
much of the observed Kok effect: a model. *New Phytologist* **214**, 570–584.

Farquhar G.D., von Caemmerer S. & Berry J.A. (1980) A biochemical model of photosynthetic
CO_2 assimilation in leaves of C_3 species. *Planta* **149**, 78–90.

Field C. (1983) Allocating Leaf Nitrogen for the Maximization of Carbon Gain: Leaf Age as a
Control on the Allocation Program. *Oecologia* **56**, 341–347.

Gamon A., Penuelas J. & Field C.B. (1992) A Narrow-Waveband Spectral Index That Tracks
Diurnal Changes in Photosynthetic Efficiency. *Remote Sensing of Environment* **41**, 35–44.

Gamon J.A., Huemmrich K.F., Wong C.Y.S., Ensminger I., Garrity S., Hollinger D.Y., …
Peñuelas J. (2016) A remotely sensed pigment index reveals photosynthetic phenology in
evergreen conifers. *PNAS* **113**, 13087–13092.

Gaston K.J. (1996) Species-range-size distributions: Patterns, mechanisms and implications.
Trends in Ecology and Evolution **11**, 197–201.

Germino M.J. & Smith W.K. (1999) Sky exposure, crown architecture, and low-temperature
photoinhibition in conifer seedlings at alpine treeline. *Plant, Cell and Environment* **22**, 407–
415.

Givnish T.J. (1988) Adaptation to Sun and Shade: A Whole-plant Perspective. *Australian*

Journal of Plant Physiology **15**, 63–92.

González-Zurdo P., Escudero A., Babiano J., García-Ciudad A. & Mediavilla S. (2016) Costs of leaf reinforcement in response to winter cold in evergreen species. *Tree Physiology* **36**, 273–86.

Griffin K.L., Griffin Z.M., Schmiege S.C., Bruner S., Boelman N.T., Vierling L.A. & Eitel J.U.H. (2022) Variation in white spruce needle respiration at the species range limits: a potential impediment to northern expansion. *Plant, Cell & Environment*, 1–15.

Griffin K.L., Schmiege S.C., Bruner S.G., Boelman N.T., Vierling L.A. & Eitel J.U.H. (2021) High Leaf Respiration Rates May Limit the Success of White Spruce Saplings Growing in the Kampfzone at the Arctic Treeline. *Frontiers in Plant Science* **12**.

Griffin K.L., Tissue D.T., Turnbull M.H., Schuster W. & Whitehead D. (2001) Leaf dark respiration as a function of canopy position in *Nothofagus fusca* trees grown at ambient and elevated CO_2 partial pressures for 5 years. *Functional Ecology* **15**, 497–505.

Harley P.C. & Baldocchi D.D. (1995) Scaling carbon dioxide and water vapour exchange from leaf to canopy in a deciduous forest. I. Leaf model parametrization. *Plant, Cell and Environment* **18**, 1146–1156.

Herrmann H.A., Schwartz J.M. & Johnson G.N. (2020) From empirical to theoretical models of light response curves - linking photosynthetic and metabolic acclimation. *Photosynthesis Research* **145**, 5–14.

Heskel M.A., Atkin O.K., Turnbull M.H. & Griffin K.L. (2013) Bringing the Kok effect to light: A review on the integration of daytime respiration and net ecosystem exchange. *Ecosphere* **4**, 1–14.

Hikosaka K. & Terashima I. (1995) A model of the acclimation of photosynthesis in the leaves of C_3 plants to sun and shade with respect to nitrogen use. *Plant, Cell and Environment* **18**, 605–618.

Hilker T., Coops N.C., Schwalm C.R., Jassal R.S., Black T.A. & Krishnan P. (2008) Effects of mutual shading of tree crowns on prediction of photosynthetic light-use efficiency in a coastal Douglas-fir forest. *Tree Physiology* **28**, 825–834.

Hirose T. & Werger M.J.A. (1987) Maximizing daily canopy photosynthesis with respect to the leaf nitrogen allocation pattern in the canopy. *Oecologia* **72**, 520–526.

Johnson D.M., Germino M.J. & Smith W.K. (2004) Abiotic factors limiting photosynthesis in

Abies lasiocarpa and *Picea engelmannii* seedlings below and above the alpine timberline.

Tree Physiology **24**, 377–386.

Kirschbaum M.U. & Farquhar G.D. (1987) Investigation of the CO_2 Dependence of Quantum

Yield and Respiration in *Eucalyptus pauciflora*. *Plant Physiology* **83**, 1032–1036.

Kitajima K. & Hogan K.P. (2003) Increases of chlorophyll a/b ratios during acclimation of

tropical woody seedlings to nitrogen limitation and high light. *Plant, Cell and Environment*

26, 857–865.

Kok B. (1948) A critical consideration of the quantum yield of *Chlorella*-photosynthesis.

Enzymologia **13**, 1–56.

Körner C. (1989) The nutritional status of plants from high altitudes - A worldwide comparison.

Oecologia **81**, 379–391.

Körner C. (2012) *Alpine treelines: Functional Ecology of the Global High Elevation Tree Limits*.

Springer, Basel.

Kuuluvainen T. (1992) Tree Architectures Adapted to Efficient Light Utilization: Is There a

Basis for Latitudinal Gradients? *Nordic Society Oikos* **65**, 275–284.

Kuznetsova A., Brockhoff P. & Christensen R. (2017) lmerTest Package: Tests in Linear Mixed

Effects Models. *Journal of Statistical Software* **82**, 1–26.

Lambers H. & Oliveira R.S. (2019) Physiological Ecology and the Distribution of Organisms. In

Plant Physiological Ecology. pp. 2–5. Springer, Cham.

Lenth R. (2020) emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.4.5. https://CRAN.R-project.org/package=emmeans.

Lewis J.D., McKane R.B., Tingey D.T. & Beedlow P. A. (2000) Vertical gradients in

photosynthetic light response within an old-growth Douglas-fir and western hemlock

canopy. *Tree Physiology* **20**, 447–456.

Lichtenthaler H.K. (1987) Chlorophylls and carotenoids: Pigments of photosynthetic

biomembranes. *Methods in Enzymology* **148**, 350–382.

Little Jr. E.L. (1999) Digital representation of "Atlas of United States Trees." *U.S. Geological*

Survey Professional Paper 1650.

Lowry E. & Lester S.E. (2006) The biogeography of plant reproduction: Potential determinants

of species' range sizes. *Journal of Biogeography* **33**, 1975–1982.

Magney T.S., Bowling D.R., Logan B.A., Grossmann K., Stutz J., Blanken P.D., … Frankenberg

C. (2019) Mechanistic evidence for tracking the seasonality of photosynthesis with solar-
induced fluorescence. *Proceedings of the National Academy of Sciences of the United States*
of America **116**, 11640–11645.

Magney T.S., Eitel J.U.H., Griffin K.L., Boelman N.T., Greaves H.E., Prager C.M., … Vierling
L.A. (2016) LiDAR canopy radiation model reveals patterns of photosynthetic partitioning
in an Arctic shrub. *Agricultural and Forest Meteorology* **221**, 78–93.

Maguire A.J., Eitel J.U.H., Vierling L.A., Johnson D.M., Griffin K.L., Boelman N.T., …
Meddens A.J.H. (2019) Terrestrial lidar scanning reveals fine-scale linkages between
microstructure and photosynthetic functioning of small-stature spruce trees at the forest-
tundra ecotone. *Agricultural and Forest Meteorology* **269**–**270**, 157–168.

Martin T.A., Hinckley T.M., Meinzer F.C. & Sprugel D.G. (1999) Boundary layer conductance,
leaf temperature and transpiration of Abies amabilis branches. *Tree Physiology* **19**, 435–
443.

Mavi H.S. & Tupper G.J. (2004) Solar radiation and its role in plant growth. In
Agrometeorology: Principles and applications of climate studies in agriculture. pp. 13–42.
Food Products Press, Binghampton, NY.

Niinemets Ü. (2007) Photosynthesis and resource distribution through plant canopies. *Plant, Cell*
& Environment **30**, 1052–1071.

Niinemets Ü., Keenan T.F. & Hallik L. (2015) A worldwide analysis of within-canopy variations
in leaf structural, chemical and physiological traits across plant functional types. *New*
Phytologist **205**, 973–993.

Niinemets U., Kull O. & Tenhunen J.D. (1998) An analysis of light effects on foliar morphology,
physiology and light interception in temperate deciduous woody species of contrasting
shade tolerance. *Tree Physiology* **18**, 681–696.

Nilsen J. (1983) Light Climate in Northern Areas. In *Plant Production in the North*. (eds A.
Kaurin, O. Junttila & J. Nilsen), pp. 62–72. Norwegian University Press, Tromso, Norway.

Patterson A.E., Arkebauer R., Quallo C., Heskel M.A., Li X., Boelman N. & Griffin K.L. (2018)
Temperature response of respiration and respiratory quotients of 16 co-occurring temperate
tree species. *Tree Physiology* **00**, 1–14.

Peltoniemi M.S., Duursma R.A. & Medlyn B.E. (2012) Co-optimal distribution of leaf nitrogen
and hydraulic conductance in plant canopies. *Tree Physiology* **32**, 510–519.

Penuelas J., Filella I. & Gamon J.A. (1995) Assessment of photosynthetic radiation-use

efficiency with spectral reflectance. *New Phytologist*, 291–296.

Poorter H., Niinemets Ü., Poorter L., Wright I.J. & Villar R. (2009) Causes and consequences of

variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist* **182**, 565–588.

de Pury D.G.G. & Farquhar G.D. (1997) Simple scaling of photosynthesis from leaves to

canopies without the errors of big-leaf models. *Plant, Cell and Environment* **20**, 537–557.

R Core Team (2022) R: A language and environment for statistical computing.

Reich P.B. & Oleksyn J. (2004) Global patterns of plant leaf N and P in relation to temperature

and latitude. *Proceedings of the National Academy of Sciences of the United States of*

America **101**, 11001–11006.

Reich P.B., Walters M.B., Ellsworth D.S., Vose J.M., Volin J.C., Gresham C. & Bowman W.D.

(1998) Relationships of leaf dark respiration to leaf nitrogen, specific leaf area and leaf life-

span: a test across biomes and functional groups. *Oecologia* **114**, 471–482.

Rogers A., Medlyn B.E., Dukes J.S., Bonan G., von Caemmerer S., Dietze M.C., … Zaehle S.

(2017) A roadmap for improving the representation of photosynthesis in Earth system

models. *New Phytologist* **213**, 22–42.

Ruban A. V (2015) Evolution under the sun: optimizing light harvesting in photosynthesis.

Journal of Experimental Botany **66**, 7–23.

Sands P. (1995) Modelling Canopy Production. I. Optimal Distribution of Photosynthetic

Resources. *Australian Journal of Plant Physiology* **22**, 593–601.

Scartazza A., Di Baccio D., Bertolotto P., Gavrichkova O. & Matteucci G. (2016) Investigating

the European beech (*Fagus sylvatica* L.) leaf characteristics along the vertical canopy

profile: leaf structure, photosynthetic capacity, light energy dissipation and photoprotection

mechanisms. *Tree Physiology* **36**, 1060–1076.

Schimel J.P., Bilbrough C. & Welker J.M. (2004) Increased snow depth affects microbial

activity and nitrogen mineralization in two Arctic tundra communities. *Soil Biology and*

Biochemistry **36**, 217–227.

Schimel J.P. & Stuart Chapin F. (1996) Tundra plant uptake of amino acid and NH_4^+ nitrogen in

situ: Plants compete well for amino acid N. *Ecology* **77**, 2142–2147.

Schnieder C.A., Rasband W.S. & Eliceiri K.W. (2012) NIH Image to ImageJ: 25 years of image

analysis. *Nature Methods* **9**, 671–675.

Schulze E.D., Kelliher F.M., Körner C., Lloyd J. & Leuning R. (1994) Relationships among
maximal stomatal conductance, ecosystem surface conductance, carbon assimilation rate
and plant nitrogen nutrition: A global ecology scaling exercise. *Annual Review of Ecology*
and Systematics **25**, 629–662.

Schuster W.S.F., Griffin K.L., Roth H., Turnbull M.H., Whitehead D. & Tissue D. T (2008)
Changes in composition, structure and aboveground biomass over seventy-six years (1930 -
2006) in Black Rock Forest, Hudson Highlands, southeastern New York State. *Tree*
Physiology **28**, 537–549.

Sellers P.J., Berry J.A., Collatz G.J., Field C.B. & Hall F.G. (1992) Canopy reflectance,
photosynthesis, and transpiration. III. A reanalysis using improved leaf models and a new
canopy integration scheme. *Remote Sensing of Environment* **42**, 187–216.

Sharp R.E., Matthews M.A. & Boyer J.S. (1984) Kok effect and the quantum yield of
photosynthesis: light partially inhibits dark respiration. *Plant Physiology* **75**, 95–101.

Slaughter C.W. & Viereck L.A. (1986) Climatic Characteristics of the Taiga in Interior Alaska.
In *Forest Ecosystems in the Alaskan Taiga: A synthesis of structure and function*. (eds K.
Van Cleve, F.S. Chapin III, P.W. Flanagan, L.A. Viereck & C.T. Dyrness), pp. 9–21.
Springer, New York.

Smith W.K., Vogelmann T.C., DeLucia E.H., Bell D.T. & Shepherd K.A. (1997) Leaf Form and
Photosynthesis: Do leaf structure and orientation interact to regulate internal light and
carbon dioxide? *BioScience* **47**, 785–793.

ter Steege H. (1993) HEMIPHOT, a programme to analyze vegetation indices, light and light
quality from hemispherical photographs. Tropenbos Foundation, Wageningen.

ter Steege H. (1997) winphot, Version 5.0. A programme to analyse vegetation indices, light and
light quality from hemispherical photographs. Tropenbos-Guyana Programme.

ter Steege H. (2018) Hemiphot.R: Free R scripts to analyse hemispherical photographs for
canopy openness, leaf area index and photosynthetic active radiation under forest canopies.

Tcherkez G., Gauthier P., Buckley T.N., Busch F.A., Barbour M.M., Bruhn D., … Cornic G.
(2017) Tracking the origins of the Kok effect, 70 years after its discovery. *New Phytologist*
214, 506–510.

Valladares F. & Niinemets U. (2008) Shade Tolerance, a Key Plant Feature of Complex Nature
and Consequences. *Annual Review of Ecology, Evolution, and Systematics* **39**, 237–57.

Viereck L.A., Van Cleve K. & Dyrness C.T. (1986) Forest Ecosystem Distribution in the Taiga
Environment. In *Forest Ecosystems in the Alaskan Taiga: a synthesis of structure and*
function. (eds K. Van Cleve, F.S. Chapin III, P.W. Flanagan, L.A. Viereck & C.T. Dyrness),
pp. 22–43. Springer, New York.
Whitehead D., Griffin K.L., Turnbull M.H., Tissue D.T., Engel V.C., Brown K.J., … Walcroft
A.S. (2004) Response of total night-time respiration to differences in total daily
photosynthesis for leaves in a *Quercus rubra* L. canopy: Implications for modelling canopy
CO_2 exchange. *Global Change Biology* **10**, 925–938.
Wong C.Y.S. & Gamon J.A. (2015a) Three causes of variation in the photochemical reflectance
index (PRI) in evergreen conifers. *New Phytologist* **206**, 187–195.
Wong C.Y.S. & Gamon J.A. (2015b) The photochemical reflectance index provides an optical
indicator of spring photosynthetic activation in evergreen conifers. *New Phytologist* **206**,
196–208.
Wong S.C., Cowan I.R. & Farquhar G.D. (1978) Leaf Conductance in Relation to Assimilation
in *Eucalyptus pauciflora* Sieb. ex Spreng: influence of irradiance and partial pressure of
carbon dioxide. *Plant Physiology* **62**, 670–674.
Yamori W. & von Caemmerer S. (2009) Effect of rubisco activase deficiency on the temperature
response of CO_2 assimilation rate and rubisco activation state: Insights from transgenic
tobacco with reduced amounts of rubisco activase. *Plant Physiology* **151**, 2073–2082.
Yin X. (1993) Variation in foliar nitrogen concentration by forest type and climatic gradients in
North America. *Canadian journal of forest research* **23**, 1587–1602.
Zweifel R., Böhm J.P. & Häsler R. (2002) Midday stomatal closure in Norway spruce -
Reactions in the upper and lower crown. *Tree Physiology* **22**, 1125–1136.

957 TABLES

959 **Table 1.** Means ± 1 standard error for characteristics of trees from the Forest Tundra Ecotone
960 (FTE), Alaska (n=18) ,and Black Rock Forest (BRF), New York (n = 6), including diameter at
961 breast height (DBH; cm), tree height (m), dripline area (m^2), stems per hectare (means only, as
962 reported in Whitehead et al. (2004) for BRF and from Jensen pers. comm. for the FTE).

Location	DBH (cm)	Tree Height (m)	Dripline area (m^2)	Stems ha^{-1}
BRF	23.10 (1.99)	9.91 (0.73)	27.34 (2.00)	760
FTE	16.71 (0.94)	9.32 (0.57)	3.52 (0.37)	407

Parameter	INTERCEPTS		SLOPES		r_m^2	r_c^2
	BRF	FTE	BRF	FTE		
Photosynthesis						
A_{1500}	**9.282 (0.730)*** a**	**9.542 (1.389)*** a**	**0.066 (0.024)** a**	-0.075 (0.054)+ b	0.4	0.5
Φ	**0.055 (0.003)*** a**	**0.039 (0.004)*** b**	**-2e-04 (1e-04)** a**	-3e-04 (2e-04)+ a	0.67	0.82
LSP	**366.132 (29.038)*** a**	**540.489 (51.805)*** b**	**6.912 (0.855)*** a**	**4.185 (2.012)* a**	0.57	0.69
log(LCP)	**2.744 (0.116)*** a**	**4.170 (0.223)*** b**	**0.038 (0.004)*** a**	0.014 (0.009) b	0.75	0.79
V_{cmax}	**43.123 (3.494)*** a**	**68.472 (6.867)*** b**	**0.316 (0.112)** a**	**-0.796 (0.270)** b**	0.17	0.31
J_{max}	**73.760 (5.499)*** a**	**108.241 (10.866)*** b**	**0.531 (0.178)** a**	**-1.092 (0.427)* b**	0.17	0.31
Respiration						
R_D	**0.72 (0.211)** a**	**2.321 (0.411)*** b**	**0.055 (0.007)*** a**	0.006 (0.016) b	0.48	0.58
R_L	0.299 (0.228) a	**1.726 (0.403)*** b**	**0.055 (0.007)*** a**	0.020 (0.016) b	0.52	0.62
Leaf traits						
% Carbon	**47.498 (0.514)*** a**	**49.003 (0.630)*** a**	**0.020 (0.007)** a**	-0.015 (0.023) a	0.08	0.8
% Nitrogen	**1.064 (0.052)*** a**	**0.972 (0.095)*** a**	0.002 (0.001) a	4e-04 (0.004) a	0.17	0.53
C_{area}	**702.603 (58.824)*** a**	**1481.729 (139.897)*** b**	**13.332 (2.136)*** a**	2.254 (5.484) a	0.71	0.71
N_{area}	**15.084 (1.684)*** a**	**26.773 (2.842)*** b**	**0.349 (0.037)*** a**	0.176 (0.109) a	0.58	0.81
log(C:N)	**3.822 (0.053)*** a**	**3.9303 (0.100)*** a**	-0.002 (0.001) a	-9e-04 (0.004) a	0.18	0.54
SLA	**65.803 (2.053)*** a**	**34.393 (4.748)*** b**	**-0.676 (0.072)*** a**	-0.089 (0.186) b	0.76	0.77
Pigments						
Chl_{area}	**77.180 (5.705)*** a**	**89.363 (11.060)*** a**	0.325 (0.178)+ a	0.288 (0.407) a	0.21	0.44
$Chla_{area}$	**48.129 (3.576)*** a**	**54.266 (6.907)*** a**	**0.304 (0.111)* a**	0.367 (0.254) a	0.3	0.51
$Chlb_{area}$	**29.06 (2.353)*** a**	**35.186 (4.55)*** a**	0.021 (0.073) a	-0.082 (0.167) a	0.09	0.36
Car_{area}	**6.058 (0.905)*** a**	**10.722 (1.795)*** b**	**0.170 (0.030)*** a**	**0.160 (0.066)* a**	0.67	0.74
$Chl:Car_{area}$	**11.583 (0.461)*** a**	**7.973 (0.901)*** b**	**-0.116 (0.015)*** a**	-0.052 (0.033) a	0.69	0.77
$Chla:Chlb_{area}$	**1.673 (0.070)*** a**	**1.510 (0.128)*** a**	**0.009 (0.002)*** a**	**0.017 (0.005)*** a**	0.35	0.63
Spectral Indices						
PRI	**0.051 (0.006)*** a**	0.005 (0.015) b	**-9e-04 (2e-04)*** a**	8e-04 (6e-04) b	0.24	0.31

1013 Chla_{area}, area-based chlorophyll a content (μg cm^{-2}); Chlb_{area}, area-based chlorophyll b content
1014 (μg cm^{-2}); Car$_{area}$, area-based carotenoid content (μg cm^{-2}); Chl:Car$_{area}$, ratio of total chlorophyll
1015 content to carotenoids; Chl$a$:Chlb_{area}, ratio of chlorophyll a to chlorophyll b; PRI, photochemical
1016 reflectance index.

Table 3. Parameter estimates for the intercepts and slopes of linear mixed effects models with
tree as a random effect examining relationships between photosynthesis at 1500 µmol m^{-2}s^{-1}
(A_{1500}), maximum rate of carboxylation (V_{cmax}), respiration in the dark (R_D), respiration in the
light (R_L) or total pigment content (Chl+Car$_{area}$) and nitrogen per leaf area (N$_{area}$); as well as the
relationship between photochemical reflectance index (PRI) and the chlorophyll to carotenoid
ratio (Chl:Car$_{area}$). In all models tree was included as a random effect. Values in parentheses are
standard SEs. Bolded and starred values are those that are significantly different from zero
(***P<0.001; **P<0.01; *P<0.05; $^{+}$P<0.1). Significant differences between locations are
indicated by different letters (P<0.05). Marginal and conditional coefficients of determination
(r_m^2 and r_c^2, respectively) are also reported.

Model	INTERCEPTS		SLOPES		r_m^2	r_c^2
	BRF	FTE	BRF	FTE		
A_{1500} ~ N$_{area}$*location	**6.181 (1.250)*** a**	**8.130 (2.220)*** a**	**0.200 (0.049)*** a**	-0.024 (0.071) b	0.5	0.64
V_{cmax} ~ N$_{area}$*location	**28.828 (6.253)*** a**	**61.318 (12.106)*** b**	**0.934 (0.230)*** a**	-0.440 (0.383) b	0.15	0.55
R_D ~ N$_{area}$*location	-0.761 (0.410)$^{+}$ a	0.714 (0.620) a	**0.118 (0.017)*** a**	**0.055 (0.020)** b**	0.49	0.49
R_L ~ N$_{area}$*location	-0.908 (0.518)$^{+}$ a	0.386 (0.723) a	**0.113 (0.021)*** a**	**0.059 (0.023)* a**	0.42	0.42
(Chl+Car$_{area}$) ~ N$_{area}$*location	**62.309 (11.765)*** a**	**63.348 (22.613)** a**	**1.436 (0.501)** a**	**1.518 (0.712)* a**	0.39	0.52
PRI ~ Chl:Car$_{area}$*location	**-0.060 (0.015)*** a**	0.033 (0.023) b	**0.010 (0.002)*** a**	-0.0008 (0.003) b	0.43	0.54

1091
1092
Figure 1. (a) Average daily PPFD calculated from canopy photos over the study period (June-
July) at the Forest Tundra Ecotone (FTE), Alaska, and Black Rock Forest (BRF), New York.
Ambient (above canopy) PPFD is shown in orange. Boxplots show the median and first and
third quartiles. Whiskers display the range of groups with individual points representing outliers
falling outside 1.5 times the interquartile range. Different letters represent significant differences
between locations and canopy positions (P<0.05). (b) Ambient PPFD projected from canopy
photos during one day (July 4, 2017) at both locations. (c) Air temperature (°C) measured at
BRF and the FTE for June and July 2017. (d) Ambient PPFD measured from field instruments at
BRF and the FTE for June and July 2017. Green and blue shaded areas in 1c and 1d denote the
dates of the study campaigns at BRF and the FTE, respectively.

1103
Figure 2. Linear relationships between average daily photosynthetic photon flux density (PPFD)
over the measurement campaign and area-based foliar photosynthetic characteristics of white
spruce from the Forest Tundra Ecotone (FTE; blue), Alaska, and Black Rock Forest (BRF;
green), New York including photosynthesis at 1500 µmol m^{-2}s^{-1} (A_{1500}; 2a; FTE n = 38; BRF n =
33), apparent quantum yield (Φ; 2b; FTE n = 38; BRF n = 33), light saturation point (LSP; 2c;
FTE n = 38; BRF n = 33), light compensation point (LCP; 2d; FTE n = 38; BRF n = 33),
respiration in the dark (R_D; 2e; FTE n = 38; BRF n = 36) and respiration in the light (R_L; 2f; FTE
n = 37; BRF n = 29). Light and dark colors at each location (BRF or FTE) represent high and
low canopy positions, respectively. Parameter estimates of the linear mixed effects regression
models, and statistical differences between slopes and intercepts are presented in Tables 2, S1 &
S2. Regression lines are only shown for significant relationships (slope P<0.05). Also included
are boxplots by canopy position and location for each parameter. Different letters represent
significant differences between locations and canopy positions (P<0.05; Tables S1 & S2).
1117
1118
1119

1120
Figure 3. Linear relationships between average daily photosynthetic photon flux density (PPFD)
and foliar respiratory characteristics on an area basis of white spruce from the Forest Tundra
Ecotone (FTE; blue), Alaska, and Black Rock Forest (BRF; green), New York including the
maximum rate of carboxylation (V_{cmax}; 3a; FTE n = 39; BRF n = 37), and the maximum electron
transport rate (J_{max}; 3b; FTE n = 39; BRF n = 37). Light and dark colors at each location (BRF
or FTE) represent high and low canopy positions, respectively. Parameter estimates of the linear
mixed effects regression models, and statistical differences between slopes and intercepts are
presented in Tables 2, S1, S2. Regression lines are only shown for significant relationships
(slope P<0.05). Also included are boxplots by canopy position and location for each respiratory
parameter. Different letters represent significant differences between locations and canopy
positions (P<0.05; Tables S1 & S2).

1155
Figure 4. Linear relationships between average daily photosynthetic photon flux density (PPFD)
and leaf traits of white spruce from the Forest Tundra Ecotone (FTE; blue), Alaska, and Black
Rock Forest (BRF; green), New York including specific leaf area (SLA; 4a; FTE n = 37; BRF n
= 38), the ratio of chlorophyll *a* to chlorophyll *b* (Chl*a*:Chl*b*$_{area}$; 4b; FTE n = 25; BRF n = 22), %
nitrogen (4c; FTE n = 37; BRF n = 38), carotenoids (Car$_{area}$; 4d; FTE n = 25; BRF n = 22),
nitrogen per leaf area (N$_{area}$; 4e; FTE n = 37; BRF n = 38), and the ratio of chlorophylls *a* + *b* to
carotenoids (Chl:Car$_{area}$; 4c; FTE n = 25; BRF n = 22). Light and dark colors at each location
(BRF or FTE) represent high and low canopy positions, respectively. Parameter estimates of the
linear mixed effects models, and statistical differences between slopes and intercepts are
presented in Tables 2, S1 & S2. Regression lines are only shown for significant relationships
(slope P<0.05). Also included are boxplots by canopy position and location for each leaf trait.
Different letters represent significant differences between locations and canopy positions
(P<0.05; Tables S1 & S2).
1169
1170
1171
1172

1173
Figure 5. Linear relationships between photochemical reflectance index (PRI) and either
average daily photosynthetic photon flux density (PPFD) (5a; FTE n = 30; BRF n = 34) or the
ratio of chlorophylls *a* + *b* to carotenoids (Chl:Car$_{area}$; 5b) of white spruce from the Forest
Tundra Ecotone (FTE; blue), Alaska, and Black Rock Forest (BRF; green), New York. Light
and dark colors at each location (BRF or FTE) represent high and low canopy positions,
respectively. Parameter estimates of the linear mixed effects regression models, and statistical
differences between slopes and intercepts are presented in Tables 2, S1 & S2. Regression lines
are only shown for significant relationships (slope P<0.05). Also included are boxplots by
canopy position and location for PRI. Different letters represent significant differences between
locations and canopy positions (P<0.05; Tables S1 & S2).

1207
1208
Figure 6. Linear relationships between photosynthesis at 1500 µmol m^{-2}s^{-1} (A_{1500}; 6a), maximum
rate of carboxylation (V_{cmax}; 6b), respiration in the dark (R_D; 6c) and total pigment content
(Chl+Car$_{area}$; 6d) versus nitrogen per leaf area (N$_{area}$) of white spruce from the Forest Tundra
Ecotone (FTE; blue), Alaska and Black Rock Forest (BRF; green), New York. Light and dark
colors at each location (BRF or FTE) represent high and low canopy positions, respectively.
Parameter estimates of the linear mixed effects regression models, and statistical differences
between slopes and intercepts are presented in Tables 3 & S3. Regression lines are only shown
for significant relationships (slope P<0.05).

1231
1232
Figure 7. Ratios of (a) maximum rate of carboxylation to respiration at 25°C from Griffin *et al.*
(2022) (V_{cmax}/R_{25}) and (b) maximum electron transport rate to respiration at 25°C from Griffin *et*
al. (2022) (J_{max}/R_{25}) of white spruce from high and low canopy positions at the Forest Tundra
Ecotone (FTE, blue), Alaska, and Black Rock Forest (BRF, green), New York. Boxplots show
the median and first and third quartiles. Whiskers display the range of groups with individual
points representing outliers falling outside 1.5 times the interquartile range. Different letters
represent significant differences between locations and canopy positions (P<0.05; Tables S1 &
S2).

1246

1247

1248

1249

1250

1251

1252

1253

1254   SUPPLEMENTAL INFORMATION

1255

1256   **Methods S1**   *Linear mixed effects analysis of the relationships between physiological variables*

1257   *and N_{area} or PRI and $Chl:Car_{area}$*

1258

1259   The relationships between N_{area} and other physiological parameters by location presented in

1260   Table 3 and Figures 5 & 6 were assessed with a linear mixed effects model in which the

1261   interaction term allowed for the slope to vary by location (Black Rock Forest (BRF) or the Forest

1262   Tundra Ecotone (FTE)). Multiple measurements per tree were accounted for by including a

1263   random effect for each individual tree. The full model can be written as such:

$$Parameter_{t,i} = b0 + b1 * N_{area\,i} + b2 * BRF_i + b3 * N_{area\,i} * BRF_i + \gamma_t + e_i$$

1264   Where *i* represents each individual measurement taken, *t* represents each tree measured, BRF is a

1265   binary variable (e.g. BRF is 1 if the individual is from BRF, and 0 if it is not (i.e. from the FTE

1266   instead)), γ is a normally distributed random variable for different trees: $\gamma \sim N(0, \sigma)$, and *e* is the

1267   standard normally distributed random error term: $e \sim N(0, \sigma_e)$.

1268   This means that, for trees from the FTE, the equation is:

$$Parameter_{t,i} = b0 + b1 * N_{area\,i} + \gamma_t + e_i$$

1269   Whereas, for trees from BRF, the equation is:

$$Parameter_{t,i} = b0 + b1 * N_{area\,i} + b2 + b3 * N_{area\,i} + \gamma_t + e_i$$

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

Table S1. Summaries of the mixed effects models in which parameters were modeled as a
function of the interaction between canopy position and location with a random effect for each
individual tree. P-values are estimated using the Kenward-Roger approximation of the degrees of
freedom as suggested by (Halekoh & Højsgaard 2014). See table 2 for explanations of
abbreviations and units. P<0.05 are bolded.

	Canopy		Location		Canopy x Location	
	F-value	p-value	F-value	p-value	F-value	p-value
A_{1500}	0.14	0.7106	26.14	**<0.001**	13.28	**<0.001**
Φ	15.57	**<0.001**	60.94	**<0.001**	0.68	0.4129
LSP_{1500}	59.20	**<0.001**	27.38	**<0.001**	11.31	**0.0015**
log(LCP)	94.83	**<0.001**	139.98	**<0.001**	41.65	**<0.001**
V_{cmax}	1.35	0.2506	0.23	0.6381	23.37	**<0.001**
J_{max}	0.65	0.4233	1.00	0.3333	23.82	**<0.001**
R_D	55.17	**<0.001**	7.61	**0.0146**	46.38	**<0.001**
R_L	49.13	**<0.001**	10.09	**0.006**	25.33	**<0.001**
% Carbon	1.55	0.2194	1.41	0.2482	12.79	**<0.001**
% Nitrogen	0.09	0.7625	5.97	**0.0265**	1.88	0.1757
C_{area}	70.78	**<0.001**	148.02	**<0.001**	4.19	**0.0452**
N_{area}	77.16	**<0.001**	25.95	**<0.001**	19.83	**<0.001**
log(C:N)	0.05	0.8181	6.25	**0.0236**	1.58	0.2146
SLA	90.17	**<0.001**	134.52	**<0.001**	31.35	**<0.001**
Chl_{area}	2.68	0.1125	5.32	**0.0363**	0.02	0.8997
$Chla_{area}$	8.12	**0.0081**	6.27	**0.0246**	0.05	0.825
$Chlb_{area}$	0.10	0.7494	3.09	0.0997	0.0004	0.9843
Car_{area}	49.82	**<0.001**	22.34	**<0.001**	3.56	0.0703
$Chl{:}Car_{area}$	53.41	**<0.001**	31.14	**<0.001**	23.67	**<0.001**
$Chla{:}Chlb_{area}$	34.49	**<0.001**	0.37	0.5532	0.42	0.5213
PRI	7.37	**0.0094**	0.33	0.5751	20.94	**<0.001**
V_{cmax}/R_{25}	5.32	**0.0319**	91.24	**<0.001**	0.15	0.7041
J_{max}/R_{25}	3.68	0.0693	85.21	**<0.001**	0.60	0.4484

Table S2. Means ± 1 standard error for physiological, leaf and spectral traits. Several traits
report back-transformed means including the light compensation point (LCP) and the ratio of
carbon to nitrogen (C:N). Significant differences are marked by different letters (P < 0.05) with
differences between locations marked by capital letters and differences between location and
canopy positions marked by lowercase letters. See table 2 for explanations of abbreviations and
units.

	BRF	FTE	BRF low	BRF high	FTE low	FTE high
A_{1500}	10.8 (0.5) A	7.7 (0.4) B	9.8 (0.6) b	11.7 (0.6) a	8.3 (0.5) bc	7.0 (0.5) c
Φ	0.05 (0.002) A	0.03 (0.001) B	0.05 (0.002) a	0.05 (0.002) b	0.03 (0.002) c	0.03 (0.002) c
LSP_{1500}	522.0 (18.6) A	643.9 (14.1) B	427.4 (22.5) a	616.7 (22.2) bc	605.6 (17.8) b	682.2 (18.4) c
LCP	36.8 (2.1) A	90.7 (4.5) B	20.7 (1.6) a	65.1 (5.0) b	79.7 (5.3) b	103.1 (7.2) c
V_{cmax}	50.3 (2.2) A	48.5 (1.7) A	46.2 (2.7) ab	54.5 (2.7) c	54.7 (2.2) bc	42.3 (2.35) a
J_{max}	85.9 (3.4) A	80.8 (2.7) A	78.7 (4.1) ab	93.1 (4.2) c	90.1 (3.5) bc	71.5 (3.7) a
R_D	1.98 (0.14) A	2.46 (0.11) B	1.14 (0.17) a	2.82 (0.17) b	2.41 (0.13) b	2.5 (0.14) b
R_L	1.58 (0.16) A	2.22 (0.12) B	0.77 (0.19) a	2.38 (0.19) b	2.05 (0.14) b	2.39 (0.14) b
% Carbon	47.96 (0.50) A	48.63 (0.30) A	47.64 (0.51) a	48.27 (0.51) b	48.81 (0.32) ab	48.45 (0.32) ab
% Nitrogen	1.1 (0.04) A	0.98 (0.03) B	1.09 (0.05) ab	1.12 (0.05) b	1.00 (0.03) ab	0.96 (0.04) a
C_{area}	1007.2 (33.5) A	1547.8 (29.5) B	813.4 (42.8) a	1201.1 (42.1) b	1430.0 (38.2) c	1665.5 (40.9) d
N_{area}	23.1 (1.3) A	31.2 (0.9) B	18.3 (1.4) a	27.9 (1.4) bc	29.7 (1.0) b	32.7 (1.1) c
C:N	43.9 (1.8) A	49.9 (1.5) B	44.4 (2.0) ab	43.3 (2.0) a	48.9 (1.7) ab	50.9 (1.8) b
SLA	50.4 (1.2) A	32.0 (1.1) B	60.2 (1.5) d	40.5 (1.5) c	34.5 (1.4) b	29.5 (1.5) a
Chl_{area}	84.6 (3.9) A	96.0 (3.1) B	81.4 (4.8) a	87.7 (4.5) ab	93.3 (4.6) ab	98.7 (3.4) b
$Chla_{area}$	55.0 (2.5) A	62.9 (2.0) B	51.5 (3.0) a	58.4 (2.9) ab	60.0 (2.9) ab	65.9 (2.2) b
$Chlb_{area}$	29.6 (1.6) A	33.1 (1.3) A	29.8 (1.9) a	29.3 (1.8) a	33.3 (1.9) a	32.9 (1.4) a
Car_{area}	9.9 (0.8) A	14.5 (0.5) B	7.6 (0.9) a	12.1 (0.8) b	13.2 (0.7) b	15.7 (0.6) c
$Chl:Car_{area}$	9.0 (0.3) A	6.7 (0.2) B	10.7 (0.4) b	7.3 (0.3) a	7.0 (0.3) a	6.4 (0.3) a
$Chla:Chlb_{area}$	1.87 (0.05) A	1.92 (0.04) A	1.75 (0.06) a	2.00 (0.05) bc	1.81 (0.05) ab	2.02 (0.04) c
PRI	0.03 (0.004) A	0.02 (0.004) A	0.04 (0.005) b	0.02 (0.005) a	0.02 (0.005) a	0.03 (0.004) ab
V_{cmax}/R_{25}	70.2 (7.5) A	21.4 (1.4) B	23.4 (1.88) a	19.5 (1.67) a	61.7 (8.35) b	79.9 (10.81) b
J_{max}/R_{25}	119.9 (13.2) A	36.5 (2.4) B	38.7 (3.06) a	34.5 (2.98) a	105.8 (14.08) b	135.8 (18.07) b

1317   **Table S3.** Fixed effects parameter estimates and their significance for the linear mixed effects
1318   models assessing relationships between physiological variables and N_{area} or PRI and $Chl:Car_{area}$
1319   with an interaction term for location and a random effect for individual tree (taking into account
1320   multiple measurements taken on each tree at the high and low canopy positions). The full
1321   models can be seen in the Methods S1 available in the supplemental information.
1322

Model	b0	b1	b2	b3
A_{1500} ~ N_{area}*location	**0.0007**	0.7342	0.4477	**0.0118**
V_{cmax} ~ N_{area}*location	**0.00001**	0.2553	**0.0207**	**0.0031**
R_D ~ N_{area}*location	0.2539	**0.0060**	0.0516	**0.0190**
R_L ~ N_{area}*location	0.5956	**0.0131**	0.1514	0.0921
$(Chl+Car_{area})$ ~ N_{area}*location	**0.0080**	**0.0395**	0.9677	0.9253
PRI ~ $Chl:Car_{area}$*location	0.1590	0.8108	**0.0016**	**0.0073**

1328
1329