Title: Hyperinsulinemia acts through acinar cell insulin receptors to drive obesity-associated

pancreatic cancer initiation by promoting digestive enzyme production and inflammation

Authors: Anni M.Y. Zhang[1], Yi Han Xia[1], Jeffrey S.H. Lin[1], Ken H. Chu[1], Titine J.J. Ruiter[1], Jenny

C.C. Yang[1], Nan Chen[1], Justin Chhuor[1], Shilpa Patil[1], Haoning Howard Cen[1], Elizabeth J. Rideout[1],

Vincent R. Richard[3], David F. Schaeffer[2], Rene P. Zahedi[3,4,5], Christoph H. Borchers[3,6], James D.

Johnson[1] *, Janel L. Kopp[1] *

Affiliations: [1]Department of Cellular and Physiological Sciences, Life Sciences Institute, University of

British Columbia, Vancouver, Canada; [2]Department of Pathology and Laboratory Medicine, University

of British Columbia, Vancouver, Canada; [3]Segal Cancer Proteomics Centre, Lady Davis Institute,

Jewish General Hospital, McGill University, Montreal, Canada. [4]Department of Internal Medicine,

University of Manitoba, Winnipeg, Canada. [5]Manitoba Centre for Proteomics and Systems Biology,

Winnipeg, Canada. [6]Gerald Bronfman Department of Oncology, Jewish General Hospital, McGill

University, Montreal, Canada.

*Correspondence: janel.kopp@ubc.ca or james.d.johnson@ubc.ca

Keywords: pancreatic ductal adenocarcinoma, PanIN, hyperinsulinemia, obesity, insulin resistance

Funding: The project was supported by CIHR grants to E.J.R (PJT-153072), J.L.K. (MOP-142216 and

PJT-162239) and J.D.J. (PJT - 168854) and by a Lustgarten Foundation Therapeutics Focused

Research Program grant to J.D.J. and J.L.K. A.M.Y.Z was supported by Frederick Banting and

Charles Best Canada Graduate Scholarships and a Four-Year Fellowship from the University of British

Columbia. J.L.K was supported by a CIHR New Investigator Award (Msh-147794) and the MSFHR

Scholar Award (18309). E.J.R was supported by the CIHR Sex and Gender Chair Program (GSA-

171465) and the MSFHR Scholar Award (16876). C.H.B is grateful to Genome Canada for financial

support through the Genomics Technology Platform (GTP: 264PRO) and the Quebec Cancer

Consortium, Ministère de l'Économie et de l'Innovation, Quebec. C.H.B is also grateful for support

from the Segal McGill Chair in Molecular Oncology at McGill University (Montreal, Quebec, Canada),

and for support from the Warren Y. Soper Charitable Trust and the Alvin Segal Family Foundation to

the Jewish General Hospital (Montreal, Quebec, Canada).

Author Contributions: A.M.Y.Z designed, managed, and conducted the project including acquiring

and interpreting the majority of the data and she wrote the manuscript. Y.X performed the proteomics

and phospho-proteomics analyses. K.H.C and J.C provided technical assistance with pancreas

sectioning and staining (Alcian blue, H&E, Ck19, or GFP). J.S.H.L and T.J.J.R provided technical

assistance with dissection and genotyping and J.S.H.L performed the explant experiments. J.C.C.Y

provided technical assistance with dissection and Ck19 staining. N.C analysed staining (Sirius Red).

S.P assisted on dissection and sample preparation. H.H.C assisted with proteomics sample

preparation and proteomics analyses. E.J.R evaluated the manuscript and helped with interpretation

of the data in a sex-specific manner. V.R.R, R.P.Z and C.H.B co-designed and conducted proteomic

analyses. D.F.S provided histopathology analysis and expertise. C.H.B, J.L.K, and J.D.J supervised

the project, obtained funding, interpreted the data, and edited the manuscript. J.L.K is the guarantor of

this work.

Acknowledgements: The authors thank members from Johnson and Kopp labs for discussions. We

thank Christopher Wright (Vanderbilt, USA) for the *Ptf1a*CreER mice.

Conflict of interest: The authors report no relevant conflicts of interest.

Availability of data and materials: All data generated and analyzed in this study is included within

the article, additional files, or is available from the corresponding author on request.

Abstract

The rising incidence of pancreatic cancer is largely driven by the skyrocketing prevalence of obesity

and type 2 diabetes (T2D). Hyperinsulinemia is a cardinal feature of both conditions, and is

independently associated with increased cancer incidence and mortality. Our previous studies

demonstrated that genetically reducing insulin production suppressed formation of pancreatic

intraepithelial neoplasia (PanIN) pre-cancerous lesions in mice with mutant *Kras*. However, we found

that hyperinsulinemia affected many cell types in the pancreatic microenvironment. Thus, it remained

unclear whether hyperinsulinemia exerted its effects directly on the cells that give rise to PanINs or

indirectly on the tumor microenvironment, and molecular mechanisms involved were unknown. Here,

we tested whether insulin receptors (*Insr*) in $Kras^{G12D}$-expressing pancreatic acinar cells are necessary

for the effects of hyperinsulinemia on obesity-associated pancreatic cancer development. Loss of *Insr*

in $Kras^{G12D}$-expressing acinar cells did not prevent hyperinsulinemia or weight gain associated with

high fat diet (HFD) consumption in mice. However, solely reducing *Insr* in $Kras^{G12D}$-expressing acinar

cells significantly reduced formation of PanIN and tumors, in a gene dose-dependent manner.

Mechanistically, proteomic analyses showed that hyperinsulinemia acts through *Insr* to drive the

excess production of digestive enzymes in acinar cells by modulating the activity of the spliceosome,

ribosome, and secretory machinery. This resulted in increased inflammation, which was abrogated by

acinar-specific *Insr* knockout. We confirmed that insulin increased the conversion of wild-type acinar

cells into acinar-to-ductal metaplasia (ADM) in a trypsin-dependent manner. Collectively, these data

demonstrate that hyperinsulinemia acting via acinar cells insulin receptors promotes inflammatory

conditions that cooperate with Kras signaling to increase the risk of developing pancreatic cancer,

mechanistically linking obesity and pancreatic cancer.

Introduction

The 5-year survival rate of pancreatic ductal adenocarcinoma (PDAC) is less than 10% and it is

projected to become the 2nd leading cause of cancer death by 2030 (Rahib et al., 2014). Chronic

pancreatitis, family history, smoking, obesity, and T2D are risk factors for pancreatic cancer (Ilic and

Ilic, 2016). Obesity and T2D are usually accompanied by hyperinsulinemia, hyperglycemia, increased

inflammation, and dyslipidemia, which have all been proposed as underlying factors that drive the

increased PDAC morbidity and mortality in this patient population (Gallagher and LeRoith, 2020;

Zhang et al., 2021). Epidemiological studies consistently show that hyperinsulinemia is associated

with increased risk of developing PDAC and poorer survival (Pisani, 2008; Stolzenberg-Solomon et al.,

2005). Complementing clinical observations, our *in vivo* animal experiments demonstrated that

endogenous hyperinsulinemia causally contributes to PDAC development, independently of

hyperglycemia (Zhang et al., 2022; Zhang et al., 2019). Single-cell analysis revealed that

hyperinsulinemia altered gene expression in multiple cell types in the PanIN microenvironment (Zhang

et al., 2022), leaving open the question of whether the protective effects of reduced insulin production

are direct on the tumor precursor cells or whether they are mediated indirectly by local immune cells,

local fibroblasts, and/or via distant effects on adiposity (Mehran et al., 2012; Templeman et al., 2015;

Templeman et al., 2017). Consistent with a direct effect on the epithelium, insulin stimulates

proliferation in the PANC-1 and HPDE cell lines *in vitro* (Chan et al., 2014a), but this does not provide

information on the initiation of PDAC *in vivo*.

Insulin/IGF signaling, which includes KRAS/MAPK/ERK or PI3K/AKT/mTOR cascades, is

prominent in human and animal pancreatic cancer. Activating mutations in *KRAS* are detected in

~95% of PDAC clinical cases and induce PanIN pre-cancerous lesions and rare tumours in mice

(Waters and Der, 2018). Activating mutations in *PIK3CA* are also found in 3-5% of PDAC patients

(Heestand and Kurzrock, 2015; Janku et al., 2013) and can initiate PDAC in mice (Payne et al., 2015).

Additionally, *Pik3ca* silencing was protective in *Pdx1*-Cre;*Kras*[LSL-G12D];*Trp53*[LSL-R172H] mice (Sivaram et

al., 2019). Strategies that systemically reduce signaling downstream of Insr/Igf1r can suppress PDAC

(Hopkins et al., 2018; Payne et al., 2015), but they do not distinguish between the roles for insulin,

IGFs, or other upstream growth factors. Despite indirect evidence for an important role of insulin/IGF

signaling in this and other cancers, a direct and causal role for the insulin receptor alone has not been

demonstrated for any cancer. Insulin receptor protein is increased in some breast, prostate, and liver

cancers (Aljada et al., 2015; Belfiore et al., 2017; Law et al., 2008), but its role in the pancreas

remains enigmatic.

In this study, we tested the hypothesis that hyperinsulinemia-induced enhancement of PDAC

initiation is mediated through direct insulin receptor signaling in pancreatic acinar cells. We found mice

that consumed a high fat diet had a significant reduction in PanIN and tumor development when they

lacked *Insr* specifically in Kras^{G12D}-expressing acinar cells. These findings indicate that

hyperinsulinemia directly contributes to pancreatic cancer initiation through Insr in acinar cells via a

mechanism that involves increased production of digestive enzymes and subsequent pancreatic

inflammation.

Results

Effects of acinar cell-specific *Insr* loss on body weight and glucose homeostasis

We first generated *Ptf1a*^{CreER};*Insr*$^{w/w}$;nTnG or *Ptf1a*^{CreER};*Insr*$^{f/f}$;nTnG mice to assess the baseline

roles of *Insr* in acinar cells (Figure 1A). All mice were fed with HFD after 3 weeks of age to sustain

hyperinsulinemia. Tamoxifen was injected at 4 weeks of age to induce recombination of the *Insr*^{flox}

alleles and a Cre-dependent nTnG allele (Muzumdar et al., 2007), specifically in acinar cells (Figure

1B). As expected, mice consuming HFD gained weight over time. We did not find a significant

difference in body weight between male *Ptf1a*^{CreER};*Insr*$^{f/f}$;nTnG mice and *Ptf1a*^{CreER};*Insr*$^{w/w}$;nTnG mice

but observed a significant and consistent reduction in body weight in female *Ptf1a*^{CreER};*Insr*$^{f/f}$;nTnG

mice compared to *Ptf1a*^{CreER};*Insr*$^{w/w}$;nTnG mice (Figures 1C-D), perhaps as a result of a minor

pancreatic insufficiency (see below). Fasting glucose was not different between genotypes (Figures

1E-F). HFD induced similar levels of hyperinsulinemia in both genotypes (Figures 1G-H) and males,

as previously reported (Zhang et al., 2022; Zhang et al., 2019), had higher overall levels of insulin

compared to females on HFD (Figures 1G-H). These insulin levels are well above the expected fasting

insulin levels of C57Bl6J mice on a chow diet (138 +/- 24 pM) (Berglund et al., 2008) and higher than

we have previously reported in mice with reduced insulin gene dosage (Zhang et al., 2022; Zhang et

al., 2019). Pancreas weight of *Ptf1a*^{CreER};*Insr*$^{w/w}$;nTnG and *Ptf1a*^{CreER};*Insr*$^{f/f}$;nTnG mice were not

different at 12 weeks of age (Figure 1I). These data demonstrate that mice with acinar-specific *Insr*

deletion having wild type-like pancreata.

Body weight and glucose homeostasis in mice with acinar-specific *Insr* loss and Kras mutation

To test our primary hypothesis, that hyperinsulinemia drives pancreatic cancer development via

Insr cell autonomously, we generated mouse models in which $Kras^{G12D}$ expression (Kopp et al., 2012)

and loss of *Insr* were both induced in acinar cells. Our cohorts contained mice with full *Insr* gene

dosage, $Ptf1a^{CreER};Kras^{LSL-G12D};Insr^{w/w};$nTnG (PK-$Insr^{w/w}$); mice with reduced *Insr* , $Ptf1a^{CreER};Kras^{LSL-}$

$^{G12D};Insr^{w/f};$nTnG (PK-$Insr^{w/f}$); or mice without *Insr*, $Ptf1a^{CreER};Kras^{LSL-G12D};Insr^{f/f};$nTnG (PK-$Insr^{f/f}$) in

$Kras^{G12D}$-expressing acinar cells (Figure 2A). In this model, tamoxifen induced simultaneous acinar-

specific recombination to express mutant $Kras^{G12D}$, a nTnG, and deletion of $Insr^{flox}$ alleles. Multiple

studies have shown that HFD accelerates PanIN and PDAC development (Chang et al., 2017;

Dawson et al., 2013). *Insr* loss in acinar cells did not significantly affect the weight gain of male or

female mice in the context of mutant *Kras* (Figures 2B-C). Fasting glucose and insulin levels were also

similar between groups (Figures 2D-G). Thus, our model enabled us to test the role of acinar cell *Insr*

in the context of intact hyperinsulinemia and normoglycemia, which normally occurs in the obese

state.

Loss of *Insr* in $Kras^{G12D}$-expressing acinar cells reduced PanIN development

To test whether *Insr* loss affected $Kras^{G12D}$-mediated PDAC formation from acinar cells, we

examined a cohort of 9-12 PK-$Insr^{w/w}$, PK-$Insr^{w/f}$, and PK-$Insr^{f/f}$ mice for each sex and genotype and

planned to assess lesions at ~1 year of age based on our previous work (Zhang et al., 2022; Zhang et

al., 2019). However, half the male PK-$Insr^{w/w}$ mice and a few female PK-$Insr^{w/w}$ mice reached humane

endpoint earlier than we had expected (Figure 3A). This suggested that HFD promoted tumor

development in mice with full insulin gene dosage earlier than in mice with reduced insulin (Zhang et

al., 2022; Zhang et al., 2019). Through necropsy we noted that macroscopic tumors were present in

male PK-$Insr^{w/w}$ mice reaching humane endpoint (Figure 3A). Additionally, the remaining male PK-

Insr^{w/w} mice had PDAC or the parenchyma was almost completely replaced by ductal metaplasia

comprising cysts, PanIN, and ADM at 43.5 weeks of age (Figure 3A-B). Three male PK-*Insr*^{w/f} mice

also reached human endpoint prior to 43.5 weeks, but tumors were not observed at necropsy (Figure

3A). Finally, no male PK-*Insr*^{f/f} mice reached humane endpoint by 43.5 weeks of age (Figure 3A).

PDAC was noted by histology in 40-50% of PK-*Insr*^{w/f} mice and PK-*Insr*^{f/f} mice (Figures 3A-C). In sum,

male PK-*Insr*^{f/f} mice lived longer than PK-*Insr*^{w/f} mice and PK-*Insr*^{w/w} mice and had more normal

parenchyma left at 40.5 weeks of age suggesting that loss of *Insr* in Kras^{G12D}-expressing acinar cells

limits HFD-mediated promotion of acinar cell transformation.

As noted above, a few female PK-*Insr*^{w/w} mice, as well as a few PK-*Insr*^{w/f} mice also reached

humane endpoint prior to 43.5 weeks, but no macroscopic tumors were noted at necropsy (Figure 3A).

No female PK-*Insr*^{f/f} mice reached humane endpoint by 43.5 weeks of age. Histologically, we found

that the incidence of PDAC in females was *Insr* dosage-dependent (Figures 3A-C). The majority of

female mice from every genotype, retained some normal parenchyma (Figures 3B-C). This suggested

that the extent of disease in females at 10 months of age was different than males and that female

mice had not yet maximally disrupted normal parenchyma function. This is consistent with previous

reports suggesting that the timing and/or extent of lesion formation between male and female mice in

the context of HFD may differ (Chang et al., 2017). Altogether, these data strongly suggested that

limiting or eliminating insulin/Insr activity specifically in acinar cells in male and female mice reduced

the propensity of HFD and Kras activation to transform the pancreas.

We next quantified the extent of the tissue disruption and lesion formation at 43.5 weeks of age to

assess the extent of disease present in the presence or absence of *Insr*. In parallel with PanIN

quantification between groups, we stained for nuclear GFP from the nTnG lineage reporter allele

(Muzumdar et al., 2007) in all pancreata. As expected, most acinar cells and PanIN lesions were GFP

positive (Figures S1A-B) confirming that the lesions arose from Ptf1a⁺ acinar cells. Rare mice with a

GFP labeling efficiency of acinar cells below 20% were excluded from further analysis. Pancreata from

Ptf1a^{CreER};*Insr*^{w/w};nTnG and *Ptf1a*^{CreER};*Insr*^{f/f};nTnG mice were similarly comprised of acinar cells and

endocrine islets and by IHC had GFP expression widely present in acinar cells (Figures S1C-F),

consistent with their *Kras* wild-type genotype. In contrast, pancreata from male and female PK-*Insr*$^{w/w}$,

PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$ mice all contained ductal lesions with histological characteristics of

metaplastic ducts, including ADM, low-grade and high-grade PanIN, and sometimes PDAC (see

above) (Figures 3B-C). When we quantified the PanIN and/or tumor area, we found that reducing *Insr*

in Kras^{G12D}-expressing acinar cells reduced the area of PanINs plus tumors, PanIN alone, or tumor

area in a dose dependent manner in males and females (Figures 3D-F). Consistent with this

histological-based quantification, measuring the pancreatic area containing ductal metaplasia (ducts,

ADM, PanIN and PDAC) or mucinous lesions (PanIN and some tumors) with Ck19 or Alcian blue

staining, respectively, similarly showed that female PK-*Insr*$^{w/w}$ mice formed significantly more lesions

than female PK-*Insr*$^{f/f}$ mice (Figures S2A-D). Notably, male PK-*Insr*$^{w/w}$ mice had a higher Ck19$^+$, but a

similar amount of Alcian blue$^+$ area compared to other genotypes. This latter observation is likely

explained by the presence of Alcian blue negative high-grade PanIN lesions and large cysts with

predominantly normal ductal epithelium which were more prevalent in male than female, PK-*Insr*$^{w/w}$

pancreata (Figure 3B and Figures S2A, S2C, S2E-F). Indeed, most male PK-*Insr*$^{w/w}$ and some female

PK-*Insr*$^{w/w}$ pancreata were comprised of almost all Ck19$^+$ area with little acinar cell area left, while

larger areas of normal acinar cells correlated with lower Ck19$^+$ areas in PK-*Insr*$^{w/f}$ and PK-*Insr*$^{f/f}$ mice

(Figures 3B-C, Figure 3G and Figures S2G-H). Therefore, our data strongly suggested that acinar cell

Insr dose-dependently regulates oncogenic Kras-induced PanIN initiation in the context of diet-

induced hyperinsulinemia.

Loss of *Insr* in *Kras*^{G12D}-expressing acinar cells reduced PanIN initiation

To examine whether PanIN initiation was specifically affected by loss of *Insr* in acinar cells, we

examined 12-week-old male and female mice of each genotype. At 8 weeks post-tamoxifen injection,

acinar cells and PanIN lesions were also GFP positive indicating good recombination efficiency

(Figures S3A-B). There were no significant differences in pancreas weight between PK-*Insr*$^{w/w}$, PK-

Insr$^{w/f}$, and PK-*Insr*$^{f/f}$ mice in either sex (Figure 4A). This is consistent with previous data showing that

the pancreas was largely normal in young *Ptf1a*^{CreER};*Kras*$^{LSL-G12D}$ mice (Kopp et al., 2012). Pancreata

from PK-*Insr*[f/f] mice were predominantly normal with a small number of ductal lesions having

characteristics of ADM or low-grade PanIN (Figures 4B-D). In contrast, significantly more of the

pancreas was occupied by PanIN in PK-*Insr*[w/w] mice (Figures 4B-D). The significant inhibition of PanIN

formation solely through reducing *Insr* in acinar cells strongly supports a model in which the primary

mechanism by which diet-induced hyperinsulinemia promotes tumor initiation is through insulin

receptors on acinar cells.

Proteomic and phospho-proteomic analyses of pancreata from mice lacking acinar *Insr*

To investigate the underlying molecular mechanisms of *Insr* action in PanIN initiation, we

conducted unbiased total and phospho-proteomic analyses using the head of the pancreas from the

12-week-old female mice (Figure 4) with and without Kras[G12D] and/or *Insr*. For the total proteome

dataset, we obtained reliable, quantitative data on 2889 proteins across all the samples. Consistent

with our histological analyses identifying a change in PanIN, we found that the inflammation- and

PanIN-associated proteins, Reg3a, Reg3b, Reg2, Tff1, Gkn1, and Gkn2 (Chen et al., 2019; Li et al.,

2016; Steiner et al., 2022), were among the 124 proteins significantly enriched in PK-*Insr*[w/w] compared

to *Ptf1a*[CreER];*Insr*[w/w];nTnG pancreata (Figure 5A and Table S1). We also found that some proteins

associated with acinar cell function, such as Ctrc, Dbi, Pla2g1b, and Clps, were among the 122

proteins significantly down-regulated as a consequence of *Kras*[G12D] expression in acinar cells (Figure

5A). However, the majority of proteins associated with acinar cell function, like Cpa1, Spink1, and

Cela1, remain unchanged between the PK-*Insr*[w/w] and *Ptf1a*[CreER];*Insr*[w/w];nTnG genotypes (Table S1).

Importantly, we compared the differentially expressed pathways between PK-*Insr*[w/w] and

Ptf1a[CreER];*Insr*[w/w];nTnG mice to differentially expressed pathways between human PDAC and normal

pancreas tissues adjacent to human PDAC (Cao et al., 2021) (Figures S4A-B). We found that most of

the pathways were significantly enriched in both mouse and human datasets, which suggested our

proteomic and phospho-proteomic results were relevant to human samples. Interestingly, we also

found 92 proteins increased in abundance and 155 proteins had decreased abundance solely due to

loss of *Insr* in Kras wild-type acinar cells (Figure 5B), with the caveat that we analyzed a low sample

number. Nevertheless, reductions in Ctrc, Clps, Pla2g1b suggested that *Insr* may have a role in

regulating the function of wild-type acinar cells in mice fed HFD (Figure 5C).

To define the molecular mechanisms associated with *Insr* deletion in the context of mutant Kras, we

focused further analyses on comparing the PK-*Insr*$^{f/f}$ pancreata to the PK-*Insr*$^{w/w}$ pancreata. We found

that 135 proteins were enriched and 117 were depleted in PK-*Insr*$^{f/f}$ mice compared to PK-*Insr*$^{w/w}$

controls (Figure 5C). We then used these differentially expressed genes to perform a K-means

clustering analysis with the differentially abundant genes between PK-*Insr*$^{w/w}$ and PK-*Insr*$^{f/f}$ mice using

the protein abundance values in all genotypes to groups of proteins that varied by Insr status, Kras

mutation, or both (Figure 5D). As expected from our histological analyses, proteins associated with

PanIN initiation or formation in PK-*Insr*$^{w/w}$ compared to *Ptf1a*^{CreER};*Insr*$^{w/w}$;nTnG pancreata, such as

Reg3a, Reg3b, Reg2, Tff1, Gkn1, and Gkn2, (Figure 5A and 5D, cluster 1), were reduced in PK-*Insr*$^{f/f}$

mice (Figure 5C, Figure 5D, cluster 1 and Table S4). In order to investigate potential functional

enrichment of protein groups based on cell signaling, intracellular localization, and biological process,

we used the differentially abundant proteins between PK-*Insr*$^{w/w}$ and PK-*Insr*$^{f/f}$ mice (Adj. p<0.05) to

build protein-protein interaction networks using STRING (Szklarczyk et al., 2021) and assigned these

proteins to their intracellular organelle locations in a diagram using the COMPARTMENTS section of

GeneCards and/or existing knowledge of their function (Binder et al., 2014) (Figure 6A). This data

depiction (Figure 5D, cluster 2 and 3) highlighted the striking downregulation of the majority of the

proteins packaged into zymogen granules for secretion in PK-*Insr*$^{f/f}$ compared to

Ptf1a^{CreER};*Insr*$^{w/w}$;nTnG and PK-*Insr*$^{w/w}$ mice (Figure 6A-B). Acinar cells produce a large amount of

protein every day, and in *Ptf1a*^{CreER};*Insr*$^{w/w}$;nTnG mice fed HFD, 18.7% of the peptides detected in our

analyses were associated with the zymogen granules and this was similar in *PK-Insr*$^{w/w}$ pancreata

(18.7%) (Figure 6C). However, this percentage was reduced significantly to 12.2% in PK-*Insr*$^{f/f}$

pancreata, as well as to 16% in *Ptf1a*^{CreER};*Insr*$^{f/f}$;nTnG pancreata (Figure 6C). This reduction in

zymogen proteins is likely an underestimate of the effect on the total proteome, as our differential

abundance analyses were normalized to total protein content. Thus, loss of *Insr* in acinar cells results

in a coordinated decrease in the amount of digestive enzyme produced by acinar cells in the context

of HFD.

Pancreatic acinar cells represent the cellular majority in our bulk tissue proteome. With the caveat

that our bulk tissue includes multiple cell types, our STRING network analysis nevertheless highlighted

potential mechanisms downstream of *Insr* loss that could underlie this dramatic reduction in zymogen

granule content in acinar cells. This included increases in proteins with key roles in suppressing

protein synthesis at the ribosome (eEF2K) and suppressing insulin signaling (Ptpn1), as well as

increases in the first enzyme of the fatty acid beta oxidation pathway (Acox1), a protein regulating

lysosome function (Grn), and protein processing machinery at the ER and Golgi (Ilvbl, Ssr3, Ssr4,

Irag2). Many components of mitochondrial electron transport complexes were changed, as well as

endosomal and cytoskeletal proteins, many of which are involved in moving organelles within the cells

or in exocytosis, such as Snx5, Vps35l, Sycn (Figure 6A). Finally, we also found decreases in critical

parts of the spliceosome (Snrpa), the signal recognition particle complex receptor (Srprb), and

components of the large ribosome (Rpl29, Rpl36, Rpl37a). Altogether, these observations suggested

that loss of *Insr* in acinar cells and a reduction in PanIN formation is associated with a reduction in

synthesis of digestive enzymes.

Parallel phosphoproteomic analysis on the same samples confirmed and extended the findings

from our total proteomic analysis (Figure 7A-D). We identified 225 downregulated phospho-peptides

and 177 upregulated phospho-peptides by comparing the PK-*Insr*$^{f/f}$ pancreas to PK-*Insr*$^{w/w}$ controls

(Figure 7A). In general, statistically significant phospho-peptide differences were not due to underlying

differences in total protein abundance (Figure 7B). Using a similar strategy as above, we mapped the

function of the proteins to compartments and processes in the cell (Figure 7D). We found significant

decreases in phosphorylation in PK-*Insr*$^{f/f}$ compared PK-*Insr*$^{w/w}$ pancreata for proteins involved in

transcription elongation (Eloa, Top2a, Supt5h), mRNA splicing and nuclear speckle formation (Srpk1,

many Srsf proteins, Cherp, Srrm1 and Srrm2), as well as protein translation initiation (Eif4b, Eif5b, and

eIF3 complex proteins) and elongation (Eef1b, Eef1d, and many ribosomal subunits). There were also

decreases in phospho-peptides for Cavin3, which has been implicated in Akt-Erk signaling bias

(Haoning Howard Cen, 2022; Hernandez et al., 2013), and PP1 inhibitor Ppp1r2, at sites known to be

influenced by insulin signaling (Figure 7D). Finally, there was reduced phosphorylation of Larp1 at

sites known to be regulated by Raptor and Rictor. Larp1 is an RNA-binding protein that links mTorc1

to the regulation of terminal oligopyrimidine tract (TOP) mRNA that encodes for ribosomal proteins

and elongation factors (Hong et al., 2017). Altogether, our proteomics data suggested that Insulin/Insr

promotes the production of proteins in acinar cells in part through its modulation of Larp1 and many

other components controlling transcription, translation, and secretion of digestive enzymes.

Interestingly, multiple sites on the cholecystokinin (Cck) receptor (Cckar) also showed altered

phosphorylation (Figure 7C-D), including sites in the main intracellular loop that are indicative of

ligand-induced desensitization (Rao et al., 2000) and phospho-sites in the C-terminal tail that have not

been previously reported. Collectively, these unbiased total and phospho-proteomic studies delineate

the possible molecular mechanisms by which hyperinsulinemia, acting via the Insr, may promote Kras-

driven pancreatic cancer initiation.

Our histological analyses, as well as our total proteomics and phosphoproteomics suggested that

loss of the *Insr* in acinar cells prevented the formation of PanIN and prevented the induction of genes

associated with pancreatic injury, such as Reg3a, Reg3b, and Reg2. To examine whether loss of *Insr*

in acinar cells prevented the injury and/or fibrosis typically associated with Kras^{G12D} activation and

PanIN formation, we performed Sirius Red staining on 12-week-old pancreata (Figures 8A-B). In PK-

Insr$^{w/w}$ control pancreata, Sirius Red staining surrounded the metaplastic ducts, containing ADM and

PanIN, as well as the surrounding acini in nearby lobules (Figures 8A-B). In contrast, in the PK-*Insr*$^{f/f}$

pancreata, the metaplasia that did form tended to be associated with less Sirius Red staining and less

staining in the neighboring lobules (Figures 8A-C). In male PK-*Insr*$^{w/w}$ pancreata, the extent of Sirius

Red staining significantly correlated with the amount of PanINs formed (Figure S5A). In female PK-

Insr$^{w/w}$ pancreata, there were higher than expected amounts of Sirius Red given the amount of PanIN

observed, but the relationship between the values was still strong (Figure S5B). However, in the

absence of *Insr* very few PanIN formed in both sexes and this was associated with a lower amount of

Sirius Red staining, especially in females. This supported our hypothesis that inflammation was

reduced in the absence of *Insr* in Kras^{G12D}-expressing acinar cells.

Our proteomics data suggested that fewer digestive enzymes were present in the absence of Insr

in acinar cells when Kras^{G12D} expression was also present (Figure 6). Given that autoactivation of

trypsinogen in acinar cells or in the pancreatic parenchyma contributes, at least in part, to the

induction of pancreatitis (Smith and Solomon, 2014), we reasoned that the reduced presence of

enzymes in the absence of *Insr* could result in less tissue damage spreading into other lobules. This

would result in decreased ADM and PanIN formation, as in the PK-*Insr*$^{f/f}$ pancreata. To test this

hypothesis, we utilized an 3D *ex vivo* model of ADM formation from wild-type acinar cells (Figure 8D)

(Fleming Martinez and Storz, 2019; Means et al., 2005a). Wild-type acinar clusters grown in collagen

for 5 days maintained their acinus morphology, as previously shown (Means et al., 2005a). In addition,

treatment with TGF-α induced these clusters to form a duct-like lumen structure (Figure 8E) (Means et

al., 2005b). Addition of increasing concentrations of insulin alone had only a modest effect on acini

clusters (Figure 8E-F). However, increasing concentrations of insulin in combination with TGF-α

induced ring formation significantly (Figure 8E-F). Remarkably, the synergism between insulin and

TGF-α was significantly reduced by the presence of trypsin inhibitor in the culture media (Figure 8F).

Because acinar cells are the source of trypsinogen and this pro-enzyme is prone to autoactivation, our

data suggested that hyperinsulinemia-mediated promotion of enzyme production in acinar cells can

result in increased inflammation and promote initiation of PanIN lesions from acinar cells sustaining

activating mutations in *Kras* (Figure 8G).

Discussion

The purpose of this study was to test the hypothesis that hyperinsulinemia in the obese state acts

directly on acinar cells to promote pancreatic cancer initiation. Our data clearly support a model where

Insr in acinar cells plays a causal role in supporting cancer initiation in the context of diet-induced

obesity and mutant Kras. The contribution of direct insulin action on acinar cells during initiation from

normal cells explains a large portion of the effects of HFD, but our results do not preclude roles for Insr

in other local or distant cell types (Zhang et al., 2022) and at other times during the progression of the

initial precursor lesions to PDAC. Our unbiased proteomic and phospho-proteomic analyses led us to

propose a model in which HFD and hyperinsulinemia through Insr in acinar cells promote the

physiological function of acinar cells to supply digestive enzymes to breakdown lipid rich food in the

duodenum. However, the increased presence of enzymes increases the risk of autoactivated trypsin,

as well. This increased propensity for trypsin-induced injury would result in sub-clinical levels of

inflammation and acinar-to-ductal metaplasia. Thus, the physiological response of pancreas to a diet

rich in fat results in increased insulin production from beta cells, which further promotes an increased

enzyme production capacity in acinar cells. However, in the context of *Kras* mutations, this increased

production can lead to increased chances of inflammation that enhance Kras^{G12D} signaling and

promote an irreversible transformation in Kras^{G12D}-expressing acinar cells. Thus, our studies provide a

key missing link explaining the connection between obesity and hyperinsulinemia and increased

pancreatic inflammation and PanIN initiation.

Diet-induced obesity induces insulin hypersecretion, increases beta cell mass and impairs insulin

clearance, resulting in sustained hyperinsulinemia (Mehran et al., 2012; Zhang et al., 2021). Mice

used in this study, which had the full complement of all 4 insulin alleles (*Ins1* and *Ins2*), exhibited

higher fasting insulin levels (males 1000-1500 pmol/L; females 500 pmol/L) than our previous models

with insulin gene dosage reduced to 1 or 2 copies (males, 400-800 pmol/L; females 100-200 pmol/L)

(Zhang et al., 2022; Zhang et al., 2019). Consistent with our working model that endogenous

hyperinsulinemia contributes to pancreatic cancer development, more than 90% of the normal

pancreatic area in the PK-*Insr*$^{w/w}$ mice in this study was replaced by ductal metaplasia, PanIN, and

tumors at 10 months of age. Notably, 1-year-old PK-*Ins1*$^{+/+}$;*Ins2*$^{-/-}$ control mice had just ~25% of the

pancreas replaced by PanINs, while PK-*Ins1*$^{-/-}$;*Ins2*$^{+/+}$ mice were at ~1-4% by 1 year of age (Zhang et

al., 2022; Zhang et al., 2019). If only considering PDAC, ~70% of female PK-*Insr*$^{w/w}$ mice developed

PDAC in this study at 10 months of age, while only 1 female PK-*Ins1*$^{+/+}$;*Ins2*$^{-/-}$ mouse developed PDAC

after a year (Zhang et al., 2022; Zhang et al., 2019). Together, these findings suggest that simply

reducing insulin production limits PanIN initiation in the context of obesity.

All cells in the body have insulin receptors and require insulin signaling for key functions, including

nutrient uptake for storage and anabolism. While the roles of Insr in hepatocytes, myocytes, and

adipocytes are well studied (Cen et al., 2022; Cherrington, 2005; Haeusler et al., 2018), the

consequences of *Insr* loss in pancreatic acinar cells remain understudied. In this study, we specifically

deleted the *Insr* gene from acinar cells using the *Ptf1a*[CreER] allele and showed that insulin insensitivity

in acinar cells had no obvious effects on systemic glucose homeostasis or serum insulin levels

regardless of the *Kras* gene status. This suggests that the systemic regulation of glucose and insulin

homeostasis are similarly perturbed in our mice fed HFD. Our data, combined with our previous

studies (Zhang et al., 2022; Zhang et al., 2019), effectively rule out an essential role for hyperglycemia

in PanIN and PDAC formation, but do not preclude potentially important roles for hyperglycemia in the

later stages of disease (Sato et al., 2020; Vaziri-Gohar et al., 2022).

Insulin receptor signaling activates PI3K/AKT/mTOR and MAPK/ERK signaling cascades, both of

which have mitogenic effects. Effector proteins in these two pathways are frequently mutated during

tumorigenesis (Guo et al., 2020; Samuels et al., 2004; Yang et al., 2019), including in pancreatic

cancer. Indeed, activating mutations in Kras, a key mediator of insulin and insulin-like growth factor

signaling, drive the vast majority of pancreatic cancers (Waters and Der, 2018). Previous *in vitro*

evidence supported the concept that hyperinsulinemia could promote cancer cell growth through over-

activating the signaling cascades downstream of Insr protein (Chan et al., 2014b; Gallagher and

LeRoith, 2020; Godsland, 2009; Zhang et al., 2021). Our findings are the first *in vivo* studies in any

cancer type to demonstrate a direct causal role for hyperinsulinemia in the cancer cell of origin.

Mechanistically, our proteomic data demonstrated that *Insr* loss in acinar cells results in the

coordinated reduction in digestive enzymes, with or without mutant Kras. This indicates that insulin

signaling normally supports exocrine function. The reduced body weight in mice lacking acinar-cell

Insr is also consistent with sub-clinical pancreatic insufficiency and a relative reduction in the ability to

utilize ingested nutrients. Our observations are consistent with previous studies, including the

observation that amylase production is diminished by β-cell ablation using streptozotocin and restored

with insulin injection (Frier et al., 1976; Henderson et al., 1981; Soling and Unger, 1972). Mutations

that result in increased trypsin activity in the pancreas (Chang et al., 2017), as well as animal studies

using the Cck analog, caerulein, to stimulate enzyme secretion at supraphysiological levels (Guerra et

al., 2007), have demonstrated that tight control of digestive enzyme function reduces the risk of

forming PDAC. Interestingly, our phospho-proteomic data identified hyper-phosphorylation of the Ccka

receptor in Insr-knockout pancreas, linking local insulin signaling to Cck, a key endogenous regulator

of acinar cell function and pancreatic weight. Cck secretion from I-cells is stimulated by the presence

of amino acids and fat, which induces the highest I-cell secretion responses, in the small intestine

(Otsuki, 2000). Cck affects pancreatic secretions predominately through local action on nerve fibres in

the gut to induce a vago-vagal reflex circuit, resulting in acetylcholine secretion from nerves in the

pancreas. Alternatively, it also can act directly on murine acinar cells through the Ccka receptor. In

addition to mediating digestive enzyme release from acinar cells, previous studies have also shown

that injecting low levels of Cck into mice results in increased pancreatic size over time (Dembinski and

Johnson, 1980; Varga et al., 1988). This suggests that Cck helps regulate the enzyme production

capacity of the pancreas. Finally, it has been reported that HFD-associated inflammation can also

promote islet Cck expression, which was proposed to play a role in obesity-associated PanIN

formation (Incio et al., 2016). However, our data indicate that hyperinsulinemia, acting through Insr, is

the upstream driver of diet-induced inflammation via hyperactive digestive enzyme production. This

supports a working model whereby hyperinsulinemia promotes PanIN initiation via increased local

inflammation associated with elevated digestive enzyme production and/or release (Figure 8G). *Insr*

loss in acinar cells counters the increased signal for acinar cell enzyme production or cell proliferation

induced by HFD. Further studies are needed to fully understand the impact of dietary content on

acinar cells and their susceptibility to Kras^{G12D}-mediated transformation.

Our proteomic analyses revealed other key mechanisms associated with suppressed pancreatic

cancer initiation from acinar cells lacking *Insr*. For example, mRNA splicing and translation factors,

known targets of insulin signaling (Haeusler et al., 2018), are differentially abundant in the pancreas

after *Insr* loss in Kras^{G12D}-expressing acinar cells, suggesting possible mechanisms by which insulin

might regulate the production of digestive enzymes. We also found evidence that reduced insulin

signaling affected cellular metabolism. Previous studies have suggested that insulin can promote

glycolysis in wildtype acinar cells to protect them during pancreatitis (Bruce et al., 2021), however,

detailed analyses of cell-type-specific mechanisms await single-cell cell proteomic and single-cell

metabolomic characterization of this model.

We acknowledge the limitations of our study. One limitation of our study, and the field, is that

existing antibody reagents are not specific enough to perform accurate anti-Insr staining that would

allow us to determine whether acinar cells with normal morphology had escaped recombination at the

Insr floxed allele. Another limitation is that, while our proteomic analysis provided quantification of

2889 protein abundances and 8787 phospho-sites, coverage of the proteome was not complete and

biased against membrane proteins. For example, analysis of the abundance and phosphorylation of

Igf1r, a protein that could compensate in *Insr* knockout cells, will require targeted assays and/or

membrane fractionation. Insulin can bind to homodimeric Igf1 receptors and heterodimeric Insr/Igf1r

hybrids, although at a lower affinity (Belfiore et al., 2009; Belfiore et al., 2017), to mediate its pro-

tumourigenic effects. Nevertheless, it is clear that simply reducing *Insr* gene dosage was sufficient to

reduce the effects of hyperinsulinemia on PanIN initiation. While we were unable to assess the role of

Igf1r, future studies examining acinar cell-specific *Igf1r* loss in the presence or absence of concomitant

Insr loss would be necessary to delineate any contribution to PanIN formation.

In summary, our data strongly suggest that insulin receptor signaling in acinar cells contributes to

the PanIN and PDAC development. Our data illustrate the complex and interconnected molecular

mechanisms by which hyperinsulinemia, acting directly through acinar cell Insr, promotes pancreatic

tumourigenesis. We can infer that targeting insulin receptor signaling pathways, or hyperinsulinemia

itself, may be beneficial in treating and preventing pancreatic cancer.

Methods

Mice

All animal experiments were conducted at the University of British Columbia with approval of the

University of British Columbia Animal Care Committee in accordance with Canadian Council for

Animal Care guidelines. All alleles have been described previously (Bruning et al., 1998; Kopp et al.,

2012; Pan et al., 2013; Prigge et al., 2013; Søs Skovsø, 2021; Tuveson et al., 2004). $Kras^{LSL\text{-}G12D/w}$

(#008179), $Insr^{f/f}$ (#006955), and nuclear TdTomato-to-nuclear EGFP (nTnG) mice (#023035) were

purchased from Jackson Labs (Bar Harbour, USA). $Ptf1a^{CreER/w}$ mice (C57BL/6) were a gift from Chris

Wright (Vanderbilt, USA). Mice were maintained on C57BL/6 genetic background and housed at the

University of British Columbia Modified Barrier Facility which was temperature-controlled and specific

pathogen-free. They were kept on a 12:12 hr light: dark cycle with food and drinking water *ad libitum*.

To generate genetic background-matched $Ptf1a^{CreER/w}$;$Kras^{LSL\text{-}G12D/w}$;$Insr^{w/w}$;nTnG, $Ptf1a^{CreER/w}$;$Kras^{LSL\text{-}}$

$^{G12D/w}$;$Insr^{w/f}$;nTnG, and $Ptf1a^{CreER/w}$;$Kras^{LSL\text{-}G12D/w}$;$Insr^{f/f}$;nTnG mice, $Ptf1a^{CreER/w}$;$Kras^{LSL\text{-}}$

$^{G12D/w}$;$Insr^{w/f}$;nTnG mice were bred with $Insr^{w/f}$ mice. After weaning (3 weeks), the resulting litters were

fed with high fat diet (HFD), and at 4 weeks of age, recombination was induced over three consecutive

days by subcutaneous injections of tamoxifen in corn oil (20 mg/mL) of 5 mg tamoxifen/40 g body

mass. One cohort of mice was euthanized at 10 months of age for histopathology analyses and the

whole pancreas was used for histopathology analyses. Another cohort of mice was euthanized at 12

weeks of age. For mice euthanized at 12 weeks, each pancreas was cut into three pieces roughly

based on the pancreas anatomical structure: head, body, and tail. The body piece was used for

histopathological analyses, while the other pieces were snap frozen in liquid nitrogen and kept at -

80^{o}C. The head piece was processed for proteomics and phospho-proteomics analyses (see below).

Assessment of glucose homeostasis

Mouse body weight and fasting blood glucose levels were measured every 4 weeks and fasting

insulin were measured every 3 months. Before the measurements, mice were fasted for 4 hours in

clean and fresh cages during the light period. One drop of blood was collected from each mouse's

saphenous vein and a Lifescan OneTouch Ultra Mini glucometer was used to measure the fasting

blood glucose levels. About 30 µl of blood was collected with a heparinized microhematocrit capillary

tube (Fisher Scientific, 22-362566, Waltham, MA, USA) for measuring fasting insulin levels. The

collected blood was centrifuged at 10,000 rpm for 10 minutes to collect the blood serum. Then the

blood serum was kept at -20^{o}C until used to measure the fasting insulin levels with insulin ELISA (80-

INSMSU-E10; ALPCO Diagnostics, Salem, NH).

Histopathological, morphological, and immunohistochemical analyses

Pancreata were fixed in 4% paraformaldehyde for 24 hours followed by paraffin embedding. Mouse

pancreata were sectioned then stained with hematoxylin and eosin (H&E), and Alcian blue as

previously described (Lee et al., 2018; Zhang et al., 2022; Zhang et al., 2019). The stained slides were

scanned with a 20x objective using a 3DHISTECH Panoramic MIDI (Quorum Technologies Inc.

Guelph, Canada) slide scanner. Histopathological analyses were conducted in a de-identified manner

and verified separately by J.L.K and D.F.S. All histopathological analyses were performed on one of

the stained sections that displayed the maximal pancreatic cross-sectional area unless otherwise

stated. Every gland with a lumen was categorized as normal, ADM, PanIN, or neoplasia, and glands

representing more than one of these categories were scored based on their highest-grade feature.

The total pancreatic area, PanIN plus tumor area, PanIN area, and normal acinar cell area were

measured as previously described (Zhang et al., 2022; Zhang et al., 2019). Briefly, the total pancreatic

area, PanIN plus tumor area, PanIN area, normal acinar cell area was determined by masking all

pancreatic tissue, selective masking of the PanIN plus tumor area, selective masking of the only

PanIN area, and selective masking of the normal acinar cell area by Adobe Photoshop. Pixels for the

total pancreatic area or each histological feature were measured by ImageJ and this was used to

calculate the percentage area occupied by each histological feature. For Alcian blue positive area,

Adobe Photoshop 2020 Black & White function was used to highlight the blue area (red filter). The

total pixels for pancreas or Alcian blue positive area were counted using ImageJ.

Immunohistochemical (IHC) staining was performed according to published standard protocols (Lee

et al., 2018; Zhang et al., 2022). Primary antibodies were rabbit anti-cytokeratin 19 (Ck19) (Abcam,

ab133496, 1:1000) and goat anti-GFP (Abcam, ab6673, 1:200). Secondary antibodies were biotin-

conjugated donkey anti-rabbit (Jackson ImmunoResearch Laboratories, Inc., 711-065-152, 1:500) and

biotin-conjugated rat anti-goat (Vector Laboratories, MP-7404, 1:2). IHC slides were scanned with a

20x objective using a 3DHISTECH Panoramic MIDI slide scanner. Ck19 positive area was measured

with Adobe Photoshop 2020 using the Black & White function to filter and highlight the brown area

(blue filter), which was then taken as a percent of pixels in the pancreatic section.

Proteomics analyses

The whole head of pancreata collected from female mice at 12 weeks of age ($Ptf1a^{CreER}$;$Kras^{LSL\text{-}G12D}$;

$Insr^{w/w}$;nTnG (n=3), $Ptf1a^{CreER}$;$Kras^{LSL\text{-}G12D}$;$Insr^{w/f}$;nTnG (n=6), $Ptf1a^{CreER}$;$Kras^{LSL\text{-}G12D}$;$Insr^{f/f}$;nTnG

(n=6), $Ptf1a^{CreER}$;$Insr^{w/w}$;nTnG (n=3), and $Ptf1a^{CreER}$;$Insr^{f/f}$;nTnG (n=2) mice) were frozen and used for

(phospho)proteomic analyses. The frozen sample was ground into powder in a liquid nitrogen-cooled

mortar and pestle and kept on dry ice until mass-spec analysis. Proteins were extracted from

cryopulverized tumors in a buffer containing 5% sodium dodecyl sulfate (SDS) and 100 mM TRIS pH

7.8 supplemented with PhosStop phosphatase inhibitor cocktail (Roche). The protein concentration of

the lysate was determined using bicinchoninic acid assay (BCA) (Thermo Fisher Scientific), and

protein disulfide bonds were reduced and free Cysteines alkylated with 20 mM tris(2-

carboxyethyl)phosphine (TCEP) and 30 mM iodoacetamide, respectively. 250 µg of total protein was

used for proteolytic digestion using S-TRAP Mini columns (Protifi LLC, Huntington NY) (HaileMariam

et al., 2018). Resultant tryptic peptides were then vacuum concentrated and desalted using Oasis HLB

SPE cartridges (Waters). Peptides were vacuum concentrated and reconstituted in 0.1% trifluoro

acetic acid (TFA), and 10% of the total sample was reserved for measurement of the total proteome.

The remaining 90% of the sample was diluted in 80% acetonitrile with 0.1% TFA for automated

phosphopeptide enrichment using AssayMap Fe-NTA (III) immobilized metal affinity chromatography

(IMAC) cartridges and a Bravo liquid handling system using the phosphopeptide enrichment v 2.1

application (Agilent).

LC-MS/MS acquisition and data analysis

Samples for both total proteomics and phosphoproteomics were analyzed by data dependent

acquisition (DDA) using an Easy-nLC 1200 and Q Exactive Plus (both Thermo Fisher Scientific).

Samples were first loaded onto a precolumn (Acclaim PepMap 100 C18, 3 μm particle size, 75 μm

inner diameter x 2 cm) in 0.1% formic acid (buffer A), and gradient elution was performed using a 100-

min method from 3 to 40% buffer B (84% acetonitrile, 0.1% formic acid) on the analytical column

(Acclaim PepMap 100 C18, 2 μm particle size, 75 μm inner diameter x 25 cm) at a flow rate of 300

nL/min. MS scans were acquired between 350-1,500 m/z at a resolution of 70,000, with an automatic

gain control (AGC) target of 1 x 10e610E6 ions and a maximum injection time of 50 ms. The top 15

precursor ions with charge states +2, +, +3, and +4 were isolated with a window of 1.2 m/z, an AGC

target of 2 x 10e410E4 and a maximum injection time of 64 ms and fragmented using a normalized

collision energy (NCE) of 28. MS/MS were acquired at a resolution of 17,500 and the dynamic

exclusion was set to 30 s. DDA MS raw data was processed with Proteome Discoverer 2.5 (Thermo

Scientific) and searched using Sequest HT against the mouse reference proteome FASTA database

from Uniprot (downloaded October 1st 2021; 17,054 forward sequences). The enzyme specificity was

set to trypsin with a maximum of 2 missed cleavages. Carbamidomethylation of cysteine was set as a

fixed modification and oxidation of methionine, as well as phosphorylation of serine, threonine, and

tyrosine as variable modifications. The precursor ion mass tolerance was set to 10 parts per million,

and the product ion mass tolerance was set to 0.02 Da. Percolator was used to assess posterior error

probabilities and the data was filtered using a false discovery rate (FDR) of <1% on peptide and

protein level. The Minora node of Proteome Discoverer was used for label free quantitation. For the

new human comparison, Pathway enrichment was performed our mouse data and PDAC human data

(Cao et al., 2021) using Reactome (v80) (Gillespie et al., 2022). We performed differential abundance

analysis on the human data using the R package limma (Ritchie et al., 2015). For our data, we used a

cutoff of Adj. p<0.05. For the human data, we used Adj. p<1e-19 and 1e-12 for the total proteins and

phospho-proteins respectively to ensure a comparable proportion of the total number detected.

Primary mouse acinar cell isolation and three-dimensional culture

Primary pancreatic acinar cells were isolated from pancreata of wild-type female mice at the age of

6-8 weeks. The acinar cell isolation procedure was adapted from the protocol described by Martinez

and Storz (Fleming Martinez and Storz, 2019). Briefly, the mouse pancreas was harvested and

washed three times in 1x HBSS, which were then minced into 1-5mm pieces. The fragmented tissue

was then digested with 5mL (0.4mg/mL) collagenase P in 37 °C with gentle shaking for 18 minutes.

HBSS (10mL) with 5% FBS was added to terminate the digestion reaction, followed by 3 washes with

10mL HBSS (5% FBS) to remove the residual collagenase P. After each wash, the tissue was pelleted

(450g RCF, 2 min at room temperature) and the supernatant was removed. The digested tissue was

resuspended in 10mL HBSS (5% FBS) and filtered through a 100 μM cell strainer, followed by a wash

with 10mL HBSS (5% FBS). The filtrate was gently added to 20mL HBSS 30% FBS cushion to form

layers of cells. The cell mixture was centrifuged at 180g RCF at room temperature for 2 minutes to

pellet acinar cells. Appropriate number of acinar cells were then resuspended in premade collagen

solution (1mg/ml rat tail type1 collagen, 10x Waymouth's media, RPMI1640 complete media (1% FBS,

1x penicillin/streptomycin, 1μg/mL dexamethasone), adjusted to pH = ~7.8 with 1M NaOH) and plated

(50μL per well) in collagen pre-coated 96-well plates. After solidification, 100 μL of the RPMI1640

complete media with insulin (0.1nM-100nM), TGF-α (50ng/mL) or combined treatments, with or

without 0.1mg/mL soybean trypsin inhibitor was added. Media was replaced on day 1 and 3, and the

numbers of ADM events in each well were quantified on day 5. Isolated cells were maintained at 37 ^{o}C

in a humidified incubator with 5% CO_2.

Statistical analyses and data visualization

Animals were excluded from the histopathological analyses if they were found dead or if nTnG

recombination efficiency (% acinar cells or PanIN cells labeled with GFP) was lower than 20%.

Statistical analyses were conducted with GraphPad Prism 9.3.0. The Shapiro-Wilk test was used to

assess data normality. One-way ANOVA was performed unless otherwise stated. Mixed-effect

analyses were run for glucose homeostasis data (mouse body weight, fasting glucose level, and

fasting insulin level). When comparing the histopathological measurements between male and female

mice from the same genotype, a two-tailed student's t-test was run for normally distributed data and a

Mann-Whitney test was performed for non-normally distributed data. Statistical parameters, including

the exact value of sample size n (animal number), precision measures and dispersion (mean ± SEM),

and statistical significance levels, are reported in the Figures and Figure legends. A p-value <0.05 was

considered significant. In Figures, asterisks denote statistical significance (*p<0.05, **p<0.01,

***p<0.001, and ****p<0.0001).

For total proteomics data, normalization, imputation, and differential expression analyses were

performed on the protein abundances using Proteome Discoverer. Differential expression analysis

used background-based t-tests with p-values adjusted using Benjamini-Hochberg correction. The

proteins were filtered for significant comparisons (Adj. p<0.05), and heatmaps were created with

resultant protein lists. We also applied k-means clustering and extracted clusters with patterns of

interest (Gu et al., 2016). From this heatmap, a class of specific secreted proteins were identified and

values for all samples were plotted in a heatmap and sorted with hierarchical clustering. We used

STRING (v11) to generate an edge table with differentially expressed total proteins (Adj. p<0.05) with

an absolute $\log_2$(Fold Change) cut-off of 2.5 in $Ptf1a^{CreER}$;$Kras^{LSL-G12D}$;$Insr^{w/w}$;nTnG compared to

$Ptf1a^{CreER}$;$Kras^{LSL-G12D}$;$Insr^{f/f}$;nTnG. Protein-protein relationships were only included if they involved

experimental evidence or interactions annotated in other databases (Szklarczyk et al., 2021).

Cytoscape (v3.9.1) was used to visualize the network (Shannon et al., 2003), with background images

created with BioRender.com.

Phospho-proteomics data were analyzed two ways. In the first approach, we only used phospho-

peptides with measurements made in a minimum of 2/3 of all the samples and at least one sample per

group. We imputed missing phospho-protein values in samples before normalization and statistical

analysis using the same process as the total proteomics, allowing a broader assessment of the

phospho-protein landscape. These were plotted using volcano plots and against the corresponding

proteins in the total protein data set with calculated $\log_2$(Fold Change) of the median. In the second

approach, we aggregated the abundances of each sample for each phospho-site detected, and

normalized each of the detected phospho-sites to the total abundance of the corresponding protein.

We only included phospho-sites where every sample had an experimentally measured total protein

value. The second method used differential expression analysis performed using the R package limma

on the $Ptf1a^{CreER};Kras^{LSL-G12D};Insr^{w/w}$;nTnG vs $Ptf1a^{CreER};Kras^{LSL-G12D};Insr^{f/f}$;nTnG comparison (Ritchie et

al., 2015). The top differentially abundant phospho-sites (sorted by Adj. p-value) were plotted as a

heatmap and sorted with hierarchical clustering (Gu et al., 2016). Phosphosite (www.phosphosite.org)

was used to determine upstream activators, cellular location, and function of each protein, as well as

the specific kinases of the detected phospho-sites (if known). STRING was used to identify proteins

that physically interact with each other (Szklarczyk et al., 2021).

References

Aljada, A., Saleh, A. M., Al-Aqeel, S. M., Shamsa, H. B., Al-Bawab, A., Al Dubayee, M., and Ahmed, A.
A. (2015). Quantification of insulin receptor mRNA splice variants as a diagnostic tumor marker in
breast cancer. Cancer Biomark *15*, 653-661.

Belfiore, A., Frasca, F., Pandini, G., Sciacca, L., and Vigneri, R. (2009). Insulin receptor isoforms and
insulin receptor/insulin-like growth factor receptor hybrids in physiology and disease. Endocr Rev *30*,
586-623.

Belfiore, A., Malaguarnera, R., Vella, V., Lawrence, M. C., Sciacca, L., Frasca, F., Morrione, A., and
Vigneri, R. (2017). Insulin Receptor Isoforms in Physiology and Disease: An Updated View. Endocr
Rev *38*, 379-431.

Berglund, E. D., Li, C. Y., Poffenberger, G., Ayala, J. E., Fueger, P. T., Willis, S. E., Jewell, M. M.,
Powers, A. C., and Wasserman, D. H. (2008). Glucose metabolism in vivo in four commonly used
inbred mouse strains. Diabetes *57*, 1790-1799.

Binder, J. X., Pletscher-Frankild, S., Tsafou, K., Stolte, C., O'Donoghue, S. I., Schneider, R., and
Jensen, L. J. (2014). COMPARTMENTS: unification and visualization of protein subcellular
localization evidence. Database (Oxford) *2014*, bau012.

Bruce, J. I. E., Sanchez-Alvarez, R., Sans, M. D., Sugden, S. A., Qi, N., James, A. D., and Williams, J.
A. (2021). Insulin protects acinar cells during pancreatitis by preserving glycolytic ATP supply to
calcium pumps. Nat Commun *12*, 4386.

Bruning, J. C., Michael, M. D., Winnay, J. N., Hayashi, T., Horsch, D., Accili, D., Goodyear, L. J., and
Kahn, C. R. (1998). A muscle-specific insulin receptor knockout exhibits features of the metabolic
syndrome of NIDDM without altering glucose tolerance. Mol Cell *2*, 559-569.

Cao, L., Huang, C., Cui Zhou, D., Hu, Y., Lih, T. M., Savage, S. R., Krug, K., Clark, D. J., Schnaubelt,
M., Chen, L.*, et al.* (2021). Proteogenomic characterization of pancreatic ductal adenocarcinoma. Cell
184, 5031-5052 e5026.

Cen, H. H., Hussein, B., Botezelli, J. D., Wang, S., Zhang, J. A., Noursadeghi, N., Jessen, N.,
Rodrigues, B., Timmons, J. A., and Johnson, J. D. (2022). Human and mouse muscle transcriptomic
analyses identify insulin receptor mRNA downregulation in hyperinsulinemia-associated insulin
resistance. FASEB J *36*, e22088.

Chan, D. S., Vieira, A. R., Aune, D., Bandera, E. V., Greenwood, D. C., McTiernan, A., Navarro
Rosenblatt, D., Thune, I., Vieira, R., and Norat, T. (2014a). Body mass index and survival in women
with breast cancer-systematic literature review and meta-analysis of 82 follow-up studies. Ann Oncol
25, 1901-1914.

Chan, M. T., Lim, G. E., Skovso, S., Yang, Y. H., Albrecht, T., Alejandro, E. U., Hoesli, C. A., Piret, J.
M., Warnock, G. L., and Johnson, J. D. (2014b). Effects of insulin on human pancreatic cancer
progression modeled in vitro. BMC Cancer *14*, 814.

Chang, H. H., Moro, A., Takakura, K., Su, H. Y., Mo, A., Nakanishi, M., Waldron, R. T., French, S. W.,
Dawson, D. W., Hines, O. J.*, et al.* (2017). Incidence of pancreatic cancer is dramatically increased by
a high fat, high calorie diet in KrasG12D mice. PLoS One *12*, e0184455.

Chen, Z., Downing, S., and Tzanakakis, E. S. (2019). Four Decades After the Discovery of
Regenerating Islet-Derived (Reg) Proteins: Current Understanding and Challenges. Front Cell Dev
Biol *7*, 235.

Cherrington, A. D. (2005). The role of hepatic insulin receptors in the regulation of glucose production.
J Clin Invest *115*, 1136-1139.

Dawson, D. W., Hertzer, K., Moro, A., Donald, G., Chang, H. H., Go, V. L., Pandol, S. J., Lugea, A.,
Gukovskaya, A. S., Li, G.*, et al.* (2013). High-fat, high-calorie diet promotes early pancreatic neoplasia
in the conditional KrasG12D mouse model. Cancer Prev Res (Phila) *6*, 1064-1073.

Dembinski, A. B., and Johnson, L. R. (1980). Stimulation of Pancreatic Growth by Secretin, Caerulein,
and Pentagastrin. Endocrinology *106*, 323-328.

Fleming Martinez, A. K., and Storz, P. (2019). Mimicking and Manipulating Pancreatic Acinar-to-Ductal
Metaplasia in 3-dimensional Cell Culture. J Vis Exp.

Frier, B. M., Saunders, J. H., Wormsley, K. G., and Bouchier, I. A. (1976). Exocrine pancreatic function
in juvenile-onset diabetes mellitus. Gut *17*, 685-691.

Gallagher, E. J., and LeRoith, D. (2020). Hyperinsulinaemia in cancer. Nat Rev Cancer *20*, 629-644.

Gillespie, M., Jassal, B., Stephan, R., Milacic, M., Rothfels, K., Senff-Ribeiro, A., Griss, J., Sevilla, C.,
Matthews, L., Gong, C.*, et al.* (2022). The reactome pathway knowledgebase 2022. Nucleic Acids Res
50, D687-D692.

Godsland, I. F. (2009). Insulin resistance and hyperinsulinaemia in the development and progression
of cancer. Clin Sci (Lond) *118*, 315-332.

Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics *32*, 2847-2849.

Guerra, C., Schuhmacher, A. J., Cañamero, M., Grippo, P. J., Verdaguer, L., Pérez-Gallego, L.,
Dubus, P., Sandgren, E. P., and Barbacid, M. (2007). Chronic pancreatitis is essential for induction of
pancreatic ductal adenocarcinoma by K-Ras oncogenes in adult mice. Cancer Cell *11*, 291-302.

Guo, Y. J., Pan, W. W., Liu, S. B., Shen, Z. F., Xu, Y., and Hu, L. L. (2020). ERK/MAPK signalling
pathway and tumorigenesis. Exp Ther Med *19*, 1997-2007.

Haeusler, R. A., McGraw, T. E., and Accili, D. (2018). Biochemical and cellular properties of insulin
receptor signalling. Nat Rev Mol Cell Biol *19*, 31-44.

HaileMariam, M., Eguez, R. V., Singh, H., Bekele, S., Ameni, G., Pieper, R., and Yu, Y. (2018). S-Trap,
an Ultrafast Sample-Preparation Approach for Shotgun Proteomics. J Proteome Res *17*, 2917-2924.

Haoning Howard Cen, J. R., Libin Abraham, Michael R. Gold, Leonard J. Foster, James D. Johnson
(2022). Insulin-dependent and -independent dynamics of insulin receptor trafficking in muscle cells.
bioRxiv.

Heestand, G. M., and Kurzrock, R. (2015). Molecular landscape of pancreatic cancer: implications for
current clinical trials. Oncotarget *6*, 4553-4561.

Henderson, J. R., Daniel, P. M., and Fraser, P. A. (1981). The pancreas as a single organ: the
influence of the endocrine upon the exocrine part of the gland. Gut *22*, 158-167.

Hernandez, V. J., Weng, J., Ly, P., Pompey, S., Dong, H., Mishra, L., Schwarz, M., Anderson, R. G.,
and Michaely, P. (2013). Cavin-3 dictates the balance between ERK and Akt signaling. Elife *2*, e00905.

Hong, S., Freeberg, M. A., Han, T., Kamath, A., Yao, Y., Fukuda, T., Suzuki, T., Kim, J. K., and Inoki,
K. (2017). LARP1 functions as a molecular switch for mTORC1-mediated translation of an essential
class of mRNAs. Elife *6*.

Hopkins, B. D., Pauli, C., Du, X., Wang, D. G., Li, X., Wu, D., Amadiume, S. C., Goncalves, M. D.,
Hodakoski, C., Lundquist, M. R*., et al.* (2018). Suppression of insulin feedback enhances the efficacy
of PI3K inhibitors. Nature *560*, 499-503.

Ilic, M., and Ilic, I. (2016). Epidemiology of pancreatic cancer. World J Gastroenterol *22*, 9694-9705.

Incio, J., Liu, H., Suboj, P., Chin, S. M., Chen, I. X., Pinter, M., Ng, M. R., Nia, H. T., Grahovac, J., Kao,
S*., et al.* (2016). Obesity-Induced Inflammation and Desmoplasia Promote Pancreatic Cancer
Progression and Resistance to Chemotherapy. Cancer Discov *6*, 852-869.

Janku, F., Wheler, J. J., Naing, A., Falchook, G. S., Hong, D. S., Stepanek, V. M., Fu, S., Piha-Paul, S.
A., Lee, J. J., Luthra, R*., et al.* (2013). PIK3CA mutation H1047R is associated with response to
PI3K/AKT/mTOR signaling pathway inhibitors in early-phase clinical trials. Cancer Res *73*, 276-284.

Kopp, J. L., von Figura, G., Mayes, E., Liu, F. F., Dubois, C. L., Morris, J. P. t., Pan, F. C., Akiyama, H.,
Wright, C. V., Jensen, K*., et al.* (2012). Identification of Sox9-dependent acinar-to-ductal
reprogramming as the principal mechanism for initiation of pancreatic ductal adenocarcinoma. Cancer
Cell *22*, 737-750.

Law, J. H., Habibi, G., Hu, K., Masoudi, H., Wang, M. Y., Stratford, A. L., Park, E., Gee, J. M., Finlay,
P., Jones, H. E*., et al.* (2008). Phosphorylated insulin-like growth factor-i/insulin receptor is present in
all breast cancer subtypes and is related to poor survival. Cancer Res *68*, 10238-10246.

Lee, A. Y. L., Dubois, C. L., Sarai, K., Zarei, S., Schaeffer, D. F., Sander, M., and Kopp, J. L. (2018).
Cell of origin affects tumour development and phenotype in pancreatic ductal adenocarcinoma. Gut.

Li, Q., Wang, H., Zogopoulos, G., Shao, Q., Dong, K., Lv, F., Nwilati, K., Gui, X. Y., Cuggia, A., Liu, J.
L., and Gao, Z. H. (2016). Reg proteins promote acinar-to-ductal metaplasia and act as novel
diagnostic and prognostic markers in pancreatic ductal adenocarcinoma. Oncotarget *7*, 77838-77853.

Means, A. L., Meszoely, I. M., Suzuki, K., Miyamoto, Y., Rustgi, A. K., Coffey, R. J., Jr., Wright, C. V.,
Stoffers, D. A., and Leach, S. D. (2005a). Pancreatic epithelial plasticity mediated by acinar cell
transdifferentiation and generation of nestin-positive intermediates. Development *132*, 3767-3776.

Means, A. L., Meszoely, I. M., Suzuki, K., Miyamoto, Y., Rustgi, A. K., Coffey, R. J., Wright, C. V. E.,
Stoffers, D. A., and Leach, S. D. (2005b). Pancreatic epithelial plasticity mediated by acinar cell
transdifferentiation and generation of nestin-positive intermediates. Development (Cambridge,
England) *132*, 3767-3776.

Mehran, A. E., Templeman, N. M., Brigidi, G. S., Lim, G. E., Chu, K. Y., Hu, X., Botezelli, J. D., Asadi,
A., Hoffman, B. G., Kieffer, T. J*., et al.* (2012). Hyperinsulinemia drives diet-induced obesity
independently of brain insulin production. Cell Metab *16*, 723-737.

Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L., and Luo, L. (2007). A global double-fluorescent Cre
reporter mouse. Genesis *45*, 593-605.

Otsuki, M. (2000). Pathophysiological role of cholecystokinin in humans. J Gastroenterol Hepatol *15*
Suppl, D71-83.

Pan, F. C., Bankaitis, E. D., Boyer, D., Xu, X., Van de Casteele, M., Magnuson, M. A., Heimberg, H.,
and Wright, C. V. (2013). Spatiotemporal patterns of multipotentiality in Ptf1a-expressing cells during
pancreas organogenesis and injury-induced facultative restoration. Development *140*, 751-764.

Payne, S. N., Maher, M. E., Tran, N. H., Van De Hey, D. R., Foley, T. M., Yueh, A. E., Leystra, A. A.,
Pasch, C. A., Jeffrey, J. J., Clipson, L.*, et al.* (2015). PIK3CA mutations can initiate pancreatic
tumorigenesis and are targetable with PI3K inhibitors. Oncogenesis *4*, e169.

Pisani, P. (2008). Hyper-insulinaemia and cancer, meta-analyses of epidemiological studies. Arch
Physiol Biochem *114*, 63-70.

Prigge, J. R., Wiley, J. A., Talago, E. A., Young, E. M., Johns, L. L., Kundert, J. A., Sonsteng, K. M.,
Halford, W. P., Capecchi, M. R., and Schmidt, E. E. (2013). Nuclear double-fluorescent reporter for in
vivo and ex vivo analyses of biological transitions in mouse nuclei. Mamm Genome.

Rahib, L., Smith, B. D., Aizenberg, R., Rosenzweig, A. B., Fleshman, J. M., and Matrisian, L. M.
(2014). Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and
pancreas cancers in the United States. Cancer Res *74*, 2913-2921.

Rao, R. V., Holicky, E. L., Kuntz, S. M., and Miller, L. J. (2000). CCK receptor phosphorylation
exposes regulatory domains affecting phosphorylation and receptor trafficking. Am J Physiol Cell
Physiol *279*, C1986-1992.

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., and Smyth, G. K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids
Res *43*, e47.

Samuels, Y., Wang, Z., Bardelli, A., Silliman, N., Ptak, J., Szabo, S., Yan, H., Gazdar, A., Powell, S.
M., Riggins, G. J.*, et al.* (2004). High frequency of mutations of the PIK3CA gene in human cancers.
Science *304*, 554.

Sato, K., Hikita, H., Myojin, Y., Fukumoto, K., Murai, K., Sakane, S., Tamura, T., Yamai, T., Nozaki, Y.,
Yoshioka, T.*, et al.* (2020). Hyperglycemia enhances pancreatic cancer progression accompanied by
elevations in phosphorylated STAT3 and MYC levels. PloS one *15*, e0235573-e0235573.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B.,
and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular
interaction networks. Genome Res *13*, 2498-2504.

Sivaram, N., McLaughlin, P. A., Han, H. V., Petrenko, O., Jiang, Y. P., Ballou, L. M., Pham, K., Liu, C.,
van der Velden, A. W., and Lin, R. Z. (2019). Tumor-intrinsic PIK3CA represses tumor
immunogenecity in a model of pancreatic cancer. J Clin Invest *129*, 3264-3276.

Smith, J. P., and Solomon, T. E. (2014). Cholecystokinin and pancreatic cancer: the chicken or the
egg? Am J Physiol Gastrointest Liver Physiol *306*, G91-G101.

Soling, H. D., and Unger, K. O. (1972). The role of insulin in the regulation of -amylase synthesis in the
rat pancreas. Eur J Clin Invest *2*, 199-212.

Søs Skovsø, E. P., Jelena Kolic, Haoning Howard Cen, Derek A. Dionne, Xiao-Qing Dai, Rohit B.
Sharma, Lynda Elghazi, Cara E. Ellis, Katharine Faulkner, Stephanie A.M. Marcil, Peter Overby, Nilou

Noursadeghi, Daria Hutchinson, Xiaoke Hu, Hong Li, Honey Modi, Jennifer S. Wildi, J. Diego Botezelli,
Hye Lim Noh, Sujin Suk, Brian Gablaski, Austin Bautista, Ryekjang Kim, Cras-Méneur, Stephane
Flibotte, Sunita Sinha, Dan S. Luciani, Corey Nislow, Elizabeth J. Rideout, Eric N. Cytrynbaum, Jason
K. Kim, Ernesto Bernal-Mizrachi, Laura C. Alonso, Patrick E. MacDonald, James D. Johnson (2021).
Beta-cell specific Insr deletion promotes insulin hypersecretion and improves glucose tolerance prior
to global insulin resistance. biRxiv.

Steiner, S., Seleznik, G. M., Reding, T., Stopic, M., Lenggenhager, D., Ten Buren, E., Eshmuminov,
D., Endhardt, K., Hagedorn, C., Heidenblut, A. M*., et al.* (2022). De novo expression of gastrokines in
pancreatic precursor lesions impede the development of pancreatic cancer. Oncogene *41*, 1507-1517.

Stolzenberg-Solomon, R. Z., Graubard, B. I., Chari, S., Limburg, P., Taylor, P. R., Virtamo, J., and
Albanes, D. (2005). Insulin, glucose, insulin resistance, and pancreatic cancer in male smokers. JAMA
294, 2872-2878.

Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva, N. T., Legeay,
M., Fang, T., Bork, P*., et al.* (2021). The STRING database in 2021: customizable protein-protein
networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids
Res *49*, D605-D612.

Templeman, N. M., Clee, S. M., and Johnson, J. D. (2015). Suppression of hyperinsulinaemia in
growing female mice provides long-term protection against obesity. Diabetologia *58*, 2392-2402.

Templeman, N. M., Flibotte, S., Chik, J. H. L., Sinha, S., Lim, G. E., Foster, L. J., Nislow, C., and
Johnson, J. D. (2017). Reduced Circulating Insulin Enhances Insulin Sensitivity in Old Mice and
Extends Lifespan. Cell Rep *20*, 451-463.

Tuveson, D. A., Shaw, A. T., Willis, N. A., Silver, D. P., Jackson, E. L., Chang, S., Mercer, K. L.,
Grochow, R., Hock, H., Crowley, D*., et al.* (2004). Endogenous oncogenic K-ras(G12D) stimulates
proliferation and widespread neoplastic and developmental defects. Cancer Cell *5*, 375-387.

Varga, G., Papp, M., Solomon, T. E., Scarpignato, C., Toth, I. E., and Szabo, D. (1988). Caerulein in
supramaximal doses fails to stimulate pancreatic growth, but it forces secretory granulopoiesis. Acta
Physiol Hung *71*, 99-106.

Vaziri-Gohar, A., Hue, J. J., Graor, H. G., Abbas, A., Zarei, M., Hajihassani, O., Titomihelakis, G.,
Feczko, J., Rathore, M., Wang, R*., et al.* (2022). Increased glucose availability sensitizes pancreatic
cancer to chemotherapy. bioRxiv, 2022.2004.2029.490090.

Waters, A. M., and Der, C. J. (2018). KRAS: The Critical Driver and Therapeutic Target for Pancreatic
Cancer. Cold Spring Harb Perspect Med *8*.

Yang, J., Nie, J., Ma, X., Wei, Y., Peng, Y., and Wei, X. (2019). Targeting PI3K in cancer: mechanisms
and advances in clinical trials. Mol Cancer *18*, 26.

Zhang, A. M. Y., Chu, K. H., Daly, B. F., Ruiter, T., Dou, Y., Yang, J. C. C., de Winter, T. J. J., Chhuor,
J., Wang, S., Flibotte, S*., et al.* (2022). Effects of hyperinsulinemia on pancreatic cancer development
and the immune microenvironment revealed through single-cell transcriptomics. Cancer Metab *10*, 5.

Zhang, A. M. Y., Magrill, J., de Winter, T. J. J., Hu, X., Skovsø, S., Schaeffer, D. F., Kopp, J. L., and
Johnson, J. D. (2019). Endogenous Hyperinsulinemia Contributes to Pancreatic Cancer Development.
Cell Metab *30*, 403-404.

Zhang, A. M. Y., Wellberg, E. A., Kopp, J. L., and Johnson, J. D. (2021). Hyperinsulinemia in Obesity,
Inflammation, and Cancer. Diabetes Metab J *45*, 285-311.

 Figure Legends

Fig. 1. Loss of *Insr* in pancreatic acinar cells had no effect on glucose homeostasis and

pancreas size.

A, Schematic describing mouse models designed to test the role of insulin receptor signaling in

pancreatic acinar cells. On the background of the $Ptf1a^{CreER}$ mice, we generated mice having two wild-

type *Insr* alleles ($Insr^{w/w}$) or two *Insr* floxed alleles ($Insr^{f/f}$). **B,** Three-week-old mice were weaned and

provided high fat diet (HFD) for the duration of the study. At 4-weeks-old, they were injected with

tamoxifen (TM) on 3 consecutive days. Physiological measures were taken every 3 months for 10

months, mice were euthanized at 12 or 43.5 weeks of age for histopathological, proteomics, and

phospho-proteomics analyses. **C-H,** Body weight (**C-D**), fasting blood glucose (**E-F**), and fasting

insulin (**G-H**) measurements in male (**C, E, G**) and female (**D, F, H**) $Ptf1a^{CreER}$-$Insr^{w/w}$ and $Ptf1a^{CreER}$-

$Insr^{f/f}$ mice measured over >1 year (n= 10-17). **I,** The ratio of pancreatic weight to mouse body weight

for male (M) and female (F) $Ptf1a^{CreER}$-$Insr^{w/w}$ and $Ptf1a^{CreER}$-$Insr^{f/f}$ mice (n= 3-6). Values shown as

mean ± SEM.

Fig. 2. Loss of *Insr* in pancreatic acinar cells had no effect on glucose homeostasis in

$Ptf1a^{CreER}$**-induced** $Kras^{G12D}$ **pancreatic cancer model.**

A, Schematic describing a mouse model designed to test the role of insulin receptor signaling on HFD-

accelerated PDAC initiation. On the background of the $Ptf1a^{CreER}$-induced $Kras^{G12D}$ pancreatic cancer

model (PK), we generated mice having two wild-type *Insr* alleles ($Insr^{w/w}$), one *Insr* floxed allele

($Insr^{w/f}$), or two *Insr* floxed alleles ($Insr^{f/f}$). **B-G,** Body weight (**B-C**), fasting blood glucose (**D-F**), or

fasting insulin (**F-G**) measurements for PK-$Insr^{w/w}$, PK-$Insr^{w/f}$, and PK-$Insr^{f/f}$ male (**B, D, F**) and female

(**C, E, G**) mice (n= 13-33). Values shown as mean ± SEM. *p<0.05.

Fig. 3. Loss of *Insr* in acinar cells reduced PanIN and PDAC formation.

A, Percentage of mice of the indicated genotype that were found dead or reached humane endpoint

throughout the study. The star symbol indicates whether a macroscopic mass was observed in the

pancreas at necropsy, when possible. There were 8 PK-*Insr*$^{w/w}$, 20 PK-*Insr*$^{w/f}$, and 7 PK-*Insr*$^{f/f}$ male

mice and 9 PK-*Insr*$^{w/w}$ 17 PK-*Insr*$^{w/f}$, and 12 PK-*Insr*$^{f/f}$ female mice in the cohort. At 43.5 weeks, the

remaining mice were euthanized and assessed for presence of tumors. **B-C,** Representative whole

section (top) and high-magnification (bottom) H&E images of pancreatic slides from 43.5-week-old

male (**B**) and female (**C**) PK-*Insr*$^{w/w}$, PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$ mice. **D-G,** Quantification of PanIN plus

tumor area (**D**)**,** PanIN only area (**E**), tumor area (**F**), or acinar cell area (**G**) in pancreata from each

genotype and sex (male or female) (n= 5-16). Filled dots and triangles denoted mice that developed

tumors in **D**. Scale bars: 2 mm (top) and 0.1mm (bottom). Values are shown as mean ± SEM. *p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001.

Fig. 4. Loss of *Insr* in acinar cells reduced PanIN lesions initiation.

A, The ratio of pancreatic weight to mouse body weight for male (M) and female (F) PK-*Insr*$^{w/w}$, PK-

Insr$^{w/f}$, and PK-*Insr*$^{f/f}$ mice (n= 3-15). **B,** Quantification of PanIN area in pancreata from each genotype

and sex (male or female) (n= 3-14). **C-D,** Representative whole section (top) and high-magnification

(bottom) H&E images of pancreatic slides from 12-week-old male (**C**) and female (**D**) PK-*Insr*$^{w/w}$, PK-

Insr$^{w/f}$, and PK-*Insr*$^{f/f}$ mice. Scale bars: 2 mm (top) and 0.1mm (bottom). Values are shown as mean ±

SEM. *p<0.05, **p<0.01.

Fig. 5. The proteomic analysis of insulin receptors' effects on PanIN initiation

A, Volcano plot for proteins that were significantly up- or down-regulated in PK-*Insr*$^{w/w}$ mice (n=3)

compared to *Ptf1a*^{CreER}-*Insr*$^{w/w}$ mice (n=3). **B**, Volcano plot for proteins that were significantly up- or

down-regulated in *Ptf1a*^{CreER}-*Insr*$^{f/f}$ mice (n=2) compared to *Ptf1a*^{CreER}-*Insr*$^{w/w}$ mice (n=3). **C**, Volcano

plot showing proteins that were significantly up- or down-regulated in PK-*Insr*$^{f/f}$ mice compared to PK-

Insr$^{w/w}$ mice. **D,** Proteins that were differentially abundant between PK-*Insr*$^{w/w}$ and PK-*Insr*$^{f/f}$ mice were

selected and their abundance across all genotypes was used for k-means clustering. The clusters

were visualized by heatmap to show their variance across all genotypes. The proteins in clusters 1, 2,

3, and 7 are listed in the order shown in the heatmap from top to bottom.

Fig. 6. Loss of *Insr* in acinar cells reduced digestive enzymes production in pancreas.

A, Differentially abundant proteins between PK-*Insr*$^{w/w}$ and PK-*Insr*$^{f/f}$ pancreata were connected using

STRING and diagrammatically presented here in the context of Insr signaling and cellular organelles

and functions. The color in each oval reflects the fold change, while the thickness of the line around

the oval represents the Adj. p-value. Proteins depicted in white ovals were added to show the potential

relationship of the differentially abundant proteins to insulin signaling. **B**, Heatmap showing the change

of protein abundance between genotypes for proteins involved in pancreatic digestive enzymes

secretion. **C**, The percentage of summed digestive enzyme abundances in total measured proteins for

each genotype. Values are shown as mean ± SEM. *p<0.05, **p<0.01.

Fig. 7. The phospho-proteomic analysis of insulin receptors' effects on PanIN initiation

A, Volcano plots for phospho-peptides that were significantly up or down regulated in PK-*Insr*$^{f/f}$ mice

compared to PK-*Insr*$^{w/w}$ mice. **B**, Change in phospho-peptide abundance (Adj. p-values indicated in

red) compared to the change in corresponding protein's abundance between PK-*Insr*$^{f/f}$ mice and PK-

Insr$^{w/w}$ mice. **C**, TOP: Schematic of phosphorylation sites on Cckar that were differentially abundant in

any comparison. BOTTOM: The relative phospho-peptide abundance (arbitrary units) for each

detected phospho-peptide for all samples (white dots) and genotypes. **D,** Intracellular locations and

functions of differentially abundant phospho-sites and their corresponding protein. The color in each

attached phospho-site reflects the fold change, while the thickness of the border represents the Adj. p-

value. Values are shown as mean ± SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Fig. 8. Insulin receptor signaling promotes PanIN initiation via increased inflammation

associated with hyperactive digestive enzyme production.

A-B, Representative whole section (top) and high-magnification (bottom) images of pancreatic slides

from 12-week-old male (**A**) and female (**B**) mice PK-*Insr*$^{w/w}$, PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$ stained with Sirius

red. **C,** Quantification of Sirius red positive area for mice from each genotype and sex (male (M) or

female (F)) (n= 3-4). **D,** Schematic of the experimental design of the ADM formation assay using 3D

acinar cell explant system. **E,** Representative bright field images of acinar cell clusters and duct

formation (ring structures) on day 5 of treatment with or without 50ng/mL TGF-α or 100nM insulin in

the absence of trypsin inhibitor. ADM structures are indicated by red arrows. **F,** Quantification of the

fold change in ADM events in primary mouse acinar cell 3D explants after 5 days of treatment with a

combination of ± TGF-α and ± insulin and ± trypsin inhibitor. (n = 4 separate experiments, two-way

ANOVA). Fold change values were calculated as treatment/negative control (no insulin or TGF-α or

trypsin inhibitor) for each experiment. **G.** Schematic demonstrating obesity or diet-induced

hyperinsulinemia, via Insr, promoted PanIN formation through increasing digestive enzymes

production/release and its associated inflammation. Scale bars: 2 mm (A, B; top), 0.1 mm (A, B;

bottom), and 100µm (E). Values are shown as mean ± SEM. *Adj. p<0.05, **Adj. p<0.01, ***Adj.

p<0.001.

Figure S1. The *Ptf1a*^{CreER} allele labeled acinar cells and PanINs in 43.5-week-old mice.

A-B, Representative whole section (top) and high-magnification (bottom) images of

immunohistochemical staining for GFP expressed from nTnG lineage tracing allele in 43.5-week-old

male (**A**) and female (**B**) PK-*Insr*$^{w/w}$, PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$ mice. **C-D**, Representative whole section

(top) and high-magnification (bottom) images for immunohistochemical staining for GFP expressed

from nTnG lineage tracing allele in 43.5-week-old male (**C**) and female (**D**) *Ptf1a*^{CreER}-*Insr*$^{w/w}$ and

Ptf1a^{CreER}-*Insr*$^{f/f}$ mice. **E-F**, Representative whole section (top) and high-magnification (bottom) H&E

images of pancreatic slides from 43.5-week-old male (**E**) and female (**F**) *Ptf1a*^{CreER}-*Insr*$^{w/w}$ and

Ptf1a^{CreER}-*Insr*$^{f/f}$ mice. Scale bars: 2 mm (top) and 0.1mm (bottom).

Figure S2. Loss of *Insr* in acinar cells reduced ductal metaplasia and Alcian blue$^+$ PanIN

lesions.

A-B, Representative whole section (top) and high-magnification (bottom) images of

immunohistochemical staining of Ck19 for male (**A**) and female (**B**) PK-*Insr*$^{w/w}$, PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$

mice. **C-D,** Representative whole section (top) and high-magnification (bottom) images of pancreatic

slides from male (**C**) and female (**D**) PK-*Insr*$^{w/w}$, PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$ mice stained with Alcian blue.

E, Quantification of Ck19$^+$ area for mice from each genotype and sex (M or F) (n= 5-15). **F,**

Quantification of Alcian blue positive area for mice from each genotype and sex (M or F) (n= 5-16). **G-**

H, The correlation of acinar cell area and Ck19$^+$ area for male (**G**) and female (**H**) mice. The maximum

value for Ck19$^+$ area was ~20-30% due to stromal expansion in the parenchyma around the Ck19$^+$

area. Scale bars: 2 mm (top) and 0.1 mm (bottom). Values are shown as mean ± SEM. *p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001.

Figure S3. The *Ptf1a*^{CreER} allele labeled acinar cells and PanINs in 12-week-old mice.

A-B, Representative whole section (top) and high-magnification (bottom) images of

immunohistochemical staining for GFP expressed from the nTnG lineage tracing allele in 12-week-old

male (**A**) and female (**B**) PK-*Insr*$^{w/w}$, PK-*Insr*$^{w/f}$, and PK-*Insr*$^{f/f}$ pancreata. Scale bars: 1mm (top) and 0.1

mm (bottom).

Figure S4. *Kras*^{G12D} mutant mice had similar proteomics and phospho-proteomics changes as

human PDAC tumor samples.

A, Plot of the top 20 mouse pathways from Reactome pathway enrichment for proteins that were

significantly up-or down-regulated (Adj. p<0.05) in PK-*Insr*$^{w/w}$ mice (n=3) compared to *Ptf1a*^{CreER}-

Insr$^{w/w}$ mice (n=3). The FDR for the same pathways from pathway enrichment with proteins from

human data that were significantly up- or down-regulated (Adj. p<10^{-19}) in normal (n=75)

compared to tumor samples (n=140) are also shown. The size of each dot represents the

proportion of genes inputted in all the genes found in each pathway. **B,** Plot of the top 20 mouse

pathways from Reactome pathway enrichment for proteins from corresponding phospho-sites that

were significantly up- or down-regulated (Adj. p<0.05) in PK-*Insr*$^{w/w}$ mice (n=3) compared to

Ptf1a^{CreER}-*Insr*$^{w/w}$ mice (n=3). The FDR for the same pathways from pathway enrichment with

proteins from corresponding phospho-sites from human data that were significantly up- or down-

regulated (Adj. p<10^{-12}) in normal (n=75) compared to tumor samples (n=140) are also shown.

The size of each dot represents the proportion of genes inputted in all the genes found in each

pathway.

Figure S5. Loss of *Insr* in acinar cells reduced desmoplasia.

A-B, The correlation of PanIN area and Sirius red$^+$ area for male (**A**) and female (**B**) mice. *p<0.05.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1

Figure S2

Figure S4

A

M:PK-*Insr*^{w/w}
M:PK-*Insr*^{w/f}
M:PK-*Insr*^{f/f}

r^2=0.717
*p<0.05
r^2=0.223
ns
r^2=0.321
ns

PanIN area/Pancreas area %

Sirius red$^+$ area%

B

F:PK-*Insr*^{w/w}
F:PK-*Insr*^{w/f}
F:PK-*Insr*^{f/f}

r^2=0.530
ns
r^2=0.381
ns
r^2=0.0250
ns

PanIN area/Pancreas area %

Sirius red$^+$ area%

Figure S5