Title: Characteristics of intestinal microflora and dysbiosis in relation to the disease

duration in patients with Meniere's disease

Short title: Dysbiosis in patients with Meniere's disease

1. Department of Otolaryngology,St.Mrianna University School of Medicine

2. Department of Otolaryngology ,University of Miami Miller School of Medicine:

University of Miami School of Medicine

Corresponding author:Fumihiro Mochizuki[1) 2)]

Co-author: Manabu Komori[1)], Yoshiyuki Sasano[1)], Yusuke Ito[1)],Michael E Hoffer[2)],

Izumi Koizuka[1)]

Abstract

Hypothesis

Dysbiosis of the intestinal microflora has been reported in stress-induced depression and irritable colitis, a similar abnormality may also occur in stress-triggered Meniere's disease.

Background

Meniere's disease is an intractable disease characterized by paroxysms of intense rotatory dizziness, hearing loss, and other auditory symptoms. It is believed to be caused by endolymphatic hydrops of the vestibule and cochlea. The diagnosis of Meniere's disease is based on the diagnostic criteria established by the Barany Society and AAO-HNS and Journal of Vestibular Research as definite or probable cases. Endolymphatic hydrops identification using cochlear contrast-enhanced MRI (hybrid of reversed image of positive endolymph signal and native image of positive perilymph signal [HYDROPS]) is an objective test for Meniere's disease, enabling reliable diagnosis of this disease.

Methods

We investigated the gut microbiota of 10 patients (6 males and 4 females, mean age 49.6 ± 8.1 years) who met the diagnostic criteria for a unilateral definite case of Meniere's disease using the diagnostic criteria of the Journal of Vestibular Research , and who also had significant endolymphatic hydrops on the affected side on HYDROPS. Intestinal microbiota tests were performed on these 10 patients and the results were evaluated in relation to the duration of disease, results of audiometry of the affected side, and DHI (Dizziness Handicap Inventory) score.

Results

A significant negative correlation was found between the disease duration and Shannon diversity index and Faith's phylogenetic diversity, which indicated dysbiosis of the intestinal microflora. No correlation was found between the indicators of microbial diversity and the results of audiometry or the DHI on the affected side. Dysbiosis of the intestinal microbiota worsened with increasing duration of Meniere's disease. Moreover, *Akkermansia muciniphila* was not detected in any patient with Meniere's disease.

Conclusions.

48   Despite the small number of cases in this study (n = 10), the findings indicate the
49   possibility of abnormalities of the intestinal microflora in Meniere's disease.

Introduction

Meniere's disease is an intractable disorder characterized by paroxysms of intense rotatory vertigo and auditory symptoms such as aural fullness and hearing loss. Although the pathogenesis of Meniere's disease is not fully understood, it is speculated to be caused by endolymphatic hydrops of the vestibule and cochlea. The diagnostic criteria for Meniere's disease were defined by the Barany Society and AAO-HNS and Journal of Vestibular Research, etc [1]. Objective tests for evaluation of Meniere's disease traditionally included the furosemide test and cochleography, but imaging studies using cochlear contrast-enhanced MRI (hybrid of reversed image of positive endolymph signal and native image of positive perilymph signal [HYDROPS]) have made it possible to visually evaluate endolymphatic hydrops.

HYDROPS revealed that the presence of endolymphatic hydrops on imaging is not necessarily consistent with the pathophysiology of Meniere's disease. Appearance of could be observed in Meniere's disease patients both in symptomatic and asymptomatic ears. [2,3]. Meniere's disease is considered a result of multiple genes interacting with environmental factors.[2,4] Several abnormalities have been reported as histological features of Meniere's disease. These include findings of ischemia of the stria vascularis, fibrous tissue proliferation in the saccular, atrophy of the sac and loss of epithelial

integrity, hypoplasia of the vestibular aqueduct, and spiral ganglion degeneration at the

apex of the cochlea.[2,5] In addition, fluid homeostasis in the inner ear, as in the kidney,

depends on water production, transport, and absorption. Therefore, arginine-vasopressin

(AVP) localized in the inner ear, AVP-related molecules such as vasopressin receptors,

and aquaporin (AQP), a water channel, are associated with its homeostasis. In

animal,the action of AVP on inner ear tissues caused endolymphatic edema of the inner

ear [6,7,8]. In other words, these abnormalities in inner ear homeostasis may be

involved in causing endolymphatic hydrops. This AVP is known to be elevated by

stressors, but there is controversy. [9] A study of patients with refractory Meniere's

disease, in which plasma AVP was measured before and after surgery, reported that

plasma AVP was lower after surgery than before surgery, indicating better control of

Meniere's disease. [6] This author states that avoiding stress as much as possible is

important in controlling Meniere's disease.

Treatment of Meniere's disease consists of daily lifestyle guidance and medication for

mild cases and surgical approaches, such as intratympanic corticosteroids or

intratympanic gentamicin or endolymphatic sac release or destructive surgery, for

moderate to severe cases.[2]

DNA testing with next-generation sequencing has made it possible to investigate the

intestinal microbiota in fecal material in great detail. As a result, the changes in the

intestinal microbiota have been tested in relation to various disease groups. In this

regard, various studies have investigated the intestinal microbiota of patients with

depression and irritable bowel syndrome, in which stress factors are associated with the

induction of pathological conditions. Ménière's disease is a disease that repeatedly

worsens and improves. In the course of treating many patients with Meniere's disease,

the authors observed several patients with complaints of diarrhea, constipation, or other

abnormalities despite the absence of pre-existing bowel disease such as IBS during

exacerbations. We hypothesized that the gut microbiota of Ménière's disease patients

may differ from that of healthy individuals. To test this hypothesis, we investigated the

intestinal microbiota of 10 patients who met the diagnostic criteria for a solid case and

had significant endolymphatic edema on the affected side by HYDROPS, an objective

test.

Materials and Methods

Participants

Ten patients (6 males, 4 females; mean age, 49.6 ± 8.1 years) with unilateral definite

Meniere's disease diagnosed according to the diagnostic criteria of Journal of Vestibular

Research [1] and showing significant endolymphatic hydrops on the affected side on

HYDROPS of the inner ear were included in this study. The HYDROPS images are

presented in Fig 1. The patients did not have any psychiatric, diabetic, or

gastrointestinal diseases that may have caused dysbiosis.

Methods

This study was approved and conducted as an observational study by the Clinical

Trials Subcommittee (Observational Studies) of St. Marianna University School of

Medicine. The approval number is 5674. Informed consent was given to the subjects in

writing and orally. The subject's consent was obtained orally and documented in the

medical record. The information obtained was anonymized so that individual patients

could not be identified.

The 10 participants received a fecal collection kit and obtained fecal samples during a

period with no abnormal bowel movements. Intestinal microflora testing was performed

by an external contractor (SAIKINSO Co., Tokyo, Japan). The intestinal microflora was

analyzed as follows.

Fecal sampling, DNA extraction, and sequencing

Fecal samples were collected using Cykinso fecal collection kits containing guanidine

thiocyanate solution (Cykinso Inc., Tokyo. Japan) and stored at 4°C. DNA extraction

from the fecal samples was performed using an automated DNA extraction machine

(GENE PREP STAR PI-480; Kurabo Industries Ltd., Osaka). The V1-V2 region of the

16S rRNA gene was amplified using a forward primer (16S_27Fmod: TCG TCG GCA

GCG TCA GAT GTG TAT AAG AGA CAG AGR GTT TGA TYM TGG CTC AG) and

reverse primer (16S_338R: GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG

ACA GTG CTG CCT CCC GTA GGA GT) with the KAPA HiFi Hot Start Ready Mix

(Roche). To sequence the 16S amplicons using the Illumina MiSeq platform, dual index

adapters were attached using the Nextera XT Index kit. The DNA concentration of the

mixed libraries was quantified by qPCR using KAPA SYBR FAST qPCR. The master

mix (KK4601, KAPA Biosystems) was used with primer 1 (AAT GAT ACG GCG ACC

ACC) and primer 2 (CAA GCA GAA GAC GGC ATA CGA). Library preparations were

performed according to the Illumina 16S library preparation protocol (Illumina, San

Diego, CA, USA). Libraries were sequenced using the MiSeq Reagent Kit v2 (500

cycles) and 250-bp paired ends.

Taxonomy assignment based on 16S rRNA gene sequences

The paired-end reads of the partial 16S rRNA gene sequences were analyzed using

QIIME 2 (version 2020.8). The steps for data processing and assignment based on the

QIIME 2 pipeline were as follows: (1) DADA2 was used for joining paired-end reads,

filtering, and denoising; (2) taxonomic information was assigned to each ASV by using

a naive Bayes classifier in the QIIME 2 classifier with the 16S gene of the V1-V2

region data of SILVA (version 138) to determine the identity and composition of the

bacterial genera. The reliability of these tests has been reported previously [10,11].

Shannon diversity index (Shannon index) and Faith's phylogenetic diversity (FPD)

were used as indicators of dysbiosis in the gut microbiota. These indices were averaged

over 10000 reads per sample. The Dizziness Handicap Inventory (DHI) is a standard

questionnaire that quantitatively evaluates the degree of handicap in the daily life of

patients with vestibular disorders; it consists of 25 questions [12,13]. The total score

ranges from 0 (no disability) to 100 (severe disability). Hearing test is that pure tone

audiogram thresholds were measured at octave intervals between 125 Hz and 8 kHz.

The results of the audiometric tests that were compared used a quadrant method in

which the hearing levels at frequencies of 500, 1,000, and 2,000 Hz were a, b, and c dB,

respectively, and the value (dB) was calculated using the formula (a+2b+c)/4.

We performed statistical analysis of the obtained gut microbiota data, age, BMI,

duration of Meniere's disease, DHI results, and hearing test results (quadrant method)

by using Spearman's correlation coefficient values. The hearing test and DHI evaluation

performed closest to the intestinal microbiota tests were used for analyses. Excel with

add-in software Stacel4 (OMS, Tokyo, Japan) was used as statistical software.

Results

Table 1 shows the age, sex, BMI, duration of Meniere's disease, DHI results, hearing

test results, and dysbiosis index of intestinal bacteria. A significant negative correlation

was observed between disease duration and the Shannon index (r = -0.72, p = 0.02) and

FPD (r = -0.89, p = 0.0005) (Figs 2 and 3). However, no significant correlation was

found between age and Shannon index (r = 0.16, p = 0.64) or FPD (r = -0.35, p = 0.31),

BMI and the Shannon index (r = -0.21, p = 0.55) and FPD (r = -0.14, p = 0.68), the DHI

results and the Shannon index (r = -0.42, p = 0.22) and FPD (r = -0.08, p = 0.81), and

between the hearing test results and the Shannon index (r = -0.03, p = 0.94) and FPD (r

= -0.28, p = 0.42).

The TAXON BAR showing the intestinal bacterial layer in each case is shown in Figs

4 and 5. The organisms identified by phylum, class, order, family, genus, and species are

shown in Fig 4 (the description by bacterial species is too voluminous to be included),

and those identified by phylum, class, and order are shown in Fig 5. *Akkermansia*

muciniphila was not detected in the gut microbiota of any of the patients.

Discussion

The use of gut microbiota testing to analyze various disease groups is a fairly recent

approach. Recognition of the interrelationship between the brain and intestines via

intestinal bacteria, which constitute the so-called microbiome-gut-brain axis, and the

development of analytical scientific methods such as metagenomic analysis using next-

generation sequencers and metabolome analysis have together promoted rapid

accumulation of knowledge about the mechanisms underlying the microbiome-gut-brain

axis.[14]

In particular, the intestinal microbiota have been extensively investigated in patients

with irritable bowel syndrome (IBS) [15], and dysbiosis has also been reported to be

induced in the gut microbiota of depressed patients, showing a pattern similar to that of

IBS [16]. These disease groups show dysbiosis of the intestinal microbiota in

comparison with healthy individuals [17]. In animal studies, when the gut microbiota of

patients with depression was transplanted into gut microbiota-depleted mice, the

transplanted mice exhibited characteristics of depression, both at the behavioral and

physiological levels. Thus, dysbiosis of the gut microbiota was considered as a factor in

the pathogenesis of depression [18]. Stressors have some influence on the pathogenesis

of Meniere's disease, as reported in a study of hormones in Meniere's disease.[6] In fact,

multiple reports have described the effects of stress and depression on Meniere's disease.

Patients with refractory Meniere's disease are prone to depressive symptoms, with 40%

reported to show neurotic symptoms and 60% showing depressive symptoms [19].

Adrenocorticotropic hormone and blood cortisol levels have been shown to be elevated

due to stress in patients with Meniere's disease, and the levels of oxytocin and

vasopressin, which are known as stress hormones, are also elevated in patients with

Meniere's disease [20,21].

This study showed that dysbiosis of the intestinal microbiota progresses as the

duration of Meniere's disease increases. As mentioned above, dysbiosis of the intestinal

microbiota may have progressed with increasing exposure to stress, the etiological

factor, as the duration of illness increased. In this study, patients with a history of

psychiatric disorders or complications were excluded. In the present study, the longer

the duration of Meniere's disease, the stronger the dysbiosis, suggesting that a

prolonged Meniere's disease status may have caused dysbiosis. But , in this study, no

depression scoring was conducted, and it is unclear whether this dysbiosis was caused

by the presence of depressive tendencies in patients with Meniere's disease or whether

the pathophysiology of Meniere's disease itself is a factor causing the dysbiosis. In order

to clarify these issues, it is advisable to test the same patients at different times and

compare the results with the depression scoring. The microbial diversity indices showed

no correlation with DHI and hearing test results. Meniere's disease is known to cause

exacerbation and amelioration, but there may have been no association between the

degree of disability due to Meniere's disease and dysbiosis of the gut microbiota.

However, this study was conducted on a limited number of cases, and future

comparisons based on multiple examinations in the same patients and statistical

analyses with a larger number of cases are necessary in this regard. We could not find

any reports on the association between Meniere's disease and intestinal microflora, or

between Meniere's disease and IBS.

Dysbiosis of the intestinal microbiota has been verified to occur first or at the onset of

various diseases. For example, rheumatoid arthritis is triggered by an immune response

to specific intestinal bacteria [22]. Future studies should aim to evaluate a larger number

of cases and investigate whether a specific intestinal bacterium is a trigger for Meniere's

disease.

Notably, none of the patients with Meniere's disease had *Akkermansia muciniphila,* a

gram-negative enterobacterium belonging to the phylum *Verrucomicrobia* that has been

identified in the feces of healthy individuals. *Akkermansia muciniphila* is capable of

growing on mucin covering the gastrointestinal mucosa as a single source of nutrition

[23]. It is a commensal bacterium characterized by its degradative ability and accounts

for 1%-4% of the human intestinal microflora [24]. It has been suggested that

akkermansia muciniphila is associated with metabolic syndrome such as obesity and

diabetes, and *akkermansia muciniphila* is of interest as a probiotic that can ameliorate

metabolic syndrome. In relation to obesity, the intestinal mucus layer of mice fed a

high-fat diet that developed metabolic syndrome became thin and *akkermansia*

muciniphila decreased, but oral administration of *akkermansia muciniphila* restored the

thickness and permeability of the mucus layer in the intestinal tract and also improved

metabolic syndrome The metabolic syndrome was reported to be improved by oral

administration of *akkermansia muciniphila*[25]. In addition, human observational

studies have reported that *akkermansia muciniphila* is reduced in patients with type II

diabetes and atopic dermatitis compared to healthy controls[26,27]. *Akkermansia*

muciniphila has been shown to improve insulin resistance and have immunomodulatory

effects[25].

In mice, walking exercise was reported to increase *Akkermansia muciniphila* in the

intestinal microflora. [28] In humans, increased *Akkermansia muciniphila* of the

intestinal microflora has been reported with exercise and dietary restriction. [29] A

study in mice reported an increase in *Akkermansia muciniphila* and *Lactobacillus* when

melatonin was administered to improve sleep deprivation[30]. These reports indicate

that *Akkermansia muciniphila* increases or decreases with exercise, diet, and sleep in the

intestinal microflora. Daily lifestyle guidance for Meniere's disease generally includes

improving sleep, diet, and avoidance of stress [31]. Aerobic exercise is also reported to

be effective for Meniere's disease [32].

These lifestyle guidance may be consistent with some of the factors that increase

Akkermansia muciniphila in the intestinal microflora. In the future, testing of the

intestinal microflora before and after daily lifestyle guidance in patients with Meniere's

disease will clarify the relationship between *Akkermansia muciniphila* and Meniere's

disease.

Only 10 patients were evaluated in this study, and future studies should aim to

accumulate data from more cases. However, the fact that dysbiosis of the intestinal

microbiota depended on the duration of Meniere's disease and that *Akkermansia*

muciniphila was not detected in all cases suggests that the intestinal microbiota in

patients with Meniere's disease is different from that in normal individuals. Future

investigation of the intestinal microflora in Meniere's disease may lead to elucidation of

the pathogenesis and development of new treatments. Further studies will be required to

accumulate data from more cases.

270   Conclusions

Ten patients with a definite diagnosis of Meniere's disease and marked

endolymphatic hydrops on HYDROPS underwent examinations of the intestinal

microflora. A significant negative correlation was found between the duration of

Meniere's disease and dysbiosis of intestinal microbiota. None of the patients harbored

Akkermansia muciniphila. However, this study was limited by the small number of

patients. Nevertheless, the relationship between Meniere's disease and the intestinal

microbiota has not been reported previously, and these findings may help researchers

determine the pathogenesis of Meniere's disease and propose new treatments for it in

the future.

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

References

1.   Lopez-Escamez JA, Carey J, Chung WH et al . Diagnostic criteria for

Meniere's disease. J Vestib Res.2015; 25:1–7.Doi: 10.3233/VES-150549.

2.   Yupeng Liu , Jun Yang, Maoli Duan. Current status on researches of Meniere's

disease:a   review ActaOtolaryngol.2020;14:808812.

doi: 10.1080/00016489.2020.1776385. Epub 2020 Jun 21.

3.   Liu Y, Jia H, Shi J, et al. Endolymphatic hydrops detected by 3- dimensional

fluid-attenuated inversion recovery MRI following intratympanic injection of

gadolinium in the asymptomatic contralateral ears of patients with unilateral

Meniere's disease. Med Sci Monit. 2015;21:701–707. doi:

10.12659/MSM.892383.

4.   Martin-Sierra C, Gallego-Martinez A, Requena T, et al. Variable expressivity

and genetic heterogeneity involving DPT and SEMA3D genes in autosomal

dominant familial Meniere's disease. Eur J Hum Genet. 2017;25:200–207 doi:

10.1038/ejhg.2016.154. Epub 2016 Nov 23

5.   Cureoglu S, da Costa Monsanto R, Paparella MM. Histopathology of Meniere's

disease. Oper Tech Otolayngol Head Neck Surg. 2016;27:194–204. doi:

10.1016/j.otot.2016.10.003

6.   Tadashi Kitahara , Tadao Okayasu , Taeko Ito , Hiroto Fujita , Keita Ueda.

Endolymphatic Sac Drainage Surgery and Plasma Stress Hormone Vasopressin

Levels in Meniere's Disease. Front Neurol.2021;29;12:722217.doi:

10.3389/fneur.2021.722217.

7. Takeda T, Takeda S, Kitano H, Okada T, Kakigi A. Endolymphatic hydrops

induced by chronic administration of vasopressin. Hear Res. 2000; 140:1–6.

doi: 10.1016/S0378-5955(99)00180-X

8. Kakigi A, Egami N, Uehara N, Fujita T, Nibu KI, Yamasoba T. Live imaging

and functional changes of the inner ear in an animal model of Meniere's

disease. Sci Rep. 2020; 10:12271. doi: 10.1038/s41598-020-68352-0

9. Wu J, Zhou J, Dong L, Fan W, Zhang J, Wu C. A mysterious role of arginine

vasopressin levels in Ménière's disease-meta-analysis of clinical studies. Otol

Neurotol. (2017) 38:161–7. doi: 10.1097/MAO.0000000000001310

10. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et

al. Reproducible, interactive, scalable and extensible microbiome data science

using QIIME 2. Nat Biotechnol. 2019;37: 852-857. Doi: 10.1038/s41587-019-

0209-9.

11. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA

ribosomal RNA gene database project: Improved data processing and web-

based tools. Nucleic Acids Res. 2013;41: D590–D596. Doi:

10.1093/nar/gks1219.

12. Goto F, Tsutsumi T, Ogawa K. The Japanese version of the Dizziness Handicap

Inventory as an index of treatment success; exploratory factor analysis. *Acta*

Otolaryngol. (2011) 131:817–5. doi: 10.3109/00016489.2011.565423

13. Jacobson GP, Newman CW. The development of the Dizziness Handicap

Inventory. *Arch Otolaryngol Head Neck Surg.* (1990) 116:424–7. doi:

10.1001/archotol.1990.01870040046011

14. Katarzyna Socała , Urszula Doboszewska , Aleksandra Szopa,et al. The role of

microbiota-gut-brain axis in neuropsychiatric and neurological disorders.

Pharmacol Res. 2021;172:105840. doi: 10.1016/j.phrs.2021.105840.

15. Ji S, Park H, Lee D, Song YK, Choi JP, Lee SI. Post-infectious irritable bowel

syndrome in patients with Shigella infection. J Gastroenterol Hepatol. 2005;20:

381-386. Doi: 10.1111/j.1440-1746.2005.03574.x.

16. Liu Y, Zhang L, Wang X, Wang Z, Zhang J, Jiang R, et al. Similar fecal

microbiota signatures in patients with diarrhea-predominant irritable bowel

syndrome and patients with depression. Clin Gastroenterol Hepatol. 2016;14:

1602–1611.e5. doi: 10.1016/j.cgh.2016.05.033.

17.   Sanada K, Nakajima S, Kurokawa S, Barceló-Soler A, Ikuse D, Hirata A, et al.

Gut microbiota and major depressive disorder: A systematic review and meta-

analysis. J Affect Disord. 2020;266: 1-13. Doi: 10.1016/j.jad.2020.01.102.

18.   Kelly JR, Borre Y, O' Brien C, Patterson E, El Aidy S, Deane J, et al.

Transferring the blues: Depression-associated gut microbiota induces

neurobehavioural changes in the rat. J Psychiatr Res. 2016;82: 109-118. Doi:

10.1016/j.jpsychires.2016.07.019.

19.   Furukawa M, Kitahara T, Horii A, Uno A, Imai T, Ohta Y, et al. Psychological

condition in patients with intractable Meniere's disease. Acta Otolaryngol.

2013;133: 584-589. Doi: 10.3109/00016489.2012.759274.

20.   Horner KC, Cazals Y. Stress hormones in Ménière's disease and acoustic

neuroma. Brain Res Bull. 2005;66: 1-8. Doi:

10.1016/j.brainresbull.2005.04.003.

21.   van Cruijsen N, Dullaart RP, Wit HP, Albers FW. Analysis of cortisol and other

stress-related hormones in patients with Ménière's disease. Otol Neurotol.

2005;26: 1214-1219. Doi: 10.1097/01.mao.0000179528.24909.ba.

22.   Maeda   Y,Takeda   K.Host－rnicrobiota interactions in   rheumatoid

arthritis. Exp   Mol   Med. 2019：51：1-6, doi: 10.1038/s12276-019-0283-6

23.   Derrien M, Vaughan EE, Plugge CM, de Vos WM. Akkermansia muciniphila

gen. Nov., sp. Nov., a human intestinal mucin-degrading bacterium. Int J Syst

Evol Microbiol. 2004;54: 1469-1476. Doi: 10.1099/ijs.0.02873-0.

24.   Collado MC, Derrien M, Isolauri E, de Vos WM, Salminen S. Intestinal

integrity and Akkermansia muciniphila, a mucin-degrading member of the

intestinal microbiota present in infants, adults, and the elderly. Appl Environ

Microbiol. 2007;73: 7767-7770. Doi: 10.1128/AEM.01477-07.

25.   Everard A,Belzer C,Geurts L et al: Cross-talk between Akkermansia

muciniphilia and intestional epithelium controls diet-induced obesity . ProcNatl

Acad Sci USA.2013;110:9066-9071, doi: 10.1073/pnas.1219451110.

26.   Chelakkot, C., Choi, Y., Kim, D.K., Park, H.T., Ghim, J.,Kwon, Y., et al.

Akkermansia muciniphila-derived   extracellular vesicles influence gut

permeability   through the regulation of tight junctions. Exp Mol Med.2018;

50: e450. doi: 10.1038/emm.2017.282.

27.   Candela, M., Rampelli, S., Turroni, S., Severgnini, M., Consolandi, C., De

Bellis, G, et al. Unbalance of intestinal microbiota in atopic children.BMC

Microbiol 2012 12: 95 doi: 10.1186/1471-2180-12-95.

28.   Bin Wang , Yu-Xiao Jin , Jia-Li Dong ,et al. Low-Intensity Exercise Modulates

Gut Microbiota to Fight Against Radiation-Induced Gut Toxicity in Mouse

Models. Front Cell Dev Biol. 2021 21;9:706755.

doi: 10.3389/fcell.2021.706755.

29.   Zhuye Jie , Xinlei Yu , Yinghua Liu,et al. The Baseline Gut Microbiota Directs

Dieting-Induced Weight Loss Trajectories.

Gastroenterology.2021 ;160(6):2029-2042.e16. doi: 10.1053/j.gastro.

30.   Young Sook Park , Soo Hyung Kim , Jong Won Park , et al. Melatonin in the

colon modulates intestinal microbiota in response to stress and sleep

deprivation. Intest Res. 2020 Jul;18(3):325-336. doi: 10.5217/ir.2019.00093.

31.   J Nevoux , M Barbara , J Dornhoffer,et al. International consensus (ICON) on

treatment of Ménière's disease. Eur Ann Otorhinolaryngol Head Neck Dis

. 2018 ;135(1S):S29-S32. doi: 10.1016/j.anorl.2017.12.006.

32.   Masahiro Takahashi. Introducing aerobic exercise as part of a novel therapeutic

strategy for Meniere's disease. Equilibrium Res 2011; 70 204-211.

doi:org/10.3757/jser.70.204

 Table1 Detailed data on Meniere's patients

No	Age	BMI	Sex	Disease period (year)	DHI	Hearing test (dB)	Shannon's index	FPD
1	54	22.31	Male	3	46	7.5	6.34	18.41
2	39	26.78	Male	13	56	26.3	5.34	11.17
3	53	26.37	Female	0.67	16	31.3	6.58	19.69
4	60	19.81	Female	6	20	58.8	6.25	14.48
5	54	35.11	Male	7	8	26.3	6.02	15.83
6	47	25.6	Male	5	20	13.8	5.9	13.55
7	36	23.38	Female	10	54	36.3	5.77	12.48
8	61	25.65	Female	6	86	22.5	5.14	14.45
9	46	20.28	Male	0.58	62	21.3	6.7	21.12
10	46	22.91	Male	6	38	35	5.3	12.88
Average	49.6	24.82		5.725	40.6	27.91	5.934	15.406

Short arrow is cochlea, Long arrow is vestibular

Fig 1**. HYDROPS image in case 1.** The short arrows indicate the cochlea, while the

long arrow indicates the vestibule.

Fig 2.

417   r = -0.72, p = 0.02

421 **Fig 3.**

422 r = -0.89, p = 0.0005

Fig 4. Identification by phylum, class, order, family, genus, and species.

D_0__Bacteria;D_1__Bacteroidota;D_2__Bacteroidia;D_3__Bacteroidales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Lachnospirales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Oscillospirales

D_0__Bacteria;D_1__Actinobacteriota;D_2__Actinobacteria;D_3__Bifidobacteriales

D_0__Bacteria;D_1__Firmicutes;D_2__Negativicutes;D_3__Veillonellales-Selenomonadales

D_0__Bacteria;D_1__Proteobacteria;D_2__Gammaproteobacteria;D_3__Burkholderiales

D_0__Bacteria;D_1__Actinobacteriota;D_2__Coriobacteriia;D_3__Coriobacteriales

D_0__Bacteria;D_1__Firmicutes;D_2__Bacilli;D_3__Erysipelotrichales

D_0__Bacteria;D_1__Firmicutes;D_2__Bacilli;D_3__Lactobacillales

D_0__Bacteria;D_1__Firmicutes;D_2__Negativicutes;D_3__Acidaminococcales

D_0__Bacteria;D_1__Proteobacteria;D_2__Gammaproteobacteria;D_3__Enterobacterales

D_0__Bacteria;D_1__Proteobacteria;D_2__Alphaproteobacteria;D_3__Rhodospirillales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Monoglobales

D_0__Bacteria;D_1__Fusobacteriota;D_2__Fusobacteriia;D_3__Fusobacteriales

D_0__Bacteria;D_1__Desulfobacterota;D_2__Desulfovibrionia;D_3__Desulfovibrionales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Peptostreptococcales-Tissierellales

D_0__Bacteria;D_1__Proteobacteria;D_2__Gammaproteobacteria;D_3__Aeromonadales

D_0__Bacteria;D_1__Firmicutes;D_2__Bacilli;D_3__Bacillales

D_0__Bacteria;D_1__Spirochaetota;D_2__Brachyspirae;D_3__Brachyspirales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;__

D_0__Bacteria;D_1__Cyanobacteria;D_2__Vampirivibrionia;D_3__Gastranaerophilales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Clostridiales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Eubacteriales

D_0__Bacteria;D_1__Proteobacteria;D_2__Gammaproteobacteria;D_3__Pasteurellales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Clostridia UCG-014

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Christensenellales

D_0__Bacteria;D_1__Actinobacteriota;D_2__Actinobacteria;D_3__Actinomycetales

D_0__Bacteria;D_1__Patescibacteria;D_2__Saccharimonadia;D_3__Saccharimonadales

D_0__Bacteria;D_1__Firmicutes;D_2__Clostridia;D_3__Peptococcales

D_0__Bacteria;D_1__Firmicutes;D_2__Bacilli;D_3__Staphylococcales

D_0__Bacteria;D_1__Cyanobacteria;D_2__Cyanobacteriia;D_3__Chloroplast

D_0__Bacteria;D_1__Campilobacterota;D_2__Campylobacteria;D_3__Campylobacterales

D_0__Bacteria;D_1__Bacteroidota;D_2__Bacteroidia;D_3__Flavobacteriales

D_0__Bacteria;D_1__Actinobacteriota;D_2__Actinobacteria;D_3__Micrococcales

Fig 5. Identification by phylum, class, and order.