**The hidden diversity of ancient bornaviral sequences from X and P genes in**

**vertebrate genomes**

Bea Clarise B. Garcia [1], Yahiro Mukai [2,3], Keizo Tomonaga [2,3,4], & Masayuki Horie [1,5*]

[1] Laboratory of Veterinary Microbiology, Graduate School of Veterinary Science, Osaka

Metropolitan University, Izumisano, Osaka, Japan

[2] Laboratory of RNA Viruses, Institute for Frontier Life and Medical Sciences, Kyoto

University, Kyoto, Japan

[3] Laboratory of RNA Viruses, Department of Mammalian Regulatory Network,

Graduate School of Biostudies, Kyoto University

[4] Department of Molecular Virology, Graduate School of Medicine, Kyoto University

[5] Osaka International Research Center for Infectious Diseases, Osaka Metropolitan

University, Izumisano, Osaka, Japan

**\*Corresponding author**

Email: mhorie@omu.ac.jp

**Abstract**

Endogenous bornavirus-like elements (EBLs) are heritable sequences derived from

bornaviruses in vertebrate genomes that originate from transcripts of ancient

bornaviruses. EBLs have been detected using sequence similarity searches such as

tBLASTn, whose technical limitations may hinder the detection of EBLs derived from

small and/or rapidly evolving viral X and P genes. Indeed, no EBLs derived from the X

and P genes of orthobornaviruses have been detected to date in vertebrate genomes.

Here, we aimed to develop a novel strategy to detect such "hidden" EBLs. To this aim,

we focused on the 1.9 kb read-through transcript of orthobornaviruses, which encodes a

well-conserved N gene and small and rapidly evolving X and P genes. We show a series

of evidence supporting the existence of EBLs derived from orthobornaviral X and P

genes (EBLX/Ps) in mammalian genomes. Furthermore, we found that an EBLX/P is

expressed as a fusion transcript with the cellular gene, *ZNF451*, which potentially

encodes the ZNF451/EBLP fusion protein in miniopterid bat cells. This study

contributes to a deeper understanding of ancient bornaviruses and co-evolution between

bornaviruses and their hosts. Furthermore, our data suggest that endogenous viral

elements detected thus far are just the tip of the iceberg, and further studies are required

to understand ancient viruses more accurately.

**Introduction**

Ancient viral sequences can be found within the genomes of organisms in the form of endogenous viral elements (EVEs) (1). EVEs are derived from the integration of viral sequences into the host germline DNA and transfer of the integrated sequences to subsequent progenies (1, 2). As molecular fossils of ancient viruses, EVEs have expanded our knowledge of the diversity, host range, and long-term evolution of viruses (1). Furthermore, some EVEs are co-opted by hosts to function as novel gene regulatory elements, non-coding RNAs, and proteins (2).

Most EVEs are derived from retroviruses, as the viruses reverse-transcribe and integrate the viral genomes into the host genomes using virally encoded enzymes. However, non-retroviral EVEs have also been identified. Endogenous bornavirus-like elements (EBLs) are EVEs derived from bornaviruses (family *Bornaviridae*), a group of non-segmented negative-strand RNA viruses (3, 4). Although bornaviruses are RNA viruses, evidence strongly suggests that bornaviral mRNA is reverse transcribed and integrated into the host genome via the activity of the long interspersed nuclear element 1 (LINE-1) retrotransposon (3, 5). We previously revealed an abundance of EBLs in vertebrate genomes derived from the bornaviral genes N, P, M, G, and L (S1 Fig), which are referred to as EBLN, EBLP, EBLM, EBLG, and EBLL, respectively (6). These EBLs originated from ancient bornaviruses of all genera in the family *Bornaviridae*: *Orthobornavirus*, *Carbovirus*, and *Cultervirus*.

Despite the rigorous search, there has been a bias in the detection of EBLs. EBLNs comprise the majority of the detected EBLs, while no EBLs derived from the X and P genes have been reported except for a few carboviral EBLPs (3, 6-10). We proposed two hypotheses to explain the lack of EBLX and EBLP. First, X/P mRNA, which

encodes the X and P proteins, may have a low propensity for reverse-transcription

and/or integration into the host genome. However, previous studies of Borna disease

virus 1 (BoDV-1; genus *Orthobornavirus*) showed that X/P mRNA is the most abundant

transcript among the viral mRNAs (11, 12), which should therefore provide enough

chances for integration. Moreover, we have previously demonstrated that the X/P

mRNA of BoDV-1 can be reverse-transcribed and integrated into the genome of host

cells (3, 7). This led us to the second hypothesis, which states that EBLs derived from

the X and P genes are present but are difficult to detect due to technical limitations of

the sequence similarity search. EBLs have been routinely searched using tBLASTn;

however, this can be an unsuitable method for detecting small and rapidly evolving

genes (8). Indeed, the X and P genes are small and the least conserved among bornaviral

genes (Fig 1). If the second hypothesis is correct, some EBLs may remain undetected.

Therefore, we could be missing important insights into ancient bornaviruses and the co-

evolution of bornaviruses and their hosts.

In this study, we sought to develop a novel strategy to detect these hidden EBLs by

focusing on the nature of the read-through transcription of orthobornaviruses that

produce a transcript encoding a highly conserved N gene and rapidly evolving X and P

genes. We show evidence supporting the existence of several EBLX/Ps in vertebrate

genomes. Furthermore, an EBLX/P in miniopterid bats appear to be transcribed as

chimeric transcripts with a host gene, suggesting a potential co-option of EBLs by the

host.

**Results**

***Miniopterid bat EBLN is possibly derived from an N/X/P transcript***

We previously identified miEBLN-1, an EBLN derived from an ancient orthobornavirus

that retains an intact N ORF in miniopterid bats (13). The miEBLN-1 locus is present in

the genomes of miniopterid bats, specifically in those of *Miniopterus natalensis*, *M.*

*schreibersii*, and *M. fuliginosus*, but is empty (i.e., syntenic loci lacking miEBLN-1) in

the genomes of *Rhinolophus ferrumequinum* and *Phyllostomus hastatus*. We re-

examined the presence of miEBLN-1 using the current genome datasets and found the

miEBLN-1-empty locus in *Tadarida brasiliensis* (Fig 2a and S2a Fig).

Based on the nucleotide alignment, we deduced the insertion size of the miEBLN-1

locus to be 1.9 kb, consisting of a 1.2-kb miEBLN-1 region and a 0.7-kb unannotated

region. Therefore, we carried out a series of analyses to understand the source of the

0.7-kb unannotated region. First, we performed a BLASTn search (14) against the

NCBI nt database (15) using the nucleotide sequence of the 0.7-kb unannotated region

as a query. The hits consisted only of miEBLN-1 sequences (data not shown). Next, we

assessed whether the 0.7-kb unannotated region is derived from a short interspersed

nuclear element (SINE) due to the precedence of chimeric integration comprising

nucleotide sequences from an N transcript and SINE (3, 5). We performed a BLASTn

search against the miniopterid bat genomes using the nucleotide sequence of the 0.7-kb

region as a query. We only detected five BLAST hits in the genomes of *M. natalensis*

and *M. schreibersii*, which we identified as EBLs in the later sections. We also analyzed

the sequence of the 0.7-kb region by CENSOR (16), but no hits were detected. These

results indicated that the 0.7-kb fragment is not derived from a SINE.

We noticed that the insertion, including miEBLN-1, is almost similar in size to the

N/X/P mRNA of extant orthobornaviruses (Fig 2a and S2a Fig), which led us to

speculate that the insertion may be derived from an N/X/P transcript. N/X/P mRNA is

produced by read-through of the termination signal T1, resulting in a 1.9 kb mRNA

containing the transcription signal sequences S1, T1/S2, and T2 (Fig 2b) (12).

Therefore, we investigated the presence of these signal-like sequences in the 1.9-kb

miEBLN-1 insertion. We found transcription signal-like sequences at reasonable

positions with respect to extant N/X/P transcripts (Fig 2c and S3b Fig). A poly-A stretch

was also observed immediately downstream of the T2 signal, indicating that the

insertion originated from an mRNA transcript (Fig 2c and S3b Fig). Furthermore, a

tandem duplication sequence, TTGTGCTA/TTCTTGTA, was observed immediately

upstream and downstream of S1 and T2-poly-A sequences, respectively (Fig 2c and S3b

Fig). This is a potential target site duplication (TSD), which is a hallmark of LINE-1-

mediated mRNA integration (17). These results suggested that the 1.9-kb insertion,

including miEBLN-1, is derived from an ancient N/X/P transcript integrated by LINE-1.

Therefore, we tentatively refer to this locus as miEBLN/X/P-1.

***Additional EBLs possibly derived from N/X/P transcripts in miniopterid bat genomes***

As described above, orthobornaviruses express mRNAs encoding N (1.2 kb), X and P

(X/P mRNA, 0.7 kb), or N, X, and P (N/X/P mRNA, 1.9 kb) (Fig 2a). If miEBLN/X/P-1

is truly derived from N/X/P mRNA, there may be additional EBLs derived from N/X/P

or X/P mRNA. To examine this possibility, we conducted a BLASTn search against bat

genomes using the nucleotide sequence of miEBLN/X/P-1 as a query. We obtained six

hits from the genomes of *M. natalensis* and *M. schreibersii* (Fig 3a and S1 Table). While

four hits were aligned in the entire N/X/P region of miEBLN/X/P-1 (Fig 3a, red lines),

interestingly, two other hits were aligned only to the putative X/P region (Fig 3a blue

lines). Note that the N regions of four hits aligned to entire miEBLN/X/P-1 were

detected as EBLNs in a previous study (6).

We first assessed hits that aligned to the entire region of miEBLN/X/P-1. Four hits

showed 1.9 kb alignment lengths and high sequence similarities (76–84% nucleotide

identity) to miEBLN/X/P-1 (S1 Table). The flanking regions of each pair of hit

sequences were well aligned between *M. natalensis* and *M. schreibersii*, indicating that

they are orthologous between these species (S7 Table). We tentatively refer to these loci

as miEBLN/X/P-2 and miEBLN/X/P-3. We also confirmed the presence of orthologs of

these EBLN/X/Ps in the genome of *M. fuliginosus* by PCR and Sanger sequencing of

genomic DNA isolated from *M. fuliginosus* (S2b Fig and S7 Table).

Next, we examined the presence of miEBLN/X/P-2 or miEBLN/X/P-3- empty loci

by comparative genomics. We observed that the miEBLN/X/P-3 empty locus is present

in genomes of non-miniopterid bats such as *Tadarida brasiliensis* (S2b Fig and S7

Table). We did not obtain an alignment of the flanking region of miEBLN/X/P-2 with

the other bat genomes and therefore could not identify its empty locus.

We then analyzed the nucleotide sequences of miEBLN/X/P-2 and miEBLN/X/P-

3. Similar to miEBLN/X/P-1, both EBLN/X/Ps contained transcription signal-like

sequences, and poly-A stretches were observed downstream of T2-like sequences (S3b

Fig). Furthermore, tandem duplication sequences, TGTTCA/[T/C]GTTCA and

GAGGAACG/GAGGGATG, were also observed immediately upstream of S1 and

downstream of T2/poly-A in miEBLN/X/P-2 and miEBLN/X/P-3, respectively (S3b

Fig). These results suggest that miEBLN/X/P-2 and miEBLN/X/P-3 are also derived

from ancient N/X/P transcripts and were integrated into the genome of a miniopterid bat

ancestor through LINE-1 activity.

***EBLs derived from X/P-like transcripts exist in miniopterid bat genomes***

Next, we assessed the hits that aligned to the 0.7-kb putative X/P region of

miEBLN/X/P-1 (Fig 2a, teal blue lines). Two hits from *M. natalensis* and *M.*

*schreibersii* showed 0.8 kb alignment lengths and 75% nucleotide identity to the 0.7-kb

X/P region (S1 Table). The flanking regions of the hit sequences were well aligned,

indicating that the loci are orthologous between these species (Fig 3b and S4a Fig; S7

Table). We also found an orthologous locus in the genome of *M. fuliginosus* using PCR

and Sanger sequencing (Fig 3b and S4a Fig; S7 Table). Furthermore, we detected empty

loci in genomes of non-miniopterid bats, genomes such as *Myotis lucifugus* (Fig 3b and

S4a Fig; S7 Table).

The orthobornaviral X/P mRNA contains the transcription signal sequences S2/T1

in the 5'-UTR and T2 in the 3'-UTR (Fig 3c). We observed that S2/T1 and T2-like

sequences are present in the 0.7-kb insertion, which are located at positions similar to

those in extant X/P mRNA (Fig 3d and S5b Fig). Poly-A stretches were also observed

downstream of the T2-like sequences. Furthermore, the S2 and poly-A stretches were

flanked by TSDs, CTTAAGGC/CTTAAGGC (Fig 3d and S5b Fig). Taken together,

these results indicated that the insertion may have been derived from an ancient X/P

transcript and integrated into the genome of a miniopterid bat ancestor through LINE-1

activity. We tentatively refer to the insertion as the miEBLX/P-4.

***Preservation of intact ORFs in miEBLX/P-4 and characterization of the putatively***

***encoded proteins***

Most EBLs do not retain viral ORFs because of the accumulation of mutations after

integration. Remarkably, miEBLX/P-4 retains intact X- and P-like ORFs in different

reading frames, which is consistent with the extant orthobornaviral X/P mRNA (Fig 3c).

We named these ORFs as miEBLX-4 and miEBLP-4. Notably, miEBLP-4 in *M.*

*natalensis* is disrupted by a premature stop codon.

Although there is very low sequence similarity between the amino acid sequences

of these ORFs and extant orthobornaviral proteins, further analyses showed that

miEBLX-4 and miEBLP-4 are intrinsically similar to extant orthobornaviral X and P

proteins.

First, we examined the miEBLX-4 ORF. The ORF consists of 109 codons, which is

longer than that of extant orthobornaviral X (86–90 codons). We then predicted the

secondary structure and analyzed the hydrophobicity of the putative miEBLX-4 protein

with respect to extant X proteins. Both extant X and miEBLX-4 were predicted to

contain an alpha helix in the N-terminal region (Fig 4a and S6a Fig). The hydropathy

plot showed similar hydrophobicity patterns for these proteins (S7a Fig).

We also determined whether the functional domains are conserved between extant

X and miEBLX-4. Amino acid sequence alignment showed that the N-terminal region is

relatively well conserved and houses functional domains previously characterized in

BoDV-1 X (Fig 4c). This region includes the leucine-rich domain functioning both as a

CRM1-dependent nuclear export signal (18) and a mitochondria-targeting signal (19),

and is overlapped by the importin-alpha binding domain (20). The alignment also

showed that leucine and arginine residues that are critical in the importin-alpha-binding

domain (Fig 4c; orange and blue, respectively) are conserved among extant X and

miEBLX-4.

Next, we examined the miEBLP-4 ORF. The ORF consists of 233 codons, longer

than those of extant orthobornaviral P (201–208 codons). The predicted secondary

structures and hydrophobicity pattern of miEBLP-4p were similar to those of the

modern counterparts (Fig 4d, S6b Fig, and S7b Fig). Alignment of the amino acid

sequences revealed that some of the functional domains previously characterized in

BoDV-1 P may be shared between extant P and miEBLX-4 (Fig 4d). Basic and proline

residues important for the nuclear localization signal (NLS) activity (Fig 4d, blue) (21,

22) in the N-terminal NLS were prominent in both proteins.

We also explored the interaction between miEBLX-4 and miEBLP-4, which is a

notable characteristic of the orthobornaviral X and P proteins (23). We expressed Myc-

tagged miEBLX-4 (miEBLX-4-Myc) and FLAG-tagged miEBLP-4 (miEBLP-4-FLAG)

in 293T cells and then immunoprecipitated them using an anti-FLAG antibody.

miEBLX-4-Myc was co-immunoprecipitated with miEBLP-4-FLAG, indicating the

interaction between these proteins (Fig 4e).

These results showed that miEBLX-4 and miEBLP-4 share similar intrinsic

properties with the extant X and P proteins. Furthermore, together with the above data,

they strongly suggest that miEBLN/X/P-1-3 and miEBLX/P-4 are indeed derived from

ancient viral N/X/P and X/P mRNAs, respectively.

***Diverse EBLX/Ps are found in vertebrate genomes***

To further identify the presence of other EBLX/Ps, we conducted TFASTX and

TFASTY searches against vertebrate genomes using the amino acid sequences of

miEBLX-4 and miEBLP-4 as queries. TFASTX and TFASTY are sequence similarity

search programs that allow frameshifts for alignment (24). As a result, 65 unique hits

(S2-5 Tables) were detected in the genomes of mammals (class Mammalia), reptiles

(class Reptilia), amphibians (class Amphibia), and ray-finned fish (class

Actinopterygii). The majority of these hits were detected in the bat genomes, including

miEBLN/X/P-1, miEBLN/X/P-2, miEBLN/X/P-3, and miEBLX/P-4.

Comparative genomic analyses strongly suggest that at least some of the

TFASTX/Y hits are indeed derived from insertions of ancient viral X/P transcripts, as

follows. TFASTX/Y hits in the bat families Mormoopidae and Phyllostomidae

(mpEBLX/P) are orthologous, and their empty loci are present in other lineages of bats,

such as *Noctilio leporinus* (Fig 5a and S4b Fig). For a TFASTX/Y hit in the lone bat

species *Craseonycteris thonglongyai* (ctEBLX/P), we could not find its ortholog, but its

empty sites exist in other bat species (Fig 5b and S4c Fig). We further identified

orthologous TFASTX/Y hits in the rodent superfamily Octodontoidea and their empty

loci in other lineages of rodents (ocEBLX/P; Fig 5c and S4d Fig). We also identified

transcription signal-like sequences and TSDs within and flanking the above TFASTX/Y

hits, respectively (Fig 5d and S5b Fig). These results strongly suggested that at least

some of the sequences detected by TFASTX and TFASTY are bona fide EBLX/Ps.

***Possible co-option of miEBLX/P-4 in miniopterid bats***

Some EVEs, including EBLs, retain intact ORFs which encode proteins that play roles

in host-related functions (25). As described above, miEBLX/P-4 contains intact ORFs

corresponding to orthobornaviral X and P genes. Therefore, we analyzed miEBLX/P-4

in detail.

We analyzed the expression of miEBLX/P-4 by mRNA sequencing of *M.*

*schreibersii*-derived SuBK12-08 cells. miEBLX/P-4 is located within the intron region

of *ZNF451* gene. The read coverage of the miEBLX/P-4 locus was comparable to that

of *ZNF451* (Fig 6a). Interestingly, the mapping pattern revealed the existence of an

alternative splicing site for *ZNF451* that could produce a chimeric transcript comprising

ZNF451 exons 1–4 and the miEBLX/P-4 locus (Fig 6a).

The putative chimeric transcript contains two ORFs consisting of more than 100

codons in different reading frames: a chimeric ORF of ZNF451 in-frame with miEBLP-

4 or that of the miEBLX-4 (Fig 6b). The chimeric ORF potentially encodes a fusion

protein comprising the N-terminal region of ZNF451, which contains a SUMO-

interacting motif (SIM) domain, and miEBLP-4 (Fig 6b).

These results suggest that miEBLX/P-4 has been co-opted by miniopterid bats as a

chimeric gene with *ZNF451*.

**Discussion**

We previously demonstrated that the X/P mRNA of a modern orthobornavirus can be

reverse-transcribed and integrated into the host genome (3, 7); however, no EBLX and

only a few EBLPs have been reported (6). We hypothesized that the technical

limitations of the sequence similarity search hinder the detection of EBLs derived from

the X and P genes. EBLs have been routinely searched using tBLASTn, but this can be

an unsuitable method for detecting small and rapidly evolving genes such as the X and P

genes (8). In this study, our aim was to develop a novel strategy to detect these hidden

EBLs. To this aim, we focused on the 1.9-kb read-through transcript of

orthobornaviruses, consisting of a well-conserved N gene and small and rapidly

evolving X and P genes (12), to detect hidden EBLX/Ps. We show a series of evidence

implying that EBLN/X/Ps and EBLX/Ps are present in vertebrate genomes but are not

detectable by BLAST searches (Figs 2 and 3). Interestingly, miEBLX/P-4 has

maintained intact ORFs for more than 17 million years (Fig 3) and is expressed as a

fusion transcript with the cellular gene, *ZNF451*, which potentially encodes the

ZNF451/miEBLP-4 fusion protein (Fig 6). Therefore, this study revealed the presence

of hidden EBLs, which could contribute to a deeper understanding of ancient

bornaviruses and the co-evolution between bornaviruses and their hosts.

Here, we uncovered the potential co-option of miEBLX/P-4 by miniopterid bats, as

indicated by the expression of the ZNF451/miEBLP-4 chimeric transcript, which

potentially encodes a fusion protein consisting of the N-terminal region of ZNF451 and

miEBLP-4 (Fig 6). To the best of our knowledge, this is the first report of a non-

retroviral EVE being expressed as a chimeric transcript with a known host gene.

Although we could not confirm the expression of the ZNF451/miEBLP-4 fusion protein

due to the lack of antibodies, our data suggest that miEBLX/P-4 has been co-opted by

miniopterid bats. Further studies are needed to understand the biological significance of

miEBLX/P-4.

Our data strongly suggest that the genome organization and expression patterns of

the N, X, and P genes have not substantially changed over the past million years. Here,

we show that T1-read-through transcription occurred in ancient orthobornaviruses

present at least 17.3 MYA (Fig 2). Furthermore, the ORF positions of the X and P genes

of ancient orthobornaviruses were highly similar to those of modern orthobornaviruses

(Fig 3). In addition, the sequences and order of transcription signals are highly

conserved among ancient and modern orthobornaviruses (Figs 2 and 3). In BoDV-1,

read-through transcription has been suggested to be involved in the regulation of viral

gene expression (12). This hypothetical gene regulation mechanism has been

maintained for more than 17.3 million years. Interestingly, our previous study showed

that the order of S2-T1 is also conserved in some EBLNs in primates, which were

estimated to be integrated into the host genomes more than 43 MYA (3, 5, 9). Thus, this

gene regulatory mechanism may have been employed by orthobornaviruses earlier than

17.3 MYA.

The fundamental functions of viral X and P proteins may also be evolutionarily

conserved between ancient and modern orthobornaviruses. We observed similarities in

the intrinsic properties of extant orthobornaviral X and P proteins with miEBLX-4 and

miEBLP-4 proteins despite their very low amino acid identities (Fig 4, S6 Fig, and S7

Fig). Furthermore, the interaction between X and P may have been conserved in ancient

orthobornaviruses (Fig 4e). Therefore, although the evolutionary rates of X and P genes

are high, the fundamental features and functions of X and P proteins may have been

maintained for at least 17.3 million years.

For the EVE search, TFASTX/Y may be more suitable than tBLASTn, a

conventional sequence similarity search for the detection of EVEs. After integration into

the host genome, many EVEs have acquired random mutations, which caused

frameshifts in the ORFs. Such frameshifted EVEs are a kind of "fragmented" EVEs and

are thus less detectable by tBLASTn. In contrast, TFASTX/Y takes frameshifts into

account (24) and can thus detect such fragmented EVEs more sensitively. We are

currently conducting a benchmark analysis of tBLASTn and TFASTX/Y in detail,

showing that some of the EBLX/Ps detected by TFASTX/Y cannot be detected by

tBLASTn or other BLAST programs (data not shown). Therefore, TFASTX/Y can

detect fragmented EVEs more sensitively than tBLASTn, which may contribute to

identifying other hidden EVEs.

The EBLX/Ps identified in this study would represent only the tip of the iceberg on

the hidden EVEs that could exist in the genome. In this study, the majority of EBLX/Ps

were detected in bat and rodent genomes. Note that we failed to detect any EBLX/Ps in

primate genomes, which contain numerous EBLs. Considering the number of EBLs in

the genomes of primates (i.e., 24 orthobornaviral EBLs were detected in the human

genome (6)), it is plausible that hidden EBLX/Ps exist in primates that are undetectable

because of the inherent low amino acid similarities in X and P.

We may be able to detect such hidden EBLX/Ps by identifying other EBLN/X/P

integrations. However, searching for other EBLN/X/Ps by manual curation is time-

consuming. Thus, the development of an efficient searching method to identify

approximately 1.9-kb insertions with EBLN would facilitate further exploration of

EBLX/Ps. This is probably not limited to bornaviruses; there may be many hidden

EVEs that are derived from small and/or rapidly evolving viral genes. Read-through

transcription has also been observed for other mononegaviruses (26). Our search

strategy can be applied to the detection of such EVEs. Additionally, a previous study

identified virus-like insertions using a machine learning-based method, but the origins

of insertions are still unclear (27). Our approach might also contribute to understanding

the origins of those sequences.

Taken together, we identified EBLs that were undetectable by conventional

BLAST searches. Our findings provide novel insights into ancient bornaviruses and co-

evolution between bornaviruses and their hosts. Furthermore, our search strategy will

contribute to paleovirological research.

## Materials and Methods

### *Sanger sequencing of EBLs and flanking loci in M. fuliginosus*

Genomic DNA was obtained from YubFKT1 (kidney epithelial cell line) (28) in a previous study (13). miEBLN/X/P-1, miEBLN/X/P-2, miEBLN/X/P-3, and miEBLX/P-4 and corresponding flanking loci in *M. fuliginosus* were amplified by PCR using the extracted DNA with Phusion Hot Start II High-Fidelity PCR Master Mix (Thermo Fisher Scientific). The PCR conditions are available upon request. The amplicons were purified using innuPREP PCRpure Lite Kit (Analytik Jena) and sequenced by Sanger dideoxy method at Eurofins Genomics. The primer sequences are available in S6 Table. The obtained sequences were deposited to DDBJ (accession numbers LC708263-LC708266).

### *Cell lines*

SuBK 12-08 cells (29) and human embryonic kidney 293T (HEK293T) cells were cultured in low-glucose Dulbecco's Modified Eagle Medium (Nacalai Tesque) supplemented with 10% fetal bovine serum (Sigma-Aldrich) and 100 μg/mL penicillin-streptomycin (Nacalai Tesque). The cells were maintained at 37°C in a humidified chamber containing 5% $CO_2$.

### *Construction of miEBLX-4 and miEBLP-4 expressing plasmids*

Expression vectors with C-terminal FLAG or Myc tag were first constructed. pcDNA3 was digested with EcoRI and XhoI, which was assembled with oligo DNA containing Myc or FLAG tag sequence using NEBuilder HiFi DNA Assembly Master Mix (New

England Biolabs). The resultant plasmids were named pcDNA3-C-Myc or pcDNA3-C-

FLAG. Then, miEBLX-4 and miEBLP-4 were amplified by PCR using the genomic

DNA extracted from YubFKT1 with Phusion Hot Start II High-Fidelity PCR Master

Mix, which were subsequently assembled with pcDNA3-C-Myc or pcDNA3-C-FLAG

linearized with KpnI and EcoRI using NEBuilder. The oligo DNA sequences are

available in S6 Table.

***Co-immunoprecipitation of miEBLX-4 with miEBLP-4***

pcDNA3-miEBLX-4-myc and pcDNA3-miEBLP-4-FLAG plasmids (5 μg) were

transfected into 293T cells on 10-cm dish using 20 μL of Avalanche-Everyday

Transfection Reagent (EZ Biosystems) with Opti-MEM (Thermo Fisher Scientific). One

day post-transfection, the cells were washed twice with ice-cold PBS and lysed using

500 μL IP buffer (150 mM NaCl, 20 mM Tris-HCl pH 7.4, 1 mM EDTA, 1% Triton X-

100, $1\times$ protease inhibitor cocktail [Nacalai Tesque]) for 30 min at 4°C with rotation.

In parallel, 500 μg SureBeads Protein G magnetic beads (Bio-Rad) were incubated with

3 μg mouse monoclonal anti-DDDDK antibody FLA-1 (M185-3L; MBL) in IP buffer

for 1 hr at 4°C with rotation. The lysates were then centrifuged at $12000 \times$ g for 10 min

at 4°C, and then the supernatant were immunoprecipitated with the antibody-bead

complex in IP buffer for 2 hr in 4°C with rotation. After incubation, the beads were

washed thrice with ice-cold IP buffer and then transferred into new microtubes. The

proteins were finally eluted from the beads by boiling in SDS sample buffer for 5 min at

95°C.

*Western blotting*

Protein samples were separated in 12% polyacrylamide gel in Tris-glycine-SDS buffer, which were transferred to PVDF membrane (MERCK) at 10 V for 1 hr using Trans-Blot Turbo Transfer System (Bio-Rad). Then, the membrane was blocked with 5% skim milk in 0.1% Tween 20 in PBS (PBST) for 1 hr at room temperature (RT). The membrane was incubated with mouse monoclonal anti-Myc My3 (MBL; 1:5000) or anti-DDDDK FLA-1 antibody (MBL; 1:2000) diluted in 5% skim milk for 1 hr at RT, washed, and then incubated with horseradish peroxidase-conjugated donkey polyclonal anti-mouse IgG (715-035-150, Jackson ImmunoResearch; 1:2000) antibody diluted in 5% skim milk for 45 min at RT. The membrane was washed again, after which chemiluminescence reaction was conducted using Chemi-Lumi One (Nacalai Tesque) and was subsequently viewed using Amersham Imager 680 (GE Healthcare, IL, USA).

*BLAST search against bat genomes to detect EBLN/X/Ps and EBLX/Ps*

BLASTn (14) search was conducted against the WGS database (15) of bat (Chiroptera, taxid: 9397) using the nucleotide sequence of miEBLN/X/P-1 in *M. natalensis* as a query on the BLAST web server (https://blast.ncbi.nlm.nih.gov/Blast.cgi). Changes applied to the default parameters were as follows: E-value threshold = $10^{-10}$, word size = 7, no filter for low complexity regions.

*TFASTX/Y search against vertebrate genomes to detect EBLX/Ps*

Vertebrate genomes were downloaded using NCBI Datasets and were used as the database for TFASTX/Y searches. TFASTX/Y searches, included in the FASTA36 package (30), were performed against each species of the vertebrate genomes using the

putative amino acid sequences miEBLX-4 and miEBLP-4 as queries with the ktup value

option of 1. The obtained hits with E-values less than $10^{-4}$ were extracted, which were

used for the downstream analyses.

***Identification of orthologous and syntenic empty loci***

BLASTn search was conducted against the WGS database of bats (Chiroptera, taxid:

9397) or rodents (Rodentia, taxid: 9989) using the nucleotide sequences of EBL and its

flanking regions as queries on the BLAST web server. Changes applied to the default

parameters were as follows: E-value = $10^{-20}$, word size = 11. The BLAST hits which

comprised of genomic sequences from different species were retrieved, and then aligned

by MAFFT using the E-INS-i algorithm (31) in Geneious version 11.1.5. The nucleotide

accessions and region of EBLs used for search queries are available in S7 Table.

***Annotation of EBLs and TSDs***

TSD sequences were determined based on the alignment made in "*Identification of*

*orthologous and syntenic empty loci*". To identify transcription signal-like sequences,

the nucleotide sequences of EBLs and extant orthobornaviral N/X/P or X/P transcripts

were aligned by MAFFT using the E-INS-I algorithm. The transcription signal-like

sequences and poly-A stretches were determined according to the resultant alignment.

To identify conserved functional domains of the miEBLX-4p and miEBLP-4p, the

amino acid sequences of miEBLX-4p and miEBLP-4p in *M. fuliginosus* were aligned

with extant orthobornaviral X and P proteins, respectively, by MAFFT using the E-INS-

i algorithm. Based on the alignments, the conserved domains were manually analyzed.

The nucleotide accession of extant orthobornaviruses used in the analysis are

available in S8 Table.

***mRNA sequencing analysis***

Total RNA was extracted from SuBK12-08 cells using NucleoSpin RNA plus

(MACHEREY-NAGEL) according to the manual. Sequencing library was constructed

using TruSeq Stranded mRNA Library Prep Kit (Illumina). The libraries were

sequenced by NextSeq 500 using NextSeq 500/550 Mid Output Kit v2.5 (300 Cycles).

The obtained reads were preprocessed by fastp 0.23.2 (32) with the default setting,

which were mapped to the reference genome of *M. natalensis* (GCF_001595765.1) by

HISAT2 version 2.2.1 (33) with the default setting. The mapped reads and splicing

junctions were visualized using Integrated Genomics Viewer 2.13.0 (34).

The RNA-seq data were deposited to DDBJ Sequence Read Archive (DRA) under

the accession number DRR403400.

***Pairwise similarity analysis***

Amino acid sequences of the viral proteins of extant orthobornaviruses (S8 Table) were

analyzed by MAFFT alignment tool using Sequence Demarcation Tool software version

1.2 (35). Minimum value for the percent identity scale was set to 0.1. The resulting

matrices were set to 3-color mode and exported.

***In silico characterization of putative EBLX and EBLP proteins***

To predict the protein secondary structure, amino acid sequences of miEBLX-4 and

miEBLP-4 in *M. fuliginosus*, and extant orthobornaviral X and P proteins, were

analyzed by JPred version 4 (36).

To determine the protein hydropathy, the same amino acid sequences were

submitted to ProtScale (37) in ExPASy Server, using the amino acid scale developed by

Kyte & Doolittle. The window sizes were set to 9 and 15 for X and P proteins,

respectively.

**Acknowledgments**

We are grateful to Ayato Takada for kindly providing SuBK 12-08 cell. We would like

to thank Ken Maeda for kindly sharing YubFKT1. We are grateful to Takefumi Kondo

and Yukari Sando for conducting the mRNA-sequencing. The animal silhouette images

were downloaded under public domain access from https://commons.wikimedia.org/

and https://publicdomainvectors.org/.

This study was supported by JSPS KAKENHI grant numbers JP21H01199 (MH),

20H05682 (KT), JP21K19909 (KT), JSPS Core-to-Core Program JPJSCCA20190008

(KT), and MEXT KAKENHI grant number JP19H04833 (MH).

**References**

1.    Aiewsakun P, Katzourakis A. Endogenous viruses: Connecting recent and

ancient viral evolution. Virology. 2015;479-480:26-37.

2.    Jern P, Coffin JM. Effects of retroviruses on host genome function. Annu Rev

Genet. 2008;42:709-32.

3.    Horie M, Honda T, Suzuki Y, Kobayashi Y, Daito T, Oshida T, et al.

Endogenous non-retroviral RNA virus elements in mammalian genomes. Nature.

2010;463(7277):84-7.

4.       Rubbenstroth D, Briese T, Durrwald R, Horie M, Hyndman TH, Kuhn JH, et al.

ICTV Virus Taxonomy Profile: Bornaviridae. J Gen Virol. 2021;102(7).

5.       Belyi VA, Levine AJ, Skalka AM. Unexpected inheritance: multiple

integrations of ancient bornavirus and ebolavirus/marburgvirus sequences in vertebrate

genomes. PLoS pathogens. 2010;6(7):e1001030.

6.       Kawasaki J, Kojima S, Mukai Y, Tomonaga K, Horie M. 100-My history of

bornavirus infections hidden in vertebrate genomes. Proc Natl Acad Sci U S A.

2021;118(20).

7.       Horie M, Kobayashi Y, Suzuki Y, Tomonaga K. Comprehensive analysis of

endogenous bornavirus-like elements in eukaryote genomes. Philosophical transactions

of   the   Royal   Society   of   London   Series   B,   Biological   sciences.

2013;368(1626):20120499.

8.       Horie M, Tomonaga K. Paleovirology of bornaviruses: What can be learned

from molecular fossils of bornaviruses. Virus Res. 2019;262:2-9.

9.       Katzourakis A, Gifford RJ. Endogenous viral elements in animal genomes.

PLoS genetics. 2010;6(11):e1001191.

10.      Gilbert C, Meik JM, Dashevsky D, Card DC, Castoe TA, Schaack S.

Endogenous hepadnaviruses, bornaviruses and circoviruses in snakes. Proc Biol Sci.

2014;281(1791):20141122.

11.    Kojima S, Sato R, Yanai M, Komatsu Y, Horie M, Igarashi M, et al. Splicing-

Dependent Subcellular Targeting of Borna Disease Virus Nucleoprotein Isoforms.

Journal of virology. 2019;93(5).

12.    Poenisch M, Wille S, Staeheli P, Schneider U. Polymerase read-through at the

first transcription termination site contributes to regulation of borna disease virus gene

expression. Journal of virology. 2008;82(19):9537-45.

13.    Mukai Y, Horie M, Kojima S, Kawasaki J, Maeda K, Tomonaga K. An

endogenous bornavirus-like nucleoprotein in miniopterid bats retains the RNA-binding

properties of the original viral protein. FEBS letters. 2022;596(3):323-37.

14.    Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. BMC Bioinformatics. 2009;10:421.

15.    Coordinators NR. Database resources of the National Center for Biotechnology

Information. Nucleic Acids Res. 2018;46(D1):D8-D13.

16.    Jurka J, Klonowski P, Dagman V, Pelton P. CENSOR--a program for

identification and elimination of repetitive elements from DNA sequences. Comput

Chem. 1996;20(1):119-21.

17.    Morrish TA, Gilbert N, Myers JS, Vincent BJ, Stamato TD, Taccioli GE, et al.

DNA repair mediated by endonuclease-independent LINE-1 retrotransposition. Nat

Genet. 2002;31(2):159-65.

18.    Ferre CA, Davezac N, Thouard A, Peyrin JM, Belenguer P, Miquel MC, et al.

Manipulation of the N-terminal sequence of the Borna disease virus X protein improves

its mitochondrial targeting and neuroprotective potential. FASEB J. 2016;30(4):1523-

33.

19.    Poenisch M, Burger N, Staeheli P, Bauer G, Schneider U. Protein X of Borna

disease virus inhibits apoptosis and promotes viral persistence in the central nervous

systems of newborn-infected rats. Journal of virology. 2009;83(9):4297-307.

20.    Wolff T, Unterstab G, Heins G, Richt JA, Kann M. Characterization of an

unusual importin alpha binding motif in the borna disease virus p10 protein that directs

nuclear import. J Biol Chem. 2002;277(14):12151-7.

21.    Shoya Y, Kobayashi T, Koda T, Ikuta K, Kakinuma M, Kishi M. Two proline-

rich nuclear localization signals in the amino- and carboxyl-terminal regions of the

Borna disease virus phosphoprotein. Journal of virology. 1998;72(12):9755-62.

22.    Schwemmle M, Jehle C, Shoemaker T, Lipkin WI. Characterization of the

major nuclear localization signal of the Borna disease virus phosphoprotein. J Gen

Virol. 1999;80 ( Pt 1):97-100.

23.    Fujino K, Horie M, Honda T, Nakamura S, Matsumoto Y, Francischetti IM, et

al. Evolutionarily conserved interaction between the phosphoproteins and X proteins of

bornaviruses from different vertebrate species. PLoS One. 2012;7(12):e51161.

24.    Pearson WR, Wood T, Zhang Z, Miller W. Comparison of DNA sequences with

protein sequences. Genomics. 1997;46(1):24-36.

25.    Horie M. The biological significance of bornavirus-derived genes in mammals.

Current opinion in virology. 2017;25:1-6.

26.    Banerjee AK, Barik S, De BP. Gene expression of nonsegmented negative

strand RNA viruses. Pharmacology & Therapeutics. 1991;51(1):47-70.

27.    Kojima S, Yoshikawa K, Ito J, Nakagawa S, Parrish NF, Horie M, et al. Virus-

like insertions with sequence signatures similar to those of endogenous nonretroviral

RNA viruses in the human genome. Proceedings of the National Academy of Sciences.

2021;118(5).

28.    Maeda K, Hondo E, Terakawa J, Kiso Y, Nakaichi N, Endoh D, et al. Isolation

of novel adenovirus from fruit bat (Pteropus dasymallus yayeyamae). Emerging

infectious diseases. 2008;14(2):347-9.

29.    Maruyama J, Miyamoto H, Kajihara M, Ogawa H, Maeda K, Sakoda Y, et al.

Characterization of the envelope glycoprotein of a novel filovirus, lloviu virus. Journal

of virology. 2014;88(1):99-109.

30.    Pearson WR. Finding Protein and Nucleotide Similarities with FASTA. Curr

Protoc Bioinformatics. 2016;53:3 9 1-3 9 25.

31.    Katoh K, Standley DM. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80.

32.    Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. Bioinformatics. 2018;34(17):i884-i90.

33.    Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol.

2019;37(8):907-15.

34.    Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G,

et al. Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24-6.

35.    Muhire BM, Varsani A, Martin DP. SDT: a virus classification tool based on

pairwise sequence alignment and identity calculation. PLoS One. 2014;9(9):e108277.

36.    Drozdetskiy A, Cole C, Procter J, Barton GJ. JPred4: a protein secondary

structure prediction server. Nucleic Acids Res. 2015;43(W1):W389-94.

37.    Walker JM. The proteomics protocols handbook: Springer; 2005.

38.    Kumar S, Stecher G, Suleski M, Hedges SB. TimeTree: A Resource for

Timelines, Timetrees, and Divergence Times. Mol Biol Evol. 2017;34(7):1812-9.

**Figure legends**

**Figure 1. Amino acid identities between extant orthobornaviral proteins.** Pairwise

identities between the amino acid sequences of extant orthobornaviral proteins was

analyzed using Sequence Demarcation Tool (35). The animal silhouettes show the

representative hosts of viruses.

**Figure 2. Characterization of miEBLN-1 as an EBL derived from N/X/P transcript.**

**(a)** Gene orthology analysis of the miEBLN/X/P-1 locus. Multiple alignment of the

deduced syntenic loci between the indicated species is shown. Black and gray boxes

show aligned regions. Thin gray lines indicate alignment gaps. Species phylogeny and

minimum integration age of miEBLN/X/P-1 are indicated on the left based on TimeTree

(38). The full version of this alignment is available in S1 Fig. **(b)** Schematic diagram of

orthobornaviral genomic RNA, 1.9-kb N/X/P mRNA, and miEBLN/X/P-1.

Transcription signals (S1, S2, T1, and T2) are shown. **(c)** Nucleotide sequence

alignment of extant orthobornaviral N/X/P mRNA and miEBLN/X/P-1. Red letters

indicate the start and stop codons of the N gene. Light and dark teal blue letters indicate

the start codon of X gene and the stop codon of P gene, respectively. TSD, target site

duplication. The full version of this alignment is available in S2 Fig.

**Figure 3. Identification of an EBL derived from X/P transcript in miniopterid bats.**

**(a)** Schematic diagram of the BLASTn search against the bat whole genome shotgun

sequence database, using miEBLN/X/P-1 as a query. The BLAST hits are shown by

dark to light read or teal blue lines corresponding to the EBLs indicated on the right. **(b)**

Gene orthology analysis of the miEBLX/P-4 locus. Species phylogeny and deduced

minimum integration age of miEBLX/P-4 are indicated on the left based on TimeTree

(38). miEBLX-4 and miEBLP-4 ORFs are shown in light and dark teal blue,

respectively. The full version of this alignment is available in S3 Fig. **(c)** Schematic

diagram of extant orthobornaviral genomic RNA, X/P mRNA, and miEBLX/P-4.

Transcription signals (S2, T1, and T2) are indicated. **(d)** Nucleotide sequence alignment

of extant orthobornaviral X/P transcripts and miEBLX/P-4. Light and dark teal blue

letters indicate the start codon of X gene and the stop codon of P gene, respectively.

TSD, target site duplication. The full version of this alignment is available in S4 Fig.

**Figure 4. Conservation of properties between putative EBLX and EBLP and viral**

**X and P proteins. (a** and **b)** Predicted secondary structure of extant orthobornaviral X

and miEBLX-4 proteins **(a)**, and extant orthobornaviral P and miEBLP-4 proteins **(b)**.

H, α-helix. E, β-strand. The data including amino acid residues are shown in Figs. S5

and S6. **(c** and **d)** Amino acid sequence alignment of extant orthobornaviral X and

miEBLX proteins **(c)**, and extant orthobornaviral P and miEBLP-4 proteins **(d)**. Shaded

areas indicate the functional domains previously characterized in BoDV-1 X and P

proteins. The conserved amino acid residues important for the functions of viral X or P

protein are highlighted in colors. NLS, nuclear localization signal; NES nuclear export

signal; p-Ser, phosphorylated serine residue. **(e)** Co-immunoprecipitation of Myc-tagged

miEBLX-4 with FLAG-tagged miEBLP-4. FLAG-tagged miEBLP-4 was

immunoprecipitated, and then analyzed by western blotting using the indicated

antibodies. *, Non-specific band.

**Figure 5. Identification of diverse EBLX/Ps in the vertebrate genomes.**

**(a-c)** Gene orthology analysis of each EBLX/P locus. Species phylogeny and deduced

minimum integration age of each EBLX/P are indicated on the left based on TimeTree

(38). The full version of this alignment is available in S3 Fig. **(d)** Nucleotide sequence

alignment of extant orthobornaviral X/P transcripts and EBLX/Ps. Black and gray boxes

show aligned regions. Thin gray lines indicate alignment gaps. Light and dark teal blue

letters indicate the start codon of X gene and the stop codon of P gene, respectively.

TSD, target site duplication. The full version of this alignment is available in S4 Fig.

**Figure 6. Alternative splicing generates *ZNF451/miEBLX/P-4* chimeric transcript.**

**(a)** Coverage and splicing junctions of mapped RNA reads from SuBK 12-08 cells (*M.*

*schreibersii*) at the *ZNF451* locus where miEBLX/P-4 is located. The miEBLX/P-4

region, mapped read coverage graph on miEBLX/P-4, and splicing junction between

ZNF451 and miEBLX/P-4 are shown in teal blue. **(b)** Schematic figure of putative

chimeric ZNF451/miEBLP-4 protein. Shaded teal blue box SIM, SUMO interacting

motif.

**Fig. 1**

**Fig. 2**

**a**

Insert (1.9 kb)

(kb) 0    1    2    3    4   4.6

% Identitiy

*Rhinolophus ferrumequinum*
*Phyllostomus hastatus*
*Tadarida brasiliensis*
*Miniopterus fuliginosus* — miEBLN-1
*Miniopterus natalensis* — miEBLN-1
*Miniopterus schreibersii* — miEBLN-1

miEBLN/X/P-1
17.2 MYA

**b**

Genome RNA        N   X/P   M   G   L        8.9 kb

N/X/P mRNA        N   X/P   AAA(n)   1.9 kb

miEBLN/X/P-1      miEBLN-1             1.9 kb

Transcription signal    S1    S2    S1/S2 - Transcription start sites
                        T1    T2    T1/T2 - Transcription termination sites

**c**

|  | TSD | S1 | | S2 | T1 | | T2 + Poly-A | TSD |
|---|---|---|---|---|---|---|---|---|

BoDV-1 N/X/P mRNA          GAATACA------CGCAATG  TAGAAAACAATGAA-CAAACCAATAAAAAACCAAATGCGGCAAACCCCCCGCGACCTGTGATG  TAAAAAAAA(n)

PaBV-2 N/X/P mRNA          GAACACAA----CAACATG   -------TAATGAA-CATACGAATAAAAAACCATAT--------------------CGATG  TAAAAAAAA(n)

LGSV-1 N/X/P mRNA          GAACAAAA----CAAAAATG  ----------TGAA-CAAACCTATATAAAACCATAT--------------------AGATG  TAAAAAAAA(n)

CWBV N/X/P mRNA            GAACCAAATCCACAAAAATG  -------TAATGAACCATTCATATAGAAAACCGAATTC-----------------CAATG   TAAAAAAAA(n)

miEBLN/X/P-1 in *M. natalensis*   TTGTGCTAGGAACCAA------CACAATG  ----TAATATCGAA-CCAACATTTTCAAAAACATA--------------------GATG  TAAAATAAATATATACACCATTCTTGTA

**Fig. 3**

**a**

miEBLN/X/P-1 in *M. natalensis*

Query

miEBLN-1

BLASTn hits

- miEBLN/X/P-1
- miEBLN/X/P-2
- miEBLN/X/P-3
- miEBLX/P-4

N/X/P mRNA

N · X · P

**b**

(kb) 0 · 1 · Insert (0.8 kb) · 2 · 3 · 3.9

% Identitiy

*Rousettus aegyptiacus*

*Phyllostomus hastatus*

*Myotis lucifugus*

*Miniopterus fuliginosus* — miEBLP-4 / miEBLX-4

miEBLX/P-4
17.2 MYA — *Miniopterus natalensis* — miEBLX->

*Miniopterus schreibersii* — miEBLP-4 / miEBLX-4

**c**

Genome RNA — N · X · P · M · G · L — 8.9 kb

X/P mRNA — X · P · AAA(n) — 0.7 kb

miEBLX/P-4 — miEBLX-4 / miEBLP-4 — 0.8 kb

Transcription signal — S1 · S2 · T1 · T2 — S1/S2 - Transcription start sites
T1/T2 - Transcription termination sites

**d**

| | TSD | S2 | T1 | | T2 + Poly-A | TSD |
|---|---|---|---|---|---|---|
| BoDV-1 X/P mRNA | | GAACAAAC-CAATAAAAAACCAAATGCGGCAAACCCCCCGCGACCTGTGATG | | | CATAAAAAAA(n) | |
| PaBV-2 X/P mRNA | | GAACATAC-GAATAAAAAACCATAT---------------------CGATG | | | CATAAAAAAA(n) | |
| LGSV-1 X/P mRNA | | GAACAAAC-CTATATAAAACCATAT-----------------AGATG | | | TCTAAAAAAA(n) | |
| CWBV X/P mRNA | | GAACCATTCATATAGAAAACCGAA------------------TTCCAATG | | | TCTAAAAAAA(n) | |
| miEBLX/P-4 in *M. schreibersii* | CTTAAGGCGAACCAAC-ATTTACAAAAACATAG---------------ATGTCATCT | | | ATTAAAAACTAATAATAATAAATAAATAAATAAAATACTTAAGGC | |

**Fig. 4**

**Fig. 5**

**Fig. 6**

**a**

*ZNF451* gene                                               73.9 kb

*ZNF451* mRNA isoforms   X1, X2, X3, X4

miEBLX/P-4

Read coverage    560   0

Splicing junction

Putative chimeric mRNA of *ZNF451* and miEBLX/P-4

**b**

Putative chimeric protein of ZNF451 and miEBLP-4

1     105   132        364 aa

ZNF451       miEBLP-4

31-IQFVSEGPLRPVLEYIDL-49

SIM           SIM