Unraveling the hidden diversity of endogenous bornavirus-like elements derived

from ancient orthobornaviral X and P genes in mammalian genomes

Bea Clarise B. Garcia [1], Yahiro Mukai [2,3], Keizo Tomonaga [2,3], & Masayuki Horie [1,4*]

[1] Laboratory of Veterinary Microbiology, Graduate School of Veterinary Science, Osaka

Metropolitan University, Izumisano 598-8531, Osaka, Japan

[2] Laboratory of RNA Viruses, Institute for Frontier Life and Medial Sciences, Kyoto

University, Kyoto, Japan

[3] Department of Mammalian Regulatory Network, Graduate School of Biostudies,

Kyoto University

[4] Osaka International Research Center for Infectious Diseases, Osaka Metropolitan

University, Izumisano 598-8531, Osaka, Japan

*Corresponding author

Masayuki Horie, DVM, PhD

Tel: +81-72-463-5694

Email: mhorie@omu.ac.jp

Keywords: endogenous viral element, bornavirus, orthobornavirus, paleovirology,

readthrough transcript

22
23

Abstract

Endogenous bornavirus-like elements (EBLs) are heritable sequences derived from

bornaviruses in vertebrate genomes, which are strongly suggested to originate from

transcripts of ancient bornaviruses. EBLs have been detected by sequence similarity

searches, such as tBLASTn, and thus EBLs derived from small and/or rapidly evolving

viral genes, such as viral X and P genes, are difficult to be detected. Indeed, no EBLs

derived from X and P genes of orthobornaviruses were detected. Thus, although

previous studies comprehensively analyzed the presence of EBLs in vertebrate

genomes, there may still be undetectable EBLs. In this study, we developed a novel

strategy to detect such "hidden" EBLs by focusing on the nature of readthrough

transcription of orthobornavirus. We showed a series of evidence supporting that there

are EBLs derived from orthobornaviral X and P genes in mammalian genomes.

Therefore, this study contributes to a deeper understanding of the evolution of ancient

viruses and of virus-host relationships. Further, our data also suggest that endogenous

viral elements detected thus far are just the tip of the iceberg, and thus further studies

are required to understand ancient viruses more accurately.

Introduction

Ancient viruses from millions of years ago left fossils buried within the genomes of organisms in the form of endogenous viral elements (EVEs) [1]. These unconventional molecular fossils were derived via integration of viral sequences in the germline DNA of their hosts and subsequential transmission of the integrated sequences to the host offspring [1]. As fossils of ancient viruses, EVEs have facilitated the study of diversity and long-term evolution of viruses. Further, some EVEs were eventually co-opted by the hosts to function as novel gene regulatory elements, non-coding RNAs, and proteins [2].

Most of the discovered EVEs are derived from retroviruses due to reverse transcription and integration of viral genomes into the host genomes as part of the retroviral infection cycle using the virally encoded enzymes. However, non-retroviral EVEs have also been discovered. Endogenous bornavirus-like elements (EBLs) are EVEs that are derived from bornavriuses (family *Bornaviridae*), a group of non-segmented negative strand RNA viruses [3]. The family *Bornaviridae* is further divided into three genera: *Orthobornavirus*, *Carbovirus*, and *Cultervirus* [4]. Our previous study revealed that almost all the detected EBLs are divided into three clades corresponding to the classification of modern bornaviruses [5]. A series of evidence strongly suggest that EBLs have been formed by reverse-transcription and integration of viral mRNA via the activity of LINE-1, a retrotransposon [3, 6]. Bornaviruses have six genes, N, X, P, M, G, and L, among which N, P, M, G, and L-derived EBLs (EBLN, EBLP, EBLM, EBLG, and EBLL) were identified thus far [5].

There is a bias on the numbers of detected integrated viral genes (transcripts): among the identified EBLs, no EBLs derived from X or P genes were found, with the exception of a few carboviral EBLPs [5, 7]. There are two hypotheses to explain the bias. First, X/P

transcripts may be rarely reverse-transcribed and/or integrated into the host genome.

However, the amount of X/P transcript is comparable to that of the N transcript in cells

infected with Borna disease virus 1 (BoDV-1), a modern orthobornavirus [8], suggesting

that there is enough chance to be integrated into the host genome. Further, we

experimentally demonstrated that X/P mRNA of BoDV-1 could be integrated into the host

genomes [9]. The second hypothesis is that EBLX/Ps are present, but we may not be able

to detect them by the conventional sequence similarity searches due to the technical

limitation. EBLs were mainly detected by tBLASTn-based method, therefore, small and

rapidly evolving genes are difficult to be detected [7]. Indeed, X and P genes are small

and the least conserved among the genes encoded by bornaviruses (Fig. 1). If we could

detect such hidden EBLs, it would contribute to a deeper understanding of ancient viruses

and virus-host coevolution.

In this study, we developed a novel strategy to detect EBLs that were previously

undetectable by focusing on the nature of readthrough transcription of orthobornaviruses.

We here showed a series of evidence supporting that there exist several EBLX/Ps that

were undetectable in the conventional tBLASTn searches.

Results

Identification of N/X/P transcript-like EBLs in miniopterid bat genomes

We previously found miEBLN-1, which is derived from an ancient orthobornaviral N

gene and retains the intact ORF, in miniopterid bats (Fig. 2a) and the miEBLN-1 "empty"

loci in the genomes of *Rhinolophus ferrumequinum* and *Phyllostomus hastatus* [10]. To

confirm these results, we reanalyzed the miEBLN-1 empty loci using the current genome

datasets. We confirmed the above results and further identified an miEBLN-1 empty locus

in *Tadarida brasiliensis* (Fig. 2c and Fig. S1a). Interestingly, the insertion is 1.9 kb

consisting of 1.2-kb miEBLN-1 region and 0.7-kb unannotated region.

To obtain insight into the origin of the 0.7-kb region, we performed a BLASTn

search against NCBI nt database using the sequence of 0.7-kb region as a query. We did

not find any sequence hit except for the miEBLN-1 mRNA sequences (data not shown).

We and others previously found a chimeric EBL integrant consisting of N mRNA and a

retrotransposon, short interspersed nuclear element (SINE) [3, 6]. To assess whether the

0.7-kb unannotated region is SINE or not, we performed a BLASTn search against the

miniopterid bat genomes using the nucleotide sequence of 0.7-kb region as a query,

showing that there are only 5 BLAST hits in each *M. natalensis* and *M. schreibersii*

genomes that are considered to be other EBLs (see below sections). We also analyzed the

sequence by CENSOR [11], but no hit was detected in the 0.7-kb region. These results

suggest that the 1.9 kb insertion is not derived from a chimeric integrant of N mRNA and

SINE.

The 1.9-kb insertion is almost identical in length to the N/X/P mRNA of

orthobornaviruses (Fig. 2a), and thus we speculated that this insertion is derived from an

ancient N/X/P mRNA. To assess the above hypothesis, we analyzed the sequences of the

1.9-kb insertion and its flanking region. We found viral transcription signal-like

sequences (S1, T1/S2, and T2) at reasonable positions and a poly-A stretch downstream

of T2 (Fig. 2b and Fig. S2). This sequence organization is almost identical to that of the

N/X/P mRNA of orthobornaviruses (Fig. 2a). Thus, these results suggest that the 1.9-kb

insertion including miEBLN-1 is derived from an ancient N/X/P transcript. Thus, we

tentatively named this insertion miEBLN/X/P-1.

Detection of additional N/X/P transcript-like EBLs in miniopterid bat genomes

Given that miEBLN/X/P-1 is derived from an N/X/P mRNA, other N/X/P and X/P

mRNA-derived EBLs may be present in bat or even other animal genomes. To assess this

possibility, we performed a BLASTn search using EBLN/X/P-1 as a query against bat

(order Chiroptera) genomes in the NCBI WGS database. We obtained 8 BLAST hits,

which can be aligned to the whole N/X/P or X/P regions in miEBLN/X/P-1, from each of

the *M. natalensis* and *M. schreibersii* genomes (Fig. S3 and Table S1).

The two of three BLAST hits from each of the miniopterid bat genomes are alignable

to the entire region of miEBLN/X/P-1 and show high sequence similarities (76-84%

nucleotide identities) to miEBLN/X/P-1 (Table S1). The flanking region of each hit can

be aligned between *M. natalensis* and *M. schreibersii*, implying that they are orthologous

(Fig. 2d and Fig. S1b). Therefore, we tentatively named these BLAST hits miEBLN/X/P-

2 and miEBLN/X/P-3 (Table S1). We also analyzed the genome of *M. fuliginosus* by PCR

and Sangar-sequencing, showing that *M. fuliginosus* also have these EBLs (Fig. S1).

Both miEBLN/X/P-2 and miEBLN/X/P-3 retain viral transcription signal-like

sequences, S1, T1/S2, and T2 (Fig. 2b and Fig. S2). Additionally, both has a poly-A stretch

downstream   of   the   T2-like   sequence.   Further   a   tandem   duplication

(TTGTGCTA/TGTTCA)   and   (GAGGAACG/GAGGGATG)   was   also   observed

immediately upstream and downstream of S1 and T2-poly-A sequences of miEBLN/X/P-

2 and miEBLN/X/P-3, respectively, which is probably a hallmark of LINE-1-mediated

integration, called target site duplication (TSD). These results strongly suggest that

miEBLN/X/P-1, 2, and 3 are derived from ancient N/X/P mRNAs that were processed by

LINE-1.

Detection of X/P transcript-like EBLs in miniopterid bat genomes

By the above BLASTn analysis, we also found approximately 0.8-kb BLAST hits that

can be aligned only to the putative X/P region (0.7-kb region) of miEBLN/X/P-1 with

75% nucleotide identity (Fig. S3 and Table S1). These BLAST hits are orthologous

between *M. natalensis* and M. *schreibersii* as their flanking regions were well aligned

(Fig. 3c and Fig. S4a). We found that viral transcription signal-like sequences, S2/T1 and

T2, are located at the reasonable positions, and the T2-like sequences are followed by a

poly-A stretch (Fig. 3a-3b and Fig. S5). Further, a TSD-like sequence is surrounding the

BLAST hits. Further, our comparative genomic analysis identified empty loci in other

species of bats, revealing the integrant is approximately 0.8 kb (Fig. 3c and Fig. S4a).

These observations suggest that miEBLX/P-4 is derived from X/P mRNA of an

orthobornavirus.

We also found putatively intact X- and P-like ORFs in the miEBLX/P-4 of *M.*

schreibersii and *M. fuliginosus* (Fig. 3c and Fig. S4a), although the P-like ORF in *M.*

natalensis was disrupted by a premature stop codon. We therefore name these X- and P-

like ORFs as miEBLX and miEBLP, respectively. The miEBLX and miEBLP consist of

109 and 233 codons, respectively, which are slightly longer than those of

orthobornaviruses (X: 86-90 codons; P: 201-208 codons). The putative amino acid

sequences of the miEBLX and miEBLP are totally different from those of the X and P

proteins of orthobornaviruses. However, a series of *in silico* analyses revealed that the

biochemical properties of miEBLX and miEBLP are similar to those of modern

orthobornavirus X and P protein, respectively (Fig. 4a-4b and Fig. S6-S7).

Secondary structure prediction analysis showed that orthobornaviral X and miEBLX

have an alpha helix at the N-terminal region (Fig. 4a and Fig. S6a). Hydropathy plot

showed similar patterns of viral X and miEBLX (Fig. S7a). The detailed sequence

analyses revealed that the N-terminal region is the most conserved region among the viral

X and miEBLX, which houses two functional domains that were previously characterized

in BoDV-1 X (Fig. 4c). The N terminal region of BoDV-1 X protein contain leucine-rich

domain functioning both as a CRM1-dependent nuclear export signal and mitochondria-

targeting signal [12-14].

The miEBLP protein also show similar results. A predicted secondary structure and

hydropathy plot of miEBLP protein are similar to those of orthobornavirus P proteins (Fig.

4b and Fig. S6b). Hydropathy plot showed similar patterns of viral P and miEBLP (Fig.

S7b). We then aligned miEBLP with the extant P proteins and checked the presence of

functional domains that were previously characterized in BoDV-1 P (Fig. 4d). These

results suggest that the miEBLX and miEBLP are intact ORFs.

Together with the above results, our data strongly suggest that miEBLN/X/P-1, 2, 3

and miEBLX/P-4 are derived from N/X/P and X/P transcripts of ancient

orthobornaviruses.

Detection of diverse EBLX/Ps in bat and rodent genomes

To further identify EBLX/Ps in mammalian genomes, we performed a TFASTY

search [15] against mammalian genomes using the putative amino acid sequences of

miEBLX and miEBLP. TFASTY is a sequence similarity search program that allows

frame shifts, which may be more sensitive than tBLASTn for EVE search because EVEs

often contain frame shifts due to accumulation of mutations. By the TFASTY search we

identified additional EBLX/Ps in bat, rodent, and solenodon genomes (Fig. S3b-S3c and

Table S2-S3).

By comparative genomics, we confirmed that at least some of the detected EBL

candidates are inserted sequences (Fig. 3d-3f and Fig. S4b-S4d). Further, we identified

the hallmarks of EBLX/Ps in some of the TFASTY hits, such as transcription signal-like

sequences and TSDs (Fig. 3b and Fig. S5). These results strongly suggest that the

sequences detected by TFASTY, if not all, are indeed EBLX/Ps. Taken together, divergent

EBLX/Ps are present in mammalian genomes that were not detectable by the conventional

detection method using tBLASTn alone.

Discussion

Although many EBLs were identified thus far, no X and only a few P-derived EBLs were

reported. There were two hypotheses to explain this observation: (i) X/P mRNA is rarely

reverse-transcribed and integrated into host genomes, and (ii) there are X/P-derived EBLs

which are undetectable by the conventional BLAST searches due to the technical

limitation. In this study, we focused on the nature of readthrough transcription of

orthobornaviruses that produces approximately 1.9-kb N/X/P transcripts and applied this

to exploration of EBLs derived from viral X and P genes. We showed a series of evidence

that EBLX/Ps are indeed present in mammalian genomes but not detectable by the

conventional BLAST searches. Thus, this study could reveal hidden EBLX/Ps, which

contribute to a deeper understanding of ancient bornaviruses.

The EBLX/Ps identified in this study may be just the tip of the iceberg: namely, we

are probably still missing many EBLs derived from X/P genes. In this study, we identified

the 1.9-kb integrant (miEBLN/X/P-1) in the miniopterid bat genomes by chance and did

not further explore this kind of EBLs. Manual curation is time-consuming, and thus

development of efficient searching method to identify approximately 1.9-kb insertion

with EBLN would facilitate further exploration of EBLX/Ps. This is probably not limited

to bornaviruses: there may be many hidden EVEs that are derived from small and/or

rapidly evolving viral genes. Readthrough transcription was also observed in other viruses,

and thus our searching strategy can be applied to detecting further EVEs.

Taken together, we identified EBLs that were undetectable using the conventional

BLAST searches. Our findings provide novel insights into ancient bornaviruses. Further,

our searching strategy will contribute to paleovirological research.

Materials and Methods

Sangar sequencing of EBLs and flanking regions in M. fulginosus

miEBLN/X/P-1, miEBLN/X/P-2, miEBLN/X/P-3, and miEBLX/P-4 and corresponding

flanking regions in *M. fuliginosus* were PCR-amplified from DNA sample extracted from

YubFKT1 (kidney epithelial cell line) [16] using Phusion Hot Start II High-Fidelity PCR

Master Mix (Thermo Fisher Scientific). The obtained amplicons were purified using

innuPREP PCRpure Lite Kit (Analytik Jena), and then sequenced by Sanger dideoxy

method at Eurofins Genomics. The primer sequences are available in Table S4.

BLAST searches for identification of EBLN/X/Ps and EBLX/Ps

To search for EBLX/P insertions, miEBLN/X/P-1 from *M. natalensis* was used as query

for BLASTn search [17] against the WGS database [18] of bats (order Chiroptera; taxid:

9397), with changes applied to default parameters: E-value threshold = 10^{-10}, word size =

7, no filter for low complexity regions.

TFASTY search against mammalian genomes to detect EBLX/Ps

To detect EBLX/Ps in mammalian genomes, TFASTY searches were performed using the

FASTA36 package [15]. Mammalian genomes were downloaded using NCBI Datasets

[18], and used for the database for TFASTY search.

Identification of orthologous loci

To search for the orthologous loci harboring EBL insertions or EBL "empty" loci, the

sequence of the insertion and its flanking regions were used as query for BLASTn search

against the WGS contig database of Chiroptera (taxid: 9397) or Rodentia (taxid: 9989),

with changes applied to default parameters: E-value $= 10^{-20}$, word size $= 11$.

The BLAST hit sequences were retrieved, which were used for further analysis.

Genomic sequences from different species which corresponds to the locus containing the

EBL insertion were aligned by MAFFT alignment tool. The phylogenetic tree of the

species were obtained from TimeTree [19]. See Table S5 for the nucleotide accessions

and regions of EBL insertions and the loci containing the insertions.

Annotation of EBLs

Following alignment, TSD sequences flanking the EBL insertions were determined and

promptly annotated (if present). The EBLs were then aligned by MAFFT alignment tool,

wherein transcription signal sequences S2, T1, and T2 were determined and annotated,

by also aligning to extant orthobornaviral X/P transcript sequences.

Complete X- and P-like ORFs from miEBLX/P-4 were also annotated as miEBLX

and miEBLP, respetively. The amino acid sequences were then extracted and aligned with

the amino acid sequences of extant orthobornaviral X or P proteins. Functional domains

previously characterized in BoDV-1 X and P were also determined within miEBLX and

miEBLP, as well as other extant orthobornaviral X and P proteins. See Table S6 for the

nucleotide accession of extant orthobornaviruses.

All sequence alignments and annotations, as well as phylogenetic tree reconstruction,

were performed using Geneious version 11.1.5.

Pairwise similarity analysis

Amino acid sequences of the viral proteins of extant orthobornaviruses (Table S6) were

analyzed by MAFFT alignment tool [20] using Sequence Demarcation Tool software

version 1.2 [21]. Minimum value for the percent identity scale was set to 0.1. The

resulting matrices were set to 3-color mode and exported.

In silico characterization of putative EBLX and EBLP proteins

To predict the protein secondary structure, amino acid sequences of miEBLX and

miEBLP from *M. fuliginosus*, as well as extant orthobornaviral X and P proteins, were

submitted to JPred version 4 [22]. The simple results were then exported for further

analysis.

To determine the protein hydrophathy, the same amino acid sequences were

submitted to ProtScale in ExPASy Server [23], using the amino acid scale developed by

Kyte & Doolittle [24]. The window sizes were set to 9 and 15 for miEBLX and miEBLP,

respectively. The hydropathy scores were then exported for further analysis.

Acknowledgements

The animal silhouette images were downloaded from https://commons.wikimedia.org/

and https://publicdomainvectors.org/ under public domain access.

This study was supported by JSPS KAKENHI grant numbers JP21H01199 (MH)

and 20H05682 (KT) and MEXT KAKENHI grant number JP19H04833 (MH).

285   Conflict of interest

286   None declared.

288   References

1.   Aiewsakun, P. and A. Katzourakis, *Endogenous viruses: Connecting recent and*
ancient viral evolution. Virology, 2015. **479-480**: p. 26-37.

2.   Jern, P. and J.M. Coffin, *Effects of retroviruses on host genome function.* Annu
Rev Genet, 2008. **42**: p. 709-32.

3.   Horie, M., et al., *Endogenous non-retroviral RNA virus elements in mammalian*
genomes. Nature, 2010. **463**(7277): p. 84-7.

4.   Rubbenstroth, D., et al., *ICTV Virus Taxonomy Profile: Bornaviridae.* J Gen Virol,
2021. **102**(7).

5.   Kawasaki, J., et al., *100-My history of bornavirus infections hidden in vertebrate*
genomes. Proc Natl Acad Sci U S A, 2021. **118**(20).

6.   Belyi, V.A., A.J. Levine, and A.M. Skalka, *Unexpected inheritance: multiple*
integrations of ancient bornavirus and ebolavirus/marburgvirus sequences in
vertebrate genomes. PLoS Pathog, 2010. **6**(7): p. e1001030.

7.   Horie, M. and K. Tomonaga, *Paleovirology of bornaviruses: What can be learned*
from molecular fossils of bornaviruses. Virus Res, 2019. **262**: p. 2-9.

8.   Kojima, S., et al., *Splicing-Dependent Subcellular Targeting of Borna Disease*
Virus Nucleoprotein Isoforms. J Virol, 2019. **93**(5).

9.   Horie, M., et al., *Comprehensive analysis of endogenous bornavirus-like elements*
in eukaryote genomes. Philos Trans R Soc Lond B Biol Sci, 2013. **368**(1626): p.
20120499.

10.   Mukai, Y., et al., *An endogenous bornavirus-like nucleoprotein in miniopterid bats*
retains the RNA-binding properties of the original viral protein. FEBS Lett, 2022.

11.   Jurka, J., et al., *CENSOR--a program for identification and elimination of*
repetitive elements from DNA sequences. Comput Chem, 1996. **20**(1): p. 119-21.

12.   Ferre, C.A., et al., *Manipulation of the N-terminal sequence of the Borna disease*
virus X protein improves its mitochondrial targeting and neuroprotective potential.

FASEB J, 2016. **30**(4): p. 1523-33.

13.   Poenisch, M., et al., *Protein X of Borna disease virus inhibits apoptosis and*
promotes viral persistence in the central nervous systems of newborn-infected rats.
J Virol, 2009. **83**(9): p. 4297-307.

14.   Szelechowski, M., et al., *A viral peptide that targets mitochondria protects against*
neuronal degeneration in models of Parkinson's disease. Nat Commun, 2014. **5**:
p. 5181.

15.   Pearson, W.R., *Finding Protein and Nucleotide Similarities with FASTA.* Curr
Protoc Bioinformatics, 2016. **53**: p. 3 9 1-3 9 25.

16.   Maeda, K., et al., *Isolation of novel adenovirus from fruit bat (Pteropus*
dasymallus yayeyamae). Emerg Infect Dis, 2008. **14**(2): p. 347-9.

17.   Camacho, C., et al., *BLAST+: architecture and applications.* BMC Bioinformatics,
2009. **10**: p. 421.

18.   Coordinators, N.R., *Database resources of the National Center for Biotechnology*
Information. Nucleic Acids Res, 2018. **46**(D1): p. D8-D13.

19.   Kumar, S., et al., *TimeTree: A Resource for Timelines, Timetrees, and Divergence*
Times. Mol Biol Evol, 2017. **34**(7): p. 1812-1819.

20.   Katoh, K. and D.M. Standley, *MAFFT multiple sequence alignment software*
version 7: improvements in performance and usability. Mol Biol Evol, 2013.
30(4): p. 772-80.

21.   Muhire, B.M., A. Varsani, and D.P. Martin, *SDT: a virus classification tool based*
on pairwise sequence alignment and identity calculation. PLoS One, 2014. **9**(9):
p. e108277.

22.   Drozdetskiy, A., et al., *JPred4: a protein secondary structure prediction server.*
Nucleic Acids Res, 2015. **43**(W1): p. W389-94.

23.   Duvaud, S., et al., *Expasy, the Swiss Bioinformatics Resource Portal, as designed*
by its users. Nucleic Acids Research, 2021. **49**(W1): p. W216-W227.

24.   Kyte, J. and R.F. Doolittle, *A simple method for displaying the hydropathic*
character of a protein. J Mol Biol, 1982. **157**(1): p. 105-32.

Figure legends

Figure 1. Amino acid identities within extant orthobornaviral proteins. Pairwise

sequence identities between amino acid sequences of extant orthobornaviral proteins are

shown. The accession numbers used in this analysis was shown in Table S6.

Figure 2. Characterization of EBLs derived from N/X/P transcripts. (A) A schematic

figure of orthobornviral genomic RNA, its N/X/P mRNA, transcription signals and

miEBLN-1 locus. S1/S2 and T1/T2 show transcription initiation and termination signals,

respectively. **(B)** A nucleotide alignment of orthobornaviral N/X/P transcripts and

miEBLN/X/P loci. Red letters represent the start and stop codons of viral N gene. Blue

letters indicate the start codon of X gene and stop codon of P gene. TSD, target site

duplication. **(C and D)** Gene orthology analyses for miEBLN/X/P-1 and 3. Nucleotide

alignment of each miEBLN/X/P and their flanking loci is shown. The corresponding

species and their phylogenetic relationships are indicated on the left. The minimum age

of each EBLN/X/P is also shown.

Figure 3. Characterization of EBLs derived from X/P transcripts. (A) A schematic

figure of orthobornaviral genomic RNA, its X/P mRNA, transcription signals, and

miEBLX/P loci. **(B)** A nucleotide alignment of orthobornaviral X/P transcripts and

miEBLX/P loci. Light blue and blue letters represent the start codon of X gene and stop

codon of P gene, respectively. TSD, target site duplication. **(C - F)** Gene orthology

analyses for EBLX/Ps. Nucleotide alignment of each EBLX/P and its flanking loci is

shown. The corresponding species and their phylogenetic relationships are indicated on

the left. The minimum age of each EBLN/X/P is also shown.

Figure 4. *In silico* **characterization of putative miEBLX and miEBLP proteins. (A**
and **B)** Secondary structure prediction of putative EBLX and EBLP proteins. H and E
represent helical and beta extended secondary structure, respectively. **(C)** Multiple amino
acid sequence alignment of orthobornaviral X and putative EBLX proteins. **(D)** Multiple
amino acid sequence alignment of orthobornaviral P and putative EBLP proteins.

Fig. 1

Fig. 2

A

Orthobornaviral gRNA

N/X/P mRNA   AAA(n)   1.9 kb

miEBLN/X/P-1   miEBLN-1   1.9 kb

Transcription signal

B

	TSD	S1		S2	T1		T2 + Poly-A	TSD

BoDV-1 N/X/P   GAATACA------CGCAATG   TAGAAAAACAATGAA-CAAACCAATAAAAAAACCAAATGCGGCAAACCCCCCGCGACCTGTGATG   TAAAAAAAA(n)

PaBV-2 N/X/P   GAACACAA-----CAACATG   -------TAATGAA-CATACGAATAAAAAACCATAT--------------------CGATG   TAAAAAAAA(n)

LGSV-1 N/X/P   GAACAAAA----CAAAAATG   -------TGAA-CAAACCTATATAAAAACCATAT--------------------AGATG   TAAAAAAAA(n)

CWBV N/X/P   GAACCAAATCCACAAAAATG   -------TAATGAACCATTCATATAGAAAACCGAATTC-------------------CAATG   TAAAAAAAA(n)

miEBLN/X/P-1 in *M. natalensis*   TTGTGCTAGGAACCAA------CACAATG   TAATATCGAA-CCAACATTTTCAAAAACATA---------------------GATG   TAAAATAAATATATACACCA----TTCTTGTA

miEBLN/X/P-2 in *M. natalensis*   TGTTCAGGAGCCAA-------CACAATG   -----TTATATCGAA-ACAACATTTCAAAAACATA--------------------GATA   TAAATATTGCATATATTTCCACAAATGTTCA

miEBLN/X/P-3 in *M. natalensis*   GAGGAACGGGAACCAA------CACAATG   -----TAACATCGAA-CCAACATTTACAAAAACATA--------------------GACA   TAAAAAATAATAGTAATAAAAG--GAGGGATG

C

Insert (1.9 kb)

(kb) 0   1   2   3   4   4.6

% Identitiy

Rhinolophus ferrumequinum

Phyllostomus hastatus

Tadarida brasiliensis

Miniopterus fuliginosus   miEBLN-1

Miniopterus natalensis   miEBLN-1

Miniopterus schreibersii   miEBLN-1

miEBLN/X/P-1
17.2 mya

D

Insert (1.9 kb)

(kb) 0   1   2   3   4   5   5.3

% Identitiy

Molossus molossus

Murina aurata

Myotis lucifugus

Miniopterus fuliginosus

Miniopterus natalensis

Miniopterus schreibersii

miEBLN/X/P-3
17.2 mya

Fig. 3

Fig. 4