The hidden diversity of ancient bornaviral sequences from X and P genes in

vertebrate genomes

Bea Clarise B. Garcia [1], Yahiro Mukai [2,3], Keizo Tomonaga [2,3], & Masayuki Horie [1,4*]

[1] Laboratory of Veterinary Microbiology, Graduate School of Veterinary Science, Osaka

Metropolitan University, Izumisano 598-8531, Osaka, Japan

[2] Laboratory of RNA Viruses, Institute for Frontier Life and Medial Sciences, Kyoto

University, Kyoto, Japan

[3] Department of Mammalian Regulatory Network, Graduate School of Biostudies,

Kyoto University

[4] Osaka International Research Center for Infectious Diseases, Osaka Metropolitan

University, Izumisano 598-8531, Osaka, Japan

*Corresponding author

Masayuki Horie, DVM, PhD

Tel: +81-72-463-5694

Email: mhorie@omu.ac.jp

Keywords: endogenous viral element, bornavirus, orthobornavirus, paleovirology,

readthrough transcript

Abstract

Endogenous bornavirus-like elements (EBLs) are heritable sequences derived from

bornaviruses in vertebrate genomes that originate from transcripts of ancient

bornaviruses. EBLs have been detected using sequence similarity searches such as

tBLASTn, whose technical limitations may hinder the detection of EBLs derived from

small and/or rapidly evolving viral X and P genes. Indeed, no EBLs derived from the X

and P genes of orthobornaviruses have been detected to date in vertebrate genomes.

Here, our aim was to develop a novel strategy to detect such "hidden" EBLs. To that

aim, we focused on the 1.9 kb readthrough transcript of orthobornaviruses, which

encodes a well-conserved N gene and small and rapidly evolving X and P genes. We

show a series of evidence supporting the existence of EBLs derived from

orthobornaviral X and P genes (EBLX/Ps) in mammalian genomes. Furthermore, we

found that an EBLX/P is expressed as a fusion transcript with the cellular gene,

ZNF451, which potentially encodes the ZNF451EBLP-4 fusion protein in miniopterid

bat cells. This study contributes to a deeper understanding of ancient bornaviruses and

co-evolution between bornaviruses and their hosts. Furthermore, our data suggest that

endogenous viral elements detected thus far are just the tip of the iceberg, and further

studies are required to understand ancient viruses more accurately.

Introduction

Ancient viral sequences can be found within the genomes of organisms in the form of endogenous viral elements (EVEs) [1]. EVEs are derived from the integration of viral sequences into the host germline DNA and transfer of the integrated sequences to subsequent progenies [2]. As molecular fossils of ancient viruses, EVEs have expanded our knowledge of the diversity, host range, and long-term evolution of viruses. Furthermore, some EVEs are co-opted by hosts to function as novel gene regulatory elements, non-coding RNAs, and proteins [2].

Most EVEs are derived from retroviruses, as the viruses reverse-transcribe and integrate the viral genomes into the host genomes using virally encoded enzymes. However, non-retroviral EVEs have also been identified. Endogenous bornavirus-like elements (EBLs) are EVEs derived from bornaviruses (family *Bornaviridae*), a group of non-segmented negative-strand RNA viruses [3, 4]. Although bornaviruses are RNA viruses, evidence strongly suggests that bornaviral mRNA is reverse transcribed and integrated into the host genome via the activity of the long interspersed nuclear element 1 (LINE-1) retrotransposon [3, 5]. We previously revealed an abundance of EBLs in vertebrate genomes derived from the bornaviral genes N, P, M, G, and L, which are referred to as EBLN, EBLP, EBLM, EBLG, and EBLL, respectively [6]. These EBLs originated from ancient bornaviruses of all genera in the family *Bornaviridae*: *Orthobornavirus*, *Carbovirus*, and *Cultervirus*.

Despite the rigorous search, there has been a bias in the detection of EBLs. EBLNs comprise the majority of the detected EBLs, while no EBLs derived from the X and P genes have been reported except for a few carboviral EBLPs [3, 6-8]. We proposed two hypotheses to explain the lack of EBLX and EBLP. First, X/P mRNA, which encodes

the X and P proteins, may have a low propensity for reverse-transcription and/or

integration into the host genome. However, our previous studies of Borna disease virus

1 (BoDV-1; genus *Orthobornavirus*) showed that X/P mRNA is the most abundant

transcript among the viral mRNAs [9], which should therefore provide enough chances

for integration. Moreover, we have previously demonstrated that the X/P mRNA of

BoDV-1 can be reverse-transcribed and integrated into the genome of host cells [3, 7].

This led us to the second hypothesis, which states that EBLs derived from the X and P

genes are present but are difficult to detect due to technical limitations of the sequence

similarity search. EBLs have been routinely searched using tBLASTn; however, this can

be an unsuitable method for detecting small and rapidly evolving genes [8]. Indeed, the

X and P genes are small and the least conserved among bornaviral genes (Fig. 1). If the

second hypothesis is correct, some EBLs may remain undetected. Therefore, we could

be missing important insights into ancient bornaviruses and the co-evolution of

bornaviruses and their hosts.

In this study, we sought to develop a novel strategy to detect these hidden EBLs by

focusing on the nature of the readthrough transcription of orthobornaviruses that

produce a transcript encoding a highly conserved N gene and rapidly evolving X and P

genes. We show evidence supporting the existence of several EBLX/Ps in vertebrate

genomes. Furthermore, an EBLX/P in miniopterid bats appear to be transcribed as

chimeric transcripts with a host gene, suggesting a potential co-option of EBLs by the

host.

Results

Miniopterid bat EBLN is possibly derived from an N/X/P transcript

We previously identified miEBLN-1, an EBLN derived from an ancient orthobornavirus

that retains an intact N ORF in miniopterid bats [10]. The miEBLN-1 locus is present in

the genomes of miniopterid bats, specifically in those of *Miniopterus natalensis*, *M.*

schreibersii, and *M. fuliginosus*, but is empty (i.e., syntenic loci lacking miEBLN-1) in

the genomes of *Rhinolophus ferrumequinum* and *Phyllostomus hastatus*. We re-

examined the presence of miEBLN-1 using the current genome datasets and found the

miEBLN-1-empty locus in *Tadarida brasiliensis* (Fig. 2a and Fig. S1a). Based on the

nucleotide alignment, we deduced the insertion size of the miEBLN-1 locus to be 1.9

kb, consisting of a 1.2-kb miEBLN-1 region and a 0.7-kb unannotated region.

We carried out a series of analyses to understand the source of the 0.7-kb

unannotated region. First, we performed a BLASTn search against the NCBI nt

database using the nucleotide sequence of the 0.7-kb unannotated region as a query. The

hits consisted only of miEBLN-1 sequences (data not shown). Next, we assessed

whether the 0.7-kb unannotated region is derived from a short interspersed nuclear

element (SINE) due to the precedence of chimeric integration comprising nucleotide

sequences from an N transcript and SINE [3, 5]. We performed a BLASTn search

against the miniopterid bat genomes using the nucleotide sequence of the 0.7-kb region

as a query. We only detected five BLAST hits in the genomes of *M. natalensis* and *M.*

schreibersii, which we identified as EBLs in the later sections. We also analyzed the

sequence of the 0.7-kb region by CENSOR [11], but no hits were detected. These results

indicated that the 0.7-kb fragment is not derived from a SINE.

We noticed that the insertion, including miEBLN-1, is almost similar in size to the

N/X/P mRNA of extant orthobornaviruses (Fig. 2b), which led us to speculate that the

insertion may be derived from an N/X/P transcript. N/X/P mRNA is produced by

readthrough of the termination signal T1, resulting in a 1.9 kb mRNA containing the

transcription signal sequences S1, T1/S2, and T2 (Fig. 2b) [12]. Therefore, we

investigated the presence of these signal-like sequences in the 1.9-kb miEBLN-1

insertion. We found transcription signal-like sequences at reasonable positions with

respect to extant N/X/P transcripts (Fig. 2c). A poly-A stretch was also observed

immediately downstream of the T2 signal, indicating that the insertion originated from

an mRNA transcript (Fig. 2c and Fig. S2). Furthermore, a tandem duplication sequence,

TTGTGCTA/TTCTTGTA, was observed immediately upstream and downstream of S1

and T2-poly-A sequences, respectively. This is a potential target site duplication (TSD),

which is a hallmark of LINE-1-mediated mRNA integration [13]. These results

suggested that the 1.9-kb insertion, including miEBLN-1, is derived from an ancient

N/X/P transcript integrated by LINE-1. Therefore, we tentatively refer to this locus as

miEBLN/X/P-1.

Additional EBLs possibly derived from N/X/P transcripts in miniopterid bat genomes

As described above, orthobornaviruses express mRNAs encoding N (1.2 kb), X and P

(X/P mRNA, 0.7 kb), or N, X, and P (N/X/P mRNA, 1.9 kb). If miEBLN/X/P-1 is truly

derived from N/X/P mRNA, there may be additional EBLs derived from N/X/P or X/P

mRNA. To examine this possibility, we conducted a BLASTn search against bat

genomes using the nucleotide sequence of miEBLN/X/P-1 as a query (Fig. 3a). We

obtained six hits from the genomes of *M. natalensis* and *M. schreibersii*. While four hits

were aligned in the entire N/X/P region of miEBLN/X/P-1, interestingly, two other hits

were aligned only to the putative X/P region (Fig. 3a and Table S1).

We first assessed hits that aligned to the entire region of miEBLN/X/P-1. Four hits

showed 1.9 kb alignment lengths and high sequence similarities (76–84% nucleotide

identity) to miEBLN/X/P-1 (Table S1). The flanking regions of each pair of hit

sequences were well aligned between *M. natalensis* and *M. schreibersii*, indicating that

they are orthologous between these species (Table S7). We tentatively refer to these loci

as miEBLN/X/P-2 and miEBLN/X/P-3. We also confirmed the presence of orthologs of

these EBLN/X/Ps in the genome of *M. fuliginosus* by PCR and Sanger sequencing of

genomic DNA isolated from *M. fuliginosus* (Table S7).

Next, we examined the presence of miEBLN/X/P-2 or miEBLN/X/P-3- empty loci

by comparative genomics. We observed that the miEBLN/X/P-3 empty locus is present

in genomes of non-miniopterid bats such as *Tadarida brasiliensis* (Fig. S1B and Table

S7). We did not obtain an alignment of the flanking region of miEBLN/X/P-2 with the

other bat genomes and therefore could not identify its empty locus.

We then analyzed the nucleotide sequences of miEBLN/X/P-2 and miEBLN/X/P-

3. Similar to miEBLN/X/P-1, both EBLN/X/Ps contained transcription signal-like

sequences, and poly-A stretches were observed downstream of T2-like sequences (Fig.

S2). Furthermore, tandem duplication sequences, TGTTCA/[T/C]GTTCA and

GAGGAACG/GAGGGATG, were also observed immediately upstream of S1 and

downstream of T2/poly-A in miEBLN/X/P-2 and miEBLN/X/P-3, respectively. These

results suggest that miEBLN/X/P-2 and miEBLN/X/P-3 are also derived from ancient

N/X/P transcripts and were integrated into the genome of a miniopterid bat ancestor

through LINE-1 activity.

EBLs derived from X/P-like transcripts exist in miniopterid bat genomes

Next, we assessed the hits aligned to the 0.7-kb putative X/P region of miEBLN/X/P-1.

Two hits from *M. natalensis* and *M. schreibersii* showed 0.8 kb alignment lengths and

75% nucleotide identity to the 0.7-kb X/P region (Table S1). The flanking regions of the

hit sequences were well aligned, indicating that the loci are orthologous between these

species (Table S7). We also found an orthologous locus in the genome of *M. fuliginosus*

using PCR and Sanger sequencing (Table S7). Furthermore, we detected empty loci in

genomes of non-miniopterid bats, genomes such as *Myotis lucifugus* (Fig. 3b and Fig.

S3a).

The orthobornaviral X/P mRNA contains the transcription signal sequences S2/T1

in the 5'-UTR and T2 in the 3'-UTR. We observed that S2/T1 and T2-like sequences are

present and located at positions similar to those in extant X/P mRNA. Additionally,

poly-A stretches were observed downstream of the T2 sequences. Furthermore, the S2

and poly-A stretches were flanked by TSDs, CTTAAGGC/CTTAAGGC (Fig. 3a, b and

Fig. S5). Taken together, these results indicated that the locus may have been derived

from an ancient X/P transcript and integrated into the genome of a miniopterid bat

ancestor through LINE-1 activity. We tentatively refer to it as the miEBLX/P-4.

Preservation of intact ORFs in miEBLX/P-4 and characterization of the putatively

encoded proteins

Most EBLs do not retain viral ORFs because of the accumulation of mutations after

integration. Remarkably, however, miEBLX/P-4 retains intact X- and P-like ORFs in

different reading frames, which is consistent with the extant orthobornaviral X/P mRNA

(Fig. 3c and Fig. S4a). Therefore, we named these ORFs as miEBLX-4 and miEBLP-4.

Notably, miEBLP-4 in *M. natalensis* is disrupted by a premature stop codon.

Although there is very low sequence similarity between the amino acid sequences

of these ORFs and extant orthobornaviral proteins, further analyses showed that

miEBLX-4 and miEBLP-4 are intrinsically similar to extant orthobornaviral X and P

proteins.

First, we examined the miEBLX-4 ORF. The ORF consists of 109 codons, which is

longer than the that of extant orthobornaviral X (86–90 codons). We then predicted the

secondary structure and analyzed the hydrophobicity of the putative miEBLX-4 protein

with respect to extant X proteins. Both extant X and miEBLX-4 were predicted to

contain an alpha helix in the N-terminal region (Fig. 4a and Fig. S6a). The hydropathy

plot showed similar hydrophobicity patterns for these proteins (Fig. S7a).

We also determined whether the functional domains are conserved between extant

X and miEBLX-4. Amino acid sequence alignment showed that the N-terminal region is

relatively well conserved and houses functional domains previously characterized in

BoDV-1 X (Fig. 4c). This region includes the leucine-rich domain functioning both as a

CRM1-dependent nuclear export signal [14] and a mitochondria-targeting signal [15],

and is overlapped by the importin-alpha binding domain [16]. The alignment also

showed that leucine and arginine residues that are critical in the importin-alpha-binding

domain (Fig. 4d; orange and blue, respectively) are conserved among extant X and

miEBLX-4.

Next, we examined the miEBLP-4 ORF. The ORF consists of 233 codons; it is

therefore longer than those of extant orthobornaviral P (201–208 codons). Extant P and

putative miEBLP-4 proteins also showed similar predicted secondary structures (Fig. 4b

and Fig. S6b) and hydrophobicity patterns (Fig. S7b). The predicted secondary

structures of miEBLP-4p were similar to those of the modern counterparts (Fig. 4b and

Fig. S6b). Alignment of the amino acid sequences revealed that some of the functional

domains previously characterized in BoDV-1 P may be shared between extant P and

miEBLP-4 (Fig. 4d). Proline residues (Fig. 4d, blue) in the N-terminal and C-terminal

nuclear localization signals were prominent in both proteins [17].

We also explored the interaction between miEBLX-4 and miEBLP-4, which is a

notable characteristic of the orthobornaviral X and P proteins [18]. We expressed Myc-

tagged miEBLX-4 (miEBLX-4-Myc) and FLAG-tagged miEBLP-4 (miEBLP-4-FLAG)

in 293T cells and then immunoprecipitated them using an anti-FLAG antibody.

miEBLX-4-Myc was co-immunoprecipitated with miEBLP-4-FLAG, indicating the

interaction between these proteins (Fig. 4e).

These results showed that miEBLX-4 and miEBLP-4 share similar intrinsic

properties with the extant X and P proteins. Furthermore, together with the above data,

they strongly suggest that miEBLN/X/P-1-3 and miEBLX/4 are indeed derived from

ancient viral N/X/P and X/P mRNAs, respectively.

Diverse EBLX/Ps are found in vertebrate genomes

To further identify the presence of other EBLX/Ps, we conducted TFASTX and

TFASTY searches against vertebrate genomes using the amino acid sequences of

miEBLX-4 and miEBLP-4 as queries. TFASTX and TFASTY are sequence similarity

search programs that allow frameshifts for alignment [19]. As a result, 65 unique hits

(Table S2–S5) were detected in the genomes of mammals (class Mammalia), reptiles

(class Reptilia), amphibians (class Amphibia), and ray-finned fish (class

Actinopterygii). The majority of these hits were detected in the bat genomes, including

miEBLN/X/P-1, miEBLN/X/P-2, miEBLN/X/P-3, and miEBLX/P-4.

Comparative genomic analyses strongly suggest that at least some of the

TFASTX/Y hits are indeed derived from insertions of ancient viral X/P transcripts, as

follows. TFASTX/Y hits in the bat families Mormoopidae and Phyllostomidae

(mpEBLX/P) are orthologous, and their empty loci are present in other lineages of bats,

such as *Noctilio leporinus* (Fig. 5a and Fig. S3b). For a TFASTX/Y hit in the lone bat

species *Craseonycteris thonglongyai* (ctEBLX/P), we could not find its ortholog, but its

empty sites exist in other bats (Fig. 5b and Fig. S3c). We further identified orthologous

TFASTX/Y hits in the rodent superfamily Octodontoidea and their empty loci in other

lineages of rodents (ocEBLX/P; Fig. 5c and Fig. S3d). We also identified transcription

signal-like sequences and TSDs within and flanking the above TFASTX/Y hits,

respectively (Fig. 5d and Fig. S4). These results strongly suggested that at least some of

the sequences detected by TFASTX and TFASTY are bona fide EBLX/Ps.

Interestingly, the insertion sizes of bat mpEBLX/P and ctEBLX/P were both 0.8

kb, which is similar to that of miEBLX/P-4. In contrast, that of rodent ctEBLX/P is 0.7

kb, which is closer to the size of extant orthobornaviral X/P transcripts (see discussion).

Possible co-option of EBLX/P in miniopterid bats

Some EVEs, including EBLs, retain intact ORFs that encode proteins that play roles in

host-related functions [20]. As described above, miEBLX/P-4 contains intact ORFs

corresponding to orthobornaviral X and P genes. Therefore, we analyzed miEBLX/P-4

in detail. We first examined the location of miEBLX/P-4 in the *M. natalensis* genome.

miEBLX/P-4 is located within the intron region of *ZNF451* gene (Fig. 6).

Next, we analyzed the expression of miEBLX/P-4 by mRNA sequencing of *M.*

schreibersii-derived SubK12-08 cells. The read coverage of the miEBLX/P-4 locus was

comparable to that of ZNF451 (Fig. 6A). Interestingly, the mapping pattern revealed the

existence of an alternative splicing site for *ZNF451* that could produce a chimeric

transcript comprising ZNF451 exons 1–4 and the miEBLX/P-4 locus (Fig. 6A).

The putative chimeric transcript contains two ORFs consisting of more than 100

codons in different reading frames: a chimeric ORF of ZNF451 in-frame with miEBLP-

4 or that of the miEBLX-4. The chimeric ORF potentially encodes a fusion protein

comprising the N-terminal region of ZNF451, which contains a SUMO-interacting

motif (SIM) domain, a short intronic region of *ZNF451*, and miEBLP-4 (Fig. 6).

These results suggest that miEBLX/P-4 has been co-opted by miniopterid bats as a

chimeric gene with *ZNF451*.

Discussion

We previously demonstrated that the X/P mRNA of a modern orthobornavirus can be

reverse-transcribed and integrated into the host genome [7]; however, no EBLX and

only a few EBLPs have been reported [6]. We hypothesized that the technical

limitations of the sequence similarity search hinder the detection of EBLs derived from

the X and P genes. EBLs have been routinely searched using tBLASTn, but this can be

an unsuitable method for detecting small and rapidly evolving genes such as the X and P

genes [8]. In this study, our aim was to develop a novel strategy to detect these hidden

EBLs. To that aim, we focused on the 1.9-kb readthrough transcript of

orthobornaviruses, consisting of a well-conserved N gene and small and rapidly

evolving X and P genes [12], to detect hidden EBLX/Ps. We show a series of evidence

implying that EBLN/X/Ps and EBLX/Ps are present in vertebrate genomes but are not

detectable by BLAST searches. Interestingly, miEBLX/P-4 has maintained intact ORFs

for more than 17 million years and is expressed as a fusion transcript with the cellular

gene, *ZNF451*, which potentially encodes the ZNF451/miEBLP-4 fusion protein.

Therefore, this study revealed the presence of hidden EBLs, which could contribute to a

deeper understanding of ancient bornaviruses and the co-evolution between

bornaviruses and their hosts.

Here, we uncovered the potential co-option of miEBLX/P-4 by miniopterid bats, as

indicated by the expression of the ZNF451/miEBLP-4 chimeric transcript, which

potentially encodes a fusion protein consisting of the N-terminal region of ZNF451 and

miEBLP-4. To the best of our knowledge, this is the first report of a non-retroviral EVE

being expressed as a chimeric transcript with a known host gene. Although we could not

confirm the expression of the ZNF451/miEBLP-4 fusion protein due to the lack of

antibodies, our data suggest that miEBLX/P-4 co-opted by miniopterid bats. Further

studies are needed to understand the biological significance of EBLs.

Our data strongly suggest that the genome organization and expression patterns of

the N, X, and P genes have not substantially changed over the past million years. Here,

we show that T1-readthrough transcription occurred in ancient orthobornaviruses

present at least 17.3 MYA (Fig. 2). Furthermore, the ORF positions of the X and P genes

of ancient orthobornaviruses were highly similar to those of modern orthobornaviruses

(Fig. 3). In addition, the sequences and order of transcription signals are highly

conserved among ancient and modern orthobornaviruses (Figs. 2 and 3). In BoDV-1,

readthrough transcription has been suggested to be involved in the regulation of viral

gene expression [12]. This hypothetical gene regulation mechanism has been

maintained for more than 17.3 million years. Interestingly, our previous study showed

that the order of S2-T1 is also conserved in some EBLNs shared among Old World

monkeys, which were estimated to be integrated into the host genomes more than 43

MYA [3]. Thus, this gene regulatory mechanism may have been employed by

orthobornaviruses earlier than 17.3 MYA.

The fundamental functions of viral X and P proteins may also be evolutionarily

conserved between ancient and modern orthobornaviruses. We observed similarities in

the intrinsic properties of extant orthobornaviral X and P proteins with miEBLX-4p and

miEBLP-4p despite their very low amino acid identities (Fig. 4). Furthermore, the

interaction between X and P may have been conserved in ancient orthobornaviruses

(Fig. 4). Although the evolutionary rates of X and P genes are high, the fundamental

features and functions of X and P proteins may have been maintained for at least 17.3

million years. Some orthobornaviruses are known to be fetal zoonotic pathogens [21].

Thus, although the biological functions of the X protein and the interaction between X

and P proteins are not fully understood, these proteins can be good targets for the

development of antivirals.

For the EVE search, TFASTX/Y, a conventional sequence similarity search for the

detection of EVEs, may be more suitable than tBLASTn. After integration into the host

genome, many EVEs have acquired random mutations, which caused frameshifts in the

ORFs. Such frameshifted EVEs are a kind of "fragmented" EVEs and are thus less

detectable by tBLASTn. In contrast, TFASTX/Y takes frameshifts into account and can

thus detect such fragmented EVEs more sensitively. We are currently conducting a

benchmark analysis of tBLASTn and TFASTX/Y in detail, showing that some of the

EBLX/Ps detected by TFASTX/Y cannot be detected by tBLASTn or other BLAST

programs (data not shown). Therefore, TFASTX/Y can detect fragmented EVEs more

sensitively than tBLASTn, which may contribute to identifying other hidden EVEs.

The EBLX/Ps identified in this study would represent only the tip of the iceberg on

the hidden EVEs that could exist in the genome. In this study, the majority of EBLX/Ps

were detected in bat and rodent genomes. Note that we failed to detect any EBLX/Ps in

primate genomes, which contain numerous EBLs. Considering the number of EBLs in

the genomes of primates (i.e., 24 orthobornaviral EBLs were detected in the human

genome [6]), it is plausible that hidden EBLX/Ps exist in primates that are undetectable

because of the inherent low amino acid similarities in X and P.

We may be able to detect such hidden EBLX/Ps by identifying other EBLN/X/P

integrations. However, searching for other EBLN/X/Ps by manual curation is time-

consuming. Thus, the development of an efficient searching method to identify

approximately 1.9-kb insertions with EBLN would facilitate further exploration of

EBLX/Ps. This is probably not limited to bornaviruses; there may be many hidden

EVEs that are derived from small and/or rapidly evolving viral genes. Readthrough

transcription has also been observed for other mononegaviruses. Our search strategy can

be applied to the detection of such EVEs.

Taken together, we identified EBLs that were undetectable by conventional

BLAST searches. Our findings provide novel insights into ancient bornaviruses.

Furthermore, our search strategy will contribute to paleovirological research.

Materials and Methods

Sanger sequencing of EBLs and flanking loci in M. fuliginosus

Genomic DNA was obtained from YubFKT1 (kidney epithelial cell line) [22] in a

previous study [10]. miEBLN/X/P-1, miEBLN/X/P-2, miEBLN/X/P-3, and miEBLX/P-

4 and corresponding flanking locus in *M. fuliginosus* were PCR-amplified from the

DNA extracted from YubFKT1 using Phusion Hot Start II High-Fidelity PCR Master

Mix (Thermo Fisher Scientific). The amplicons were then purified using innuPREP

PCRpure Lite Kit (Analytik Jena) and sequenced by Sanger dideoxy method at Eurofins

Genomics. The primer sequences are available in Table S6. The obtained sequences

were deposited to DDBJ (accession numbers LC708263-LC708266).

Cell lines

SuBK 12-08 cells [23] and human embryonic kidney 293T (HEK293T) cells were

cultured in low-glucose Dulbecco's Modified Eagle Medium (Nacalai Tesque)

supplemented with 10% fetal bovine serum (Sigma-Aldrich) and 100 μg/mL penicillin-

streptomycin (Nacalai Tesque). The cells were then maintained at 37°C in a humidified

chamber containing 5% CO_2.

Construction of miEBLX-4 and miEBLP-4 expressing plasmids

Expression vectors with C-terminal FLAG or Myc tag were first constructed. pcDNA3

was digested with EcoRI and XhoI, which was assembled with oligo DNA containing

Myc or FLAG tag sequence using NEBuilder HiFi DNA Assembly Cloning Kit (New

England Biolabs). The resultant plasmids were named pcDNA3-C-Myc or pcDNA3-C-

FLAG. miEBLX-4 and miEBLP-4 sequences were then PCR-amplified from the

YubFKT1 DNA sample, and were subsequently assembled with pcDNA3-C-Myc or

pcDNA3-C-FLAG linearized with KpnI and EcoRI using NEBuilder. The oligo DNA

sequences are available in Table S6.

Co-immunoprecipitation of miEBLX-4 with miEBLP-4

The constructed plasmids (5 μg) were transfected into seeded 293T cells on 10-cm dish

using 20 μL of Avalanche-Everyday Transfection Reagent (EZ Biosystems), with Opti-

MEM (Thermo Fisher Scientific) as the transfecting medium. One day post-

transfection, the cells were washed twice with ice-cold PBS and lysed using 500 μL IP

buffer (150 mM NaCl, 20 mM Tris-HCl pH 7.4, 1 mM EDTA, 1% Triton X-100, 1x

protease inhibitor cocktail) for 30 min in 4°C with rotation. In parallel, 500 μg

SureBeads Protein G magnetic beads (Bio-Rad) were incubated with 3 μg mouse

monoclonal anti-DDDDK antibody FLA-1 (M185-3L; MBL) in IP buffer for 1 hr in

4°C with rotation. The lysates were then centrifuged at 12000 x g for 10 min at 4°C,

after which the FLAT-tagged proteins were immunoprecipitated from the lysate

supernatant by mixing with the antibody-bead complex in IP buffer for 2 hr in 4°C with

rotation. After incubation, the beads were washed thrice with ice-cold IP buffer and then

transferred into new microtubes. The proteins were finally eluted from the beads by

boiling in SDS sample buffer (組成) for 5 min in 95°C.

Western blotting

Protein samples were loaded in 12% polyacrylamide gel and run under electrophoresis

in Tris-glycine-SDS buffer. The proteins were then semi-dry transferred to PVDF

membrane at 10 V for 1 hr. Following blot transfer, the membrane was blocked with 5%

skim milk in 0.1% Tween 20 in PBS (PBST) for 1 hr at room temperature (RT). The

membrane was incubated with mouse monoclonal anti-Myc My3 (M192-3; MBL) or

anti-DDDDK antibody diluted in 5% skim milk for 1 hr at RT, washed, and then

incubated with horseradish peroxidase-conjugated donkey polyclonal anti-mouse IgG

(715-035-150; Jackson ImmunoResearch) antibody diluted in 5% skim milk for 45 min

at RT. The membrane was washed again, after which chemiluminescence reaction was

conducted using Chemi-Lumi One (Nacalai Tesque) and was subsequently viewed using

Amersham Imager 680 (GE Healthcare, IL, USA).

BLAST search against bat genomes to detect EBLN/X/Ps and EBLX/Ps

BLASTn [24] search was conducted using the nucleotide sequence of miEBLN/X/P-1

in *M. natalensis* as query, against the WGS database [25] of bat (Chiroptera, taxid:

9397) on the BLAST web server (https://blast.ncbi.nlm.nih.gov/Blast.cgi). Changes

applied to default parameters were as follows: E-value threshold = 10^{-10}, word size = 7,

no filter for low complexity regions.

TFASTX/Y search against vertebrate genomes to detect EBLX/Ps

Vertebrate genomes were downloaded using NCBI Datasets and were used as the

database for TFASTX/Y search. TFASTX/Y searches, included in FASTA36 package

[26], were performed against each vertebrate genome using the putative amino acid

sequences miEBLX-4 and miEBLP-4 with the ktup value of 1.

Identification of orthologous and empty loci

BLASTn search was conducted using the nucleotide sequences of EBL and its flanking

regions as query, against the WGS database of bats (Chiroptera, taxid: 9397) or rodents

(Rodentia, taxid: 9989) on the BLAST web server. Changes applied to default

parameters were as follows: E-value = 10^{-20}, word size = 11. The BLAST hits which

comprised of genomic sequences from different species were retrieved, and then aligned

by MAFFT using the E-INS-i algorithm [27] in Geneious version 11.1.5. The nucleotide

accessions and loci region of EBLs used for search query are available in Table S7.

Annotation of EBLs and their TSDs

TSD sequences were determined based on the above loci alignment. To identify

transcription signal-like sequences, the nucleotide sequences of EBLs and extant

orthobornaviral N/X/P or X/P transcripts were aligned by MAFFT using the E-INS-I

algorithm. The transcription signal-like sequences and poly-A stretches were then

determined according to the resulting alignment.

To identify conserved functional domains of the miEBLX-4p and miEBLP-4p, the

amino acid sequences of miEBLX-4p and miEBLP-4p in *M. fuliginosus* were aligned

with extant orthobornaviral X and P proteins, respectively, by MAFFT using the E-INS-

i algorithm. The conserved domains were manually analyzed based on the previously

characterized functional domains in BoDV-1 X and P and the alignments.

The nucleotide accession of extant orthobornaviruses used in the analysis are

available in Table S8. Multiple sequence alignments and annotations were conducted

using Geneious version 11.1.5.

mRNA-seq analysis

Total RNA sample from Subk12-08 cells was obtained by NucleoSpin RNA plus

(MACHEREY-NAGEL) according to the manual. Sequencing library was constructed

using TruSeq Stranded mRNA Library Prep Kit (Illumina). The libraries were

sequenced by NextSeq 500 using NextSeq 500/550 Mid Output Kit v2.5 (300 Cycles).

The obtained reads were preprocessed by fastp 0.23.2 [28] with the default setting,

which were mapped to the reference genome of *M. natalensis* (GCF_001595765.1) by

HISAT2 version 2.2.1 [29] with the default setting ref. The mapped reads and splicing

junctions were visualized using Integrated Genomics Viewer 2.13.0 [30].

Pairwise similarity analysis

Amino acid sequences of the viral proteins of extant orthobornaviruses (Table S8) were

analyzed by MAFFT alignment tool using Sequence Demarcation Tool software version

1.2 [31]. Minimum value for the percent identity scale was set to 0.1. The resulting

matrices were set to 3-color mode and exported.

In silico characterization of putative EBLX and EBLP proteins

To predict the protein secondary structure, amino acid sequences of miEBLX-4 and

miEBLP-4 in *M. fuliginosus*, and extant orthobornaviral X and P proteins, were

submitted to JPred version 4 [32]. The simple results were then exported.

To determine the protein hydrophathy, the same amino acid sequences were

submitted to ProtScale [33] in ExPASy Server, using the amino acid scale developed by

Kyte & Doolittle. The window sizes were set to 9 and 15 for X and P proteins,

respectively. The hydropathy scores were then exported.

Acknowledgements

This study was supported by JSPS KAKENHI grant numbers JP21H01199 (MH) and

20H05682 (KT) and MEXT KAKENHI grant number JP19H04833 (MH).

The animal silhouette images were downloaded under public domain access from

https://commons.wikimedia.org/ and https://publicdomainvectors.org/.

Conflict of interest

None declared.

References

1. Aiewsakun P, Katzourakis A: **Endogenous viruses: Connecting recent and ancient viral evolution**. *Virology* 2015, **479-480**:26-37.
2. Jern P, Coffin JM: **Effects of retroviruses on host genome function**. *Annu Rev Genet* 2008, **42**:709-732.
3. Horie M, Honda T, Suzuki Y, Kobayashi Y, Daito T, Oshida T, Ikuta K, Jern P, Gojobori T, Coffin JM *et al*: **Endogenous non-retroviral RNA virus elements in mammalian genomes**. *Nature* 2010, **463**(7277):84-87.
4. Rubbenstroth D, Briese T, Durrwald R, Horie M, Hyndman TH, Kuhn JH, Nowotny N, Payne S, Stenglein MD, Tomonaga K *et al*: **ICTV Virus Taxonomy Profile: Bornaviridae**. *The Journal of general virology* 2021, **102**(7).
5. Belyi VA, Levine AJ, Skalka AM: **Unexpected inheritance: multiple integrations of ancient bornavirus and ebolavirus/marburgvirus sequences in vertebrate genomes**. *PLoS Pathog* 2010, **6**(7):e1001030.
6. Kawasaki J, Kojima S, Mukai Y, Tomonaga K, Horie M: **100-My history of bornavirus infections hidden in vertebrate genomes**. *Proc Natl Acad Sci U S A* 2021, **118**(20).
7. Horie M, Kobayashi Y, Suzuki Y, Tomonaga K: **Comprehensive analysis of endogenous bornavirus-like elements in eukaryote genomes**. *Philosophical transactions of the Royal Society of London Series B, Biological sciences* 2013, **368**(1626):20120499.
8. Horie M, Tomonaga K: **Paleovirology of bornaviruses: What can be learned from molecular fossils of bornaviruses**. *Virus research* 2019, **262**:2-9.
9. Kojima S, Sato R, Yanai M, Komatsu Y, Horie M, Igarashi M, Tomonaga K: **Splicing-Dependent Subcellular Targeting of Borna Disease Virus Nucleoprotein Isoforms**. *Journal of virology* 2019, **93**(5).
10. Mukai Y, Horie M, Kojima S, Kawasaki J, Maeda K, Tomonaga K: **An endogenous bornavirus-like nucleoprotein in miniopterid bats retains the RNA-binding properties of the original viral protein**. *FEBS letters* 2022, **596**(3):323-337.

11. Jurka J, Klonowski P, Dagman V, Pelton P: **CENSOR--a program for identification and elimination of repetitive elements from DNA sequences**. *Comput Chem* 1996, **20**(1):119-121.

12. Poenisch M, Wille S, Staeheli P, Schneider U: **Polymerase read-through at the first transcription termination site contributes to regulation of borna disease virus gene expression**. *Journal of virology* 2008, **82**(19):9537-9545.

13. Morrish TA, Gilbert N, Myers JS, Vincent BJ, Stamato TD, Taccioli GE, Batzer MA, Moran JV: **DNA repair mediated by endonuclease-independent LINE-1 retrotransposition**. *Nat Genet* 2002, **31**(2):159-165.

14. Ferre CA, Davezac N, Thouard A, Peyrin JM, Belenguer P, Miquel MC, Gonzalez-Dunia D, Szelechowski M: **Manipulation of the N-terminal sequence of the Borna disease virus X protein improves its mitochondrial targeting and neuroprotective potential**. *FASEB J* 2016, **30**(4):1523-1533.

15. Poenisch M, Burger N, Staeheli P, Bauer G, Schneider U: **Protein X of Borna disease virus inhibits apoptosis and promotes viral persistence in the central nervous systems of newborn-infected rats**. *Journal of virology* 2009, **83**(9):4297-4307.

16. Wolff T, Unterstab G, Heins G, Richt JA, Kann M: **Characterization of an unusual importin alpha binding motif in the borna disease virus p10 protein that directs nuclear import**. *J Biol Chem* 2002, **277**(14):12151-12157.

17. Shoya Y, Kobayashi T, Koda T, Ikuta K, Kakinuma M, Kishi M: **Two proline-rich nuclear localization signals in the amino- and carboxyl-terminal regions of the Borna disease virus phosphoprotein**. *Journal of virology* 1998, **72**(12):9755-9762.

18. Fujino K, Horie M, Honda T, Nakamura S, Matsumoto Y, Franeischetti IM, Tomonaga K: **Evolutionarily conserved interaction between the phosphoproteins and X proteins of bornaviruses from different vertebrate species**. *PLoS one* 2012, **7**(12):e51161.

19. Pearson WR, Wood T, Zhang Z, Miller W: **Comparison of DNA sequences with protein sequences**. *Genomics* 1997, **46**(1):24-36.

20. Horie M: **The biological significance of bornavirus-derived genes in mammals**. *Current opinion in virology* 2017, **25**:1-6.

21. Rubbenstroth D, Schlottau K, Schwemmle M, Rissland J, Beer M: **Human bornavirus research: Back on track!** *PLoS Pathog* 2019, **15**(8):e1007873.

22. Maeda K, Hondo E, Terakawa J, Kiso Y, Nakaichi N, Endoh D, Sakai K, Morikawa S, Mizutani T: **Isolation of novel adenovirus from fruit bat**

(Pteropus dasymallus yayeyamae). *Emerging infectious diseases* 2008,
14(2):347-349.

23.   Maruyama J, Miyamoto H, Kajihara M, Ogawa H, Maeda K, Sakoda Y, Yoshida
R, Takada A: **Characterization of the envelope glycoprotein of a novel**
filovirus, lloviu virus. *Journal of virology* 2014, **88**(1):99-109.

24.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden
TL: **BLAST+: architecture and applications**. *BMC Bioinformatics* 2009,
10:421.

25.   Coordinators NR: **Database resources of the National Center for**
Biotechnology Information. *Nucleic Acids Res* 2018, **46**(D1):D8-D13.

26.   Pearson WR: **Finding Protein and Nucleotide Similarities with FASTA**. *Curr*
Protoc Bioinformatics 2016, **53**:3 9 1-3 9 25.

27.   Katoh K, Standley DM: **MAFFT multiple sequence alignment software**
version 7: improvements in performance and usability. *Mol Biol Evol* 2013,
30(4):772-780.

28.   Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ**
preprocessor. *Bioinformatics* 2018, **34**(17):i884-i890.

29.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL: **Graph-based genome**
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat*
Biotechnol 2019, **37**(8):907-915.

30.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G,
Mesirov JP: **Integrative genomics viewer**. *Nat Biotechnol* 2011, **29**(1):24-26.

31.   .

32.   Drozdetskiy A, Cole C, Procter J, Barton GJ: **JPred4: a protein secondary**
structure prediction server. *Nucleic Acids Res* 2015, **43**(W1):W389-394.

33.   Walker JM: **The proteomics protocols handbook**: Springer; 2005.

Figure legends

Figure 1. Amino acid identities between extant orthobornaviral proteins. Pairwise identities between the amino acid sequences of extant orthobornaviral proteins was analyzed using Sequence Demarcation Tool ref.

Figure 2. Characterization of miEBLN-1 as an EBL derived from N/X/P transcript. **(A)** Gene orthology analysis of the miEBLN/X/P-1 locus. Multiple alignment of the deduced syntenic loci between the indicated species is shown. Black and gray boxes show aligned regions. Thin gray lines indicate gaps. Species phylogeny and minimum integration age of miEBLN/X/P-1 are indicated on the left based on TimeTree ref. **(B)** Schematic diagram of orthobornaviral genomic RNA, 1.9-kb N/X/P mRNA, and miEBLN/X/P-1. Transcription signals (S1, S2, T1, and T2) are shown. **(C)** Nucleotide sequence alignment of extant orthobornaviral N/X/P mRNA and miEBLN/X/P-1. Red letters indicate the start and stop codons of the N gene. Light and dark teal blue letters indicate the start codon of X gene and the stop codon of P gene, respectively. TSD, target site duplication.

Figure 3. Identification of an EBL derived from X/P transcript in miniopterid bats. **(A)** Schematic diagram of the BLASTn search against the bat whole genome shotgun sequence database, using miEBLN/X/P-1 as query. The BLAST hits are shown by dark to light read or teal lines corresponding to the EBLs indicated on the right. **(B)** Gene orthology analysis of the miEBLX/P-4 locus. Species phylogeny and deduced minimum integration age of miEBLX/P-4 are indicated on the left based on TimeTree. **(C)** Schematic diagram of extant orthobornaviral genomic RNA, X/P mRNA, and

miEBLX/P-4. Transcription signals (S2, T1, and T2) are indicated. **(D)** Nucleotide

sequence alignment of extant orthobornaviral X/P transcripts and miEBLX/P-4. Light

and dark teal letters indicate the start codon of X gene and the stop codon of P gene,

respectively. TSD, target site duplication.

Figure 4. Conservation of properties between putative EBLX and EBLP and viral

X and P proteins. (A and **B)** Predicted secondary structure of extant orthobornaviral X

and miEBLX-4 proteins **(A)**, and extant orthobornaviral P and miEBLP-4 proteins **(B)**.

H, α-helix. E, β-strand. **(C** and **D)** Amino acid sequence alignment of extant

orthobornaviral X and miEBLX proteins **(C)**, and extant orthobornaviral P and

miEBLP-4 proteins **(D)**. Shaded areas indicate the functional domains previously

characterized in BoDV-1 X and P proteins. The conserved amino acid residues

important for the functions of viral X or P protein are highlighted in colors. **(E)** Co-

immunoprecipitation of Myc-tagged miEBLX-4 with FLAG-tagged miEBLP-4. FLAG-

tagged miEBLP-4 was immunoprecipitated, and then analyzed by western blotting

using the indicated antibodies. *, Non-specific band.

Figure 5. Identification of diverse EBLX/Ps in the vertebrate genomes.

(A-C) Gene orthology analysis of each EBLX/P locus. Species phylogeny and deduced

minimum integration age of each EBLX/P are indicated on the left based on TimeTree.

(D) Nucleotide sequence alignment of extant orthobornaviral X/P transcripts and

EBLX/Ps. Light teal and dark teal letters indicate the start codon of X gene and the stop

codon of P gene, respectively. TSD, target site duplication.

Figure 6. Alternative splicing generates *ZNF451/miEBLX/P-4* chimeric transcript.

(A) Coverage and splicing junctions of mapped RNA reads from SubK 12-08 cells (*M.*

schreibersii) at the *ZNF451* locus where miEBLX/P-4 is located. The miEBLX/P-4

region, mapped read coverage graph on miEBLX/P-4, and splicing junction between

ZNF451 and miEBLX/P-4 are shown in teal. **(B)** Schematic diagram of putative

ZNF451/EBLX/P-4 chimeric transcript. The transcript contains two ORFs in different

frames: miEBLX-4 alone and ZNF451 fused with miEBLP-4.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

A

Znf451 gene   73.9 kb

Znf451 mRNA isoforms   X1 X2 X3 X4

miEBL/X/P-4

Read coverage   560   0

Splicing junction

Putative *miEBL/XP-4* mRNA

B

Putative chimeric mRNA of
Znf451 and *miEBLX/P-4*

miEBLX-4
ZNF451   miEBLP-4-ext   2.1 kb