Subthalamic beta bursts correlate with dopamine-dependent motor symptoms in 106 Parkinson's patients

Roxanne Lofredi, MD[1,2], Liana Okudzhava, MSc[1], Friederike Irmen, PhD[1], Christof Brücke, MD/PhD[1], Julius Huebl, MD[1], Joachim K. Krauss, MD[3], Gerd-Helge Schneider, MD[4], Katharina Faust, MD[4], Wolf-Julian Neumann, MD[1,5], Andrea A. Kühn, MD[1,5,6,7,8]

[1]*Department of Neurology, Charité-Universitätsmedizin Berlin, Berlin, Germany*

[2] *Berlin Institute of Health (BIH), Berlin, Germany*

[3]*Department of Neurosurgery, Medizinische Hochschule Hannover, Hannover, Germany*

[4]*Department of Neurosurgery, Charité-Universitätsmedizin Berlin, Berlin, Germany*

[5]*Bernstein Center for Computational Neuroscience, Humboldt-Universität, Berlin, Germany*

[6] *NeuroCure, Exzellenzcluster, Charité-Universitätsmedizin Berlin, Berlin, Germany*

[7] *DZNE, German center for neurodegenerative diseases, Berlin, Germany*

[8] *Berlin School of Mind and Brain, Humboldt-Universität zu Berlin*

Corresponding author

Andrea A. Kühn
Department of Neurology - Campus Mitte,
Charité Universitätsmedizin Berlin,
Charitéplatz 1
10117 Berlin, Germany

Word Count: Abstract: 152 Words; **Main Text:** 2987 Words

Running Title

Beta bursts correlate with motor impairment

Financial disclosure

Dr. Lofredi, Dr. Neumann, Dr. Irmen, Dr. Brücke, Dr. Huebl and Ms Okudzhava report no conflicts of interest, outside the submitted work. Prof. Kühn reports personal fees from Medtronic, personal fees from Boston Scientific, personal fees from Abbott, personal fees from Ipsen Pharma, personal fees from Teva, outside the submitted work. Dr. Krauss reports personal fees from Medtronic, personal fees from Boston scientific, outside the submitted work. Dr. Schneider and Dr. Faust report personal fees from Medtronic, personal fees from Boston Scientific, personal fees from Abbott, outside the submitted work.

Funding Sources

Dr. Lofredi is participant in the BIH Charité Clinician Scientist Program funded by the Charité – Universitätsmedizin Berlin, and the Berlin Institute of Health at Charité (BIH). The work was supported by Deutsche Forschungsgemeinschaft (Project-ID 424778381 – TRR 295 Retune).

Abstract

Background: Pathologically increased beta power has been described as a biomarker for Parkinson's disease (PD) and related to prolonged bursts of subthalamic beta synchronization.

Methods: Here, we investigate the association between subthalamic beta dynamics and motor impairment in a cohort of 106 Parkinson's patients in the ON- and OFF- medication state, suing two different methods of beta burst determination.

Results: We report a frequency-specific correlation of low beta power and burst duration with motor impairment OFF dopaminergic medication. Furthermore, reduction of power and burst duration correlated significantly with symptom alleviation through dopaminergic medication. Importantly, qualitatively similar results were yielded with two different methods of beta burst definition.

Conclusions: Our findings validate the robustness of previous results on pathological changes in subcortical oscillations both in the frequency- as well as in the time-domain in the largest cohort of PD patients to date with important implications for next-generation adaptive deep brain stimulation control algorithms.

Introduction

Intracerebral recordings from patients undergoing deep brain stimulation (DBS) surgery for Parkinson' disease (PD) have revealed increased subthalamic oscillatory beta band activity (13-35 Hz) as a pathophysiological hallmark of the PD motor state[1–4]. Both dopaminergic medication as well as DBS are associated with a decrease of this pathologically enhanced activity, specifically in the low beta sub-band (LBeta=13-20 Hz) [5–8]. In the dopamine-depleted state, several studies have reported a significant correlation between parkinsonian symptom severity and beta synchronization[7,9], even months after chronic stimulation[10,11]. Thus, beta activity is considered a stable biomarker of motor signs in PD and has been used as feedback signal for therapeutic demand in experimental studies on adaptive closed-loop DBS (aDBS) with similar or better efficacy / side-effect profile than conventional DBS [12–15]. The beneficial effects of aDBS have been linked to a shortening of pathologically prolonged periods of beta synchronization, so-called beta bursts. In contrast, the overall decrease of beta burst duration with conventional DBS[16] possibly relies on suppression of both short and long beta bursts [17,18]. Suppressing short beta bursts, however, might be disadvantageous as they are considered part of physiological signaling within the motor circuit [19]. Accordingly, a shift towards shorter beta bursts is observed with dopaminergic medication[2,20,21], which has been correlated with clinical improvement in one study so far[22]. With increasing interest for the pathophysiological role of beta bursts in PD, novel methods of burst definition have recently been proposed [23]. These have been validated by investigating DBS-induced changes but not used for capturing dopamine-related dynamics of beta bursts yet [16,24].

The present study aims to test the robustness of previous results on a dopamine-dependent correlation between subthalamic beta burst dynamics and motor symptoms in Parkinson's disease by i) validating these findings in the largest cohort to date of > 100 PD patients and ii) comparing an established with a newly developed method of beta burst assessment.

Methods and Materials

Patients and Surgery. For this study, we have identified archival local field potential (LFP) data from 106 PD-patients (63.5±8 years, 42 female, clinical subtype: 47% akinetic-rigid; 20% tremor-dominant; 33% equivalent) who underwent bilateral implantation of subthalamic DBS-electrodes. In 95 cases, assessment of motor impairment using the UPDRS-III (denominated UDPRS in the following) was available both under regular dopaminergic medication (ON

state) and at least 12 hours after dopaminergic drug withdrawal (OFF state) either pre-operatively (n=28) or during the recording sessions (n=67). Motor impairment was significantly more pronounced in the OFF than in the ON medication state (UPDRS-OFF: 32.7 ± 12.7, UPDRS-ON: 18.3 ± 9.3, P=.002). All patients gave written informed consent, which was approved by the local ethics committee in accordance with the standards set by the Declaration of Helsinki. Three types of DBS-macroelectrodes were used: Medtronic 3389 (n=80), Boston Vercise with circular electrodes (n=11) and Boston Vercise Cartesia (n=15), in which mid-contacts are segmented. Intraoperative microelectrode recordings, test stimulation and postoperative imaging ensured correct DBS-electrode placement. A subset from a previous study (n=45) that assessed beta power in the OFF medication state of PD patients have been included in the present cohort[9].

Recordings. Subthalamic LFP recordings of at least 200 seconds were performed in the ON- and OFF medication state in the 1-7 days following DBS-surgery. Patients were comfortably seated and instructed not to move during the recording. Signals were amplified (50.000x) using a D360 amplifier (Digitimer, Hertfordshire, UK) and recorded at a sampling frequency of 1 kHz through a 1401 A-D converter (CED, Cambridge, UK) onto a computer using Spike2 software. When electrode contacts were circular, bipolar LFPs were recorded between adjacent contact pairs. In directional leads, recordings were referenced to the lowermost contact of the electrode, subsequently summed and re-referenced offline to approximate the bipolar recordings derived from adjacent pairs of circular contacts, for example: (contact 1/2 + contact 1/3 + contact 1/4)-(contact 1/5 + contact 1/6 + contact1/7).

Signal Processing. Artifact-free recording segments were identified visually and analyzed using custom MATLAB code (The Mathworks, Natick, Massachusetts) based on SPM12 (http://www.fil.ion.ucl.ac.uk/spm/) and FieldTrip (http://fieldtrip.fcdonders.nl/). Continuous recordings were filtered (high pass=5 Hz, low pass=98 Hz and notch filter=48-52 Hz) and transferred into the frequency domain using Morlet wavelets with 10 cycles and a temporal resolution of 200 Hz. Power spectra were derived through averaging over time and normalized to the total sum of the recording and further expressed as a percentage of total power. Normalization allows comparison across participants as arbitrary amplitude differences due to electrode impedance and distance to source are removed. Frequency bands of interest in this study where predefined as alpha (8-12 Hz), low beta (LBeta, 13-20 Hz) and high beta (HBeta, 21-35 Hz) band. Spectral peak frequencies and the channel displaying the highest spectral peak in the respective frequency bands of interest (alpha=8-12 Hz, Lbeta=13-20 Hz, HBeta=21-35

Hz) were automatically detected, see Figure 1. For correlative analyses with motor impairment, power values in each frequency band were calculated for the channel with highest peak frequency amplitude in the OFF medication state, separately for each hemisphere and averaged across hemispheres. There was no significant shift in peak frequency from the OFF to the ON medication state (P>.05) and no difference in band specific power between male and female subjects (P>.05) in either frequency band.

Figure 1: Exemplary trace of power spectra in the ON- and OFF-medication state. Note the multiple spectral peaks can be detected within the same patient or even the same recording channel in the alpha, low and high beta band. Specifically the detected peaks in the low beta range decrease in the ON medication state, while high beta activity remains prominent. Peak frequency with highest amplitude is labeled with a black dot, separately for each respective frequency band of interest. Shown is the left hemisphere of subject 31.

Beta Burst Analysis. For burst determination, the bipolar channel with highest peak in the respective frequency band was selected in the OFF medication state and used for both medication states. Two methods were used separately for burst determination and compared qualitatively. The first method was recently established and qualifies beta bursts when beta activity exceeds the overall noise floor within a recording [16,23]. For this approach, the raw signal was bandpass-filtered around the frequency band of interest (here low or high beta band) and envelope peaks of the squared signal were linearly connected to generate an amplitude envelope. A beta burst was detected when the amplitude crossed a threshold defined as four times the median of averaged troughs from envelopes of five overlapping 6 Hz bands in the low gamma band (55-65 Hz) from the same recording. The second method was used in previous

publications by the authors [2,25,26], qualifiying beta bursts as events in which beta activity is signifcantly higher than the overall beta amplitude distribution. Here, the wavelet amplitude was separately averaged across the respective frequency band of interest and smoothed with a moving average Gaussian smoothing kernel (175 ms). Each power amplitude trace was z-scored (X-μ/δ) over the entire recording, separately for the ON- and OFF-medication state. The threshold for bursts was defined as the 75th percentile of the normalized signal amplitude distribution. A separate threshold was calculated in the ON and OFF medication state and the average of both was applied for burst determination in both medication states. In both methods, beta burst duration was defined as time spent above the defined threshold. The distribution of burst durations was considered by categorization into 7 time windows of 100 ms starting from 100 ms to > 700 ms in duration. Additionally, burst amplitude was defined as maximal amplitude reached during a beta burst. As for power analyses, obtained values were averaged across hemispheres for correlative analyses with motor impairment.

Statistics. Nonparametric Monte Carlo permutation tests with 5000 permutations were used for statistical analyses as they do not rely on assumptions about the underlying data distribution. For correlative analyses, Spearman's correlation (denoted as Rho) was calculated and tested for significance using permutation. All results are indicated as mean ± standard deviation and reported significant at an α level of .05. Multiple comparisons were controlled by using detection of the false discovery rate (FDR) where appropriate.

Results

Correlation of beta power and motor impairment. Relative power spectra were averaged across contact pairs separately for the ON- and OFF-medication state. Each frequency bin was tested for difference across medication states, revealing a significant decrease (P<.001) of relative power between 10-19 Hz with dopaminergic medication when averaged across all channels or 10-20 Hz when only considering the channel with most pronounced Lbeta peak per hemisphere in the OFF medication state (see Figure 2A). While significant correlation with symptom severity in the OFF medication state was found in a slightly broader spectrum (9-22 Hz), only frequencies within the Lbeta band (13-19 Hz) were significantly associated to symptom alleviation with dopaminergic medication (see Figure 2B). To remain comparable with previous studies, the following analyses were performed with a predefined Lbeta band from 13-20 Hz. When averaged across the Lbeta band, there was a significant correlation between motor impairment as assessed by the UPDRS-III and Lbeta power in the OFF-medication state

(Rho=0.21, P=.03, see Figure 2C). While this correlation was not significant in the ON-medication state (P>.05), the dopamine-related motor improvement (UPDRS OFF-ON) was significantly associated with the reduction in Lbeta band power (Rho=0.36, P=.0008, see Figure 2C) as well. There was no significant correlation of motor impairment or improvement with averaged Hbeta power or averaged alpha power.

Figure 2: Low beta power in the OFF-medication state and its reduction with dopamine correlates with symptom severity. (A) Averaged power spectra across contact pairs in the ON (grey) and OFF (blue) medication state. Note the decrease in relative power with dopaminergic medication that includes the low beta and sub-beta band (underlined in grey, P<.0001). The effect is more pronounced if only the channels with highest low beta activity are considered instead of averaging across channels (lower row). (B) Matching the frequency bins that show a significant modulation with dopamine, amplitude of frequencies between 9-22 Hz show a significant correlation with motor symptoms as assessed by the UPDRS-III. However, symptom alleviation with dopamine is best reflected by amplitude changes from 13-19 Hz, commonly referred to as low beta band. In blue are shown Rho-values for each frequency bin, the scattered grey line shows the correspondent P-value, significant areas are underlined in grey. (C) There is a significant correlation between symptom severity and averaged low beta power (13-20 Hz) in the dopamine-depleted sate (Rho=0.208, P=.03). Likewise, the reduction in low beta power with dopamine correlates with symptom alleviation (Rho=0.36, P=.0084, right column).

Correlation of beta burst duration and motor impairment. When using a baseline as established in Anderson et al., 2020, averaged Lbeta burst duration was significantly prolonged after 12-hour withdrawal of dopaminergic medication (OFF=579.1± 469.1 ms; ON=358.6 ± 230.2 ms; P<.0001), see Figure 3A. In the OFF-medication state, the averaged Lbeta burst duration correlated significantly with motor impairment (Rho=0.18, P=.05) as did the shortening of mean Lbeta burst duration with motor improvement (Rho=0.26, P=.008), see Figure 3B. There was no significant correlation between LBeta burst duration and motor impairment in the ON-

medication state. When burst time bins were tested separately, motor impairment in the OFF state was positively correlated with the amount of bursts above 500 ms length (500-600 ms: Rho=0.29, P=.005; 600-700 ms: Rho=0.26, P=.01) and negatively correlated with short beta bursts (100-200 ms: Rho=-0.18, P=.05; 200-300 ms: Rho=-0.2, P=.03), see Figure 3C. Accordingly, changes in short and long bursts may best reflect symptom alleviation with medication (100-200 ms : Rho=-0.19, P=.04; 200-300 ms: Rho=-0.2, P=.03; > 700 ms : Rho=0.26, P=.02), see Figure 3C. However, these timebin-specific results did not survive FDR-correction for multiple comparisons. There was no significant correlation between motor symptoms and Lbeta burst amplitude or Hbeta burst properties in either medication state.

Figure 3: Low beta burst duration decreases with dopamine and correlates with symptom severity. (A) There is a significant shortening of averaged low beta burst duration in the ON-medication state (P<.001). (B) When averaged, the mean LBeta burst duration correlates with motor impairment in the OFF medication state (left column, Rho=.18, P=.05) and symptom alleviation with medication (right column, Rho=.26, P=.008). (C) While the amount of short LBeta bursts correlates negatively, the amount of long Lbeta bursts correlates positively with motor impairment (left column). A similar distribution is seen for the correlation with dopamine-related symptom alleviation and change in amount of short vs long Lbeta bursts (right column). However, the time-bin related results do not remain significant after FDR-correction. Shown are P-values per time-bin before FDR-correction. In boxplots, central marks indicate the median and edges the 25th and 75th percentiles of the distribution. *** p<0.001.

Comparison of different methods for burst detection. As expected, beta bursts where significantly shorter when using the common threshold method based on the 75th percentile of the amplitude distribution (OFF=304.9± 42.3 ms vs 579.1± 469.1 ms, P<.001; ON=289.4 ± 36.7 ms vs 358.6 ± 230.2 ms; P=.001). However, there was a significant correlation between

averaged burst duration between both methods (OFF: Rho=.56, P<.001; ON: Rho=.63, P<.001), specifically with regard to the amount of very short and very long Lbeta bursts in both medication states (OFF: 100-200 ms: Rho=.46, P<.001; >700 ms: Rho=.53, P<.001; ON: 100-200 ms: Rho=.52, P<.001; >700 ms: Rho=.53, P<.001).

In the OFF medication state, common threshold based results were qualitatively similar to the baseline-based results with a frequency-specific correlation between mean Lbeta burst duration and motor impairment (Rho=.27, P=.008) that may rely on the amount of long bursts (>700 ms: Rho=.22, P=.02) and negatively associated with the amount of very short bursts (100-200 ms: Rho=.18, P=.05; uncorrected). However, dopamine-related changes were less well captured by the common threshold based method. While a significant reduction of averaged burst duration with medication was observed (OFF=304.9± 42.3 ms; ON=289.4 ± 36.7 ms; P<.0001), its amount did not correlate with symptom alleviation, neither when averaged across all bursts nor when considered separately for each time bin. For a summary of common threshold based results, see Supp. Fig 1.

Discussion

In the present study, we investigated the association between subthalamic beta power and bursts with motor symptoms as well as their change with dopaminergic medication in a large cohort of 106 PD-patients. We show a significant correlation between motor symptom severity in the OFF medication state and oscillatory activity from ~10-20 Hz. Moreover, the medication-related reduction of Lbeta power (13-20 Hz) was paralleled by a dopamine-related symptom alleviation. Similar to Lbeta power, averaged Lbeta burst duration significantly correlated with motor impairment in the OFF medication state, as did dopamine-related reduction of LBeta burst duration with symptom alleviation, when using the noise floor as baseline. All observed effects were frequency-specific for the Lbeta band. We thus provide further evidence for temporal dynamics of Lbeta activity as a robust biomarker of motor symptoms in a large cohort of PD patients, relatively independent of the exact method used for burst determination.

Frequency-specific prolongation of low beta bursts in the dopamine-depleted state.

Our results are in line with previous studies, demonstrating a correlation of parkinsonian symptoms in the OFF medication state [9] and a specific dopamine-sensitivity of activity mainly covering the Lbeta band [27], which has also been described for DBS-related effects[8]. A recent study combining magnetoencephalography, subthalamic vs. pallidal LFP and computational

modelling, has attributed the difference of high- and low-beta to the hyperdirect- and indirect-pathway [28] . The findings support the notion that PD symptoms may arise from hyperactivity of the indirect pathway, while hyperdirect cortico-subthalamic communication is implicated in cognitive control and motor inhibition[29,30].

Regarding the more recently emerged concept of pathologically prolonged beta bursts in the OFF medication state, frequency-specificity for the Lbeta band has been explicitly reported only in one study investigating the association between beta bursts and bradykinesia [25]. In two other studies, beta bursts were determined around beta peak frequencies, the average of which lied in the Lbeta band as well (Tinkhauser et al., 2017: 19.9 Hz; Lofredi et al., 2019: 14.3 Hz). Not all studies have produced this distinction of high- and low beta activity, e.g. a more recently published study reported a frequency-specific correlation of high beta burst duration with motor symptoms [20]. The precise differences remain unknown, but it is important to note that potentially electrode impedance and spatial coverage may have an impact on the recorded frequency bands in LFP [31]. All of the aforementioned studies however reported results on <30 PD-patients. The present study provides further evidence that in a large cohort, the best across patient correlation with symptom severity is obtained with burst properties in the Lbeta band.

Qualitatively similar results with different methods for burst determination.

Methodologically, a vivid scientific controversy has risen around the potential pitfalls of burst determination. Earlier studies established thresholds for bursts based on amplitude distribution either by considering only amplitudes above a specific percentile[21] or median deviation [32] that were applied by following studies [2,20,25,33–35]. Others tried to train discriminators on artificially generated time series for burst detection, which is a promising direction, but might need further knowledge on the underlying amplitude distribution before assuring to generate physiologically meaningful results [17]. More recently, a novel method of burst determination, using a physiological baseline based on the recording-specific noise floor was proposed [23] and used in several studies investigating DBS-associated beta burst dynamics in PD-patients with chronically implanted pulse generators [16,24]. However, this method had not been used to assess dopamine-related beta burst dynamics in externalized recordings of PD patients yet. Here, we obtain similar burst durations in the OFF medication state than these studies, that were moreover qualitatively similar, in both used methodological approaches. This provides further evidence for a stable, hardware-independent, medication- as well as stimulation sensitive and

thus pathophysiological relevant correlation between temporal dynamics of Lbeta activity and motor state in PD patients.

Limitations of beta power and beta bursts as a biomarker.

Although subthalamic Lbeta power and burst duration appear as robust biomarkers across PD-patients, the explained variance of motor symptom severity remains relatively low, with rho-values ranging around 0.4-0.5 in previous studies and 0.18-0.36 in the present study with the largest cohort to date. This low explanatory power might relate to the following aspects: First, subthalamic LBeta power and burst duration might reflect similar aspects of the parkinsonian motor state as burst duration and amplitude are highly correlated. Thus, combining both does not significantly increase the explained variance. Second, Lbeta power and burst properties lose their explanatory value for symptom severity when considering the ON medication state and additional features beyond beta, such as movement- or dyskinesia-related low-frequency or gamma activity may be more performative in this state[33,36]. Third, an individualized investigation of combined spectral changes assessed on a case-by-case basis, may outperform the information content of spectral features based on group results for individual patients. Thus, while our study provides clear evidence for the pathophysiological relevance of LBeta in PD, other frequency bands may be better suited to act as control signals in individual patients. Also, cortical signals might be advantageous when compared to subcortical activity patterns in decoding specific motor states [37,38]. In the same vein, multisite recordings could be necessary if the entirety of motor- and non-motor symptoms have to be captured, given that PD is considered a network disease and pathological interregional connections have been described, specifically for motor cortical areas [39–41]. Interestingly, a first pioneering study investigating symptom correlations from combined chronic ECoG and LFP recordings in freely moving patients found different combinations of recording source (Motor cortex vs. STN vs. Motor cortex – STN coupling) and frequency band (alpha, beta, gamma) for each of the five patients to be most predictive of symptom severity [42].

Outlook.

In conclusion, our study corroborates and extends previous findings on the pathophysiological relevance of beta burst dynamics in PD. Moreover, our findings highlight the importance of understanding temporal brain circuit dynamics to inspire the design of bidirectional control algorithms for on-demand adaptive neuromodulation. The fact that the shift from very short to pathologically prolonged bursts can be observed in such a large cohort of PD-patients increases

the confidence that an aDBS-approach, where stimulation would be triggered by beta amplitude threshold crossings of specific lengths, can be viable and advantageous in clinical settings. Large multicenter trials are currently in development with manufacturers offering first generations of commercially available sensing enabled devices. Our results suggest that aDBS trials should implement temporal threshold parameters beyond the single or dual power threshold paradigms to consider temporal burst dynamics as a key hallmark of pathological basal ganglia activity in PD.

Supp. Figures

Supp. Figure 1: Low beta dynamics using the common threshold method based on the 75th percentile of the amplitude distribution. (A) There is a significant shortening of averaged low beta burst duration in the ON-medication state (P<.001). (B) The increasing burst duration OFF-medication relies on a shift from very short bursts (100-200 ms) to bursts with prolonged duration exceeding 700 ms. Shown is the percentage change from the ON- to the OFF-medication state. (C) Motor impairment shows a significant positive correlation with the

averaged Lbeta burst duration. (D) When tested separately, the amount of very long bursts in the OFF-medication state correlates positively with symptom severity (Rho=0.22, P=0.02), while the amount of very short Lbeta bursts show a trend towards a negative correlation with motor impairment. However, both results do not remain significant after FDR-correction. In boxplots, central marks indicate the median and edges the 25th and 75th percentiles of the distribution. *** p<0.001.

Acknowledgements

We would like to thank all patients for their participation.

Author Contributions

Roxanne Lofredi:	conception and design of the study, acquisition and analysis of data and drafting a significant portion of the manuscript or figures.
Liana Okudzhava:	analysis of data.
Friederike Irmen:	acquisition and analysis of data.
Christof Brücke:	acquisition of data.
Julius Hübl:	acquisition of data.
Joachim K. Krauss:	acquisition of data.
Gerd- Helge Schneider:	acquisition of data.
Katharina Faust:	acquisition of data.
Gerd- Helge Schneider:	acquisition of data.
Wolf-Julian Neumann:	acquisition and analysis of data.
Andrea A. Kühn:	conception and design of the study, drafting a significant portion of the manuscript.

References

1. Geng, X. *et al.* Comparison of oscillatory activity in subthalamic nucleus in Parkinson's disease and dystonia. *Neurobiol. Dis.* **98**, 100–107 (2017).

2. Lofredi, R. *et al.* Pallidal beta bursts in Parkinson's disease and dystonia. *Mov. Disord.* **34**, (2019).

3. Silberstein, P. *et al.* Patterning of globus pallidus local field potentials differs between Parkinson's disease and dystonia. *Brain* **126**, 2597 (2003).

4. Wang, D. D. *et al.* Pallidal Deep-Brain Stimulation Disrupts Pallidal Beta Oscillations

and Coherence with Primary Motor Cortex in Parkinson's Disease. *J. Neurosci.* **38**, 4556–4568 (2018).

5. Eusebio, A. *et al.* Deep brain stimulation can suppress pathological synchronisation in parkinsonian patients. *J. Neurol. Neurosurg. Psychiatry* **82**, 569–573 (2011).

6. Kühn, A. A. *et al.* High-Frequency Stimulation of the Subthalamic Nucleus Suppresses Oscillatory β Activity in Patients with Parkinson ' s Disease in Parallel with Improvement in Motor Performance. *J. Neurosci.* **28**, 6165–6173 (2008).

7. Kühn, A. A., Kupsch, A., Schneider, G.-H. & Brown, P. Reduction in subthalamic 8-35 Hz oscillatory activity correlates with clinical improvement in Parkinson's disease. *Eur. J. Neurosci.* **23**, 1956–1960 (2006).

8. Oswal, A. *et al.* Deep brain stimulation modulates synchrony within spatially and spectrally distinct resting state networks in Parkinson ' s disease. 1482–1496 (2016) doi:10.1093/brain/aww048.

9. Neumann, W.-J. *et al.* Subthalamic synchronized oscillatory activity correlates with motor impairment in patients with Parkinson's disease. *Mov. Disord.* **31**, 1748–1751 (2016).

10. Neumann, W.-J. *et al.* Long term correlation of subthalamic beta band activity with motor impairment in patients with Parkinson's disease. *Clin. Neurophysiol. Off. J. Int. Fed. Clin. Neurophysiol.* **128**, 2286–2291 (2017).

11. Trager, M. H. *et al.* Subthalamic beta oscillations are attenuated after withdrawal of chronic high frequency neurostimulation in Parkinson's disease. *Neurobiol. Dis.* **96**, 22–30 (2016).

12. Little, S. *et al.* Bilateral adaptive deep brain stimulation is effective in Parkinson's disease. *J. Neurol. Neurosurg. Psychiatry* **87**, 717–721 (2016).

13. Little, S. *et al.* Adaptive deep brain stimulation in advanced Parkinson disease. *Ann. Neurol.* **74**, 449–457 (2013).

14. Piña-Fuentes, D. *et al.* Toward adaptive deep brain stimulation for dystonia. *Neurosurg. Focus* **45**, E3 (2018).

15. Pina-Fuentes, D. *et al.* Adaptive DBS in a Parkinson's patient with chronically implanted DBS: A proof of principle. *Movement disorders : official journal of the Movement Disorder Society* vol. 32 1253–1254 (2017).

16. Kehnemouyi, Y. M. *et al.* Modulation of beta bursts in subthalamic sensorimotor circuits predicts improvement in bradykinesia. *Brain* **144**, 473–486 (2021).

17. Schmidt, S. L., Peters, J. J., Turner, D. A. & Grill, W. M. Continuous deep brain

stimulation of the subthalamic nucleus may not modulate beta bursts in patients with Parkinson's disease. *Brain Stimul.* **13**, 433–443 (2020).

18. Tinkhauser, G. *et al.* The modulatory effect of adaptive deep brain stimulation on beta bursts in Parkinson's disease. *Brain* **140**, 1053–1067 (2017).

19. Shin, H., Law, R., Tsutsui, S., Moore, C. I. & Jones, S. R. The rate of transient beta frequency events predicts behavior across tasks and species. *Elife* **6**, 1–31 (2017).

20. Sure, M., Vesper, J., Schnitzler, A. & Florin, E. Dopaminergic Modulation of Spectral and Spatial Characteristics of Parkinsonian Subthalamic Nucleus Beta Bursts   . *Frontiers in Neuroscience*   vol. 15 1450 (2021).

21. Tinkhauser, G. *et al.* Beta burst dynamics in Parkinson's disease OFF and ON dopaminergic medication. *Brain* **140**, 2968–2981 (2017).

22. Tinkhauser, G. *et al.* Beta burst dynamics in Parkinson ' s disease OFF and ON dopaminergic medication. 2968–2981 (2018) doi:10.1093/brain/awx252.

23. Anderson, R. W. *et al.* A novel method for calculating beta band burst durations in Parkinson's disease using a physiological baseline. *J. Neurosci. Methods* **343**, 108811 (2020).

24. Anidi, C. *et al.* Neuromodulation targets pathological not physiological beta bursts during gait in Parkinson's disease. *Neurobiol. Dis.* **120**, 107–117 (2018).

25. Lofredi, R. *et al.* Beta bursts during continuous movements accompany the velocity decrement in Parkinson's disease patients. *Neurobiol. Dis.* **127**, 462–471 (2019).

26. Lofredi, R. *et al.* Dopamine-dependent scaling of subthalamic gamma bursts with movement velocity in patients with Parkinson's disease. *Elife* **7**, (2018).

27. Priori, A. *et al.* Rhythm-specific pharmacological modulation of subthalamic activity in Parkinson's disease. *Exp. Neurol.* **189**, 369–379 (2004).

28. Oswal, A. *et al.* Neural signatures of hyperdirect pathway activity in Parkinson's disease. *Nat. Commun.* **12**, 5185 (2021).

29. Lofredi, R. *et al.* Subthalamic stimulation impairs stopping of ongoing movements. *Brain* **144**, 44–52 (2021).

30. Neumann, W.-J. *et al.* Functional segregation of basal ganglia pathways in Parkinson's disease. *Brain* **141**, 2655–2669 (2018).

31. Buzsáki, G. Rhythms of the brain. in (Oxford University Press, 2006). doi:http://dx.doi.org/10.1093/acprof:oso/9780195301069.001.0001.

32. Feingold, J., Gibson, D. J., DePasquale, B. & Graybiel, A. M. Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys

performing movement tasks. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13687–13692 (2015).

33. Lofredi, R. *et al.* Dopamine-dependent scaling of subthalamic gamma bursts with movement velocity in patients with Parkinson's disease. *Elife* **7**, (2018).

34. Torrecillos, F. *et al.* Modulation of Beta Bursts in the Subthalamic Nucleus Predicts Motor Performance. *J. Neurosci.* **38**, 8905–8917 (2018).

35. Neumann, W.-J. *et al.* Pallidal and thalamic neural oscillatory patterns in tourette's syndrome. *Ann. Neurol.* **84**, 505–514 (2018).

36. Alegre, M. *et al.* Subthalamic activity during diphasic dyskinesias in Parkinson's disease. *Mov. Disord.* **27**, 1178–1181 (2012).

37. Swann, N. C. *et al.* Gamma Oscillations in the Hyperkinetic State Detected with Chronic Human Brain Recordings in Parkinson's Disease. *J. Neurosci.* **36**, 6445–6458 (2016).

38. Merk, T. *et al.* Electrocorticography is superior to subthalamic local field potentials for movement decoding in Parkinson's disease. *bioRxiv* 2021.04.24.441207 (2021) doi:10.1101/2021.04.24.441207.

39. Swann, N. C. *et al.* Adaptive deep brain stimulation for Parkinson's disease using motor cortex sensing. *J. Neural Eng.* **15**, (2018).

40. de Hemptinne, C. *et al.* Therapeutic deep brain stimulation reduces cortical phase-amplitude coupling in Parkinson's disease. *Nat. Neurosci.* **18**, 779–786 (2015).

41. Litvak, V. *et al.* Movement-related changes in local and long-range synchronization in Parkinson's disease revealed by simultaneous magnetoencephalography and intracranial recordings. *J. Neurosci.* **32**, 10541–10553 (2012).

42. Gilron, R. *et al.* Long-term wireless streaming of neural recordings for circuit discovery and adaptive stimulation in individuals with Parkinson's disease. *Nat. Biotechnol.* **39**, 1078–1085 (2021).