Sonic Hedgehog Activates Prostaglandin Signaling to Stabilize Primary Cilium Length

Shariq S. Ansari[1], Miriam E. Dillard[1], Yan Zhang[1], Mary Ashley Austria[1,2], Naoko Boatwright[3], Elaine L. Shelton[3,4], Amanda Johnson[5], Brandon M. Young[6], Zoran Rankovic[6], John D. Schuetz[7], Camenzind G. Robinson[5], and Stacey K. Ogden[1]

[1]Department of Cell and Molecular Biology, [5]Cell and Tissue Imaging Center, [6]Department of Chemical Biology and Therapeutics, [7]Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN 38105

[3]Department of Pediatrics, Monroe Carell Jr. Children's Hospital at Vanderbilt and Vanderbilt University Medical Center, [4]Department of Pharmacology, Vanderbilt University School of Medicine, Nashville, TN 37232

[2]Rhodes College Summer Plus Program, Memphis, TN 38112

Correspondence: Stacey.Ogden@stjude.org

ABSTRACT

Sonic Hedgehog (SHH) signaling is an essential driver of embryonic patterning that, when

corrupted, leads to developmental disorders and cancers. SHH effector responses are

organized through nonmotile primary cilia that grow and retract with the cell cycle and in

response to distinct extracellular cues. Destabilization of primary cilium length corrupts SHH

pathway regulation, which places significant pressure on SHH to maintain ciliary architecture.

Herein, we investigate whether SHH signaling promotes ciliary length control. We reveal a

signal crosstalk circuit induced by SHH activation of Phospholipase A2 ($cPLA_2$) that drives

ciliary EP_4 signaling to stabilize primary cilium length. We demonstrate that chemical or genetic

blockade of SHH-EP_4 crosstalk leads to destabilized primary cilium cyclic AMP (cAMP) control,

reduced ciliary length, and attenuated SHH pathway induction. Accordingly, we find that $Ep_4^{-/-}$

mice display shortened neuroepithelial primary cilia and altered SHH-dependent neuronal cell

fate specification. Thus, SHH initiates a signaling crosstalk circuit that maintains primary cilium

length for a robust downstream signaling response.

INTRODUCTION

During embryonic patterning, Sonic Hedgehog (SHH) signaling proteins provide instructional

cues that guide cell fate decisions to organize tissues. Disruption of SHH pathway activation

leads to developmental disorders and, consistent with regulated signaling being important for

tissue homeostasis post-development, aberrant SHH activity drives cancer (Barakat et al.,

2010). As such, sophisticated regulatory mechanisms are in place to ensure appropriate

pathway control in both basal and activated signaling states. Tight regulation of pathway activity

requires localization of SHH pathway components to a specialized sensory organelle called the

primary cilium (Caspary et al., 2007; Huangfu and Anderson, 2006). Each cell possesses a

single primary cilium that is anchored to the basal body and allows for organization of numerous

signal transduction cascades to instruct diverse cellular responses (Anvarian et al., 2019;

Wheway et al., 2018). Despite the membrane of the primary cilium being contiguous with the

plasma membrane, it maintains a distinct membrane lipid composition through action of lipid

metabolic enzymes that localize near the ciliary base (Arensdorf et al., 2017; Findakly et al.,

2021; Garcia et al., 2018; Raleigh et al., 2018). Further, a cytoplasmic diffusion barrier controls

entry of soluble proteins into the ciliary body, ensuring that primary cilia offer protected

environments for interpretation and transduction of myriad extracellular signals (Garcia et al.,

2018; Shi et al., 2017; Ye et al., 2013).

The SHH receptor Patched (PTCH), signal transducer Smoothened (SMO), and GLI2/GLI3

transcriptional effectors all function at or cycle through primary cilia in ligand-regulated manners

(Arensdorf et al., 2016; Petrov et al., 2017). In the absence of SHH, SMO retention in primary

cilia is restricted by PTCH-mediated depletion of SMO-activating sterols from ciliary membrane

(Kinnebrew et al., 2019; Kinnebrew et al., 2021; Kong et al., 2019). In this off state,

phosphorylation of GLI2 and GLI3 transcription factors by cyclic AMP (cAMP)-dependent protein

kinase (PKA) in primary cilia tags them for partial degradation to remove their transcriptional

activation domains (Bangs and Anderson, 2017; Haycraft et al., 2005; Tuson et al., 2011). SHH

binding to PTCH attenuates PTCH-mediated ciliary sterol elimination, which leads to SMO sterol

binding and retention in primary cilia where it signals to block PKA-promoted GLI truncation.

This allows for accumulation of full-length GLI2/3 proteins that are subsequently activated for

nuclear translocation and target gene induction (Niewiadomski et al., 2014; Umberger and

Ogden, 2021).

Current models suggest dependency of SHH signaling on primary cilia evolved due to the

specialized lipid composition of ciliary membrane allowing for tight control over availability of

SMO-activating sterols (Kinnebrew et al., 2019; Kinnebrew et al., 2021; Kong et al., 2019;

Raleigh et al., 2018). Further, the small size of primary cilia allows for rapid modulation of

second messengers such as cAMP (Hansen et al., 2020; Truong et al., 2021). Consistent with

this functionality and the key role of PKA in regulating GLI that cycles through primary cilia, SHH

signal output is highly sensitive to changes in ciliary cAMP concentration (Mukhopadhyay et al.,

2013; Truong et al., 2021; Vuolo et al., 2015). Pathway activation lowers primary cilium cAMP to

reduce PKA activity by stimulating ciliary exit of $G\alpha_s$-coupled GPR161 (Mukhopadhyay et al.,

2013). In addition, SMO can activate heterotrimeric G proteins of the $G\alpha_i$ family to inhibit

adenylyl cyclase (AC), suggesting that high-level activation of SHH signaling can lead to

sustained reduction in primary cilium cAMP (Barzi et al., 2011; Ogden et al., 2008; Riobo and

Manning, 2007).

Paradoxically, a growing body of evidence supports that ciliary length stability requires high

ciliary cAMP. Studies in zebrafish demonstrate that Multidrug Resistance Protein 4 Transporter

(ABCC4), which exports the prostaglandin signaling molecule PGE_2, is required for motile cilium

elongation. Ciliary length is regulated by PGE_2 activation of the ciliary G protein coupled

receptor (GPCR) E-type prostanoid receptor 4 (EP_4) (Jin et al., 2014; Marra et al., 2019). Active

EP_4 stimulates $G\alpha_s$ heterotrimeric G proteins, which raise intraciliary cAMP to promote

anterograde intraflagellar transport (IFT) for motile cilium length maintenance (Besschetnova et

al., 2010; Hansen et al., 2020; Hansen et al., 2022; Jin et al., 2014). Primary cilium length

control likely occurs through similar mechanisms because reduced ABCC4 activity in cultured

fibroblasts leads to primary cilium shortening and increasing ciliary cAMP leads to primary cilium

elongation (Besschetnova et al., 2009; Jin et al., 2014). This raises the possibility that ciliary

cAMP reduction in response to high-level activation of SHH signaling could slow IFT, shorten

primary cilia, and blunt the downstream signal response. The fail-safes that prevent ciliary

shortening in SHH-stimulated cells have not been described.

We previously reported that SHH activates cytosolic phospholipase A2α (cPLA$_2$) through SMO

activation of Gβγ to produce arachidonic acid. Arachidonic acid drives a feed-forward signal that

enhances SMO ciliary enrichment such that inhibition of cPLA$_2$ attenuates agonist-induced SMO

ciliary accumulation and blunts signaling to GLI transcription factors (Arensdorf et al., 2017).

Notably, arachidonic acid can be metabolized to generate the EP$_4$ agonist PGE$_2$. This suggests

that, in addition to enhancing SMO ciliary enrichment, SHH-mediated production of arachidonic

acid may connect the SHH pathway with PGE$_2$/EP$_4$-regulated primary cilium length control.

Herein, we investigate this hypothesis and reveal a crosstalk circuit between SHH and

prostaglandin signaling pathways. We report that SHH-stimulated production of arachidonic acid

drives ciliary length homeostasis through production of PGE$_2$ and activation of EP$_4$. This

provides a mechanism by which ciliary cAMP levels are stabilized in SHH-stimulated cells to

sustain anterograde IFT and prevent primary cilium shortening. We show that genetic ablation

or small molecule inhibition of SHH-to-EP$_4$ crosstalk destabilizes primary cilium cAMP

concentration, slows IFT, shortens primary cilia, reduces SMO ciliary enrichment, and

attenuates SMO signaling to GLI. Consistent with EP$_4$ signaling being crucial for regulated SHH

pathway activation *in vivo*, we find that mice lacking EP$_4$ have shortened primary cilia and

exhibit neural tube patterning defects indicative of compromised SHH signaling. As such, we

propose the SHH pathway evolved to link with prostaglandin signaling through cPLA$_2$ activation

to stabilize primary cilium length for optimal function of the organelle from which it instructs

downstream signaling.

RESULTS

cPLA$_2$ contributes to ciliary length control downstream of SHH.

To determine whether SHH activation of cPLA$_2$ leads to secretion of PGE$_2$ that regulates

primary cilium morphology, we turned to murine kidney inner medullary collecting duct (IMCD3)

cells. IMCD3 cells are SHH responsive, produce long primary cilia, and provide an established

model for examination of ciliary biology (Breslow et al., 2018; May et al., 2021; Ott and

Lippincott-Schwartz, 2012; Zhou et al., 2014). IMCD3 cells were stimulated with control

conditioned media or conditioned media containing the amino-terminal signaling domain of SHH

for ~24 hours, and then media was collected to test for accumulation of PGE$_2$. Enzyme-linked

immunosorbent assays (ELISA) of conditioned media from control and ligand-treated IMCD3

cells showed that SHH stimulated release of PGE$_2$ and that release was blocked by

pretreatment with the inverse SMO agonist cyclopamine. SHH-stimulated PGE$_2$ secretion was

also blocked by pre-treatment with the specific cPLA$_2$ inhibitor giripladib (GIRI) (Figure 1A)

(Arensdorf et al., 2017; Duvernay et al., 2015). To determine whether PGE$_2$ secretion occurred

in an ABCC4-dependent manner, we measured release from control or *Abcc4* knockout NIH-

3T3 cells following SHH stimulation. Similar to what was observed with IMCD3 cells, SHH

stimulation of control NIH-3T3 cells induced secretion of PGE$_2$ into cell culture medium. This

secretion was blocked by inhibiting SMO with cyclopamine or by inhibiting the cyclooxygenase

(COX) responsible for initiating arachidonic acid to PGE$_2$ conversion (Figure 1A', Coxib).

Consistent with ABCC4 secreting PGE$_2$ that is generated in response to SHH pathway

activation, *Abcc4* knockout NIH-3T3 cells failed to release prostaglandin in response to SHH

treatment (Figure 1A', light gray vs. dark gray). As such, arachidonic acid produced downstream

of SHH-stimulated cPLA$_2$ provides a substrate for PGE$_2$ production and secretion.

Having established that SHH-mediated $cPLA_2$ activation stimulates PGE_2 release from IMCD3

and NIH-3T3 cells, we next evaluated the effect of reducing $cPLA_2$-generated arachidonic acid

on ciliary length. We treated IMCD3 cells with vehicle or GIRI in the absence and presence of

SHH, and measured lengths of acetylated α-tubulin-marked primary cilia. We generated

maximum projections of Z stack images, and then traced the length of each acetylated α-tubulin

marked cilium from base to the tip using the line profile tool in LAS X software. GIRI treatment of

IMCD3 cells shortened primary cilia significantly, independent of SHH stimulation status.

Primary cilium length averaged ~5 μm in the absence of GIRI and shortened by approximately

40% to ~3 μm at the highest GIRI concentration used (Figure 1B-C). Ciliary lengths of GIRI-

treated cells were rescued by addition of supplemental PGE_2 or its precursor arachidonic acid

(Figure 1D-G). Notably, supplemental arachidonic acid augmented ciliary length in SHH-

stimulated cells in the absence of GIRI, further supporting a link between increased arachidonic

acid availability and primary cilium elongation (Figure 1G, column 2 vs. 3-4). Taken together,

these results support that ciliary shortening observed in GIRI-treated cells was a specific effect

of compromised arachidonic acid-to-PGE_2 metabolism.

To determine whether the ciliary morphology changes that occurred following $cPLA_2$ inhibition

correlated with reduced SMO ciliary accumulation in IMCD3 cells, we stimulated with SHH

conditioned media in the absence or presence of GIRI and quantified SMO ciliary signal

intensity (Supplementary Figure 1A-E'). Consistent with what we previously reported for NIH-

3T3 cells (Arensdorf et al., 2017), $cPLA_2$ inhibition disrupted SMO ciliary retention in SHH-

treated cells (Supplementary Figure 1A-E, red). SHH-stimulated ciliary exit of GPR161

(magenta) and retention of the primary cilium resident protein ARL13B (green) were unaffected

by GIRI treatment (Supplementary Figure 1A-D, E'), suggesting specificity in the ciliary functions

affected by $cPLA_2$ inhibition.

EP_4 activation promotes ciliary length homeostasis for SHH signaling to GLI.

Based on established connections between prostaglandin signaling and motile cilia elongation

(Jin et al., 2014; Marra et al., 2019), we hypothesized PGE_2 produced downstream of SHH

signals through ciliary EP_4 to influence primary cilium length. We therefore sought to determine

whether EP_4 localized to primary cilia of IMCD3 cells. EP_4 was evident in primary cilia of both

IMCD3 cells and control mouse embryonic fibroblasts (MEF) and was undetectable in cilia of

$Ep_4^{-/-}$ MEFs. These results support specificity of the EP_4 antibody signal and demonstrate that

EP_4 localizes to primary cilia in both MEFs and IMCD3 cells (Figure 2A-C').

To test for EP_4 contributions to primary cilium length maintenance, average ciliary lengths were

calculated in control and SHH-treated IMCD3 cells exposed to vehicle or the high-affinity EP_4

inhibitor, L161,982 (Cherukuri et al., 2007). Primary cilia of IMCD3 cells treated with increasing

concentrations of L161,982 shortened in a dose-dependent manner in the absence of SHH.

Notably, ciliary shortening of cells treated with low-dose L161,982 was significantly enhanced by

SHH treatment (Figure 2D), demonstrating that an active SHH pathway disrupts primary cilium

length control if EP_4 activity is compromised. Supplementing culture media of L161,982-treated

cells with high concentrations of the EP_4 ligand PGE_2 partially rescued ciliary length of SHH-

stimulated cells, supporting that primary cilium shortening following L161,982 treatment was a

specific effect of on-target inhibitor activity toward EP_4 (Figure 2E).

Having determined that EP$_4$ contributes to ciliary length maintenance in SHH-stimulated cells,

we next wanted to understand how EP$_4$ inhibition affected SHH signaling output. Thus, SHH-

stimulated SMO ciliary signal intensity was quantified in control, GIRI-treated, and L161,982-

treated IMCD3 cells. Similar to what we observed following cPLA$_2$ inhibition with GIRI, inhibition

of EP$_4$ with L161,982 attenuated ciliary accumulation of endogenous SMO following SHH

exposure (Figure 3A-B and Supplementary Figure 1). Ciliary shortening and reduced SMO

accumulation relative to control were also observed in $Ep4^{-/-}$ MEFs (Figure 3C-E), further

supporting on-target activity of L161,982 to EP$_4$. To determine whether reduced SMO ciliary

enrichment following EP$_4$ inhibition compromised activation of GLI transcriptional effectors,

qPCR analyses of SHH transcriptional targets *Gli1* and *Ptch1* were performed in IMCD3 and

NIH-3T3 cells. Inhibition of either cPLA$_2$ with GIRI or EP$_4$ with L161,982 significantly reduced

the ability of SHH to activate a transcriptional response (Figure 3F-F'). Thus, compromised

PGE$_2$/EP$_4$ signaling shortens cilia and attenuates SHH pathway activity.

EP$_4$ signaling stimulates ciliary cAMP recovery following SHH pathway activation.

We hypothesized that the ciliary length reduction and SHH signal attenuation observed following

EP$_4$ inhibition resulted from altered ciliary cAMP regulation (Figure 4A). To address this

hypothesis, we first tested whether reduced ciliary cAMP would lead to shortening of IMCD3

primary cilia by knocking down expression of ciliary adenylyl cyclases AC3, AC5, or AC6

(Supplementary Figure 2A-B). Consistent with these ACs localizing to primary cilia, their

knockdown did not prevent forskolin-stimulated accumulation of whole cell cAMP by cytoplasmic

ACs (Supplementary Figure 1C). To specifically measure IMCD3 cell ciliary cAMP following AC

knockdown, we used a primary cilium-localized fluorescent sensor system to track ciliary cAMP

changes in real time (Supplementary Figure 2D-F'). The sensor revealed that knockdown of

each of the ciliary ACs attenuated ciliary cAMP increase by the AC stimulating drug forskolin

(Figure 4B). Consistent with reduced ciliary cAMP leading to ciliary shortening, we observed a

significant decrease in average IMCD3 cell primary cilium length following knockdown of *Adcy3*,

Adcy5, or *Adcy6* (Figure 4C). Consistent with reduced primary cilium length compromising the

SHH signal response, *Gli1* induction was attenuated following ciliary AC knockdown (Figure

4D), suggesting that sustained depletion of ciliary cAMP can compromise ligand-stimulated

SHH pathway induction.

Despite the established link between ciliary length destabilization and reduced SHH pathway

activation (Caspary et al., 2007; He et al., 2014), our observation that decreased ciliary cAMP

correlated with reduced *Gli1* induction was curious. This is because reduced ciliary AC

localization following loss of the ciliary AC chaperone protein ANKMY2 has been shown to

induce GLI gain-of-function phenotypes *in vivo* (Somatilaka et al., 2020). To query this

discrepancy, we knocked down *Ankmy2* in IMCD3 cells to probe effects of combined ciliary AC

depletion on primary cilium cAMP concentration and length control. Consistent with published

results, *Ankmy2* knockdown reduced ciliary enrichment of endogenous AC3 and AC5/6

(Somatilaka et al., 2020) and blocked the ability of forskolin to trigger rapid ciliary cAMP

accumulation (Supplementary Figure 3A-D). To determine how ciliary AC depletion impacted

primary cilium length and SHH pathway induction, we measured primary cilia and performed

Gli1 qPCR analyses in control and *Ankmy2* knockdown cells treated with control or SHH

conditioned media. Notably, *Ankmy2* knockdown triggered a 23% reduction in IMCD3 cell

primary cilium length and attenuated SHH-stimulated *Gli1* induction (Figure 4E-F). These results

suggest the inability of cells to raise ciliary cAMP in SHH-stimulated cells compromises the

ligand-induced transcriptional response.

To further evaluate this hypothesis, we next probed effects of ciliary AC depletion on the SHH

response in highly SHH-responsive NIH-3T3 cells. Due to baseline primary cilia lengths in NIH-

3T3 cells being shorter than those of IMCD3 cell cilia, length reductions occurring in response to

ciliary AC depletion were difficult to appreciate. Nevertheless, we observed that knockdown of

Adcy5 or *Ankmy2* led to modest, but statistically significant, reduction in primary cilium lengths

in NIH-3T3 cells (Supplementary Figure 3E-F). *Adcy3* knockdown did not alter NIH-3T3 ciliary

length, suggesting AC5 may be the primary modulator of cAMP-mediated primary cilium length

regulation in this cell type (Supplementary Figure 3F). Accordingly, and consistent with

published results (Somatilaka et al., 2020), we observed that knockdown of *Ankmy2* significantly

reduced SHH-stimulated *Gli1* induction in NIH-3T3 cells (Supplementary Figure 3G). Consistent

with *Ankmy2* knockdown leading to a pan-AC ciliary depletion, individual knockdown of either

Adcy3 or *Adcy5* attenuated the SHH transcriptional response to a lesser extent than *Ankmy2*

knockdown (Supplementary Figure 3G). We did not detect endogenously expressed *Adcy6*

transcript in NIH-3T3 cells, so did not test knockdown of *Adcy6* in this cell type. These results,

combined with the results presented above, lead us to conclude that primary cilium AC

depletion is detrimental to the ligand-induced SHH signal response.

Having established that lowering ciliary cAMP by AC depletion appreciably and significantly

shortened primary cilia and attenuated SHH signaling in IMCD3 and NIH-3T3 cells, we next

sought to determine whether cAMP dynamics downstream of SMO activation were impacted by

EP_4. Thus, IMCD3 cells were stimulated with the AC activator forskolin, and ciliary cAMP was

monitored in control and SMO stimulated cells using the fluorescent cAMP system introduced

above. As anticipated, forskolin treatment stimulated a rapid increase in ciliary cAMP

fluorescence intensity ratios (Figure 5A, red and Supplementary Figure 2F-F', blue). Consistent

with the ability of activated SMO to couple with $G\alpha_i$ and induce exit of $G\alpha_s$-coupled GPR161

from primary cilia, co-stimulation of IMCD3 cells with forskolin and the direct SMO agonist SAG

reduced the level to which forskolin could raise ciliary cAMP. Despite this, ciliary cAMP in cells

treated with forskolin + vehicle and forskolin + SAG equilibrated similarly after an initial lag

(Figure 5A, red vs blue). We hypothesized cAMP equilibration was the result of EP_4 activation

downstream of SMO-stimulated PGE_2 release. Accordingly, ciliary cAMP equilibration in SAG-

stimulated cells was blocked by direct EP_4 inhibition with L161,982 (Figure 5A, green vs. blue).

We conclude EP_4 activation is required to equilibrate primary cilium cAMP levels in cells with an

active SHH signaling pathway.

Previous reports indicate that activation of EP_4 raises cAMP to promote anterograde IFT

(Besschetnova et al., 2010; Jin et al., 2014). To determine if this activity was compromised by

EP_4 inhibition in SHH-stimulated cells, we calculated anterograde velocity of the ciliary

intraflagellar transport protein IFT88 in control and L161,982-treated cells. We observed

increased anterograde IFT in vehicle-treated, SHH-stimulated cells (Figure 5B, lane 1 vs. 2).

Strikingly, anterograde IFT88 velocity was significantly reduced following treatment with the EP_4

inhibitor L161,982 in both control and SHH-stimulated IMCD3 cells (Figure 5B). Thus, EP_4

activity is required to maintain anterograde IFT in control cells and enhance its velocity in SHH-

stimulated cells.

If cAMP equilibration that stabilizes primary cilium length in SHH-stimulated cells occurs through

SHH-to-EP_4 crosstalk, we reasoned that providing supplemental ciliary cAMP might restore

ciliary length to rescue SHH pathway induction in L161,982-treated cells. To affect ciliary cAMP

in a controlled manner, we adapted a published optogenetic system (Truong et al., 2021) for

use in IMCD3 cells. In this system, bacterial photoactivatable adenylyl cyclase (bPAC) is

expressed in the cell cytoplasm or targeted to primary cilia through fusion with the primary

cilium-localized protein ARL13B, and then activated to specifically raise cytoplasmic or ciliary

cAMP by blue light stimulation (Supplementary Figure 4A). IMCD3 cells were engineered to

stably express cytoplasmic or ciliary bPAC-GFP, and then exposed to blue light to quantify

primary cilium length changes (Supplementary Figure 4B-C"). IMCD3 cells expressing

cytoplasmic or ciliary bPAC proteins showed similar average primary cilium lengths in the

absence of blue light and cilia of cyto-bPAC-expressing cells did not change length upon blue

light exposure, despite its ability to increase cytoplasmic cAMP levels (Supplementary Figure

4C-C'). Conversely, cilia-bPAC-expressing cells increased their average primary cilium length

by ~30% upon blue light exposure (Supplementary Figure 4C, C"). To support that elongation

resulted from increased primary cilium cAMP, we measured primary cilium lengths in IMCD3

cells treated with forskolin, which stimulates both ciliary and non-ciliary AC. We observed an

increase in average primary cilium length in forskolin-treated cells that was similar to that

observed following cilia-bPAC activation (Supplementary Figure 4D-D'). Thus, small molecule or

optogenetic increase in ciliary cAMP can increase primary cilium length.

Having validated feasibility of using ciliary bPAC to modulate primary cilium cAMP in IMCD3

cells, we next determined whether shortened cilia of EP_4-inhibited cells could be reversed by

increasing ciliary cAMP. Consistent with the hypothesis that reduced primary cilia lengths in

SHH-stimulated L161,982-treated cells resulted from failed cAMP equilibration, blue light

stimulation of cilia-bPAC partially rescued average primary cilium length in EP_4-inhibited, SHH-

stimulated cells. Blue light stimulation increased average ciliary lengths of L161-982-treated,

ciliary-bPAC-expressing cells by ~25% in the absence and presence of SHH (Figure 5C). To

test whether ciliary length increase correlated with rescue of SHH pathway activity, we

quantified SMO ciliary enrichment in control and blue light-stimulated, cilia-bPAC-expressing

cells after L161,982 treatment. Consistent with the ability of blue light-induced cAMP to partially

rescue ciliary length following EP_4 inhibition, SHH-stimulated SMO ciliary enrichment was also

partially restored in blue light-stimulated, L161,982-treated cilia-bPAC cells (Figure 5D, column

3 vs. 6). Intriguingly, SHH-stimulated enrichment of SMO in vehicle-treated cells was enhanced

by blue light stimulation, supporting that augmenting cAMP production in cells with an active

SHH pathway may enhance anterograde IFT to facilitate increased SMO ciliary enrichment

(Figure 5D, column 2 vs. 5). Thus, EP_4-regulated ciliary cAMP control impacts both primary

cilium length and SMO ciliary accumulation.

EP_4 is required for ciliary length control and SHH signaling during neural tube development.

Having established a cell biological connection between SHH and EP_4 that stabilizes primary

cilium length, we next sought to determine whether EP_4 loss compromised ciliary length control

to impact SHH signaling *in vivo*. Thus, we analyzed developing neural tubes of *$Ep4^{-/-}$* mice to

assess primary cilium length and evaluate SHH-regulated ventral neural tube cell fate

specification. *$Ep4^{-/-}$* mice are present at normal Mendelian ratios *in utero* but die as neonates

due to heart failure resulting from highly penetrant patent ductus arteriosus (Segi et al., 1998).

Thus, we collected E9.5/29 somite stage *$Ep4^{+/-}$* and *$Ep4^{-/-}$* embryos to perform scanning electron

microscopy for visualization of apical primary cilia of cells lining the neural tube lumen (Figure

6A). Measurement of apical primary cilia revealed statistically significant length reductions in

$Ep4^{-/-}$ embryos compared to cilia of heterozygous littermate controls (Figure 6B). Whereas

primary cilia in control neural tubes averaged ~0.9 µm, cilia in knockout embryos showed an

331 approximate 30% length reduction to an average length of ~0.6 µm. Thus, EP_4 contributes to

332 primary cilium length control in the developing neural tube.

334 During neural tube development, SHH signals in a graded manner from the notochord and

335 neural tube floor plate to instruct transcriptional programs that establish ventral neural tube cell

336 fates (Ericson et al., 1997; Patten and Placzek, 2000; Ribes and Briscoe, 2009). To test whether

337 EP_4 loss led to alteration of SHH-dependent neural tube progenitor domain specification, we

338 examined expression of the ventral SHH transcriptional target *Olig2,* which is activated by high

339 SHH, and the intermediate fate marker *Pax6* (Ericson et al., 1997; Ribes and Briscoe, 2009).

340 Because specification of ventral progenitor domains is highly sensitive to SHH signal alteration,

341 monitoring their induction allows for reliable reporting of SHH signaling activity (Ribes and

342 Briscoe, 2009). Comparison of E9.5/27 somite stage knockout and heterozygous littermate

343 control neural tubes revealed that EP_4 loss led to reduction in the *Olig2*-postive motor neuron

344 progenitor cell population, indicative of reduced SHH signaling response in *Ep4$^{-/-}$* mice.

345 Coincident with reduction in the *Olig2*-expressing progenitor domain, we observed expansion of

346 the *Pax6*-expressing progenitor population (Figure 6C-D). Taken together with the above, these

347 results support that EP_4 loss shortens primary cilia and compromises SHH-directed cell fate

348 specification *in vivo*.

350 **DISCUSSION**

351 Prostaglandin signaling is important for motile cilia development and beat frequency and

352 contributes to primary cilium elongation (Gayner and McCaffrey, 1998; Haxel et al., 2001; Jin et

353 al., 2014; Marra et al., 2019). The study presented here reveals the SHH pathway, which

354 requires functional primary cilia to initiate downstream effector responses, links with the PGE_2

pathway to ensure ciliary length homeostasis. We delineate a SHH-to-PGE$_2$/EP$_4$ signal

crosstalk circuit and demonstrate that SHH-stimulated PGE$_2$ production leads to activation of

Gα_s-coupled EP$_4$. EP$_4$ signaling equilibrates ciliary cAMP after an initial dip following SHH/SAG-

stimulated ciliary exit of Gα_s-coupled GPR161 and activation of Gα_i-coupled SMO. We propose

that cAMP equilibration maintains anterograde IFT that is necessary for SMO ciliary enrichment

and primary cilium length stability (Figure 7A-B). Genetic or pharmacological interruption of the

SHH-to-EP$_4$ signal crosstalk cascade short-circuits ciliary cAMP equilibration in SHH-stimulated

cells, slows anterograde IFT, shortens primary cilia, and attenuates SHH signal output (Figure

7C). *In vivo* loss of EP$_4$ leads to short neuroepithelial primary cilia and neural tube defects

indicative of compromised SHH signaling. These results support an essential link between SHH

and EP$_4$ ciliary signaling pathways during embryonic development.

Although the requirement for functional primary cilia in regulation of SHH signaling is well

established, the complex relationship between SHH and cAMP modulation in cilia has yet to be

clearly defined. To initiate a transcriptional response, SHH must lower ciliary cAMP to halt PKA-

instructed GLI processing into its repressor form (Mukhopadhyay et al., 2013; Niewiadomski et

al., 2014). Consistent with this biology, genetic loss of the AC chaperone *Ankmy2* depletes

primary cilia of AC activity, attenuates PKA-directed GLI repressor conversion, and allows for

unregulated accumulation of full length GLI proteins that induce gain-of-function phenotypes *in*

vivo. Although ciliary length reduction was not noted in *Ankmy2* knockout mice, mechanistic *in*

vitro analyses revealed that *Ankmy2* knockout NIH-3T3 cells were, nonetheless, compromised

in their ability to induce SHH-stimulated transcriptional responses (Somatilaka et al., 2020). This

suggests the ability to precisely tune ciliary cAMP levels up or down is essential for optimal

ligand-stimulated SHH pathway activity. Our data suggest SHH-promoted EP$_4$ signaling is a key

mechanism by which up-tuning occurs, and that this crosstalk is necessary to prevent slowing of

anterograde IFT and primary cilium shortening that can occur in response to sustained ciliary

cAMP depletion. However, the requirement for SHH to activate $G\alpha_s$-coupled EP_4 to stabilize

primary cilium length prompts the question of how GLI escapes PKA-mediated repressor

conversion upon ciliary cAMP re-elevation. We hypothesize GLI stabilization is maintained in

SHH-stimulated cells through direct PKA inhibition by active SMO. Recent reports demonstrate

that SHH-stimulated phosphorylation of the intracellular carboxyl-terminal tail of SMO exposes a

PKA inhibitory domain (PKI) that prevents GLI phosphorylation (Arveseth et al., 2021; Happ et

al., 2022). As such, we propose SHH coordinates a sophisticated ciliary signal response

through a network of interconnected GPCR activities.

Notably, GPCR signal crosstalk downstream of Hedgehog (Hh) family ligands that equilibrates

cAMP may be a conserved mechanism by which pathway activity is controlled. In *Drosophila*,

Hh-activated Smo stimulates $G\alpha_i$ to lower cAMP and halt PKA-regulated truncation of the GLI

ortholog Cubitus interruptus (Ci) (Ogden et al., 2008). However, a *Drosophila*-specific positive

regulatory role for PKA-mediated phosphorylation of Smo is required for high-level pathway

induction, and could be disrupted by Smo activation of $G\alpha_i$ (Jia et al., 2004). A potential

mechanism by which Smo solves this problem was revealed by the discovery that, in some cell

types, Smo stimulation leads to $G\alpha_s$ activation to enhance pathway responsiveness to low-

threshold Hh stimulation (Praktiknjo et al., 2018). It is not yet clear whether $G\alpha_s$ activation

downstream of Hh occurs through Smo signal bias or by Smo crosstalk with $G\alpha_s$-coupled GPCR

signaling partners. Nevertheless, the observation that Smo can lead to $G\alpha_s$ activation in flies

supports an evolutionarily conserved ability of Hedgehog family members to affect exquisite

control of cAMP modulators to ensure appropriate downstream signaling activity.

Table 1: Reagents and Resources

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Smoothened	Santa Cruz Biotechnology	sc-166685, RRID:AB2239686
Acetylated α-Tubulin	Cell Signaling	5335S, RRID:AB_0544694
ARL13B	Antibodies Incorporated	75-287, RRID:AB_2341543
EP4	Santa Cruz Biotechnology	sc-55596, RRID:AB_2174775
AC3	Proteintech	19492-1-AP RRID:AB_10638445
AC5/6	FabGennix	PAC-501AP
PAX6	Developmental Studies Hybridoma Bank	RRID:AB_528427
OLIG2	Millipore	AB9610, RRID:AB_570666
GPR161	Saikat Mukhopadhyay	N/A
AlexaFluor 488	Life Technologies	A11029 (Mouse), RRID:AB_2534088; A11034 (Rabbit), RRID:AB_2576217
AlexaFluor 555	Life Technologies	A21424 (Mouse), RRID:AB_141780; A21429 (Rabbit), RRID:AB_2535850
AlexaFluor 647	Life Technologies	A21236 (Mouse), RRID:AB_2535805; A21245 (Rabbit), RRID:AB_2535813
Chemicals, Peptides, and Recombinant Proteins		
Arachidonic Acid	Cayman Chemical Company	90010-50

Giripladib	This Paper	N/A
L161,982	Tocris Chemicals	2514
Prostaglandin E2	Sigma Chemicals	P5640
Cyclopamine	LC Laboratories	C-8700
Celecoxib	Cayman Chemical Company	10008672
Forskolin	Sigma Chemicals	F6886-10MG
SAG	Selleck Chemical Company	S7779-2MG
Lipofectamine™ 3000	Thermo Fisher Scientific	L3000008
Lipofectamine™ RNAiMAX	Thermo Fisher Scientific	13778150
IBMX	Sigma-Aldrich	I5879-1G
DMSO	Sigma-Aldrich	D2650
Hygromycin	Life Technologies	10687010
PowerUp SYBR Green Master Mix	Applied Biosystems	A25742
Commercial Assay Kits		
Prostaglandin E2 Parameter Assay Kit	R&D Systems	KGE004B
BCA Protein Assay Kit	Thermo Scientific™ Pierce™	23227
RNeasy Plus Mini Kit	Qiagen	74136
High-Capacity cDNA Reverse Transcription Kit	Applied Biosystems	4368813
Ratio-metric Cilia-Targeted cADDis cAMP	Montana Molecular	D0211G
Direct cAMP ELISA kit	Enzo Life Sciences	ADI900066
Oligonucleotides		
Ptch1 Forward: TCACCTCCATCAGCAATGTC	Thermo Fisher Scientific	N/A
Ptch1 Reverse: GAATACCACCACCACAGCAG	Thermo Fisher Scientific	N/A

Gli1 Forward: CCAAGCCAACTTTATGTCAGGG	Thermo Fisher Scientific	N/A
Gli1 Reverse: AGCCCGCTTCTTTGTTAATTTGA	Thermo Fisher Scientific	N/A
Adcy3 Forward: GCTTTCTTTGTCTTCTCCTTC	Thermo Fisher Scientific	N/A
Adcy3 Reverse: ATGATAGCACACAGGTAGAG	Thermo Fisher Scientific	N/A
Adcy5 Forward: GTCTCAGAAAGTGGCAATTC	Thermo Fisher Scientific	N/A
Adcy5 Reverse: CATTCAGGTAGTTGAGTGTG	Thermo Fisher Scientific	N/A
Anmky2 Forward: GCTGCCAAAACATAAGAGACAG	Thermo Fisher Scientific	N/A
Anmky2 Reverse: CTTCTTTCTCCCTTCCACAG	Thermo Fisher Scientific	N/A
Adcy6 Forward: ACGTTGCCATGGAGATGAAA	Thermo Fisher Scientific	N/A
Adcy6 Reverse: TCAATGTCCGCAAACAGGAT	Thermo Fisher Scientific	N/A
Ppia Forward: AGCACTGGAGAGAAAGGATT	Thermo Fisher Scientific	N/A
Ppia Reverse: ATTATGGCGTGTAAAGTCACCA	Thermo Fisher Scientific	N/A
Btf3 Forward: GAACAACATCTCTGGTATTGAAGA	Thermo Fisher Scientific	N/A
Btf3 Reverse: AATGGTGAAGGTGTTTGCTG	Thermo Fisher Scientific	N/A
Adcy3 siRNA	Thermo Fisher Scientific	Assay ID: S98223
Adcy5 siRNA	Thermo Fisher Scientific	Assay ID: S104765

Adcy6 siRNA	Thermo Fisher Scientific	Assay ID: s104033
Anmky2 siRNA	Thermo Fisher Scientific	Assay ID: MSS235827
Experimental Models: Cell Lines		
IMCD3	ATCC	CRL-2123
IMCD3 Flp-In Cells	Maxence Nachury Lab	N/A
IMCD3 EF1α-Cyto-bPAC-GFP	This Paper	N/A
IMCD3 EF1α-Arl13b-Cilia-bPAC-GFP	This Paper	N/A
NIH-3T3	ATCC	CRL-1658
Abcc4$^{-/-}$ NIH-3T3	John Schuetz lab	N/A
Ep4$^{-/-}$ Mouse Embryonic Fibroblast	This paper	N/A
Wild Type Mouse Embryonic Fibroblast	This paper	N/A
Experimental Models: Mouse Model		
Ep4$^{+/-}$ Mice	Elaine Shelton	(Segi et al., 1998)
Ep4$^{-/-}$ Mice	Elaine Shelton	(Segi et al., 1998)
Recombinant DNA		
pgLAP5-EF1α dTATA-bPAC-GFP	Jeremy Reiter	(Truong et al., 2021)
pgLAP5-EF1α dTATA-Arl13b-bPAC-GFP	Jeremy Reiter	(Truong et al., 2021)
pEF1α-mIFT88-EGFP	Addgene	149697
5HT6-mCherry-G-GECO1.0	Addgene	47500
pcDNA-DEST47-Arl13b-GFP	Addgene	40872
Software and Algorithms		
GraphPad Prism v. 8.2.0	GraphPad Software	http://www.graphgad.com/

ImageJ	NIH software	https://imagej.nih.gov/ij/
LAS X	Leica	
JACoP	Open Source	https://imagej.nih.gov/ij/plugins/track/jacop2.html
KymographDirect v. 2.0	Open Source	https://sites.google.com/site/kymographanalysis/
NIS-Elements Viewer v. 4.50.00	Nikon	https://www.microscope.healthcare.nikon.com/products/software/nis-elements/viewer
Inkscape	Open Source	https://inkscape.org/

RESOURCE AVAILABILITY

Requests for resources and reagents should be directed to Stacey K. Ogden

(Stacey.Ogden@stjude.org). Cell lines and novel reagents will be available upon request

pursuant to SJCRH material transfer agreement.

METHODS

Mammalian Cell culture

Wild type and Flp-In IMCD3 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM):

F12 supplemented with 10% fetal bovine serum (FBS) and 1% Penicillin Streptomycin solution

(Gibco). Wild type MEFs, NIH-3T3, $Ep4^{-/-}$ MEF, and $Abcc4^{-/-}$-NIH-3T3, were supplemented with

10% fetal calf serum. Cells were passaged routinely with 0.25% Trypsin/EDTA solution and

maintained under standard incubation conditions (5% CO_2, 37°C) in a humidified incubator. All

cell lines were routinely tested for mycoplasma contamination. To promote ciliogenesis, cell

lines were serum starved for 2 hours in serum-free media (DMEM or DMEM: F12,

supplemented with 0.1 mM nonessential amino acids, 2mM L-Glutamine, 1mM sodium

pyruvate, and 1% Pen-Strep). After starvation, media was replaced with low serum (0.5%)

media and cells were incubated for 18 or 36 hours.

Chemicals

L161,982 (Tocris), Cyclopamine (LC Laboratories) and Celecoxib (Coxib, Cayman Chemical

Company) were reconstituted in DMSO and stored at -20°C. Arachidonic acid (Cayman

Chemical Company) was stored in ethanol at -20°C. Prostaglandin E_2 (Sigma Chemicals) and

Forskolin (Sigma Chemicals) were reconstituted in ethanol and stored at 4°C. Giripladib (GIRI)

was synthesized and stored according to published procedures (Duvernay et al., 2015; Thotala

et al., 2013).

Immunofluorescence and ciliary length measurement

For cultured cells ciliary analysis, cells were plated onto coverslips (Corning) and pretreated

with either vehicle or drug in serum-free media. After 2 hours, media was changed to low serum

media containing drug and SHH-N conditioned media (100 µL/mL), and then incubated for ~18

hours. After incubation, cells were washed with PBS, and then fixed in 4% paraformaldehyde for

12 minutes. Cells were then washed with wash buffer (PBS with 0.1% Triton X-100) three times

for 5 minutes and incubated in blocking buffer for 60 minutes at room temperature (PBS with 2%

BSA, 0.1% Triton X-100, 1% goat serum). Primary antibody incubations were performed

overnight at 4°C. The following antibodies and dilutions were used: anti-SMO (Santa Cruz,

1:500), anti-acetylated α-tubulin (Cell Signaling, 1:1000), anti-ARL13B (Antibodies Incorporated,

1:500), anti-EP$_4$ (Santa Cruz, 1:100), anti-AC5/6 (FabGennix, 1:75), anti-AC3 (Proteintech,

1:200) diluted in blocking buffer. Secondary antibody incubations were performed with

AlexaFluor 488, 555, or 647 conjugated secondary antibodies (Life Technologies, 1:1000) and

DAPI for 60 minutes at room temperature. Following antibody incubation, samples were washed

three times in wash buffer followed by a rinse in PBS, prior to mounting with ProLong Diamond

(Life Technologies). Images were collected using Leica TCS SP8 STED 3x confocal microscope

with a 63x oil-immersion objective and processed using LAS X (Leica). For all

immunofluorescence experiments, multiple cells (≥100) were examined over at least two

independent experiments. Representative images are shown.

For ciliary length measurement, a maximum projection of the acquired Z stack images was

created. Each cilium was manually traced from the base to the tip using the line profile tool in

the LAS X software. For all experiments, >100 cilia were measured for each condition across a

minimum of two independent experiments. For quantification graphs, all data are included. For

microscopy images, representative cilia are shown.

In vivo analysis

Data and materials generated from animals were obtained in accordance with SJCRH/IACUC

approved protocol 608-100616-10/19 and Vanderbilt protocol M1600016-01. All animal

husbandry and procedures were performed in accordance with protocols approved by St. Jude

Children's Research Hospital and Vanderbilt University Institutional Animal Care and Use

Committees.

$Ep4^{+/-}$ and $Ep4^{-/-}$ embryos in the C57BL/6 background were harvested and processed for

immunohistochemistry at E9.5. Pregnant dams were harvested, uterine horns removed, and

embryos were dissected in 1X PBS, then rinsed three times in 1X PBS. Embryos were fixed

overnight at 4°C in 4% PFA. The following day, embryos were rinsed three times in 1X PBS and

transferred to 30% sucrose to cryo-protect. The following day, embryos were frozen in O.C.T.

Compound (Tissue-Tek) on dry ice. Embryos were sectioned transverse at 15 µm thickness on

a Leica Microm CM1950 cryo-stat. Sections were briefly dried, then washed in 1X TBST, then

blocked with 2% BSA, 1% goat serum, 0.1% Triton-X-100 in 1X PBS. Antibodies were diluted in

blocking buffer and incubated overnight on sections at room temperature. The following

antibodies and dilutions were used: mouse anti-PAX6 (1:25; DSHB), and rabbit anti-OLIG2

(1:300; Millipore). Primary antibody was removed, sections were washed with 1X TBST three

times, then incubated for 3 hours in secondary antibodies (Invitrogen) at a 1:500 dilution.

Sections were washed three times in 1X TBST, then rinsed with water, dried, and cover slips

were applied with ProLong Gold mounting media with DAPI. Sections were imaged on a Leica

DMi8 widefield microscope. A minimum of four embryos per genotype were analyzed.

To quantify the *Olig2* and *Pax6* expressing domains, at least three sections per embryo in the

thoracic region were randomly chosen, and their area was measured using the Segmented Line

Tool in ImageJ. The percentage area of each domain was calculated with respect to the entire

Dorsal Ventral extent of the neural tube. The data are presented as mean ± SD, and statistical

comparisons were made using Student's t-test. GraphPad software was used for statistical

analysis. p-values less than 0.05 were considered statistically significant.

For electron microscopy of neural tube primary cilia, samples were fixed in 2.5% glutaraldehyde,

2% paraformaldehyde in 0.1M cacodylate buffer containing 2 mM $MgCl_2$. Following fixation,

samples were dissected to expose the interior of the neural tube lumen. Dissected samples

were buffer washed, post-fixed in 0.1% aqueous osmium tetroxide for 1 hour, washed with

ddH_2O, dehydrated with an ascending ethanol series, and critical point dried in liquid CO_2 using

an Autosamdri 931 (Tousimis). Dried samples were sputter-coated under planetary rotation with

15 nm iridium and imaged in a ThermoFisher Scientific Teneo scanning electron microscope at

2 kV using the Everhart-Thornley and T1 detectors.

cAMP measurement

cAMP in whole cell lysates was measured using the Direct cAMP ELISA kit (Enzo). Cells were

seeded in 6-well plates at a density of 3.5×10^5 cells per well. The next day, cells were starved

for 2 hours in serum free media followed by overnight incubation in low serum media to promote

ciliation. On the day of treatment, cells were treated with 100 µM IBMX diluted in low serum

media and scraped into lysis buffer (0.1 M HCl, Enzo). Samples were analyzed per the

manufacturer's protocol. cAMP concentrations were calculated using a 4-parameter logistic

(4PL) and cAMP concentrations were normalized to the total protein concentrations, determined

by a BCA protein assay (Thermo Fisher Scientific).

Ciliary cAMP assay

IMCD3 cells were seeded at 18,000 cells/well in an 8-well chamber slide (Nunc™ Lab-Tek™ II

Chamber Slide™ System) and transduced the next day with the ratiometric cilia-targeted

cADDis BacMam (Montana Molecular, D0211G) per manufacturer's recommendation. Cells

were infected with 20 µl of BacMam sensor stock in a total of 250 µl of media containing 2 mM

Sodium Butyrate (Molecular Montana) for 30 minutes at room temperature in dark followed by 5

hours at 37°C. BacMam was removed and replaced with low serum media containing 1 mM

Sodium Butyrate for 16-24 hours with or without vehicle, GIRI, or L-161,982 treatment. Prior to

imaging, cells were incubated in PBS for 20 minutes at room temperature. Positive agonist FSK

(100 µM) was added 30 seconds after the recording was initiated and images were acquired on

a Nikon A1R confocal on live cell imaging mode (63x, epifluorescence) every 10 seconds for 8

minutes. For experiments with SAG, the signal was recorded for basal conditions followed by

incubation with forskolin for 1.5 minutes and then treated with either SAG or vehicle and

recorded every 10 seconds for 8 minutes.

For knockdown experiments, IMCD3 cells were seeded in 8-well chamber slides (Nunc™ Lab-

Tek™ II Chamber Slide™ System) followed by siRNA treatment the next day with either the

control siRNA or *Adcy3, Adcy5*, *Adcy6* or *Ankmy2* siRNA. Cells were then transduced the

following day with the ratiometric cilia-targeted cADDis BacMam and the experiment was carried

out as described above.

Generation of optogenetic bPAC cells

IMCD3 cell lines stably expressing Cyto-bPAC-GFP and Cilia-bPAC-GFP were generated using

the Flp-In system following manufacturer's instructions (Invitrogen). IMCD3 Flp-In cells (Breslow

and Nachury, 2015) were transfected using Lipofectamine 3000 with the appropriate plasmids

and selected with 40 mg/mL hygromycin (Gibco). Single colonies were expanded, and protein

expression was confirmed by fluorescence microscopy for GFP.

Optogenetic stimulation

bPAC-expressing IMCD3 Flp-In cells were plated onto coverslips, grown to confluency, and

pretreated with either vehicle or L161-982 (10 μM) in serum-free media. After 2 hours, media

was changed to low serum media containing vehicle or drug and SHH-N conditioned media

(100 μL/mL), and then incubated for ~18 hours. Prior to immunofluorescence analysis, cells

were transferred to a custom-made LED humidified incubator with 5% CO_2 for continuous blue

light 450 nm stimulation at ~0.4 mW/cm^2 for 3 hours (Zhang et al., 2019). Following this,

immunofluorescence staining was performed as described above.

Quantitative Reverse Transcriptase Polymerase Chain Reaction (qRT-PCR)

Total RNA was extracted from cells using the RNeasy Mini Kit (Qiagen) according to

manufacturer's protocol. One thousand nanograms of RNA was used to synthesize

complementary DNA (cDNA) using High-Capacity cDNA Reverse Transcription Kit (Applied

Biosystems). qRT-PCR reactions were performed on a QuantStudio 7 Flex PCR machine using

PowerUp Sybr Green Master Mix (Applied Biosystems). Corresponding changes in the

expression levels of selected genes were calculated using the ΔΔCt method relative to

housekeeping genes, *Ppia* and *Btf3* (Arensdorf et al., 2017).

Cell Line Generation

EP$_4$ knockout cells were generated from *Ep4$^{-/-}$* embryos (Segi et al., 1998). Embryos (13.5 dpc)

were freshly isolated and placed in individual 10 cm culture dishes where they were eviscerated.

Heads were removed and used for genotyping to identify homozygous null embryos and wild

type littermates using conventional PCR. Embryo bodies were transferred to 15 mL conical

tubes containing 5 mL fresh PBS and dissociated by aspirating through a 16G needle into a 10

mL syringe and expelling the contents. Warm media (DMEM+10%FBS+1%pen/strep) was then

added to each conical tube to bring the total volume to 10 mL. This solution was transferred to a

15 cm culture dish containing 25 mL additional media and placed in a 37°C incubator. After 24h,

media and non-adherent tissue pieces were removed, and fresh media was added. Cells were

grown to 80-90% confluence, after which they were passaged and frozen down after passage

number two. *Abcc4$^{-/-}$* NIH-3T3 cells were generated as previously described (Wijaya et al.,

2020).

PGE$_2$ ELISA

IMCD3 cells were cultured in a 12-well culture plate and were pre-treated for 2 hours with either

Cyclopamine (10 µM) or Giripladib (GIRI, 4 µM) in serum free medium. Following pretreatment,

medium was changed to low serum containing either vehicle, Cyclopamine, or GIRI in control or

SHH-N conditioned media, and then incubated for 36 hours. PGE_2 levels were determined using

by ELISA (R&D Systems) according to the manufacturer's instructions. The absorbance of each

well was measured at 450 nm with correction at 540 nm.

siRNA Transfection

Transfections using Lipofectamine RNAiMAX were performed following the manufacturer's

protocols. Briefly, IMCD3 or NIH-3T3 cells were plated in a 6-well tissue culture plate at a

density of 175,000 or 100,000 cells per well. The following day, cells were treated with 25 pmol

of either the scrambled siRNA or *Adcy3, Adcy5*, *Adcy6* or *Ankmy2* siRNA. Cells were analyzed

48 hours post transfection.

IFT88 Velocity

To analyze intraciliary trafficking, Kymographs for anterograde IFT88 velocities were generated

in IMCD3 cells as previously described (Besschetnova et al., 2009). Briefly, 120,000 cells were

grown on a coverslip in a 24 well plate. Cells were transfected with IFT88-GFP and 5HT6-

mCherry plasmids the following day. Cells were pretreated with the EP4 inhibitor L161,982 for 2

hours and then switched to low serum media containing L161,982 either in the presence or

absence of SHH-N conditioned media and incubated overnight. On the day of the experiment,

the coverslips were flipped onto a 2-well chamber slide (Nunc™ Lab-Tek™ II Chamber Slide™

System) and the cells were imaged every 200 ms for 30 s on a Nikon A1R confocal microscope

with a 60x oil objective and 37 °C, 5% CO_2 heat stage (Live Cell Instrument). The resulting

videos were analyzed by ImageJ macros KymographClear and KymographDirect to generate

violin plots (Mangeol et al., 2016).

QUANTIFICATION AND STATISTICAL ANALYSIS

Quantification of ciliary lengths and SMO ciliary accumulation values were calculated from three

independent experiments with at least 100 cilia per condition per experiment. A maximum

fluorescence intensity projection from a sum of at least 10 slices was generated before length

and ciliary intensity quantifications. SMO ciliary intensity was normalized to acetylated α-tubulin

intensity. At least three randomly chosen fields of view were selected per condition per

experiment, and all cilia within the field were quantified using LAS X. For ciliary SMO signal

intensity determination, each cilium surface area was traced from the base to the tip using

spline profile function. The average fluorescence measurement adjacent to the cilium was

subtracted from ciliary fluorescence as background. The intensity values along the cilium were

exported and analyzed via GraphPad Prism. Colocalization of AC3 and AC6 within the primary

cilium marker ARL13B was carried out using the ImageJ plugin JACoP. Manders' coefficients

(M1 and M2) estimate the co-occurrence fraction of a fluorescent signal on one channel

(ARL13B) with a fluorescent signal of another channel (AC3/AC5/6). Manders' coefficients

range from 0 to 1. The former corresponds to non-overlapping images and the latter reflects a

complete co-localization between both images.

For quantification of cAMP, the red fluorescence and green fluorescence intensities over time

were processed on the Nikon NIS-Elements imaging software by defining a mask for each

cilium and determining the ratio of green fluorescence to red fluorescence. The data were

normalized to the initial time point for each experiment (set to 100%), and the inverse of each

value was plotted using GraphPad Prism.

Statistical analyses were performed using Prism 8 (GraphPad Software). Results are presented

as mean ± standard deviation. For comparisons between two groups, a two tailed unpaired t-

test was used unless otherwise specified. For multiple group comparisons, either one-way or

two-way (depending on the number of variables) ANOVA followed by multiple comparison post-

hoc testing was performed as indicated using Prism 8. Significance is depicted as $*p < 0.05$, $**p$

< 0.01, $***p < 0.001$, $****p < 0.0001$, ns = not significant.

ACKNOWLEDGEMENTS

We thank M. Hatley and members of the Ogden lab for reagents, advice, thoughtful discussion,

and comments on the manuscript. We thank S. Narumiya (Kyoto University) and the Center for

Animal Resources and Development (Kumamoto University) for providing $Ptger4^{-/-}$ frozen sperm

used to generate the $Ep4^{-/-}$ mouse line. J. Reiter (UCSF) provided bPAC expression vectors, S.

Mukhopadhyay (UTSW) provided anti-GPR161, and M. Nachury (UCSF) provided Flp-In IMCD3

cells. We thank J. Temirov, J. Messing, A. Carisey, and the St. Jude CTIC for assistance with

confocal and electron microscopy. Diagrams were created with BioRender.com.

FUNDING

This work was supported by National Institute of Health grants R35GM122546 (S.K.O.) and

R01HD099777 (E.L.S.) and by ALSAC of St. Jude Children's Research Hospital. The CTIC is

supported by SJCRH and NCI P30 CA021765. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the funding agencies.

AUTHOR CONTRIBUTIONS

Conceptualization, S.S.A. and S.K.O.; Methodology, S.S.A., M.A.A., B.M.Y., M.E.D., Y.Z.,

E.L.S., A.J., C.G.R., and S.K.O.; Validation, S.S.A., M.A.A., M.E.D., and Y.Z.; Analysis, S.S.A.,

M.A.A., M.E.D., Y.Z., and A.J.; Investigation, S.S.A., M.A.A., M.E.D., and Y.Z.; Resources,

J.D.S., Z.R., C.G.R., N.B., E.L.S., and S.K.O.; Writing – Original Draft, S.S.A. and S.K.O.;

Writing – Review & Editing, S.S.A., M.E.D., J.D.S., E.L.S., and S.K.O.; Visualization, S.S.A.;

Supervision, S.K.O. All authors approved the submitted manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Anvarian, Z., K. Mykytyn, S. Mukhopadhyay, L.B. Pedersen, and S.T. Christensen. 2019. Cellular signalling by primary cilia in development, organ function and disease. *Nat Rev Nephrol*. 15:199-219.

Arensdorf, A.M., M.E. Dillard, J.M. Menke, M.W. Frank, C.O. Rock, and S.K. Ogden. 2017. Sonic Hedgehog Activates Phospholipase A2 to Enhance Smoothened Ciliary Translocation. *Cell Rep*. 19:2074-2087.

Arensdorf, A.M., S. Marada, and S.K. Ogden. 2016. Smoothened Regulation: A Tale of Two Signals. *Trends Pharmacol Sci*. 37:62-72.

Arveseth, C.D., J.T. Happ, D.S. Hedeen, J.F. Zhu, J.L. Capener, D. Klatt Shaw, I. Deshpande, J. Liang, J. Xu, S.L. Stubben, I.B. Nelson, M.F. Walker, K. Kawakami, A. Inoue, N.J. Krogan, D.J. Grunwald, R. Huttenhain, A. Manglik, and B.R. Myers. 2021. Smoothened transduces Hedgehog signals via activity-dependent sequestration of PKA catalytic subunits. *PLoS Biol*. 19:e3001191.

Bangs, F., and K.V. Anderson. 2017. Primary Cilia and Mammalian Hedgehog Signaling. *Cold Spring Harb Perspect Biol*. 9.

Barakat, M.T., E.W. Humke, and M.P. Scott. 2010. Learning from Jekyll to control Hyde: Hedgehog signaling in development and cancer. *Trends Mol Med*. 16:337-348.

Barzi, M., D. Kostrz, A. Menendez, and S. Pons. 2011. Sonic Hedgehog-induced proliferation requires specific Galpha inhibitory proteins. *J Biol Chem*. 286:8067-8074.

Besschetnova, T.Y., E. Kolpakova-Hart, Y. Guan, J. Zhou, B.R. Olsen, and J.V. Shah. 2010. Identification of signaling pathways regulating primary cilium length and flow-mediated adaptation. *Curr Biol*. 20:182-187.

Besschetnova, T.Y., B. Roy, and J.V. Shah. 2009. Imaging intraflagellar transport in mammalian primary cilia. *Methods Cell Biol*. 93:331-346.

Breslow, D.K., S. Hoogendoorn, A.R. Kopp, D.W. Morgens, B.K. Vu, M.C. Kennedy, K. Han, A. Li, G.T. Hess, M.C. Bassik, J.K. Chen, and M.V. Nachury. 2018. A CRISPR-based screen for Hedgehog signaling provides insights into ciliary function and ciliopathies. *Nat Genet*. 50:460-471.

Breslow, D.K., and M.V. Nachury. 2015. Analysis of soluble protein entry into primary cilia using semipermeabilized cells. *Methods Cell Biol*. 127:203-221.

Caspary, T., C.E. Larkins, and K.V. Anderson. 2007. The graded response to Sonic Hedgehog depends on cilia architecture. *Dev Cell*. 12:767-778.

Cherukuri, D.P., X.B. Chen, A.C. Goulet, R.N. Young, Y. Han, R.L. Heimark, J.W. Regan, E. Meuillet, and M.A. Nelson. 2007. The EP4 receptor antagonist, L-161,982, blocks prostaglandin E2-induced signal transduction and cell proliferation in HCA-7 colon cancer cells. *Exp Cell Res*. 313:2969-2979.

Duvernay, M.T., A. Matafonov, C.W. Lindsley, and H.E. Hamm. 2015. Platelet Lipidomic Profiling: Novel Insight into Cytosolic Phospholipase A2alpha Activity and Its Role in Human Platelet Activation. *Biochemistry*. 54:5578-5588.

Ericson, J., P. Rashbass, A. Schedl, S. Brenner-Morton, A. Kawakami, V. van Heyningen, T.M. Jessell, and J. Briscoe. 1997. Pax6 controls progenitor cell identity and neuronal fate in response to graded Shh signaling. *Cell*. 90:169-180.

Findakly, S., V. Daggubati, G. Garcia, S.A. LaStella, A. Choudhury, C. Tran, A. Li, P. Tong, J.Q. Garcia, N. Puri, J.F. Reiter, L. Xu, and D.R. Raleigh. 2021. Sterol and oxysterol synthases near the ciliary base activate the Hedgehog pathway. *J Cell Biol*. 220.

Garcia, G., 3rd, D.R. Raleigh, and J.F. Reiter. 2018. How the Ciliary Membrane Is Organized Inside-Out to Communicate Outside-In. *Curr Biol*. 28:R421-R434.

Gayner, S.M., and T.V. McCaffrey. 1998. Muscarinic ciliostimulation requires endogenous prostaglandin production. *Am J Rhinol*. 12:203-207.

Hansen, J.N., F. Kaiser, C. Klausen, B. Stuven, R. Chong, W. Bonigk, D.U. Mick, A. Moglich, N.
Jurisch-Yaksi, F.I. Schmidt, and D. Wachten. 2020. Nanobody-directed targeting of
optogenetic tools to study signaling in the primary cilium. *Elife*. 9.
Hansen, J.N., F. Kaiser, P. Leyendecker, B. Stuven, J.H. Krause, F. Derakhshandeh, J. Irfan,
T.J. Sroka, K.M. Preval, P.B. Desai, M. Kraut, H. Theis, A.D. Drews, E. De-Domenico, K.
Handler, G.J. Pazour, D.J.P. Henderson, D.U. Mick, and D. Wachten. 2022. A cAMP
signalosome in primary cilia drives gene expression and kidney cyst formation. *EMBO*
Rep. 23:e54315.
Happ, J.T., C.D. Arveseth, J. Bruystens, D. Bertinetti, I.B. Nelson, C. Olivieri, J. Zhang, D.S.
Hedeen, J.F. Zhu, J.L. Capener, J.W. Brockel, L. Vu, C.C. King, V.L. Ruiz-Perez, X. Ge,
G. Veglia, F.W. Herberg, S.S. Taylor, and B.R. Myers. 2022. A PKA inhibitor motif within
SMOOTHENED controls Hedgehog signal transduction. *Nat Struct Mol Biol*. 29:990-999.
Haxel, B.R., D. Schafer, L. Klimek, and W.J. Mann. 2001. Prostaglandin E2 activates the ciliary
beat frequency of cultured human nasal mucosa via the second messenger cyclic
adenosine monophosphate. *Eur Arch Otorhinolaryngol*. 258:230-235.
Haycraft, C.J., B. Banizs, Y. Aydin-Son, Q. Zhang, E.J. Michaud, and B.K. Yoder. 2005. Gli2
and Gli3 localize to cilia and require the intraflagellar transport protein polaris for
processing and function. *PLoS Genet*. 1:e53.
He, M., R. Subramanian, F. Bangs, T. Omelchenko, K.F. Liem, Jr., T.M. Kapoor, and K.V.
Anderson. 2014. The kinesin-4 protein Kif7 regulates mammalian Hedgehog signalling
by organizing the cilium tip compartment. *Nat Cell Biol*. 16:663-672.
Huangfu, D., and K.V. Anderson. 2006. Signaling from Smo to Ci/Gli: conservation and
divergence of Hedgehog pathways from Drosophila to vertebrates. *Development*. 133:3-
14.
Jia, J., C. Tong, B. Wang, L. Luo, and J. Jiang. 2004. Hedgehog signalling activity of
Smoothened requires phosphorylation by protein kinase A and casein kinase I. *Nature*.
432:1045-1050.
Jin, D., T.T. Ni, J. Sun, H. Wan, J.D. Amack, G. Yu, J. Fleming, C. Chiang, W. Li, A. Papierniak,
S. Cheepala, G. Conseil, S.P. Cole, B. Zhou, I.A. Drummond, J.D. Schuetz, J. Malicki,
and T.P. Zhong. 2014. Prostaglandin signalling regulates ciliogenesis by modulating
intraflagellar transport. *Nat Cell Biol*. 16:841-851.
Kinnebrew, M., E.J. Iverson, B.B. Patel, G.V. Pusapati, J.H. Kong, K.A. Johnson, G. Luchetti,
K.M. Eckert, J.G. McDonald, D.F. Covey, C. Siebold, A. Radhakrishnan, and R. Rohatgi.
2019. Cholesterol accessibility at the ciliary membrane controls hedgehog signaling.
Elife. 8.
Kinnebrew, M., G. Luchetti, R. Sircar, S. Frigui, L.V. Viti, T. Naito, F. Beckert, Y. Saheki, C.
Siebold, A. Radhakrishnan, and R. Rohatgi. 2021. Patched 1 reduces the accessibility of
cholesterol in the outer leaflet of membranes. *Elife*. 10.
Kong, J.H., C. Siebold, and R. Rohatgi. 2019. Biochemical mechanisms of vertebrate hedgehog
signaling. *Development*. 146.
Mangeol, P., B. Prevo, and E.J. Peterman. 2016. KymographClear and KymographDirect: two
tools for the automated quantitative analysis of molecular and cellular dynamics using
kymographs. *Mol Biol Cell*. 27:1948-1957.
Marra, A.N., B.D. Adeeb, B.E. Chambers, B.E. Drummond, M. Ulrich, A. Addiego, M. Springer,
S.J. Poureetezadi, J.M. Chambers, M. Ronshaugen, and R.A. Wingert. 2019.
Prostaglandin signaling regulates renal multiciliated cell specification and maturation.
Proc Natl Acad Sci U S A. 116:8409-8418.
May, E.A., M. Kalocsay, I.G. D'Auriac, P.S. Schuster, S.P. Gygi, M.V. Nachury, and D.U. Mick.
2021. Time-resolved proteomics profiling of the ciliary Hedgehog response. *J Cell Biol*.
220.

Mukhopadhyay, S., X. Wen, N. Ratti, A. Loktev, L. Rangell, S.J. Scales, and P.K. Jackson.
2013. The ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic
hedgehog pathway via cAMP signaling. *Cell*. 152:210-223.
Niewiadomski, P., J.H. Kong, R. Ahrends, Y. Ma, E.W. Humke, S. Khan, M.N. Teruel, B.G.
Novitch, and R. Rohatgi. 2014. Gli protein activity is controlled by multisite
phosphorylation in vertebrate Hedgehog signaling. *Cell Rep*. 6:168-181.
Ogden, S.K., D.L. Fei, N.S. Schilling, Y.F. Ahmed, J. Hwa, and D.J. Robbins. 2008. G protein
Galphai functions immediately downstream of Smoothened in Hedgehog signalling.
Nature. 456:967-970.
Ott, C., and J. Lippincott-Schwartz. 2012. Visualization of live primary cilia dynamics using
fluorescence microscopy. *Curr Protoc Cell Biol*. Chapter 4:Unit 4 26.
Patten, I., and M. Placzek. 2000. The role of Sonic hedgehog in neural tube patterning. *Cell Mol
Life Sci*. 57:1695-1708.
Petrov, K., B.M. Wierbowski, and A. Salic. 2017. Sending and Receiving Hedgehog Signals.
Annu Rev Cell Dev Biol. 33:145-168.
Praktiknjo, S.D., F. Saad, D. Maier, P. Ip, and D.R. Hipfner. 2018. Activation of Smoothened in
the Hedgehog pathway unexpectedly increases Galphas-dependent cAMP levels in
Drosophila. *J Biol Chem*. 293:13496-13508.
Raleigh, D.R., N. Sever, P.K. Choksi, M.A. Sigg, K.M. Hines, B.M. Thompson, D. Elnatan, P.
Jaishankar, P. Bisignano, F.R. Garcia-Gonzalo, A.L. Krup, M. Eberl, E.F.X. Byrne, C.
Siebold, S.Y. Wong, A.R. Renslo, M. Grabe, J.G. McDonald, L. Xu, P.A. Beachy, and
J.F. Reiter. 2018. Cilia-Associated Oxysterols Activate Smoothened. *Mol Cell*. 72:316-
327 e315.
Ribes, V., and J. Briscoe. 2009. Establishing and interpreting graded Sonic Hedgehog signaling
during vertebrate neural tube patterning: the role of negative feedback. *Cold Spring Harb
Perspect Biol*. 1:a002014.
Riobo, N.A., and D.R. Manning. 2007. Pathways of signal transduction employed by vertebrate
Hedgehogs. *Biochem J*. 403:369-379.
Segi, E., Y. Sugimoto, A. Yamasaki, Y. Aze, H. Oida, T. Nishimura, T. Murata, T. Matsuoka, F.
Ushikubi, M. Hirose, T. Tanaka, N. Yoshida, S. Narumiya, and A. Ichikawa. 1998. Patent
ductus arteriosus and neonatal death in prostaglandin receptor EP4-deficient mice.
Biochem Biophys Res Commun. 246:7-12.
Shi, X., G. Garcia, 3rd, J.C. Van De Weghe, R. McGorty, G.J. Pazour, D. Doherty, B. Huang,
and J.F. Reiter. 2017. Super-resolution microscopy reveals that disruption of ciliary
transition-zone architecture causes Joubert syndrome. *Nat Cell Biol*. 19:1178-1188.
Somatilaka, B.N., S.H. Hwang, V.R. Palicharla, K.A. White, H. Badgandi, J.M. Shelton, and S.
Mukhopadhyay. 2020. Ankmy2 Prevents Smoothened-Independent Hyperactivation of
the Hedgehog Pathway via Cilia-Regulated Adenylyl Cyclase Signaling. *Dev Cell*.
54:710-726 e718.
Thotala, D., J.M. Craft, D.J. Ferraro, R.P. Kotipatruni, S.R. Bhave, J.J. Jaboin, and D.E.
Hallahan. 2013. Cytosolic phospholipaseA2 inhibition with PLA-695 radiosensitizes
tumors in lung cancer animal models. *PLoS One*. 8:e69688.
Truong, M.E., S. Bilekova, S.P. Choksi, W. Li, L.J. Bugaj, K. Xu, and J.F. Reiter. 2021.
Vertebrate cells differentially interpret ciliary and extraciliary cAMP. *Cell*. 184:2911-2926
e2918.
Tuson, M., M. He, and K.V. Anderson. 2011. Protein kinase A acts at the basal body of the
primary cilium to prevent Gli2 activation and ventralization of the mouse neural tube.
Development. 138:4921-4930.
Umberger, P.A., and S.K. Ogden. 2021. SPOP and CUL3 Modulate the Sonic Hedgehog Signal
Response Through Controlled Degradation of GLI Family Transcription Factors. *Front
Cell Dev Biol*. 9:710295.

Vuolo, L., A. Herrera, B. Torroba, A. Menendez, and S. Pons. 2015. Ciliary adenylyl cyclases
801   control the Hedgehog pathway. *J Cell Sci*. 128:2928-2937.
802   Wheway, G., L. Nazlamova, and J.T. Hancock. 2018. Signaling through the Primary Cilium.
803   *Front Cell Dev Biol*. 6:8.
804   Wijaya, J., B.T. Vo, J. Liu, B. Xu, G. Wu, Y. Wang, J. Peng, J. Zhang, L.J. Janke, B.A. Orr, J.
805   Yu, M.F. Roussel, and J.D. Schuetz. 2020. An ABC Transporter Drives Medulloblastoma
806   Pathogenesis by Regulating Sonic Hedgehog Signaling. *Cancer Res*. 80:1524-1537.
807   Ye, F., D.K. Breslow, E.F. Koslover, A.J. Spakowitz, W.J. Nelson, and M.V. Nachury. 2013.
808   Single molecule imaging reveals a major role for diffusion in the exploration of ciliary
809   space by signaling receptors. *Elife*. 2:e00654.
810   Zhang, P., B. Fan, P. Yang, J. Temirov, J. Messing, H.J. Kim, and J.P. Taylor. 2019. Chronic
811   optogenetic induction of stress granules is cytotoxic and reveals the evolution of ALS-
812   FTD pathology. *Elife*. 8.
813   Zhou, X., L.X. Fan, K. Li, R. Ramchandran, J.P. Calvet, and X. Li. 2014. SIRT2 regulates
814   ciliogenesis and contributes to abnormal centrosome amplification caused by loss of
815   polycystin-1. *Hum Mol Genet*. 23:1644-1655.

 FIGURE LEGENDS

Figure 1: PGE$_2$ generated downstream of SHH contributes to ciliary length control. (A-A').

PGE$_2$ is secreted downstream of SMO and cPLA$_2$ activation. PGE$_2$ was measured in IMCD3 **(A)**

or WT and *Abcc4$^{-/-}$* NIH-3T3 **(A')** cell culture media by ELISA. **(A).** Cells were pretreated with

vehicle control (DMSO), the inverse SMO agonist cyclopamine (10 µM), or cPLA$_2$ inhibitor

giripladib (GIRI, 2 µM), and then exposed to SHH. PGE$_2$ was measured in cell culture

supernatant ~36 hours post treatment. The experiment was repeated twice with 3 biological

replicates per experiment. Pooled data are shown. **(A').** Cells were pretreated with vehicle

(DMSO), cyclopamine (10 µM), or COX2 inhibitor Celecoxib (Coxib,10 µM) and then stimulated

with control or SHH conditioned media. PGE$_2$ was measured in the supernatant ~36 hours after

treatment. The experiment was performed twice with 3 biological replicates per experiment. A

representative experiment is shown. **(B-C).** cPLA$_2$ inhibition shortens primary cilia. **(B)**. IMCD3

cells were treated with SHH conditioned media in the presence of GIRI (4 µM), or vehicle control

(DMSO). The ciliary axoneme is marked by acetylated α-tubulin (magenta). DAPI (blue) marks

the nucleus. Scale bar = 10 µm. **(C)**. Lengths of primary cilia were quantified in IMCD3 cells

treated with SHH conditioned media in the absence or presence of GIRI. Average primary cilium

length was determined by measuring cilia of ≥100 cells/condition across three independent

experiments. **(D-E).** PGE$_2$ supplementation rescues ciliary length in GIRI-treated IMCD3 cells.

Cells were treated with 4 µM GIRI in the presence of PGE$_2$ media supplementation (40 µM) or

vehicle control. Scale bar = 10 µm. Ciliary length quantification is shown in **E. (F-G).** Arachidonic

acid supplementation rescues primary cilium length following cPLA$_2$ inhibition. IMCD3 cells were

treated with SHH conditioned media in the presence of vehicle or 4 µM GIRI in media

supplemented with increasing concentrations of arachidonic acid (12.5 µM or 25 µM). Scale bar

= 10 µm. Ciliary length quantification is shown in **G**. For all experiments, statistical significance

was calculated using one-way ANOVA. A p value of less than 0.05 was considered statistically

significant with significance indicated as follows: *<0.05, **<0.01, ***<0.001, ****<0.0001, and

ns, p > 0.05. Data are represented as mean ± SD.

Figure 2: EP$_4$ is required for primary cilium length homeostasis. (A). EP$_4$ localizes to

primary cilia of IMCD3 cells. Endogenous EP$_4$ is shown in green and acetylated α-tubulin

(magenta) marks the primary cilium. Scale bar = 5 µm. **(B-C').** EP$_4$ localizes to primary cilia in

wild type MEFs. Ciliary EP$_4$ signal is absent in *Ep$_4$*$^{-/-}$ cells. Line scans indicate the degree of

localization between EP$_4$ (green) and acetylated α-tubulin (magenta) within the primary cilium in

control **(B')** and *Ep$_4$*$^{-/-}$ MEFs **(C').** Intensity profiles are presented as arbitrary units (AU). Scale

bars = 5 µm. **(D).** EP$_4$ inhibition shortens primary cilia in IMCD3 cells. Cells were pretreated with

vehicle control (DMSO) or EP$_4$ inhibitor L161,982 (10 µM or 20 µM) for two hours prior to

addition of SHH. **(E).** PGE$_2$ partially rescues primary cilia length following EP$_4$ inhibition. IMCD3

cells were treated with SHH conditioned media in the presence of vehicle or 20 µM L161,982 in

media supplemented with increasing concentrations of PGE$_2$ (25 or 50 µM). At least 150

cilia/condition were measured over three experiments. Statistical significance was calculated

using one-way ANOVA. For all experiments, a p value of less than 0.05 was considered

statistically significant with significance indicated as follows: *<0.05, **<0.01, ***<0.001,

****<0.0001, and ns, p > 0.05. Data are represented as mean ± SD.

Figure 3: SHH signaling is attenuated by inhibition of cPLA$_2$ or EP$_4$. (A-B). SMO ciliary

enrichment is reduced by EP$_4$ inhibition. SMO (**A,** red) enrichment in primary cilia was examined

in IMCD3 cells. Primary cilia are marked with acetylated α-tubulin (magenta). Cells were

pretreated with vehicle (DMSO), GIRI (4 µM), or L161,982 (20 µM) for 18 hours. Scale bar = 5

µm. **(B).** Average SMO ciliary signal intensity was quantified for each condition. **(C-E).** EP$_4$ loss

shortens primary cilia and attenuates SMO ciliary enrichment. Ciliary length and SMO (red)

ciliary enrichment were examined in wild type and $Ep4^{-/-}$ MEFs 18 hours post treatment with

control or SHH conditioned media. Cilia are marked by acetylated α-tubulin (magenta, **C**). Scale

bar =10 µm. A representative experiment is shown. Quantification of average primary cilium

length **(D)** and ciliary SMO signal intensity **(E)** were determined as above by measuring cilia of

≥150 cells/condition across three independent experiments. **(F-F')**. qRT-PCR analyses of *Gli1*

and *Ptch1* expression in IMCD3 and NIH-3T3 cells were performed. Cells were pretreated with

vehicle (DMSO), GIRI (4 µM), or L161,982 (10 µM) for 2 hours and then cultured in control or

SHH conditioned media plus inhibitor for 18 hours. Fold-change in expression was determined

using the $2^{-\Delta\Delta Ct}$ method. Average fold change was calculated across two independent

experiments with three biological replicates per experiment. All data are pooled. For all

experiments, statistical significance was determined using a one-way ANOVA. A p value of less

than 0.05 was considered statistically significant. Significance is denoted as follows: *<0.05,

<0.01, *<0.001, ****<0.0001, and ns, p > 0.05. Error bars indicate SD.

Figure 4: Ciliary adenylyl cyclases contribute to primary cilium length control and SHH-

stimulated transcriptional activation. (A). A model for SHH and EP$_4$ effects on primary cilium

cAMP control. **(B-D).** Knockdown of ciliary AC3, AC5, or AC6 disrupts ciliary cAMP control,

shortens primary cilia, and compromises *Gli1* induction. **(B).** IMCD3 cells were treated with

scrambled, *Adcy3*, *Adcy5*, or *Adcy6* siRNA for 48 hours and ciliary cAMP was measured using

the cADDis cAMP sensor. cADDis activity was recorded as cells were exposed to vehicle

(ethanol, black) or forskolin (FSK,100 µM, blue). Fluorescence ciliary intensity was recorded

over 8 minutes in live cell imaging mode. An average of ~5 cilia were recorded for each

condition. The experiment was performed three times. A representative experiment is shown.

Significance was determined by calculating the area under the curve followed by Student's t-test

analysis. **(C)**. Ciliary AC knockdown shortens primary cilia. Knockdown experiments were

repeated at least twice with 75-100 primary cilia measured per experiment per condition. All

data were pooled. Significance was determined by one-way ANOVA. **(D)**. *Gli1* fold change was

determined in control and SHH-stimulated IMCD3 cells following *Adcy3*, *Adcy5*, or *Adcy6*

knockdown. Fold change was assessed by qRT-PCR ~48 hours after treatment with AC-specific

siRNA or scrambled control following 2 hours of serum starvation and overnight treatment with

SHH conditioned media. Expression was determined using the $2^{-\Delta\Delta Ct}$ method. Significance was

determined by a one-way ANOVA. **(E-F)**. Knockdown of *Ankmy2* shortens primary cilia and

attenuates ligand-stimulated *Gli1* induction. **(E).** Knockdown experiments were repeated at least

twice with 75-100 primary cilia measured per experiment per condition. All data were pooled.

Significance was determined by Student's t-test **(F).** *Gli1* fold change was determined in control

or SHH-stimulated IMCD3 cells following *Ankmy2* knockdown, as described above. The

experiment was performed twice in triplicate and all data pooled. Significance was determined

by a one-way ANOVA. For all experiments, a p value of less than 0.05 was considered

statistically significant. Significance is denoted as follows: *<0.05, **<0.01, ***<0.001,

****<0.0001, and ns, p > 0.05. Error bars indicate SD.

Figure 5: EP$_4$ stabilizes ciliary cAMP in SHH-stimulated cells to maintain anterograde IFT

and SMO ciliary accumulation. (A). EP$_4$ signaling equilibrates SMO-induced reduction of

primary cilium cAMP. cADDis-expressing IMCD3 cells were pretreated overnight with vehicle or

L161,982 (10 µM). The following morning, cADDis activity was monitored by live imaging to

track the ciliary cAMP response as cells were treated with forskolin (FSK, 100 µM) for 1.5

minutes prior to addition of the SMO agonist SAG (1 µM) or vehicle control. Fluorescence ciliary

intensity was recorded over 10 minutes in live cell imaging mode. An average of ~6 cilia were

recorded for each condition and the experiment was performed twice. A representative

experiment is shown. **(B)**. Anterograde IFT velocity was calculated in IMCD3 cells by tracking

IFT88-GFP movement in the presence and absence of SHH, L161,982 (10 μM), or vehicle

control. IFT velocity was calculated for 30 cilia per condition across 4 experiments. Velocity is

shown as a violin plot with SD indicated. **(C-D).** Average ciliary length and SMO ciliary intensity

were quantified in IMCD3-bPAC cells exposed to control or SHH conditioned media in the

absence or presence of 10 μM L161,982 in control or blue light-exposed cells. Significance was

determined by one-way ANOVA. For all experiments, a p value of less than 0.05 was

considered statistically significant. Significance is denoted as follows: *<0.05, **<0.01,

<0.001, *<0.0001, and ns, p > 0.05. Data are represented as mean ± SD. For all ciliary

analyses, 50-100 cells per condition were analyzed over at least 3 independent experiments.

Figure 6: EP$_4$ loss *in vivo* reduces primary cilium length and alters SHH-dependent neural

tube patterning. (A). Scanning electron microscopy images of cardiac level neural tube

sections from E9.5/27 somite stage *Ep4$^{+/-}$* and *Ep4$^{-/-}$* embryos show apical plasma membranes

of cells lining the neural tube lumen. Scale bar = 2 μm. Cilia are shaded in pink **(B)**. Average

primary cilium length was calculated by measuring cilia of ~30 cells/embryo lining the neural

tube lumen. Primary cilia of *Ep$_4$$^{-/-}$* embryos show a statistically significant length reduction.

Statistical significance was calculated using one-way ANOVA. Error bars indicate SD**.**

Significance is denoted as follows: ****<0.0001, and ns, p > 0.05. **(C-D).** *Ep$_4$$^{-/-}$* embryos show

altered ventral progenitor domains. Cardiac level sections of E9.5/29 somite stage neural tubes

from the indicated genotypes were immuno-stained for neural progenitor domain markers *Olig2*

(magenta) and *Pax6* (green). *Ep$_4$$^{-/-}$* neural tubes show disrupted ventral progenitor specification

as indicated by reduced *Olig2*-labeled cellular populations. For all experiments, four embryos

per genotype were analyzed. Scale bar = 100 μm. Progenitor domain area is shown in **(D)**.

Statistical significance was calculated using a Student's t-test. Error bars indicate SD**.**

Significance is denoted as follows: *<0.05. Sections shown in **(C)** are indicated as colored dots

in each graph.

Figure 7: A model for SHH-to-EP4 signal crosstalk. (A). In the absence of SHH, the SHH

receptor PTCH prevents ciliary accumulation and signaling of the GPCR SMO. In this basal

condition, GPR161 signaling and basal PGE_2 secretion and activation of EP_4-$G\alpha_s$ maintain

sufficient ciliary cAMP for anterograde IFT and length homeostasis. **(B).** SHH binding to PTCH

leads to SMO de-repression and GPR161 ciliary exit. SMO activates $G\alpha_i\beta\gamma$ to increase

production of arachidonic acid by $\beta\gamma$ stimulation of $cPLA_2\alpha$. Arachidonic acid is metabolized to

PGE_2, which is secreted to activate ciliary EP_4 for $G\alpha_s$ activation. This ensures ciliary cAMP

levels remain sufficiently high in SHH-stimulated cells to prevent anterograde IFT slowing that

leads ciliary shortening. **(C).** Inhibition of $cPLA_2$ or EP_4 reduces SHH-to-EP4 signal crosstalk,

reduces primary cilium length, and blunts SMO ciliary accumulation and high-level signaling.

 SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1: cPLA$_2$ is required for SMO ciliary accumulation but not for

GPR161 ciliary exit. (A-D). IMCD3 cells were treated with vehicle **(A-B)** or GIRI **(C-D**, 4 µM**)** in

the absence or presence of SHH as indicated. GPR161 is magenta, SMO is red, ARL13B

(green) marks primary cilia. Scale bar = 3 µm. **(E-E').** Quantification of ciliary signal intensity for

SMO **(E)** and GPR161 **(E')** in cells treated with SHH, GIRI, or vehicle control. The experiment

was performed twice with at least 30 cilia imaged per condition per experiment and all data were

pooled. Significance was determined using a one-way ANOVA. Significance is indicated as

follows: *<0.05, **<0.01, ***<0.001, ****<0.0001, and ns, p > 0.05.

Supplementary Figure 2: Ciliary AC controls primary cilium cAMP. (A). AC3 and AC5/6

localize to primary cilia in IMCD3 cells. ARL13B (magenta in merge and white in insets) marks

primary cilia. Ciliary ACs are shown in green (merge) and in white (insets). Scale bar = 5µm.

(B). *Adcy3*, *Adcy5,* or *Adcy6* knockdown in IMCD3 cells was validated by qRT-PCR ~48 hours

after treatment with AC-specific siRNA or scrambled control. Fold-change in expression was

determined using the $2^{-\Delta\Delta Ct}$ method. Experiments were performed 3 times in triplicate and all

data pooled. Statistical significance was calculated using Student's t-test. **(C).** Knockdown of

ciliary AC does not alter forskolin (FSK)-stimulated whole cell cAMP elevation. The indicated

ciliary AC were knocked down as above. Cells were stimulated with 100 µM forskolin ~48 hours

after siRNA treatment and whole cell cAMP was measured using the direct cAMP ELISA assay.

Assays were performed at least 3 times and all data pooled. Statistical significance was

calculated using Student's t-test. **(D-F').** Live imaging of cADDis-expressing IMCD3 cells shows

changes in fluorescence intensity of a ciliary cAMP sensor (green) and stable fluorescence of

the ciliary reference marker (magenta) following exposure to vehicle **(D)** or forskolin (FSK, 100

µM) **(E)**. Increasing cAMP decreases green fluorescence. Scale bar = 2 µm. **(F-F').** Relative

cAMP shift is shown as native **(F)** or inverse **(F')** of the fluorescent intensity ratio of the cAMP

sensor to reference marker. Inverse ratios are shown in the main figures to clearly illustrate

cAMP increase. Significance was determined by calculating the area under the curve followed

by Student's t-test analysis. Significance is indicated as follows: *<0.05, **<0.01, ***<0.001,

****<0.0001, and ns, p > 0.05.

Supplementary Figure 3: *Ankmy2* knockdown impacts IMCD3 ciliary AC enrichment and

SHH-stimulated transcriptional activation. (A). qRT-PCR verification of *Ankmy2* knockdown

in IMCD3 cells ~48 hours after treatment with *Ankmy2* siRNA or scrambled control. Fold-change

in expression was determined using the $2^{-\Delta\Delta Ct}$ method. **(B-C).** Knockdown of the ciliary AC

chaperone *Ankmy2* reduces primary cilium AC occupancy. Colocalization of AC3 **(B)** and AC5/6

(C) to ARL13B in primary cilia of IMCD3 cells in scrambled and *Ankmy2* siRNA-treated cells is

shown. Both M1 and M2 scores of AC3 and AC5/6 to ARL13B are significantly reduced in cilia

of the *Ankmy2* siRNA treated cells compared to control. Statistical significance was calculated

using Student's t-test. **(D).** *Ankmy2* knockdown reduces forskolin-induced ciliary cAMP increase.

Forskolin (FSK) induced changes in primary cilium cAMP were monitored in *Ankmy2*

knockdown cells using the CADDis sensor. Forskolin-mediated ciliary cAMP accumulation is

blocked by *Ankmy2* knockdown. Significance was determined by calculating the area under the

curve followed by Student's t-test analysis. **(E).** *Adcy3*, *Adcy5,* and *Anmky2* knockdown in NIH-

3T3 cells was validated by qRT-PCR ~48 hours after treatment with *Adcy3, Adcy5,* or *Ankmy2*

siRNA or scrambled control. Fold-change in expression was determined using the $2^{-\Delta\Delta Ct}$

method. Student's t-test was used to calculate statistical significance. **(F-G).** Knockdown of

Adcy3, Adcy5, or *Ankmy2* shortens primary cilia and reduces SHH target gene induction in NIH-

3T3 cells. **(F).** Average lengths of ARL13B marked primary cilia of NIH-3T3 cells were

determined as described above. Knockdown experiments were repeated three times with 75-

100 primary cilia measured per condition per experiment. **(G).** *Gli1* fold change was determined

in control or SHH-stimulated NIH-3T3 cells following *Adcy3, Adcy5,* or *Ankmy2* knockdown.

Significance was determined by one-way ANOVA. For all experiments, significance is denoted

as follows: *<0.05, **<0.01, ***<0.001, ****<0.0001, and ns, p > 0.05. Data are represented as

mean ± SD. qPCR analyses were performed twice in triplicate and all data were pooled.

1009

Supplementary Figure 4: Validation of optogenetic cAMP modulation in IMCD3 cells. (A).

Schematic of optogenetic cAMP modulation. Ciliary bPAC is activated by blue light to increase

ciliary cAMP**. (B).** Immunofluorescence imaging of IMCD3 cells expressing Cyto-bPAC-GFP or

Cilia-bPAC-GFP (green). Acetylated α-tubulin marks cilia (magenta) and DAPI (blue) marks

nuclei. Scale bars = 10 µm. **(C-C").** Average ciliary length was quantified in IMCD3 Cyto-bPAC,

and IMCD3 Cilia-bPAC cells in control conditions or following 180 minutes of blue light

exposure. **(C').** Total cellular cAMP was measured as in Supplementary Figure 2C. Error bars

indicate SD. Significance was determined by a one-way ANOVA in **(C)** and a Student's t-test in

(C'). **(D-D').** Ciliary lengths were measured in vehicle and forskolin treated cells. Average

primary cilium length was calculated by measuring cilia of ≥100 cells/condition across three

independent experiments. All data were pooled. **(D').** IMCD3 cells were treated with vehicle or

100 µM forskolin. The ciliary axoneme is marked by ARL13B (magenta) and DAPI (blue) marks

nuclei. Scale bar = 5 µm. Significance was determined using a one-way ANOVA. For all

experiments significance is indicated as follows: *<0.05, **<0.01, ***<0.001, ****<0.0001, and

ns, p > 0.05.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

SHH Pathway OFF

GPR161 signaling to Gα$_s$

No SMO signaling to Gα$_i$ or GLI

Basal PGE$_2$/EP$_4$ signaling to Gα$_s$

Basal [cAMP]

B

C

SHH Pathway ON

GPR161 exits cilia

SMO signaling to Gα$_i$ active

High SMO ciliary enrichment

High SMO signaling to GLI

Increased EP$_4$ signaling to Gα$_s$

[cAMP] equilibration

SHH Pathway ON +cPLA$_2$ or EP$_4$ inhibition

GPR161 exits cilia

SMO signaling to Gα$_i$ active

Decreased SMO ciliary enrichment

Attenuated SMO signaling to GLI

EP$_4$ signaling to Gα$_s$ inhibited

Disrupted [cAMP] equilibration

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

A Cilia-bPAC / Cilia-bPAC — No Light / Blue Light — ↑cAMP

B IMCD3-Cyto-bPAC / IMCD3-Cilia-bPAC — Ac-α-Tub Cyto-bPAC-GFP DAPI / Ac-α-Tub Cilia-bPAC-GFP DAPI

C Cilia length (μm) — ns, ns, **** — IMCD3 Cyto-bPAC No Light, IMCD3 Cyto-bPAC Blue Light, IMCD3 Cilia-bPAC No Light, IMCD3 Cilia-bPAC Blue Light

C' cAMP (pmol/ml) — **** — Cyto-bPAC - No Light, Cyto-bPAC - Blue Light

C'' IMCD3 Cyto-bPAC / IMCD3 Cilia-bPAC — No Light / Blue Light — ARL-13B / DAPI

D IMCD3 Cilia Length (μm) — ** — Vehicle, Forskolin

D' Vehicle / Forskolin — IMCD3 - WT — ARL-13B / DAPI