The bacterial lysate Lantigen B reduces the expression of ACE2 on primary

oropharyngeal cells

Caterina Pizzimenti[1], Antonella D'Agostino [1], Paola Pirrello[1], Gilberto Rinaldi[1,2], Giovanni

Melioli[2,3]

[1]Department of Research and Development, Bruschettini Ltd, Via Isonzo 6, 16147, Genova,

Italy

[2]Department of Medical Affair, Bruschettini Ltd, Via Isonzo 6, 16147, Genova, Italy

[3]Laboratory Medicine, Albaro Site, Alliance Medical Diagnostic Ltd, Via Boselli 30, 16146,

Genova, Italy

Corresponding author

Giovanni Melioli, MD

Laboratory Medicine, Albaro Site

Alliance Medical Diagnostic Ltd,

Via Boselli 30, 16146, Genova, Italy

Email: covid.lab@alliancemedical.it

Abstract

Background: Vercelli and coworkers recently observed that a well-established bacterial

lysate (OM-85, Vifor Pharma; CH) was able to downregulate the expression of Angiotensin-

Converting Enzyme 2 (ACE2) on epithelial cells. This downregulation was also associated

with a reduced infectivity of cells, resulting in a reduced viral titre. We evaluated whether

another bacterial lysate (Lantigen B, Bruschettini Ltd; Italy) may have similar activities.

However, while OM-85 is given per os and has a systemic effect after absorption at the gut

level, Lantigen B is given locoregionally. Thus, the concentration that the bacterial lysate can

reach at the mucosal level seems to be promising.

Methods: Oropharyngeal cells were collected from healthy donors. After 24 hours of

treatment in vitro with doses of Lantigen B comparable to those that are reached in vivo, the

expression of ACE2 was evaluated by direct fluorescence and flow cytometry.

Results: A reduction in the number of ACE2-positive cells was observed in 80% of treated

samples. Only a few donors had poor expression of ACE2, and in these donors, the

downregulation was less evident or absent.

Conclusions: These results suggest that Lantigen B, at pharmacological doses, could be an

interesting drug to reduce ACE2 expression on oropharyngeal cells, thus contributing to the

prophylaxis of COVID-19 in humans.

Trial registration

Not applicable

Keywords

SARS-CoV-2; ACE2; COVID-19; Bacterial lysates; Lantigen B

Background

The cell surface molecule angiotensin-converting enzyme 2 (ACE2) is the main receptor for

SARS-CoV-2 spike (S) protein attachment [1, 2]. The binding of the spike protein to the

ACE2 receptor, together with other mechanisms, allows the fusion of the virus and the cell

membrane, thus infecting the target cell. ACE2 has been shown in many different cell types,

including nasal, intestinal, kidney, testis, gallbladder, and heart cells, and, to a lesser extent,

tongue, tonsil, and the olfactory region cells [3, 4]

The COVID-19 pandemic has pushed research to the limit, and in a few months, not

only advanced protocols for virus identification but also active vaccines have been developed

and introduced in humans [5, 6]. At present, few monoclonal antibodies are active against all

viral variants [7], and novel drugs have been proposed or are under development [8].

In a recent interesting article [9], the authors showed that a bacterial lysate developed

decades ago (OM-85) has the capability of reducing the expression of ACE2 both in cell lines

and in primary cells from human airways. This finding suggests that the administration of

OM-85, which would reduce the expression of ACE2, could to some extent inhibit

respiratory tract target cell infection, acting as a prophylactic treatment to be used in

association with vaccines and other future therapies.

Bacterial lysates have been shown to have the capacity to reduce the number of

respiratory tract infections (or exacerbations) in treated patients [10, 11]. The detailed

mechanisms are not completely described, although the capacity for activating dendritic cells,

inducing a specific and functionally effective immune response, and recruiting innate

immune-competent cells has been observed in the past [12, 13]. In addition, it has been

shown that epithelial cells are able to recognize certain bacterial lysates (such as PMBL),

resulting in epithelial cell activation, proliferation and tight junction sealing [14]. Lantigen B

is a bacterial lysate that was originally developed in the 1960s and has been empirically used

in recent decades to reduce the number of respiratory tract infections during the winter

season. A recent clinical trial has clearly demonstrated the clinical activity of Lantigen B

[15]. Lantigen B is administered under the tongue; thus, the drug enters into direct contact

with the epithelial cells of the mouth and the oropharynx. If ACE2 expression modulation is a

characteristic of bacterial lysates (and not only of OM-85), a possible effect of Lantigen B on

the same protein could be hypothesized. In this report, we show that epithelial cells of the

oropharynx express ACE2 and that Lantigen B is able to reduce the expression of ACE2 after

a 24-hour treatment at doses comparable to those used in vivo.

Methods

Epithelial cells of the oropharynx were collected from a group of researchers and employees

(14 males and 16 females, age ranging from 35 to 73) at Bruschettini Ltd, the Italian firm that

produces Lantigen B. All donors gave verbal informed consent. Donors were first screened

for the presence of SARS-CoV-2 antigens (using a rhino-pharyngeal swab and a standard

antigenic lateral flow cassette). All donors were negative for SARS-CoV-2 infection and thus

were recruited in the different sessions of the study. Epithelial cells were then collected by a

sterile swab of the oropharynx of the donors. The cells were washed in saline and then

resuspended in RPMI 1640 with 10% foetal bovine serum and gentamycin. The vitality of the

cells was scored by trypan blue exclusion.

The expression of ACE2 on epithelial cells was measured by a specific anti-ACE2

monoclonal antibody (R&D Systems, labelled with Alexa Fluor 488) and flow cytometry

(Cytoflex, Beckman Coulter). An isotypic control (FITC-labelled anti-CD83, Beckman

Coulter) was used to evaluate the possible nonspecific binding to epithelial cells. To evaluate

whether the modulation of ACE2 can be considered in the context of a more general

mechanism, another surface molecule (CD54 – ICAM-1, receptors of rhinoviruses) [16] was

used. These experiments were repeated three times using three different cohorts consisting of

six donors. Polystyrene six micrometre microspheres (Polysciences, USA) were used to

calibrate the flow cytometer and to evaluate the cell size.

Modulation of ACE2 by Lantigen B. Cells collected from donors were washed with

saline and resuspended in RPMI (Euroclone, Pero, Italy) as described, and different

experimental conditions were provided. First, cells were incubated overnight in RPMI with

FBS (HyClone, Sigma, USA) and gentamycin in the absence of any stimulus. Second, the

cells were incubated with Lantigen B diluent containing polysorbate 80, chlorhexidine

diacetate and sodium methyl parahydroxybenzoate. This condition was considered the

experimental negative control. Third, the cells were diluted 1:1 with the drug used in humans.

The drug contained the particulate and soluble fraction obtained by the chemical lysis of six

different microbes of the respiratory tract (*Staphylococcus aureus*, *Klebsiella pneumoniae*,

Streptococcus pneumoniae, *Streptococcus pyogenes*, *Moraxella catarrhalis* and *Haemophilus*

influenzae). After 24 hours of culture, the cells were collected, stained with the anti-ACE2

monoclonal antibody, and analysed on a cytofluorimeter. Vitality was evaluated for the same

cells using trypan blue dye exclusion. The test was performed on 10 different donors in three

different sessions.

Results

Expression of ACE2 on normal oropharyngeal cells. As described, the first series of

experiments was focused on the identification of ACE2 on normal oropharyngeal cells

immediately after cell collection with an oropharyngeal swab. Indeed, only in the presence of

clear ACE2 expression could any attempt to downregulate this expression be started. Figure 1

shows the expression of ACE2 in the FITC channel and the same population treated with an

isotypic control. As described, the assay was performed on cells from 18 different donors.

ACE2 was expressed on the surface of oropharyngeal cells with a certain heterogeneity. The

median fraction of ACE2-positive cells was 60% (ranging from 52 to 80%). Few samples

were characterized by a reduced percentage of ACE2-positive cells (ranging from 37 to

47%). The intensity of fluorescence was more homogeneous. Of note, epithelial cells were

large; the diameter measured using 6 micrometre plastic beads on the forward scatter channel

ranged between 18 and 56 micrometres. Under these conditions, the very large surface area

caused a strong signal in the FITC channel. Additionally, autofluorescence was high but

significantly lower than that of ACE2 mAb-stained cells.

Modulation of ACE2 on normal oropharyngeal cells after incubation with Lantigen B.

Using the same conditions of the first experiments, donor oropharyngeal cells were tested for

the presence of ACE2 after 24 hours of incubation in the presence and absence of Lantigen B.

Treatment with Lantigen B resulted in a reduction in the number of ACE2-positive cells in

74% of samples (median reduction 44%, ranging from 18% to 69.5%). A reduction in the

fluorescence intensity (representative of the number of surface molecules) was observed in

65% of samples (median reduction 12.4%, ranging from 4.2 to 43%). Figure 2 shows the

results of the two different donors. Notably, not all samples where the percentage of ACE2-

positive cells was reduced also showed a reduction in fluorescence intensity.

CD54 (ICAM-1) was used as a control, being a rhinovirus family receptor. Despite

being well expressed on the surface of nearly all oropharyngeal cells, its expression intensity,

as well as the number of positive cells, was unmodified by the administration of Lantigen B

in this experimental setting.

Discussion

A recent article by Vercelli and coworkers showed that a bacterial lysate (OM-85) has the

capability of reducing the expression of ACE2 on epithelial cells after in vitro culture. In the

same experiments, it was demonstrated that OM-85 inhibited SARS-CoV-2 infection of

epithelial cells. This result is interesting because it can be hypothesized that drugs with this

capability can be used in the context of the prophylaxis of COVID-19. Even more

interestingly, it has been demonstrated that vaccination induces a strong systemic

immunological response lasting months. However, in the same patients, the locoregional

response at the mucosal level is shorter [17, 18]. This finding could be one of the

explanations for the inability of vaccination to prevent reinfections, while severe disease is

better controlled by systemic immunity. Thus, already available drugs that can reduce the

infection of cells, representing the first barrier, are extremely useful. However, these results

were obtained using high concentrations of the bacterial lysate (from 0.24 to 1.92 mg/mL),

doses that seem to be difficult to reach in vivo, considering that a single pill of OM-85

contains 7 mg of the active principle.

Lantigen B is administered at the mucosal level, and this could be one added value of

this treatment. Indeed, the assays performed in vitro on freshly isolated oropharyngeal

epithelial cells indicate that a significant reduction in expression is achieved using

concentrations of the drug that are easily reached in vivo. In addition, the observed reduction

is observed after 24 hours, while OM-85 has effects observed after a longer period in the

experimental conditions described.

This effect seems to be strictly related to the presence of the bacterial antigens

because the excipients and preservatives have no effect when used alone. Furthermore, the

surface expression of CD54, a rhinovirus receptor, was unmodified by in vitro treatment with

Lantigen B, thus suggesting that, at least in these experimental conditions, the

downregulation of ACE2 seems to be bacterial lysate specific.

The major limitation of this study is represented by the absence of any assay of

SARS-CoV-2 infection of oropharyngeal cells and its inhibition mediated by Lantigen B.

However, even if not formally demonstrated, this seems to be highly possible at least

comparing the results of the reduction in the expression of ACE2 in these experiments with

those described by Vercelli et al.

In conclusion, Lantigen B, a bacterial lysate used for decades in the prophylaxis of

recurrent respiratory infections, has well-established activity in reducing the number of

infections in clinical trials. Our novel findings add that Lantigen B has activity that could be

extremely useful in the prophylaxis of COVID-19 (or at least in the prevention of SARS-

CoV-2 asymptomatic infections), particularly in the period that seems to be less protected by

vaccines. Thus, in addition to the well-known effects of bacterial lysates on the adaptive and

innate immune response, the finding of this capability of Lantigen B adds to the mechanism

of action of bacterial lysates.

List of abbreviations

SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus 2

ACE2: Angiotensin-Converting Enzyme 2

COVID-19: COronaVIrus Disease-2019

ICAM-1: Intercellular Adhesion Molecule 1

FITC: Fluorescein isothiocyanate

mAb: monoclonal Antibody

Declarations

Ethics approval and consent to participate

The donors involved in this study were all employees of Bruschettini ltd, were all volunteers

and had gave an verbal informed consent to collect oropharyngeal cells by a swab.

Consent for publication

Not applicable

Availability of data and materials

The experimental design and the raw data are all included in the laboratory files of

Bruschettini ltd. Data are available upon request.

Competing interests

CP, ADA, PP and GR are employees of Bruschettini ltd. GM is a consultant.

Funding

Funding was provided by Bruschettini medical affairs department

Authors' contributions

CP, ADA and PP carried out the in vitro experimental study. CR and GM designed the study.

GM wrote the manuscript.

Acknowledgements

The Authors thank Alessandra Cavalieri for the administrative support.

Authors' information

CP, ADA and PP are biologists or biotechnologists and carried with expertise in the field of

cellular biology, immunology and quality controls of Bacterial Lysates. CR is the Medical

Director of Bruschettini, expert in clinical trials on Bacterial lysates. GM is a scientist active

in the field of immunology, virology, allergy, and clinical pathology since late seventies. In

the last two decades, GM has focused on the mechanism of action of bacterial lysates and

their clinical efficacy.

References

1.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from

patients with pneumonia in China, 2019. N Engl J Med. 2020;382:727-33.

2.   Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al.

SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a

clinically proven protease inhibitor. Cell. 2020;181:271-80.e8.

3.   Hikmet F, Méar L, Edvinsson Å, Micke P, Uhlén M, Lindskog C. The protein

expression profile of ACE2 in human tissues. Mol Syst Biol. 2020;16:e9610.

4.   Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H. Tissue

distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first

step in understanding SARS pathogenesis. J Pathol. 2004;203:631-7.

5.   Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al.

Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. Nature.

2020;586:589-93.

6.   Wang Z, Schmidt F, Weisblum Y, Muecksch F, Barnes CO, Finkin S, et al. mRNA

vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. Nature.

2021;592:616-22.

7.   Touret F, Baronti C, Bouzidi HS, de Lamballerie X. In vitro evaluation of therapeutic

antibodies against a SARS-CoV-2 Omicron B. 1.1. 529 isolate. Sci Rep.

2022;12:4683

8.   Zhao L, Li S, Zhong W. Mechanism of action of small-molecule agents in ongoing

clinical trials for SARS-CoV-2: a review. Front Pharmacol. 2022;13:840639.

9.   Pivniouk V, Pivniouk O, DeVries A, Uhrlaub JL, Michael A, Pivniouk D, et al. The

OM-85 bacterial lysate inhibits SARS-CoV-2 infection of epithelial cells by

downregulating SARS-CoV-2 receptor expression. J Allergy Clin Immunol.

2022;149:923-33.e6.

10. Braido F, Tarantini F, Ghiglione V, Melioli G, Canonica GW. Bacterial lysate in the

prevention of acute exacerbation of COPD and in respiratory recurrent infections. Int

J Chron Obstruct Pulmon Dis. 2007;2:335-45.

11. Villa E, Garelli V, Braido F, Melioli G, Canonica GW. May we strengthen the human

natural defenses with bacterial lysates? World Allergy Organ J. 2010;3:S17-23.

12. Ricci R, Palmero C, Bazurro G, Riccio AM, Garelli V, Di Marco E, et al. The

administration of a polyvalent mechanical bacterial lysate in elderly patients with

COPD results in serological signs of an efficient immune response associated with a

reduced number of acute episodes. Pulm Pharmacol Ther. 2014;27:109-13.

13. Lanzilli G, Traggiai E, Braido F, Garelli V, Folli C, Chiappori A, et al.

Administration of a polyvalent mechanical bacterial lysate to elderly patients with

COPD: effects on circulating T, B and NK cells. Immunol Lett. 2013;149:62-7.

14. Migliore GS, Campana S, Pasquale CD, Carrega P, Ferlazzo G: Human airway

epithelial cells directly recognize mechanical bacterial lysates eliciting tight junction

sealing, antimicrobial peptides production and epithelial cell proliferation, vol. 20.

Dubai, UAE: XII World Congress on COPD, Asthma & Respiratory Allergy; 2018.

15. Braido F, Melioli G, Candoli P, Cavalot A, Di Gioacchino M, Ferrero V, et al. The

bacterial lysate Lantigen B reduces the number of acute episodes in patients with

recurrent infections of the respiratory tract: the results of a double blind, placebo

controlled, multicenter clinical trial. Immunol Lett. 2014;162:185-93.

16. Mincarini M, Cagnoni F, Canonica GW, Cordone G, Sismondini A, Semino C, Pietra

G, Melioli G. Quantitative flow cytometric analysis of the effects of cetirizine on the

expression of ICAM-1/CD54 on primary cultured nasal cells. Allergy. 2000;55:226-

231.

17. Azzi L, Dalla Gasperina D, Veronesi G, Shallak M, Ietto G, Iovino D, et al. Mucosal

immune response in BNT162b2 COVID-19 vaccine recipients. EBioMedicine.

2022;75:103788.

18. Sheikh-Mohamed S, Isho B, Chao GYC, Zuo M, Cohen C, Lustig Y, et al. Systemic

and mucosal IgA responses are variably induced in response to SARS-CoV-2 mRNA

vaccination and are associated with protection against subsequent infection. Mucosal

Immunol. 2022:1-10. doi: 10.1038/s41385-022-00511-0

Figures legend

Figure 1. Demonstration of the presence of the ACE2 protein on the surface of

oropharyngeal cells. Black profile represents the distribution of ACE2 on cell surface. In

grey, the isotypic negative control shows weak autofluorescence.

Figure 2. Downregulation of ACE2 on oropharyngeal cells. The analyses were performed on

oropharyngeal cells after 24-hour incubation in the presence of pharmacological doses of

Lantigen B. In black, cells from donors treated with drug excipients alone. In grey, cells from

donors after treatment with the drug. Panel A shows a representative example of the

downregulation observed in 74% of donor samples. In Panel B, a different behaviour (absent

or poor reduction in the expression of ACE2) was observed in the remaining samples.