Amplification is the Primary Mode of Gene-by-Sex Interaction in Complex Human Traits

Carrie Zhu[1,2], Matthew J. Ming[1,2], Jared M. Cole[1,2], Michael D. Edge[3], Mark Kirkpatrick[2], Arbel Harpak[1,2,+]

[1] Department of Population Health, The University of Texas at Austin, Austin, TX
[2] Department of Integrative Biology, The University of Texas at Austin, Austin, TX
[3] Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA
[+] To whom correspondence should be addressed: arbelharpak@utexas.edu

Abstract

Sexual dimorphism in complex traits is suspected to be in part due to widespread gene-by-sex interactions (GxSex), but empirical evidence has been elusive. Here, we infer the mixture of ways polygenic effects on physiological traits covary between males and females. We find that GxSex is pervasive but acts primarily through systematic sex differences in the magnitude of many genetic effects ("amplification"), rather than in the identity of causal variants. Amplification patterns account for sex differences in trait variance. In some cases, testosterone may mediate amplification. Finally, we develop a population-genetic test linking GxSex to contemporary natural selection and find evidence for sexually antagonistic selection on variants affecting testosterone levels. Taken together, our results suggest that the amplification of polygenic effects is a common mode of GxSex that may contribute to sex differences and fuel their evolution.

Introduction

Genetic effects can depend on context. If the distribution of contexts differs between groups of people, as they do for males and females, so should the average genetic effects on traits[1,2]. In particular, such gene-by-sex interaction (GxSex) may be a result of sex differences in bodily, environmental and social contexts or epistatic interaction with sex chromosomes. Sex differences in genetic effects on complex traits are clearly of high evolutionary[3–8] and translational[9–17] importance. Yet with the exception of testosterone levels[18–21], the basis of sexual dimorphism in complex traits is not well understood[13]. To date, empirical evidence for GxSex in GWAS data— whether focused on identifying large GxSex effects at individual loci or by estimating genetic correlations between the sexes for polygenic traits—has been lacking.

Here, we set out to study governing principles of GxSex in complex human traits and explain why current approaches for characterizing GxSex may be lacking for this goal. We then suggest a mode of GxSex that may have gone largely underappreciated: A systematic difference in the magnitude of effect of many variants between the sexes, which we refer to as "amplification"[22]. Amplification can happen for a large set of variants regulating a specific pathway if the pathway responds to a shared cue[23–26]. In classic hypothesis-testing approaches that test for a GxSex effect separately in each variant, the signal of amplification may be crushed under the multiple hypothesis burden. On the other hand, even state-of-the-art tools designed with complex traits in mind may miss amplification signals: They often treat genetic correlation (between GWAS estimates based on samples from two environments) as a litmus test for whether effects are the same in two groups[27–31], but correlations are scaleless and thus may entirely miss amplification effects.

We developed a new approach for flexibly characterizing a mixture of male-female genetic covariance relationships and applied it to 27 physiological traits in the UK Biobank. We found that amplification is pervasive across traits, and that considering amplification helps explain sex differences in phenotypic variance. Finally, we consider an implication of polygenic GxSex for sexually antagonistic selection: Our model confirms that variants that affect traits may be subject to sexually antagonistic selection when male and female trait optima are very different or, surprisingly, even if the trait optima are very similar. We developed a novel test for sexually antagonistic polygenic selection, which connects GxSex to signals of contemporary viability

selection. Using this test, we find subtle evidence of sexually antagonistic selection on variants
affecting testosterone levels.

Results

The limited scope of single-locus analysis.   We conducted GWASs stratified by sex
chromosome karyotype for 27 continuous physiological traits in the UKB using a sample of ~150K
individuals with two X chromosomes and another sample of ~150K individuals with XY, and a
combined sample that included both the XX and XY samples. We chose to analyze traits with
SNP heritabilities over 7.5% in the combined sample, to have higher statistical power. While there
is not a strict one-to-one relationship between sex chromosome karyotype and biological sex, we
label XX individuals as females and XY individuals as males, and view these labels as capturing
group differences in distributions of contexts for autosomal effects, rather than as a
dichotomy[14,17,32]. Throughout, we analyze GWAS on the raw measurement units as provided by
UKB. (See note on the rationale behind this choice in the section **Amplification of genetic**
effects is the primary mode of GxSex).
Among the 27 traits, we observed substantial discordance between males and females in
associations with the trait only for testosterone and waist:hip ratio (whether or not it is adjusted
for BMI; **Fig. S1**). For testosterone, as noted in previous analyses, associated genes are often in
separate pathways in males and females[18,20]. This is reflected in the small overlap of genes
neighboring top associations in our GWAS. For example, in females, the gene CYP3A7 is
involved in the hydroxylation of testosterone, resulting in its inactivation. In males, FKBP4 plays
a role in the downstream signaling of testosterone on the hypothalamus. Both genes, to our
knowledge, do not affect testosterone levels in the other sex.
For waist:hip ratio, we saw multiple associations in females only, such as variants near
ADAMTS9, a gene involved in insulin sensitivity[33]. As previous work established[18,20,21],
testosterone and waist:hip ratio are the exception, not the rule: Most traits did not display many
sex differences in top associations. For instance, arm fat-free mass, a highly heritable dimorphic
trait, showed near-perfect concordance in significant loci (**Fig. S1**). A previous study[21] examining
the concordance in top associations between males and females found few uniquely-associated
SNPs (<20) across the 84 continuous traits they studied; waist:hip ratio was an exception with

100 associations unique to one sex. Considering the evidence for the polygenicity of additive
genetic variation affecting many complex traits[34–36], it stands to reason that looking beyond lead
associations, through a polygenic prism, may aid in the characterization of non-additive effects
(such as GxSex) as well.

The limited scope of analyzing GxSex via heritability differences and genetic correlations.
We therefore turned to consider the polygenic nature of GxSex, first by employing commonly-
used approaches: comparing sex-specific SNP heritabilities and examining genetic correlations.
We used LD Score Regression (LDSC)[31,37] to estimate these for each trait. In most traits (17/27),
males and females had a genetic correlation greater than 0.9. Testosterone had the lowest
genetic correlation of 0.01, which suggests very little sharing of signals between males and
females (see similar results by Flynn et al.[20] and Sinnott-Armstrong et al.[18]).
For the majority of traits (18/27), male and female heritabilities were both greater than the
heritability in a sample that included both sexes. For instance, in arm fat-free mass (right), the
heritability in the both-sex sample was 0.232 (± 0.009), while the heritabilities for male and female
were 0.279 (± 0.012) and 0.255 (± 0.011), respectively. In particular, all body mass-related traits,
excluding BMI-adjusted waist:hip ratio, had greater sex-specific heritabilities **(Fig. 1)**.
In addition, we noticed a trend in which, as the genetic correlation decreased, the
difference between the heritabilities within each sex and in the sample combining both sexes
tended to become larger (Pearson r = -0.88, paired t-test p-value = 10^{-10}, **Fig. 1**). Nonetheless,
several traits with genetic correlation above 0.9 also present relatively large sex differences in
heritability: For example, diastolic blood pressure and arm fat-free mass (left) had differences of
5.2% (two-sample t-test p-value = $3 \cdot 10^{-6}$) and 3.4% (two-sample t-test p-value = 0.04),
respectively. These examples are incompatible with a model of pervasive uncorrelated genetic
effects driving sex-specific genetic contributions to variation in the trait (**Table 1**, second model).
We therefore considered two other alternative hypotheses under a simple additive model
of variance in a trait. Differences in heritability are either due to sex differences in genetic variance,
in environmental variance, or both. If genetic effects are similar, differences in environmental
variance alone could cause heritability differences (**Table 1**, first model). But as we show in the
Methods section, under such a model, the heritability in the combined sample cannot be smaller
than both sex-specific heritabilities.

Therefore, the observation of higher sex-specific heritabilities for most traits suggests that
the genetic variance must differ between males and females. Given the random segregation of
autosomal alleles, independent of an individual's sex chromosome karyotype, and assuming,
further, that there is little-to-no interaction of sex and genotype affecting participation in the UKB[38],
UKB allele frequencies in males and females are expected to be very similar. Thus, this
observation suggests that causal genetic effects differ between males and females for most traits
analyzed.

A last hypothesis that might tie together the observations in **Table 1** is a less appreciated
mode of GxSex, amplification. Namely, that the identity and direction of effects are largely shared
between sexes (leading to high genetic correlation), but the magnitude of genetic effects differs—
e.g., larger genetic effects on blood pressure in females—which in turn lead to differences in
genetic variance (**Table 1**, third model).

We can test the hypothesis that amplification acts systematically—across a large fraction
of causal variants—by examining the effects of polygenic scores (PGSs), genetic predictors of a
complex trait. Under this hypothesis, regardless of whether the PGS is estimated in a sample of
males, females, or a combined sample of both males and females, it should be predictive in both
sexes, since the causal variants and the direction of their effects are shared and the magnitude
is correlated (**Table 1**, third model). At the same time, in the sex for which genetic effects are
larger, the effect of the PGS is expected to be larger. To evaluate evidence for the systematic
amplification model, we estimated PGSs based on our sex-specific GWASs, and examined their
effect in both sexes. For some traits, like albumin and lymphocyte percentage, the effects of the
same PGS on trait value in males and females were statistically indistinguishable (**Fig. 2A,E,I,J**).
In a few other traits, such as diastolic blood pressure, the result was contingent on the sample in
which the PGS was estimated (**Fig. 2C,G,I,J**). However, for roughly half of the traits examined,
regardless of the sample in which the PGS was derived, the effect of the PGS was predictive in
both sexes yet significantly larger in one of the sexes (17/27 traits with t-test p-value < 0.05 using
the PGS derived from the males sample; 13/27 using the PGS derived from the females sample;
Fig. 2B,D,F,H,I,J). These observations are consistent with systematic amplification.

The results presented in **Figs. 1,2** suggested to us that various modes of polygenic GxSex
ought to be jointly evaluated. None of the hypothesized rules of thumb (**Table 1**) for interpreting
genetic correlations and sex differences in heritability worked across all traits (see also relevant

Figure 1: Heritabilities and Genetic Correlations Cannot Fully Distinguish Models of GxSex. (A) Genetic correlations between the male and females, estimated using bi-variate LD Score Regression, are shown in descending order. **(B)** The x-axis represents the relative heritability, i.e., the SNP heritability divided by the SNP heritability estimated in the sample with both sexes combined. Red asterisks show body-mass related traits with greater heritabilities in both sex-specific samples than in the sample combining both sexes.

Table 1. Polygenic Models of GxSex. We examine different models of the nature of GxSex in complex traits that link

to previous studies and motivations. Each model leads to different expectations from the analysis of heritability and

genetic correlations (**Fig. 1**). The illustrations in the third column depict examples of directions and magnitudes of

genetic effects, corresponding to each model. h_m^2, h_f^2 and h^2 denote narrow-sense heritabilities in males, females, and

a combined sample, respectively.

discussion in Khramtsova et al.[14]). This motivated us to estimate the covariance between genetic

effects in males and females directly. Another reason to treat covariance of genetic effects

themselves as the estimand of interest is that multiple, distinct GxSex patterns may exist across

subsets of genetic factors affecting a trait (**Table 1**, fourth model).

Flexible model of sex-specific genetic effects as arising from a mixture of covariance

relationships.   We set to infer the mixture of covariance relationships of genetic effects

among the sexes directly. We analyzed all traits in their raw measurement units as provided by

the UKB. In particular, we did not normalize or standardize phenotypes within each sex before

performing the sex-stratified GWAS, because sex differences in trait variance may be partly due

to amplification. Standardization would have therefore resulted in masking amplification signals

that may exist in the data. In some cases, this is indeed the purpose of standardization[39]. More

generally, while each scaling choice has it merits, we view the measurement of genetic effects in

their raw units as the most biologically interpretable.

Figure 2: Evaluating evidence for systematic amplification. (A-D) We regressed trait values in males (green) and separately in females (orange) on a polygenic score estimated in an independent sample of males. Points show mean values in one decile of the polygenic score; the fitted line and associated effect estimate and R^2 correspond to regressions on the raw, non-binned data. In some traits, like Albumin (A), the polygenic score has a similar effect on the trait in both sexes. In other traits (B,D), the estimated effect of the polygenic score differs significantly, consistent with a substantial difference in the magnitude of genetic effects of sites included in the polygenic score. **(E-H)** Same analysis as A-D, but with a polygenic score pre-estimated in an independent sample of females. **(I-J)** Summary of the ratio of the effect of the polygenic score on the trait (±2 SE) in males to the effect in females across physiological traits. See results for other traits in **Fig. S12**.

We used multivariate adaptive shrinkage (*mash*)[40], a tool that allows the inference of
genome-wide frequencies of genetic covariance relationships. Namely, we model the marginal
SNP effect estimates as sampled (with SNP-specific, sex-specific noise) from a mixture of zero-
centered Normal distributions with various prespecified covariance relationships (2x2 Variance-
Covariance matrices for male and female effects; Eq. 1 in Urbut et al.[40]). Our prespecified
covariance matrices ("hypothesis matrices") span a wide array of amplification and correlation
relationships, and use *mash* to estimate the mixture weights. Loosely, these weights can be
interpreted as the proportion of variants that follow the pattern specified by the covariance matrix
(**Fig. 3A**). Our covariance matrices ranged from -1 to 1 in between-sex correlation, and 10 levels
of relative magnitude in females relative to males, including matrices corresponding to no effect
in one or both sexes (**Fig. S2**).

We first focus on testosterone, for which previous research sets the expectation for
polygenic male-female covariance. In terms of magnitude, the vast majority of effects should have
much greater effect in males. In terms of correlation, we expect a class of genetic effects acting
through largely independent and uncorrelated pathways alongside a class of effects via shared
pathways[18]. Independent pathways include the role of hypothalamic-pituitary-gonadal axis in male
testosterone regulation and the contrasting role of the adrenal gland in female testosterone
production. Shared pathways involve sex hormone-binding globulin (SHBG), which decreases the
amount of bioavailable testosterone in both males and females. As expected, we found that
mixture weights for testosterone concentrated on greater magnitudes in males and largely
uncorrelated effects. Out of the 32% total weights on matrices with an effect in at least one sex,
98% of the weights were placed on matrices representing larger effects in males, including 20.4%
(± 0.7%) having male-specific effects (**Fig. 3, S5**).

Amplification of genetic effects is the primary mode of GxSex.   The only trait of the 27
where a large fraction (≥10%) of non-zero effects were negatively correlated was testosterone
(17%). Most effects were instead perfectly or near-perfectly correlated. For example, diastolic
blood pressure and eosinophil percentage had 66% (**Fig. 3**) and 68% (**Fig. S5**) of effects being
perfectly correlated, respectively. Overall, the low weights on matrices representing negative
correlation do not support opposite directions of effects being a major mode of GxSex (**Fig. S8**).

In some traits, such as hemoglobin A1C or diastolic blood pressure, previously considered non-
sex-specific because of high genetic correlations between sexes and a concordance in top GWAS
hits, we find evidence for substantial GxSex through amplification (**Fig. 3B,F; Fig. S5**)[20,21].
Furthermore, about half (13/27) of the traits analyzed had the majority of weights placed on
greater effects in just one of the sexes (x-axis in **Fig. 4A**). For instance, 92% of effects on BMI-
adjusted waist:hip ratio were greater in females and 92% of effects on (right) arm fat-free mass
were greater in males. Both traits had mixture weights concentrated on highly correlated effects
(**Fig. 3**). We confirmed, using a simulation study, that this summary of sex-biased amplification
indeed captures sex differences in the magnitude of genetic effects and that it is not due to
differences in the extent of estimation noise (e.g., variation in environmental factors independent
of genetic effects; **Figs. S6-7**; **Methods**).

Across traits, the difference between the fraction of male-larger effects and the fraction of
female-larger effects correlates strongly with male-to-female phenotypic variance ratio (Pearson
r = 0.873, p-value = $6 \cdot 10^{-9}$ after removing testosterone as an outlier; **Fig. 4A**). This observation is
consistent with our hypothesis of amplification leading to differences in genetic variance between
sexes and thereby contributing substantially to sex differences in phenotypic variance. Together,
these observations point to amplification, rather than uncorrelated effects, as a primary mode of
polygenic GxSex.

Another important question about the implication of pervasive amplification is whether it
is a major driver of mean phenotypic differences. The ratio between male and female phenotypic
means is correlated with the difference between male-larger and female-larger amplification
(Pearson r = 0.75; p-value = $2 \cdot 10^{-5}$ after removing testosterone and BMI-adjusted waist:hip ratio
as outliers). Though this correlation is intriguing, within-sex GWAS aims to explain individual
differences from the mean of the sex, and such GWAS results do not dictate the values of the sex
means. Further, both the ratio of mean trait values between sexes and the difference in
amplification are strongly correlated with phenotypic variance ratios (**Fig. 4A**; **Fig. S9**; see also
Karp et al.[8]), , and many different causal accounts could explain these correlations.

Finally, the pervasiveness of GxSex, alongside the mixture of covariance relationships
across the genome for many traits, may be important to consider in phenotypic prediction. We
compared the prediction accuracy of PGSs that consider the polygenic covariance structure to

Mixtures of covariance relationships
of genetic effects between males and females

Figure 3: Polygenic covariance structure between males and females. (A) Our analysis of the polygenic covariance

between males and females is based on sex-stratified GWAS. We modelled the sex-stratified GWAS estimates as

sampled with error from true effects arising from a mixture of possible covariance relationships between female and

male genetic effects. As an example, shown are illustrations for three possible relationships of the same qualitative

nature—perfectly correlated effects which are also larger in females—and the mixture weights estimated for each in

the case of diastolic blood pressure. **(B-F)** Each box shows the sum of weights placed on all covariance relationships

of the same qualitative nature, as specified by relative magnitude (horizontal axis) and correlation (vertical axis)
between male and female effects. The full set of pre-specified covariance matrices is shown in **Fig. S2**, and the weights
placed on each of them for each trait are shown in **Fig. S5**. All weights shown are percentages of non-null weights, i.e.,
the weight divided by the sum of all weights except for the one corresponding to no effect in either sex.

that of additive models, that ignore GxSex, as well as models that include GxSex but do not

consider the polygenic covariance structure (**Supplementary Materials; Fig. S13**). Indeed, for

most traits, models that consider the polygenic covariance structure outperform all other models

(18/27 traits, $p < 0.05$ for a conservative two-sample t(1)-test for 7/27 traits). Traits which showed

better prediction accuracy using the model that considered polygenic covariance structure

included many body mass-related traits such as BMI and whole body fat mass that also tended

to have higher sex-based amplification (**Fig. 4B;** Pearson $r = 0.62, p < 0.001$ between sex-

biased amplification and prediction accuracy ratio). These results point to the utility of considering

polygenic covariance structure in context-aware polygenic score prediction.

Testosterone as an amplifier.   Thus far, we treated the genetic interaction as discretely

mediated by biological sex. One mechanism that may underlie GxSex is a cue or exposure that

modulates the magnitude (and less often, the direction) of genetic effects, and varies in its

distribution between the sexes. A plausible candidate is testosterone. Testosterone may be a key

instigator since the hormone is present in distinctive pathways and levels between the sexes and

a known contributor to the development of male secondary characteristics, so therefore could

modulate genetic causes on sex-differentiated traits.

To test this idea, we first binned individuals of each sex by their testosterone levels. Then,

for each trait and within the bin, we quantified the magnitude of total genetic effect as the linear

regression coefficient of trait to a PGS for the trait (**Methods**; see **Fig. S15** for results obtained

using sex-specific PGS). For BMI, testosterone (mean per bin) and the magnitude of genetic

effect were correlated for both males and females (Pearson p-value < 0.05; **Fig. 5A**). For all body-

mass-related traits, there was a negative correlation between the magnitude of genetic effect and

testosterone levels for males and a positive correlation for females (**Fig. 5B**). Since the

relationship with testosterone remains contingent on sex, a model of testosterone as the sole

driver of the observed sex-specificity would be invalid. These observations may help explain

Figure 4. (A) Phenotypic variance strongly correlates with amplification. The x-axis summarized the "sex-biased amplification" of polygenic effects and is calculated by taking the difference between the sum of weights on matrices with male effects greater in magnitude than female effects (M>F) and the sum of weights of M<F matrices. The solid gray line shows a linear fit across traits, excluding testosterone as an outlier. **(B) Utility of polygenic GxSex model for trait prediction.** The x-axis shows the relative prediction accuracy estimated from the incremental R2 ratio of a

285 GxSex model informed by polygenic covariance patterns and an additive model. The phenotypes are ordered by the
286 relative prediction accuracy. The size of each point corresponds to the degree of sex-biased amplification as described
287 in (A). Blue points correspond to traits with a significant difference based on a two-sample t-test with p-value < 0.05
288 (**Text S8**) between the covariance-aware GxSex model and the additive model, one way or the other.

*r is the Pearson correlation

Figure 5. Amplification of total genetic effect in relation to testosterone levels. (A) The relationship between
testosterone level bins and estimated magnitude of genetic effect on traits is shown for three traits. The magnitude of
genetic effect is estimated using the slope of the regression of phenotypic values to polygenic scores in that bin. The
units on the y-axis are effect per standard deviations (SD) of the polygenic scores across all individuals in all bins. The
hollow data points are bins with overlapping testosterone ranges between males and females; these are based on
fewer individuals (~800 compared to ~2200 in other bins) and not included in the regression. **Fig. S14** show all other
traits analyzed. **(B)** The correlation for each sex (90% CI) are shown for all 27 traits. Traits are ordered in descending
order of male-female differences in Pearson correlation.

previous reports of positive correlations between obesity and free testosterone in women, and negative correlations in men[41]. We conclude that in body-mass related traits, testosterone may be modulating genetic effects in a sexually antagonistic manner.

We performed two additional analyses designed to control for possible caveats to the association of testosterone and the magnitude of polygenic effect: An association test that controls for possible confounding with age (**Fig. S17**) and a test that mitigates confounding with other variables or reverse causality (wherein the magnitude of genetic effect affecting the focal trait causally affecting testosterone levels; **Fig. S16**). The evidence for an effect of testosterone on the magnitude of polygenic effect did not remain significant in either of these tests. It is possible, however, that this was due to low statistical power (**Methods**).

Are polygenic and environmental effects jointly amplified? Our results thus far suggest that polygenic amplification across sexes is pervasive across traits; and that the ratio of phenotypic variance scales with amplification (**Fig. 4A**). An immediate question of interest is whether the same modulators that act on the magnitude of genetic effects act on environmental effects as well (see also relevant discussion by Domingue et al.[42]). Consider the example of human skeletal muscle. The impact of resistance exercise varies between males and females. Resistance exercise can be considered as an environmental effect since it upregulates multiple skeletal muscle genes present in both males and females such as IGF-1, which in turn is involved in muscle growth[43]. However, after resistance exercise at similar intensities, upregulation of such genes is sustained in males, while levels return sooner to the resting state in females (**Fig. S18**). It is plausible that modulators of the effect of IGF-1, such as insulin[44] or sex hormones[45,46], drive a difference in the magnitude of effect of core genes such as IGF-1 in a sex-specific manner. To express this intuition with a model: If amplification mechanisms are shared, then amplification may be modeled as having the same scalar multiplier effect on genetic and environmental effects (**Fig. 6A**). In the **Methods** section, we specify the details of a null model of joint amplification, which yields the prediction that the male-female ratio of genetic variances should equal the respective ratio of environmental variances (blue line in **Fig 6B**).

This expectation is qualitatively different from those of two longstanding theoretical "rules of thumb" predictions for sex differences in trait variance (**Supplementary materials**; **Fig. S19A;**

of. Zajitschek et al.[47]): The "greater male variability" and "estrus-mediated variability" models,
which provide a poor fit across the 27 physiological traits analyzed (**Fig. S19B**).

We tested the fit of the theoretical prediction under pervasive joint amplification across
traits. We used our estimates of sex-specific phenotypic variance and SNP heritabilities to
estimate the ratios of genetic and environmental variances. We note that environmental variance
is proxied here by all trait variance not due to additive genetic effects, and caution is advised with
interpretation of this proxy. Twenty of the 27 traits were consistent with the null model of pervasive
joint amplification (within 90% CI; **Fig. 6B**). This finding may suggest a sharing of pathways
between polygenic and environmental effects for these traits (**Fig. 6A**). Interesting exceptions
include diastolic blood pressure—which was the strongest outlier (p-value = $3.06 \cdot 10^{-12}$, single-
sample z-test), excluding testosterone.

Sexually antagonistic selection.   A hypothesized cause of sexual dimorphism is sexually
antagonistic selection, in which some alleles are beneficial in one sex yet deleterious in the
other[4,5,7,48,49]. Sexually antagonistic selection is difficult to study using traditional population
genetics methods because Mendelian inheritance equalizes autosomal allele frequencies
between the sexes at conception, thereby erasing informative signals. One way around this
limitation is to examine allele-frequency differences between the sexes in the current generation,
known as "selection in real time"[7,50,51]. In this section, we consider a model of sexually antagonistic
selection acting on a polygenic trait and use it to estimate the strength of contemporary viability
selection acting on the 27 traits we analyzed.
Most theoretical models of sexually antagonistic selection on a trait under stabilizing selection
usually posit either highly distinct male and female fitness optima or genetic variants affecting
traits antagonistically. Our findings on pervasive amplification suggest that variant effects on traits
tend to have concordant signs. Yet, under pervasive amplification, a somewhat surprising intuition
arises: Alleles affecting a trait may frequently experience sexually antagonistic selection—both in
the case in which trait optima for males and females are very distinct (**Fig. 7B**) and for the case
in which they are similar (**Fig. 7A**).
We developed a theoretical model of sexually antagonistic viability selection on a single
trait that builds on this intuition. The model relates sex-specific effects on a complex trait to the
divergence in allele frequency between males and females (measured as F_{ST}[52,53]) due to viability

Figure 6. Testing a model of pervasive, joint amplification of environmental and polygenic effects. **(A)** A model of equal amplification of genetic (G) and environmental (E) effect, that produces the sex differences in the distribution of the phenotype, Y. G and E both act through a core pathway that is amplified in a sex-specific manner. **(B)** The blue 1:1 line depicts the theoretical expectation under a simple model of equal amplification of genetic and environmental effects in males compared to females. Error bars show 90% confidence intervals. Traits in blue are consistent (within their 90% CI) with the theoretical prediction. **Fig. S19** shows the same data alongside the predictions under other theoretical models of male-female variance ratios.

selection "in real time", i.e., acting in the current generation between conception and the time of sampling. We derive the expected relationship for each site i,

$$F_{ST_i} \approx AV_{GxS_i}, \qquad (1)$$

where

$$V_{\{GxS\}_i} = 2p_i(1 - p_i)\left(\beta_i^m - \beta_i^f\right)^2,$$

and p_i, β_i^m and β_i^f are the allele frequency of an allele at site i, its effect on the trait in males and its effect in females, respectively. A is a constant parameter shared across all variants and can therefore be interpreted as the effect of sexually antagonistic selection on male-female divergence

Figure 7. Testing for sexually antagonistic selection. (A,B) A model of sexually antagonistic selection. Selection coefficients, s_m and s_f, are linear with the additive effect on the trait in each sex. Sexually antagonistic selection acts

such that $s_m = -s_f$. The model yields the prediction of **Eq. 1.** In (A), the effect of an allele tends to drive trait values

towards the optimum in males, and away from the optimum in females. In (B), the fitness optima are farther in males

and females; in both examples, selection on acts antagonistically (i.e., in opposite directions). **(C)** Two examples of the

weighted least-squares linear regression preformed to estimate the strength of sexually antagonistic selection on

variants associated with a trait (A in panel A and **Eq. 1**). Each point shows one SNP. Size is proportional to each point's

regression weight. **(D)** Z-scores (90% non-parametric bootstrap CI) estimated through 1000 resampling iterations of

the weighted linear regression of panel B for each trait. The two colored estimates correspond to the examples in (B).

at variants associated with the trait (**Methods**). We estimated F_{ST_i} for all sites i across subsamples

of various ancestry groups in the gnomAD dataset [54]. To estimate $V_{\{GxS\}_i}$ at each site and for each

trait, we used our sex-stratified GWAS results. Since there is large heterogeneity in uncertainty

of GxSex-genetic variance estimates, we use a variance-weighted linear regression to estimate

A (see **Methods** for the derivation of the variance of $V_{\{GxS\}_i}$ estimates and **Supplementary**

Materials for further details).

Recent work has shown that apparent sex differences in autosomal allele frequencies

within a sample are often due to a bioinformatic artifact: The mismapping of sequencing reads

from autosomes to sex chromosomes or vice versa[48,55,56]. We identified and excluded sites which

are potentially vulnerable to this artifact (**Supplementary Materials**). In **Fig. 7D,** we only show

results for gnomAD subsamples that are the closest in their genetic ancestry to our UKB sample[57]

(results for other subsamples are shown in **Fig. S20,21**). Furthermore, given the concerns of study

recruitment biases [38,55], we place higher confidence in results that replicate qualitatively across

different subsamples, even though we note that subsample-specific selection signals may be real

since sexually antagonistic selection may act heterogeneously across groups.

With these conservative criteria considered, we only find evidence for sexually

antagonistic polygenic selection on testosterone. In the non-Finnish sample, the largest of the

three samples, the null hypothesis $H_0: A = 0$ in **Eq. 1** is rejected (p-value < 0.05) only for

testosterone (Z score = 2.2). Testosterone is among the three strongest signals in the two other

samples as well, though none of the traits are statistically significant in these samples.

Discussion

Departing from previous studies that sought GxSex through single loci or heritability analyses, we modelled GxSex as a mixture of polygenic relationships across the genome. Our analysis supports pervasive context-dependency of genetic effects on complex traits, acting largely through amplification. Surprisingly, even for some traits such as red blood cell count, previously considered non-sex-specific because of high genetic correlations between sexes and a concordance in top GWAS hits, we find evidence for substantial GxSex. The strong relationships we find between amplification, environmental variance and phenotypic variance further points to its potential importance for sex differences.

We have shown that considering the polygenic covariance structure, including amplification signals, improves phenotypic prediction for most traits. Its incorporation in polygenic scores is straightforward. We therefore recommend its broad application and further building on our approach to improve clinical risk stratification and other applications of polygenic scores.

Our findings may seem at odds with previous reports of GxSex primarily consisting of sex-limited effects (i.e., no effect in one of the sexes) or antagonistic effects (differences in sign)[58]. In the **Supplementary Materials** and **Table S6**, we illustrate that these apparent discrepancies may be rooted in ascertainment biases. Therefore, limiting analyses to variants with outsized sex differences provides a clouded picture of polygenic GxSex.

Localization of GxSex signals can provide clues into the modulators underlying amplification. Here, we proposed one such modulator, testosterone, and found a correlation between testosterone levels and the magnitude of genetic effect on whole body fat mass. The opposite signs of these correlations in females and males may reflect the discrepant relationship between testosterone and these traits at the phenotypic level.

Our approach for studying GxSex in complex physiological traits can be adopted to study the moderation of polygenic effects by other environments. Starting out with sex as an environmental variable offers a methodological advantage. The study of context-dependency in humans is often complicated by study participation biases, leading to genetic ancestry structure that confounds genotype-phenotype associations[38,59–61], reverse causality between the phenotype and environment variable, collider bias, gene-by-environment correlation and other problems[62–64]. Focusing on sex as a case study circumvents many of these "usual suspects" problems: For example, problems involving the phenotype causally affecting sex are unlikely. This

is an important benchmark for future studies of environmental modulation, both because of the

methodological advantage of sex as an environmental variable and because sex is almost always

measured; so insight into sex differences in genetic effects can be incorporated straightforwardly

in future studies and in clinical risk prediction. Here, we showed that for most of the traits

considered, modeling polygenic GxSex (as opposed to individually estimating sex-specific effects

at each site; **Fig. S13** yields sex-specific predictors that outperform standard additive polygenic

scores.

Finally, we developed a model to consider how GxSex may fuel sexually antagonistic

selection in contemporary populations. Over long evolutionary timescales, the two scenarios

depicted in **Fig. 7A,B** may lead to different predictions about the long-term maintenance of GxSex

genetic variance. Regardless, in both cases, alleles that underlie GxSex may experience sexually

antagonistic selection.

We found suggestive signals of sexually antagonistic selection on variation associated

with testosterone levels (also see related results by Ruzicka et al.[51]). The signal for our inference

of selection is systematic allele frequency differences between adult males and females, which

are consistent with contemporary viability selection. The severity, age of onset and prevalence of

nearly all diseases are sexually dimorphic[65]. These signals may therefore point to a related

disease that differentially affects lifespan in the two sexes, such as immune system suppression,

diabetes, cancers, and hypertension[66–69]. Recently, high testosterone levels have been linked to

increased rates of mortality and cancer in women, but decreased rates in men[70,71]. However, the

testosterone result is also consistent with other accounts, such as testosterone having opposing

effects on propensity to participate in a study in the two sexes. Further validation is therefore

required to better test hypotheses of sexually antagonistic selection, for example in studies with

no recruitment biases (or at least distinct recruitment biases).

In this work, we have shown that amplification of the magnitude of polygenic effects may

be important to consider as a driver of sex differences and their evolution. Our approach included

the flexible modelling of genetic effect covariance among the sexes, as well as various

subsequent analyses exploring the implications of these covariance structures. We hope this

study can inform future work on the context-specificity of genetic effects on complex traits.

Limitations of the Study

Study participation in large biobanks like the UK Biobank (UKB) differs by sex[72]; and work by Piratsu et al. further argued that allele frequency differences between males and females may reflect sex-specific recruitment biases[55]. However, a recent study by Benonisdottir and Kong found no evidence of sex-specific genetic associations with UKB participation[38], and another by Kasimatis et al. showed that many apparent associations of autosomal genotypes and biological sex in the UKB were instead primarily due to a bioinformatic artifact—the mis-hybridization of autosomal genotyping probes with sex chromosomes[48]. Even still, subtle recruitment biases affecting male and female participation differently remain a possible caveat to our conclusions. For the analysis of natural selection in particular, while the replication of signals of selection in multiple samples may lend credence to our inference, medical datasets based on recruitment of participants via referring physicians, participation biases may still plausibly be shared across studies.

Acknowledgments

We thank the Harpak Lab and Edge Lab members, Ziyue Gao, Tom Juenger, Jonathan Pritchard, Molly Przeworski, Guy Sella, Jeff Spence and Elliot Tucker-Drob for helpful comments on the manuscript. We also thank Brian Dilkes, Andrés Bendesky and Jim Fleet for insightful discussions. We thank Abin Abraham and Tony Capra for their help in implementing code from Kasimatis et al.[48], and Michelle Traglia and Lauren Weiss for useful discussions of the relationship between the results reported here and Traglia et al.[58]. This work was supported by NIH GM116853-07 to M.K. and NIH GM137758 to M.D.E. This study has been conducted using the UK Biobank resource under application Number 61666, as approved by the University of Texas at Austin Institutional Review Board, protocol 2019-02-0125.

Author Contributions

C.Z., M.J.M. and A.H. designed the experiments. C.Z. and M.J.M. performed the experiments. C.Z., M.J.M. and A.H. wrote the paper with assistance from all authors. J.M.C., M.D.E and M.K. provided expertise and feedback.

Declaration of Interests

497   The authors declare no competing interests.

Methods

METHOD DETAILS

UK Biobank sample characteristics.   The UK Biobank is an extensive database that contains a wide variety of phenotypic and genotypic information of around half a million participants aged 40-69 at recruitment[73].

In this study, we considered 337,111 individuals who passed quality control (QC) checks, which included the removal of samples identified by the UK Biobank with sex chromosome aneuploidy or self-reported sex differing from sex determined from genotyping analysis. We excluded related individuals (3rd-degree relatives or closer) as identified by the UK Biobank in data field 22020. To reduce potential population structure confounding, we further limited our sample to individuals identified by the UK Biobank as "White British" in data field 22006. These

are individuals who both self-identified as White and as British and were additionally very tightly
clustered in the genetic principal component space[73,74]. Individuals who had withdrawn from the
UK Biobank by the time of this study were removed. For each phenotype, we also removed
individuals who had missing data for the specified phenotype. These procedures left us with
between 255,426 to 336,551 individuals in the analysis for each trait.

Expectations for sex-specific heritabilities with no GxSex.   In the section "The limited
scope of analyzing GxSex via heritability differences and genetic correlations," we report our
observation that, for most traits examined, sex-specific heritabilities (i.e., estimated independently
from sex-stratified GWAS) were both higher than the heritability in the combined sample. Here,
we explain why this observation is inconsistent with a simple model in which genetic effects are
the same across the sexes.

Under a simple additive model of variance in a trait Y within each sex Z,

$$Var[Y|Z] = Var[G|Z] + Var[E|Z], \qquad (2)$$

540 where Y, G, E represent the trait value, additive effect, and environmental effect (including all non-
541 genetic context aside from sex), respectively. Under this model, the sex-specific heritability h_z^2 is

$$h_z^2 = \frac{Var[G|Z]}{Var[G|Z] + Var[E|Z]}. \qquad (3)$$

Therefore, sex differences in heritability are either due to sex differences in genetic
variance, in environmental variance, or both. If genetic effects are equal, differences in
environmental variance alone could cause heritability differences (**Table 1**, first model). But as
we show below, the heritability in the combined sample cannot be smaller than both sex-specific
heritabilities.

We assume as before that allele frequencies are highly similar between males and
females. Since genetic effects are equal, this implies

$$Var[G|Z = m] \approx Var[G|Z = f].$$

550 For the environmental variance, we have that

$$Var[E] = \mathbb{E}_Z\big[Var[E|Z]\big] + Var_Z\big[\mathbb{E}[E|Z]\big] = \mathbb{E}_Z\big[Var[E|Z]\big] + 0 = \qquad (4)$$

$$\mathbb{P}(Z = m)Var[E|Z = m] + \mathbb{P}(Z = f)Var[E|Z = f] \leq \max_{z \in \{m,f\}} Var[E|Z = z].$$

The first equality follows from the law of total variance. In the second equality, we have

assumed that there are no mean sex differences in the environmental effects (or, in practice in

our analysis and as routine in other analyses, that mean phenotypic sex differences have been

subtracted out), giving

$$\mathbb{E}[E|Z = m] = \mathbb{E}[E|Z = f] = \mathbb{E}[E].$$

Eq. 4 shows that the combined environmental variance cannot be greater than the larger of the

two sex-specific environmental variances. It follows that if the genetic variance is equal in both

sexes, then the heritability in the combined sample cannot be smaller than both of the sex-specific

heritabilities,

$$h^2 = \frac{Var[G]}{Var[G] + Var[E]} \geq \frac{Var[G]}{Var[G] + \max_{z \in \{m,f\}} Var[E|Z]} = \min_{z \in \{m,f\}} h_Z^2. \qquad (5)$$

Multivariate adaptive shrinkage (mash).   We used multivariate adaptive shrinkage (*mash*) to

examine correlation and differences in magnitude of SNP effects between males and females [40].

mash is an adaptive shrinkage method[75] that improves upon previous methods of estimating and

comparing effects across multiple conditions by flexibly allowing for a mixture of effect covariance

patterns between conditions and requiring only summary statistics from each condition (including

a point estimate of the effect and corresponding standard error for each SNP and condition). The

method adapts to patterns of sparsity, sharing, and correlation among the conditions to compute

improved effect estimates.

In this study, we set two conditions, male and female, and provided effect estimates and

corresponding standard errors from our male-specific and female-specific GWAS. *mash* learns

from the data by estimating mixture proportions of various predefined covariance matrices

representing different patterns in effects. Using maximum likelihood, mash assigns low weights

to matrices that capture fewer patterns in the data, and higher weights to those that capture more.

Mixture weights for covariance structure between male and female effects.   To interpret

patterns of SNP effects between males and females, we inputted 66 hypothesis-based covariance

matrices (**Fig. S2**) spanning a range of correlations and relative magnitudes of effects between
males and females. We used a random subset of all SNPs for mash to learn the covariance
mixture weights. In order for the random subset to contain approximately independent SNPs and
capture the weight of SNPs with no effect (**Fig. S2**), we created a subset of SNPs for each trait
by taking a random SNP from each of 1703 approximately independent LD blocks estimated for
Europeans[76]. *mash* can also generate data-driven covariance matrices that capture SNP effects
in the data, but we did not use this feature since the data-driven matrices had negligible
differences from our hypothesized matrices (in terms of $\ell2$ norm) and were less interpretable.
For each trait, we repeat this weight-learning step 100 times, sampling the SNPs from the 1703
LD blocks without replacement to fit the mash model and generate mixture proportions. We then
take the average proportion for each covariance matrix as an estimate of its weight, effectively
treating each of the 100 samples as i.i.d. draws.

Choice of SNPs used to estimate male-female effect covariance.   We   examined   the
effect of using a random subset taken from different p-value thresholds [1, 5e-2, 1e-5, 5e-8] while
selecting from LD blocks. By doing so, we can examine differences in the distribution of weights
across the p-value thresholds. We performed this test on height, BMI, testosterone, and BMI-
adjusted waist:hip ratio. For each trait, weight placed on the no-effect matrix decreased as we
reduced the p-value threshold (**Fig. S4A**). Patterns of weights for non-null effect matrices varied
across the traits (**Fig. S4B,C**). Since *mash* considers the proportion of null effects and sex-
specific, SNP-specific noise; together with the fact that for complex traits, less significant
associations may still reflect valuable signal, we decided on using the whole set of SNPs to
sample from when estimating mixture proportions.

Simulating equal genetic effects and heterogeneous estimation noise among the sexes.
602   To ensure that *mash* was not mistaking sex differences in estimation noise (e.g. via
differences in the extent of environmental variance) to be differences in the magnitude of genetic
effects, we performed a simulation study. In short, samples of males and females were generated
under the model given by **Eq. 2**. Genetic effects were set as equal, but the environmental variance
differed among the sexes. We then perform a GWAS on both samples and input the simulated
GWAS results into *mash*, and test whether the estimated mixture weights spuriously suggest the

presence of GxSex. We performed this simulation on a grid of parameters, including heritabilities

in males set to either 5% or 50%, female to male environmental variance ratio of 1, 1.5 or 5; and

100, 1,000 or 10,000 causal SNPs.

First, we created a sample of 300K individuals with randomly assigned sex. We then

sampled genotypes for all individuals consisting of 20K SNPs by sampling from the observed

distribution of allele frequencies from UK Biobank's imputed data[77], assuming linkage equilibrium.

From the 20K SNPs, we portioned out the predetermined number of causal SNPs and assigned

effect sizes by sampling from a Standard Normal distribution. We estimated the male

environmental variance for each causal SNP using the equation,

$$Var[E|Z = m] = \frac{Var[G|Z = m](1 - h_m^2)}{h_m^2} = \frac{(\sum_{i=0} \beta_i^2 2p_i(1 - p_i))(1 - h_m^2)}{h_m^2} \quad (6)$$

where $Var[E||Z = m]$ is the simulated environmental variance for males, $G|Z = m$ is a vector of

the genetic effects in males, h_m^2 is the heritability in males and β_i and p_i are the effect size and

allele frequency at site i, which are equal for males and females. We multiplied $Var[E||Z = m]$

by the predetermined environmental variance ratio to obtain the environmental variance for

females $Var[E||Z = f]$. Afterwards, for each individual j with sex z_j, we sampled the

environmental effect E_j as

$$E_j \sim N(0, Var[E|Z = z_j].$$

623   Phenotypes were then set using the following additive model,

$$y_j = \sum_{i=0} \beta_i x_{ij} + E_j \quad (7)$$

where y_j is the phenotypic value for individual j and x_{ij} is the number of effect allele copies at the

i^{th} causal SNP for the j^{th} individual. With the phenotype, genotype and environmental effect set,

we obtained the estimated effect sizes, $\{\widehat{\beta_i}\}$, using least squares simple linear regression for all

20K SNPs and used the estimated effect sizes and corresponding standard errors as input into

mash.

For nearly all parameters, out of the weights on matrices other than the null matrix, the

vast majority was placed on the matrix for perfect correlation, equal magnitude (**Fig S6**). As the

number of causal SNPs increased, the weight on the no-effect covariance matrix decreased

accordingly. These results suggest that *mash* was not grossly mistaking differences in
environmental variance as amplification.

Simulating sex-biased amplification. To evaluate whether *mash* accurately
captures sex-biased amplification of genetic effects (a measure we have used in the x-axis of **Fig.**
4A,B), we followed the same simulation procedure described in the Section "Simulating equal
genetic effects and heterogeneous estimation noise among the sexes". However, instead of using
equal genetic effects in males and females, we sampled genetic effects from pre-specified
covariance matrices (**Fig. S7** left-hand panel). We set the female to male environmental variance
ratio as 1.2 and the heritability as 0.5. We generated data from (A) a model in which all genetic
effects are sampled from a matrix where male and female effects are equal, (B) a model in which
86% of the genetic effects are sampled from a matrix where effects between the sexes are equal,
and 14% of the effects are sampled from a matrix where the female effect size magnitude is 4
times that of males, and (C) a model in which 86% of effects are sampled from a matrix where
effects between sexes are equal, and 14% of effects are sampled from a matrix of only female-
specific effects. After simulating sex-specific GWAS on the three models, we input the results into
mash to estimate mixture weights. We repeated this simulation procedure 100 times for each
model.

For model (A), the equal effect matrix received 78% of the weight, and the difference
between male-larger and female-larger magnitude was 1% (**Fig. S7**). For model (B), 67% of the
weight was placed on the matrix for equal effects. The weight difference between male-larger and
female-larger magnitude was 13%. In model (C), 69% of the weight was on the matrix for equal
effects, and the difference between male-larger and female-larger magnitude was 16%. These
simulation results therefore suggest some overestimation of the proportion of SNPs with
magnitude differences. However, the measure of "sex-biased amplification" matched that of the
pre-specified generative models up to an error of 2%. Therefore, the simulations suggest "sex-
biased amplification" is measured accurately in our estimation procedure.

Testosterone as an amplifier. We tested a model of testosterone as a modulator of
magnitude differences in males and females. We first split individuals by sex and for each sex,
created 10 bins of testosterone levels. We adjusted one of the 10 bins to have testosterone levels

overlap between males and females. The overlapping testosterone bin was based on fewer
individuals (~800) compared to the other bins (~2200). For each trait, each of the sexes, and
within each bin, we performed a simple linear regression of trait values to the PGS for the trait
(using a PGS based on both-sex summary statistics (**Supplementary Materials**)). We interpret
the estimated coefficient for the effect of the PGS as a proxy for the magnitude of polygenic effect.
Finally, we summarized the relationship between testosterone level and magnitude of polygenic
effect across bins using the Pearson correlation between the two.

To mitigate the possible effects of confounding (of testosterone and magnitude of
polygenic effect) or reverse causation (the magnitude of polygenic effect on the focal trait causally
affecting testosterone levels) we employed a version of Mendelian Randomization[78,79] of the same
analysis (**Fig. S16**). Namely, we replaced testosterone levels of each individual with their PGS for
testosterone. Here, given the near-zero genetic correlation between males and females, we used
our sex-specific PGS for each sex; otherwise, the analysis is unchanged.

We also examined whether participants' age may have confounded the relationship
between testosterone and polygenic effect. In this analysis, instead of using the polygenic effect
as the response variable across bins, we used the polygenic effect residualized for mean age in
the bin and examined the effect of an individual's polygenic score on the residual (**Fig. S17**).

Model of Shared Amplification. Here, we suggest a null model in which amplification is
shared between genetic and environmental effects. We then suggest a prediction that the model
yields and explain how we tested this prediction across traits (**Fig. 6**).

If an amplifier is shared, it may be modeled as having the same scalar multiplier effect on
genetic and environmental effects. Consider the within-sex additive model of **Eq. 1** in the section
"The limited scope of analyzing GxSex via heritability differences and genetic correlations" above.
For a phenotype value Y_z in sex $z \in \{m, f\}$

$$Y_z = c + G_z + E_z, \tag{8}$$

688 Where c is a constant, E_z is the environmental effect and

$$G_z = \sum_{site\ i} x_i \beta_i^z \tag{9}$$

689   is the polygenic effect where β_i^Z is the effect of an allele at site i (say the minor allele) in sex Z

690   and x_i is the number of copies of the allele. We assume here for simplicity that male genetic

691   effects relate to female effects solely through a shared polygenic amplification constant, α,

$$\beta_i^m = \alpha\beta_i^f \quad \forall i; \; \alpha > 0. \tag{10}$$

692   Allele frequencies are once again assumed to be close to equal between males and

693   females, since due to random segregation of alleles during meiosis, genotype frequencies at

694   autosomal sites are independent of sex; and further assuming no substantial interaction between

695   genotype and sex affecting participation in UKB[38]. Consequently, differences in polygenic effect

696   distributions between males and females are solely based on GxSex, and thus:

$$Var[G_m] = \alpha^2 Var[G_f]. \tag{11}$$

697   The model we would like to test is one where the amplification of environmental effects

698   can also be simplified to the same scalar multiplier,

$$E_m = \alpha E_f, \text{ and} \tag{12}$$

$$Var[E_m] = \alpha^2 Var[E_f].$$

699   Hence, with equal amplification,

$$\frac{Var[G_m]}{Var[G_f]} = \frac{Var[E_m]}{Var[E_f]} \tag{13}$$

701   To test the model of shared amplification between environmental and polygenic effects

702   (**Eq. 8**) we obtained the genetic and environmental variance for males and females based on the

703   following relationships,

$$Var[G_z] = h^2 Var[Y_z] \tag{14}$$

704   and

$$Var[E_z] = (1 - h^2)Var[Y_z], \tag{15}$$

where $Var[G_z], Var[E_z]$, and $Var[G_z]$ are the additive genetic, environmental, and phenotype

variances, respectively. Estimates of the sex-specific heritabilities, h_z^2, were obtained from

previous estimates using LD Score Regression (**Supplementary Materials**).

Representing male genetic or environmental variance as x, and the corresponding female

variance as y, we derived standard errors for the ratio of male to female variance using the 2$^{\text{nd}}$-

order Taylor approximation for the standard error of a ratio of estimators of x and y,

$$SE\left[\frac{\hat{x}}{\hat{y}}\right] = \sqrt{Var\left[\frac{\hat{x}}{\hat{y}}\right]} \cong \frac{E[\widehat{x}]}{E[\widehat{y}]}\sqrt{\frac{Var[\hat{x}]}{E[\hat{x}]^2} + \frac{Var[\hat{y}]}{E[\hat{y}]^2} - \frac{2Cov[\hat{x},\hat{y}]}{E[\hat{x}]E[\hat{y}]}} \approx \frac{\hat{x}}{\hat{y}}\sqrt{\frac{SE[\hat{x}]^2}{\hat{x}^2} + \frac{SE[\hat{y}]^2}{\hat{y}^2}} \tag{16}$$

711   assuming independence between $\hat{x}$ and $\hat{y}$ since they are statistics of independent sampling

712   distributions (independent samples of males and females). The standard errors of the genetic and

713   environmental variance were estimated using the law of total variance for a product of two random

714   variables. For $\hat{a}$ and $\hat{b}$, unbiased estimators of the two parameters a and b, respectively, we get

$$SE[\hat{a}\hat{b}] = \sqrt{SE[\hat{a}]^2 SE[\hat{b}]^2 + E[\hat{a}]^2 SE[\hat{b}]^2 + E[\hat{b}]^2 SE[\hat{a}]^2}.$$

715   Plugging in the point estimate $\hat{a}$ for $E[\hat{a}] = a$ and the point estimate $\hat{b}$ for $E[\hat{b}] = b$,

$$\widehat{SE}[\hat{a}\hat{b}] = \sqrt{SE[\hat{a}]^2 SE[\hat{b}]^2 + \hat{a}^2 SE[\hat{b}]^2 + \hat{b}^2 SE[\hat{a}]^2}. \tag{17}$$

716   In this case, a represents the phenotypic variance, and b represents either h_z^2 for

717   estimation of genetic variance or $(1 - h_z^2)$ for estimation of environmental variance. Lastly, to

718   obtain the standard error of the phenotypic variance, we used bootstrapping with 100 samples

719   (with replacement) of estimates of the phenotypic variance in sex z,

$$\widehat{SE}[\hat{a}] = \sqrt{\frac{\sum_{i=1}^{100}\left(Var[Y]_i - \widehat{Var}[Y_z]_i\right)^2}{100-1}}$$

720   Finally, for each trait, we estimated $\tilde{Z}$, the ratio of the two male-female ratios

721   (environmental and genetic, y and x axes in **Fig. 6**, respectively), and its standard error, $SE[\tilde{Z}]$,

722   using the same method as in **Eq. 16**. Under the null hypothesis of equal environmental and

723   genetic amplification (**Eq. 8**),

$$H_0: E[Z] = 0, \tag{18}$$

724   where

$$Z = \frac{\tilde{Z}-1}{SE[\tilde{Z}]}.$$

In **Fig. 6,** we approximated 90% confidence intervals on Z by treating it as a Z score, i.e.,

further treating Z as a Standard Normal.

A Model of Sexually antagonistic Selection.   We developed a model relating sex

differences in additive effects on a trait at a biallelic locus (β_m and β_f) and divergence in allele

frequencies. Our model resembles that of Cheng and Kirkpatrick[7] who developed a similar model

relating allele-frequency differences and sex bias in gene expression. In short, we modelled

sexually antagonistic, post-conception viability selection on a focal complex trait. We assumed

allele frequencies in adult males, p_m, and adult females, p_f, are at equilibrium, i.e. do not change

in consecutive generations. Under these conditions, we derive the relationship

$$F_{ST} \approx AV_{GxSex},$$

735   where F_{ST}[52] is the fixation index with respect to the male and female subpopulations, i.e., the

736   proportion of heterozygosity in the population that is due to allelic divergence between the sexes.

737   V_{GxSex} is defined as

$$V_{GxSex} := 2p(1-p)\left(\beta_m - \beta_f\right)^2, \tag{19}$$

738   where p is the allele frequency in zygotes. A is a parameter that, importantly, is shared across all

739   variants affecting the trait and can be thought of as the intensity of sexually antagonistic selection

740   acting on genetic variation for the trait in question.

741   In our model, allele frequencies at the autosomal locus are assumed to be equal in males

742   and female zygotes. F_{ST} at adulthood takes the form

$$F_{ST} := \frac{Var_z[p_z]}{\bar{p}(1-\bar{p})} = \frac{E[p_z^2] - \bar{p}^2}{\bar{p}(1-\bar{p})} = \frac{p_m^2 + p_f^2 - \left(\frac{p_m + p_f}{2}\right)^2}{\bar{p}(1-\bar{p})} = \frac{\left(p_m - p_f\right)^2}{4\bar{p}(1-\bar{p})}, \tag{20}$$

743   where

$$\bar{p} = \frac{p_m + p_f}{2}.$$

745   If we further assume a near-1:1 sex ratio such that $\bar{p} \approx p$,

$$F_{ST} \approx \frac{\left(p_m - p_f\right)^2}{4p(1-p)}. \tag{21}$$

Sexually antagonistic selection acting on viability will cause divergence in allele

frequencies between adult males and females. We write the relative viabilities of the homozygote

for the reference allele, the heterozygote and the homozygote for the effect allele as $1 :: 1 +$

$d_z S_z :: 1 + S_z$ for each sex $z \in \{m, f\}$. The selection coefficient S_z and dominance coefficient d_z

can be frequency-dependent, in which case these coefficients take their values at equilibrium. We

can write the additive selection coefficient of the effect allele as

$$s_z = [p + (1 - 2p)d_z]S_z. \tag{22}$$

752 Assuming that zygotes are at Hardy-Weinberg equilibrium, the allele frequency in each

753 sex at adulthood is

$$p_z \approx p + p(1 - p)s_z, \tag{23}$$

754 where we neglected terms of order s_z^2 [80]. Plugging **Eq. 23** into **Eq. 21**, the divergence between

755 males and females post-selection is

$$F_{ST} \approx \frac{1}{4}p(1 - p)\left(s_m - s_f\right)^2. \tag{24}$$

756 We model the strength of viability selection acting on males and females as linear with the

757 additive effect on a focal trait in each sex,

$$s_z = a_z\beta_z, \tag{25}$$

758 and recalling the simplifying assumption that allele frequencies are at equilibrium under sexually

759 antagonistic viability selection at the locus, such that selection favoring an allele in one sex is

760 balanced by selection against that allele in the other sex,

$$s_f = -s_m. \tag{26}$$

761 If $\beta_m = \beta_f$, then **Eq. 24** simplifies to

$$F_{ST} \approx p(1 - p)\left(a_f\beta_f\right)^2 = \frac{a_f^2}{2}V_G. \tag{27}$$

762 where

$$V_G = 2p(1 - p){\beta_f}^2. \tag{28}$$

763 is the additive genetic variance. However, when β_m does not strictly equal β_f, **Eq. 25, 26** together

764 imply

$$\beta_m + \beta_f = \frac{\beta_m + \beta_f}{\beta_m - \beta_f}(\beta_m - \beta_f) = \frac{\frac{S_m}{a_m} - \frac{S_m}{a_f}}{\frac{S_m}{a_m} + \frac{S_m}{a_f}}(\beta_m - \beta_f) = \frac{a_f - a_m}{a_f + a_m}(\beta_m - \beta_f). \tag{29}$$

765 Finally, using **Eq. 25**,

$$s_m - s_f = a_m\beta_m - a_f\beta_f = \frac{1}{2}[(a_m + a_f)(\beta_m - \beta_f) + (a_m - a_f)(\beta_m + \beta_f)], \tag{30}$$

766 which together with **Eq. 29** gives

$$s_m - s_f = \frac{1}{2}\left[(a_m + a_f) + \frac{(a_m - a_f)(a_f - a_m)}{a_f + a_m}\right](\beta_m - \beta_f) = \frac{2a_m a_f}{a_m + a_f}(\beta_m - \beta_f). \tag{31}$$

767   We denote the heritability due to GxSex at the locus as $V_{GxSex} := 2p(1-p)(\beta_m - \beta_f)^2$ and

768   the parameter relating this contribution to the differentiation in allele frequencies as

$$A := 2\left(\frac{a_m a_f}{a_m + a_f}\right)^2, \tag{32}$$

769   and plug **Eq. 31** into **Eq. 24**, we get

$$F_{ST} \approx A V_{GxSex}. \tag{33}$$

770   as given by **Eq. 3** in **Results**.

772   **Estimating the potential for sexually antagonistic selection on standing variation (A).** For

773   each trait and gnomAD subsample (**Supplementary Materials**), we estimated A using weighted

774   least squares linear regression of our estimate of F_{ST} ($\widehat{F_{ST}}$) to our estimate of V_{GxSex} ($\hat{V}_{GxSex}$), with

775   weight w inversely proportional to our site-specific estimate of noise in the estimate of V_{GxSex},

$$w = \frac{1}{Var[\widehat{\hat{V}_{GxSex}}]}. \tag{34}$$

776   To simplify the estimation of $Var[\hat{V}_{GxSex}]$, we treated the allele frequency p as perfectly

777   estimated, and as independent of the allele frequency in the GWAS sample—as different data

778   are used in the GWAS (UK Biobank) and in the allele frequency estimation (gnomAD). Under

779   these assumptions,

$$Var[\widehat{\hat{V}_{GxSex}}] = Var[2p(1-p)\widehat{D^2}] = [2p(1-p)]^2 Var[(\hat{\beta}_m - \hat{\beta}_f)^2], \tag{35}$$

780   and thus the task at hand is estimating $Var[(\hat{\beta}_m - \hat{\beta}_f)^2]$. Using the law of total variance,

$$Var[(\hat{\beta}_m - \hat{\beta}_f)^2] = Var_{\hat{\beta}_f}\left[E_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2|\hat{\beta}_f\right]\right] + E_{\hat{\beta}_f}\left[Var_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2|\hat{\beta}_f\right]\right]. \tag{36}$$

781   We begin with the argument of the first term,

$$E_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2|\hat{\beta}_f\right] = E_{\hat{\beta}_m}[\hat{\beta}_m^2 - 2\hat{\beta}_m\hat{\beta}_f + \hat{\beta}_f^2|\hat{\beta}_f] = \mu_m^2 + \sigma_m^2 - 2\mu_m\hat{\beta}_f + \hat{\beta}_f^2, \tag{37}$$

782   where we denote

$$\mu_z = E[\hat{\beta}_z]; \tag{38}$$
$$\sigma_z^2 = Var[\hat{\beta}_z]$$

783   for each sex $z \in \{m, f\}$. Plugging **Eq. 37** into the first term of **Eq. 36**,

$$Var_{\hat{\beta}_f}\left[E_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2\big|\hat{\beta}_f\right]\right] = Var_{\hat{\beta}_f}[\mu_m^2 + \sigma_m^2] + Var_{\hat{\beta}_f}[\hat{\beta}_f^2 - 2\mu_m\hat{\beta}_f] = \tag{39}$$

$$0 + Var_{\hat{\beta}_f}[\hat{\beta}_f^2 - 2\mu_m\hat{\beta}_f] = Var_{\hat{\beta}_f}[\hat{\beta}_f^2] + 4Var_{\hat{\beta}_f}[\mu_m\hat{\beta}_f] - 4\mu_m Cov_{\hat{\beta}_f}[\hat{\beta}_f^2, \hat{\beta}_f],$$

784   where the first and second step follow from the fact that $\mu_m^2 + \sigma_m^2$ is a constant. We can take note

785   of the fact that $\hat{\beta}_z$ is Normally distributed around β_z, and in particular that it has no skewness.

786   Therefore,

$$Cov_{\hat{\beta}_z}[\hat{\beta}_z^2, \hat{\beta}_z] = E[\hat{\beta}_z^3] - E[\hat{\beta}_z]E[\hat{\beta}_z^2] = (\mu_z^3 + 3\mu_z\sigma_z^2 + \gamma_z\sigma_z^3) - \mu_z(\mu_z^2 + \sigma_z^2) = 2\mu_z\sigma_z^2, \tag{40}$$

787   where $\gamma_z = 0$ is the skewness of $\hat{\beta}_z$. We can also note that

$$Var_{\hat{\beta}_z}[\hat{\beta}_z^2] = Var_{\hat{\beta}_z}[(\sigma_z b_z + \mu_z)^2], \tag{41}$$

788   where we defined

$$b_z = \frac{\hat{\beta}_z - \mu_z}{\sigma_z},$$

789   and therefore b_z is a Standard Normal and therefore b_z^2 is Chi-squared with one degree of

790   freedom. **Eq. 41** now gives

$$Var_{\hat{\beta}_z}[\hat{\beta}_z^2] = Var_{\hat{\beta}_z}[\sigma_z^2 b_z^2 + 2\sigma_z\mu_z b_z] \tag{42}$$

$$= Var_{\hat{\beta}_z}[\sigma_z^2 b_z^2] + Var[2\sigma_z\mu_z b_z] + Cov[\sigma_z^2 b_z^2, 2\sigma_z\mu_z b_z]$$

$$= Var[b_z^2]\sigma_z^4 + 4Var[b_z]\mu_z^2\sigma_z^2 + 0 = 2\sigma_z^4 + 4\mu_z^2\sigma_z^2.$$

791   Plugging **Eq. 40** and **Eq. 42** into **Eq. 39**, we find

$$Var_{\hat{\beta}_f}\left[E_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2\big|\hat{\beta}_f\right]\right] = 2\sigma_f^4 + 4\mu_f^2\sigma_f^2 + 4\mu_m^2\sigma_f^2 - 8\mu_m\mu_f\sigma_f^2. \tag{43}$$

792   We now turn to the second term of **Eq. 36**. First,

$$Var_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2\big|\hat{\beta}_f\right] = Var[\hat{\beta}_m^2 + 2\hat{\beta}_m\hat{\beta}_f|\hat{\beta}_f] \tag{44}$$

$$= Var[\hat{\beta}_m^2] + 4\sigma_m^2\hat{\beta}_f^2 - 4\hat{\beta}_f Cov[\hat{\beta}_m, \hat{\beta}_m^2].$$

793   **Eq. 40** and **42** again give us

$$Var_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2\big|\hat{\beta}_f\right] = 2\sigma_m^4 + 4\mu_m^2\sigma_m^2 + 4\sigma_m^2\hat{\beta}_f^2 - 8\mu_m\sigma_m^2\hat{\beta}_f, \tag{45}$$

794   which then gives

$$E_{\hat{\beta}_f}\left[Var_{\hat{\beta}_m}\left[(\hat{\beta}_m - \hat{\beta}_f)^2\big|\hat{\beta}_f\right]\right] = 2\sigma_m^4 + 4\mu_m^2\sigma_m^2 + 4\sigma_m^2(\sigma_f^2 + \mu_f^2) - 8\mu_m\mu_f\sigma_m^2. \tag{46}$$

795   Plugging **Eq. 43** and **Eq. 46** into **Eq. 36**, we obtain

$$Var[(\hat{\beta}_m - \hat{\beta}_f)^2] = \tag{47}$$

$$= 2(\sigma_m^4 + \sigma_f^4) + 4\sigma_m^2\sigma_f^2 + 4(\mu_m^2\sigma_m^2 + \mu_f^2\sigma_f^2) + 4(\sigma_m^2\mu_f^2 + \sigma_f^2\mu_m^2)$$
$$- 8\mu_m\mu_f(\sigma_m^2 + \sigma_f^2).$$

796   Finally, we estimate μ_z with the GWAS-derived point estimate of the effect $\hat{\beta}_z$ and σ_z with

797   its standard error, $\hat{\sigma}_z = [\hat{\beta}_z]$. Plugging back into **Eq. 35**, we obtain

$$Var\widehat{[V_{GxSex}]} = [2p(1-p)]^2[2(\hat{\sigma}_m^4 + \hat{\sigma}_f^4) + 4\hat{\sigma}_m^2\hat{\sigma}_f^2 + 4(\hat{\beta}_m^2\sigma_m^2 + \hat{\beta}_f^2\sigma_f^2) + 4(\hat{\sigma}_m^2\hat{\beta}_f^2 + \hat{\sigma}_f^2\hat{\beta}_m^2) \quad (48)$$
$$- 8\hat{\beta}_m\hat{\beta}_f(\hat{\sigma}_m^2 + \hat{\sigma}_f^2)].$$

798   Using **Eq. 33**, we estimate F_{st} with the estimator

$$\widehat{F_{st}} = n_{st}/d_{st}, \quad (49)$$

799   where

$$n_{st} = \left(\widehat{p_m} - \widehat{p_f}\right)^2 - SE(\widehat{p_m})^2 - SE\left(\widehat{p_f}\right)^2, \quad (50)$$

$$d_{st} = 4\hat{p}(1-\hat{p}) - SE(\widehat{p_m})^2 - SE\left(\widehat{p_f}\right)^2,$$

801   and noting that

$$E[\widehat{F_{st}}] \approx \frac{E[n_{st}]}{E[d_{st}]} = \frac{(p_m - p_f)^2 - Var(p_m) + Var(p_f) + E\{SE(\widehat{p_m})^2\} + E\left[SE\left(\widehat{p_f}\right)^2\right]}{4p(1-p) + Var(p_m)^2 + Var(p_f)^2 - E\{SE(\widehat{p_m})^2\} - E\left[SE\left(\widehat{p_f}\right)^2\right]} = F_{st}, \quad (51)$$

where in the first equality we approximated the expectation of a ratio with the ratio of expectations.

Therefore, **Eq. 49** provides an approximately unbiased estimator of F_{st} despite the absence of

genotype frequencies.

To perform this estimation of A on the GWAS and F_{st} data, we used paired v and V_{GxSex}

points for all sites which passed all previous stages of filtering. Weights were set by **Eq. 34** and

follow **Eq. 48** where $\hat{\beta}_m$ and $\hat{\beta}_f$ are the GWAS effect estimates as above, and $\hat{\sigma}_m$ and $\hat{\sigma}_f$ are the

GWAS standard errors (SE) estimates for the effect size of each site per trait.

To minimize the possibility of LD between sites used in the analysis as much as possible,

we used the approximately independent LD blocks in Europeans[76] as in Section "Mixture weights

for covariance structure between male and female effects". Namely, we subdivided the genome

into 1703 approximately independent LD blocks as before. We iterated over the 1703 blocks and

sampling one site per block in a given iteration, using a sample of (up to) 1703 post-filtering sites

to perform the weighted linear regression of F_{ST} on $V_{G\times Sex}$. The slope of this regression was used

as an estimate of A. We perform this estimation procedure 1,000 times and take an average of

Z scores (slope point estimates divided by their SE) as the final estimate of A. In each replicate,

we sample with replacement m LD blocks from the m LD blocks which had at least one site within

them post-filtering (Supplementary Materials); we then sample one site per resampled block. In

Fig. 7D, each point is the mean of the 1,000 samples of one site per LD block and 90% confidence

intervals show the range between the 5th and 95th percentile of the 1000 replicates.

In the main text, we focus on the results performed this estimation for Ashkenazi Jewish,

Finnish, and Non-Finnish European populations as the other ancestry group-stratified

subsamples in *gnomAD* are further diverged from the UKB White British sample and therefore

our GWAS estimates are expected to be less portable[57,81]. We also performed a similar analysis

using UKB data to measure differentiation in allele frequencies between males and females,

rather than an independent dataset (*gnomAD*) as in the main text. Since individual level data was

available in this case, we replaced F_{st} with L_{ST}, a measure developed by Ruzicka et al.[51]. L_{st} can

be thought of as site-specific F_{st} controlled for major axes of population structure differentiating

males and females (**Fig. S21)**.

References

1. van Doorn, G. S. Intralocus sexual conflict. *Ann N Y Acad Sci* **1168**, 52–71 (2009).

2. Arnqvist, G. & Rowe, L. *Sexual Conflict*. (Princeton University Press, 2005). doi:10.1515/9781400850600.

3. Barson, N. J. *et al.* Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature* **528**, 405–408 (2015).

4. Kidwell, J. F., Clegg, M. T., Stewart, F. M. & Prout, T. Regions of stable equilibria for models of differential selection in the two sexes under random mating. *Genetics* **85**, 171–83 (1977).

5. Connallon, T., Cox, R. M. & Calsbeek, R. Fitness consequences of sex-specific selection. *Evolution* **64**, 1671–82 (2010).

6. Harrison, P. W. *et al.* Sexual selection drives evolution and rapid turnover of male gene expression. *Proceedings of the National Academy of Sciences* **112**, 4393–4398 (2015).

7. Cheng, C. & Kirkpatrick, M. Sex-Specific Selection and Sex-Biased Gene Expression in Humans and Flies. *PLoS Genet* **12**, e1006170 (2016).

8. Karp, N. A. *et al.* Prevalence of sexual dimorphism in mammalian phenotypic traits. *Nat Commun* **8**, 15475 (2017).

9. Schroderus, E. *et al.* Intra- and Intersexual Trade-Offs between Testosterone and Immune System: Implications for Sexual and Sexually Antagonistic Selection. *Am Nat* **176**, E90–E97 (2010).

10. Power, R. A. *et al.* Fecundity of Patients With Schizophrenia, Autism, Bipolar Disorder,
Depression, Anorexia Nervosa, or Substance Abuse vs Their Unaffected Siblings. *JAMA*
Psychiatry **70**, 22 (2013).

11. Mokkonen, M. & Crespi, B. J. Genomic conflicts and sexual antagonism in human health:
insights from oxytocin and testosterone. *Evol Appl* **8**, 307–325 (2015).

12. Harper, J. A., Janicke, T. & Morrow, E. H. Systematic review reveals multiple sexually
antagonistic polymorphisms affecting human disease and complex traits. *Evolution (N Y)*
75, 3087–3097 (2021).

13. Oliva, M. *et al.* The impact of sex on gene expression across human tissues. *Science*
(1979) **369**, eaba3066 (2020).

14. Khramtsova, E. A., Davis, L. K. & Stranger, B. E. The role of sex in the genomics of human
complex traits. *Nat Rev Genet* **20**, 173–190 (2019).

15. Clayton, J. A. & Collins, F. S. Policy: NIH to balance sex in cell and animal studies. *Nature*
509, 282–283 (2014).

16. Clayton, J. A. Studying both sexes: a guiding principle for biomedicine. *The FASEB Journal*
30, 519–524 (2016).

17. Nature journals raise the bar on sex and gender reporting in research. *Nature* **605**, 396–
396 (2022).

18. Sinnott-Armstrong, N., Naqvi, S., Rivas, M. & Pritchard, J. K. Gwas of three molecular traits
highlights core genes and pathways alongside a highly polygenic background. *Elife* **10**, 1–
35 (2021).

19.   Carole Hooven. *T: The Story of Testosterone, the Hormone that Dominates and Divides*
871   *Us*. (Henry Holt and Co., 2021).

872   20.   Flynn, E. *et al.* Sex-specific genetic effects across biomarkers. *Eur J Hum Genet* **29**, 154–
873   163 (2021).

874   21.   Bernabeu, E. *et al.* Sex differences in genetic architecture in the UK Biobank. *Nat Genet*
875   **53**, 1283–1289 (2021).

876   22.   Muir, W., Nyquist, W. E. & Xu, S. Alternative partitioning of the genotype-by-environment
877   interaction. *Theoretical and Applied Genetics* **84**, 193–200 (1992).

878   23.   Robertson, A. The Sampling Variance of the Genetic Correlation Coefficient. *Biometrics*
879   **15**, 469 (1959).

880   24.   Falconer, D. S. The Problem of Environment and Selection. *Am Nat* **86**, 293–298 (1952).

881   25.   Fry, J. D. The Mixed-Model Analysis of Variance Applied to Quantitative Genetics:
882   Biological Meaning of the Parameters. *Evolution (N Y)* **46**, 540 (1992).

883   26.   Yamada, Y. Genotype by Environment Interaction and Genetic Correlation of the same
884   Trait under Different Environments. *The Japanese Journal of Genetics* **37**, 498–509 (1962).

885   27.   Brown, B. C., Asian Genetic Epidemiology Network Type 2 Diabetes Consortium, Ye, C.
886   J., Price, A. L. & Zaitlen, N. Transethnic Genetic-Correlation Estimates from Summary
887   Statistics. *Am J Hum Genet* **99**, 76–88 (2016).

888   28.   Galinsky, K. J. *et al.* Estimating cross-population genetic correlations of causal effect sizes.
889   *Genet Epidemiol* **43**, 180–188 (2019).

29.   Ni, G., Moser, G., Schizophrenia Working Group of the Psychiatric Genomics Consortium,
Wray, N. R. & Lee, S. H. Estimation of Genetic Correlation via Linkage Disequilibrium Score
Regression and Genomic Restricted Maximum Likelihood. *Am J Hum Genet* **102**, 1185–
1194 (2018).

30.   Shi, H., Mancuso, N., Spendlove, S. & Pasaniuc, B. Local Genetic Correlation Gives
Insights into the Shared Genetic Architecture of Complex Traits. *The American Journal of*
Human Genetics **101**, 737–751 (2017).

31.   Bulik-Sullivan, B. K. *et al.* An atlas of genetic correlations across human diseases and traits.
Nat Genet **47**, 1236–1241 (2015).

32.   DiMarco, M., Zhao, H., Boulicault, M. & Richardson, S. S. Why "sex as a biological variable"
conflicts with precision medicine initiatives. *Cell Rep Med* **3**, 100550 (2022).

33.   Lumish, H. S., O'Reilly, M. & Reilly, M. P. Sex Differences in Genomic Drivers of Adipose
Distribution and Related Cardiometabolic Disorders: Opportunities for Precision Medicine.
Arteriosclerosis, thrombosis, and vascular biology vol. 40 45–60 Preprint at
https://doi.org/10.1161/ATVBAHA.119.313154 (2020).

34.   Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. *Cell* **169**, 1177–1186 (2017).

35.   Shi, H., Kichaev, G. & Pasaniuc, B. Contrasting the Genetic Architecture of 30 Complex
Traits from Summary Association Data. *Am J Hum Genet* **99**, 139–53 (2016).

36.   Sella, G. & Barton, N. H. Thinking About the Evolution of Complex Traits in the Era of
Genome-Wide Association Studies. *Annu Rev Genomics Hum Genet* **20**, 461–493 (2019).

37. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity
in genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).

38. Benonisdottir, S. & Kong, A. The Genetics of Participation: Method and Analysis. *bioRxiv*
(2022) doi:10.1101/2022.02.11.480067.

39. Lynch, M. & Walsh, B. *Genetics and Analysis of Quantitative Traits*. (Sinauer Associates,
1998).

40. Urbut, S. M., Wang, G., Carbonetto, P. & Stephens, M. Flexible statistical methods for
estimating and testing effects in genomic studies with multiple conditions. *Nat Genet* **51**,
187–195 (2019).

41. Pasquali, R. Obesity and androgens: facts and perspectives. *Fertil Steril* **85**, 1319–40
(2006).

42. Domingue, B. W., Kanopka, K., Mallard, T. T., Trejo, S. & Tucker-Drob, E. M. Modeling
Interaction and Dispersion Effects in the Analysis of Gene-by-Environment Interaction.
Behav Genet **52**, 56–64 (2022).

43. Liu, D. *et al.* Skeletal muscle gene expression in response to resistance exercise: sex
specific regulation. *BMC Genomics* **11**, 659 (2010).

44. Lutz, S. Z. *et al.* Sex-Specific Associations of Testosterone With Metabolic Traits. *Front
Endocrinol (Lausanne)* **10**, (2019).

45. Kraemer, W. J., Ratamess, N. A. & Nindl, B. C. Recovery responses of testosterone,
growth hormone, and IGF-1 after resistance exercise. *J Appl Physiol* **122**, 549–558 (2017).

46.   Roberts, B. M., Nuckols, G. & Krieger, J. W. Sex Differences in Resistance Training: A
932   Systematic Review and Meta-Analysis. *J Strength Cond Res* **34**, 1448–1460 (2020).

933   47.   Zajitschek, S. R. *et al.* Sexual dimorphism in trait variability and its eco-evolutionary and
934   statistical implications. *Elife* **9**, (2020).

935   48.   Kasimatis, K. R. *et al.* Evaluating human autosomal loci for sexually antagonistic viability
936   selection in two large biobanks. *Genetics* **217**, (2021).

937   49.   Ryan, M. *The Genetics of Political Behavior*. (Routledge, 2020).

938   50.   Ruzicka, F., Holman, L. & Connallon, T. Polygenic signals of sexually antagonistic selection
939   in contemporary human genomes. *bioRxiv* (2021) doi:10.1101/2021.09.20.461171.

940   51.   Ruzicka, F. *et al.* The search for sexually antagonistic genes: Practical insights from studies
941   of local adaptation and statistical genomics. *Evol Lett* **4**, 398–415 (2020).

942   52.   Wright, S. The genetical structure of populations. *Ann Eugen* **15**, 323–54 (1951).

943   53.   Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population
944   Structure. *Evolution (N Y)* **38**, 1358 (1984).

945   54.   Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in
946   141,456 humans. *Nature* **581**, 434–443 (2020).

947   55.   Pirastu, N. *et al.* Genetic analyses identify widespread sex-differential participation bias.
948   *Nat Genet* **53**, 663–671 (2021).

949   56.   Bissegger, M., Laurentino, T. G., Roesti, M. & Berner, D. Widespread intersex
950   differentiation across the stickleback genome – The signature of sexually antagonistic
951   selection? *Mol Ecol* **29**, 262–271 (2020).

952   57.   Privé, F. *et al.* Portability of 245 polygenic scores when derived from the UK Biobank and
953   applied to 9 ancestry groups from the same cohort. *The American Journal of Human*
954   *Genetics* **109**, 12–23 (2022).

955   58.   Traglia, M., Bout, M. & Weiss, L. A. Sex-heterogeneous SNPs disproportionately influence
956   gene expression and health. *PLoS Genet* **18**, e1010147 (2022).

957   59.   Berg, J. J. *et al.* Reduced signal for polygenic adaptation of height in UK Biobank. *Elife* **8**,
958   (2019).

959   60.   Sohail, M. *et al.* Polygenic adaptation on height is overestimated due to uncorrected
960   stratification in genome-wide association studies. *Elife* **8**, (2019).

961   61.   Young, A. I., Benonisdottir, S., Przeworski, M. & Kong, A. Deconstructing the sources of
962   genotype-phenotype associations in humans. *Science* **365**, 1396–1400 (2019).

963   62.   Coop, G. & Przeworski, M. Lottery, luck, or legacy. A review of "The Genetic Lottery: Why
964   DNA matters for social equality". *Evolution (N Y)* **76**, 846–853 (2022).

965   63.   Mills, M. C. & Tropf, F. C. Sociology, Genetics, and the Coming of Age of Sociogenomics.
966   *Annu Rev Sociol* **46**, 553–581 (2020).

967   64.   Coop, G. Reading tea leaves? Polygenic scores and differences in traits among groups.
968   Preprint at (2019).

969   65.   Ober, C., Loisel, D. A. & Gilad, Y. Sex-specific genetic architecture of human disease. *Nat*
970   *Rev Genet* **9**, 911–922 (2008).

971   66.   Khan, S. S. *et al.* Association of Body Mass Index With Lifetime Risk of Cardiovascular
972   Disease and Compression of Morbidity. *JAMA Cardiol* **3**, 280 (2018).

973   67.   Vazquez, G., Duval, S., Jacobs, D. R. & Silventoinen, K. Comparison of Body Mass Index,
974   Waist Circumference, and Waist/Hip Ratio in Predicting Incident Diabetes: A Meta-
975   Analysis. *Epidemiol Rev* **29**, 115–128 (2007).

976   68.   Ning, Y., Wang, L. & Giovannucci, E. L. A quantitative analysis of body mass index and
977   colorectal cancer: findings from 56 observational studies. *Obesity Reviews* **11**, 19–30
978   (2010).

979   69.   Brown, C. D. *et al.* Body Mass Index and the Prevalence of Hypertension and Dyslipidemia.
980   *Obes Res* **8**, 605–619 (2000).

981   70.   Arthur, R. S., Dannenberg, A. J. & Rohan, T. E. The association of prediagnostic circulating
982   levels of cardiometabolic markers, testosterone and sex hormone-binding globulin with risk
983   of breast cancer among normal weight postmenopausal women in the UK Biobank. *Int J*
984   *Cancer* **149**, 42–57 (2021).

985   71.   Wang, J. *et al.* Sex-specific associations of circulating testosterone levels with all-cause
986   and cause-specific mortality. *Eur J Endocrinol* **184**, 723–732 (2021).

987   72.   Fry, A. *et al.* Comparison of Sociodemographic and Health-Related Characteristics of UK
988   Biobank Participants With Those of the General Population. *Am J Epidemiol* **186**, 1026–
989   1034 (2017).

73.   Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.
Nature **562**, 203–209 (2018).

74.   Haworth, S. *et al.* Apparent latent structure within the UK Biobank sample has implications
for epidemiological analysis. *Nat Commun* **10**, 333 (2019).

75.   Stephens, M. False discovery rates: a new deal. *Biostatistics* kxw041 (2016)
doi:10.1093/biostatistics/kxw041.

76.   Berisa, T. & Pickrell, J. K. Approximately independent linkage disequilibrium blocks in
human populations. *Bioinformatics* **32**, 283–5 (2016).

77.   Sudlow, C. *et al.* UK Biobank: An Open Access Resource for Identifying the Causes of a
Wide Range of Complex Diseases of Middle and Old Age. *PLoS Med* **12**, e1001779 (2015).

78.   Davies, N. M., Holmes, M. v & Davey Smith, G. Reading Mendelian randomisation studies:
a guide, glossary, and checklist for clinicians. *BMJ* k601 (2018) doi:10.1136/bmj.k601.

79.   Davey Smith, G. & Ebrahim, S. 'Mendelian randomization': can genetic epidemiology
contribute to understanding environmental determinants of disease?*. *Int J Epidemiol* **32**,
1–22 (2003).

80.   Gillespie, J. H. *Population Genetics: A Concise Guide*. (Johns Hopkins University Press,
2004).

81.   Wang, Y. *et al.* Theoretical and empirical quantification of the accuracy of polygenic scores
in ancestry divergent populations. *Nat Commun* **11**, 3865 (2020).

1009