Potential future climate change effects on global reptile distribution and diversity

Matthias F. Biber[1,*], Alke Voskamp[2], Christian Hof[1]

[1]Technical University of Munich, Terrestrial Ecology Research Group, Department for Life Science Systems, School of Life Sciences, 85354 Freising, Germany

[2]Senckenberg Biodiversity and Climate Research Centre (SBiK-F), 60325 Frankfurt, Germany

Running head: Global reptiles under climate change

Abstract

Aim: Until recently, complete information on global reptile distributions has not been widely available. Here, we provide the first comprehensive climate impact assessment for reptiles at a global scale.

Location: Global, excluding Antarctica

Time period: 1995, 2050, 2080

Major taxa studied: Reptiles

Methods: We modeled the distribution of 6296 reptile species and assessed potential global as well as realm-specific changes in species richness, the change in global species richness across climatic space and species-specific changes in range extent, overlap and position, under future climate change. To assess the future climatic impact on 3768 range-restricted species, which could not be modeled, we compared the future change in climatic conditions between both modeled and non-modeled species.

Results: Reptile richness was projected to decline significantly over time, globally but also for most zoogeographic realms, with the strongest decrease in Brazil, Australia and South Africa. Species richness was highest in warm and most regions, which were projected to shift further to climate extremes in the future. Range extents were projected to decline considerably in the future, with a low overlap between projected current and future ranges. Shifts in range centroids differed among realms and taxa, with a dominating global poleward shift. Non-modeled species were likely to be significantly stronger affected by climatic changes than modeled species.

Main conclusions: With ongoing future climate change, reptile richness is likely to decrease significantly across most parts of the world. This effect, as well as considerable impacts on species' range extent, overlap and position, were visible across lizards, snakes and turtles alike. Together with other anthropogenic impacts, such as habitat loss and harvesting, this is a cause for concern. Given the historical lack of information on global reptile distributions, this calls for a re-assessment of global conservation efforts towards reptile species, with specific focus on anticipated future climatic changes.

KEYWORDS: bioclim, biodiversity, environmental niche model, global warming, ISIMIP, lizard, snake, species distribution model, species richness, turtle

Introduction

Emissions from anthropogenic activities have led to an increase in global surface temperature of around 1°C in the last 100 years. This has already led to changes in weather and climate extremes in every region across the globe (IPCC, 2021). Unless emissions are vastly reduced in the coming decades, global warming will continue and exceed 1.5°-2°C compared to pre-industrial levels by the end of the 21st century (IPCC, 2021).

Climate change has already had adverse effects on biodiversity and ecosystem functioning and these effects are likely to worsen as warming proceeds in future (IPBES, 2019; IPCC, 2022). Climate change impacts on ecological processes scales from genes to entire ecosystems, can affect organisms, populations or entire communities and vary between physiological, morphological, phenological and distributional shifts (Bellard *et al.*, 2012; Scheffers *et al.*, 2016). Especially changes in species abundance and distribution due to climate change have already been frequently observed (Bowler *et al.*, 2017; Lenoir *et al.*, 2020), with many species shifting their range towards higher latitudes and elevations (Chen *et al.*, 2011). However, some species also respond to climate change by idiosyncratic range shifts (Gibson-Reinemer & Rahel, 2015).

In the past most climate change impact assessments on vertebrate biodiversity have focused on endotherms (birds & mammals). Reptiles, though accounting for a third of global terrestrial vertebrate diversity, have been largely ignored (Pacifici *et al.*, 2015), and previous assessments of climate change impacts on reptiles species have either used only a subset of species (Warren *et al.*, 2018; Newbold, 2018) or have not been of global extent (Araújo *et al.*, 2006). Global biodiversity assessments further often either consider overall effects on a single taxon (Baisero *et al.*, 2020; Voskamp *et al.*, 2021) or compare multiple taxa (Hof *et al.*, 2018; Newbold, 2018; Warren *et al.*, 2018; Thuiller *et al.*, 2019), but only very rarely compare different taxonomic groups within one taxon (but see, for example Hof *et al.* 2011).

Reptiles have in the past also often been neglected when assessing global conservation priorities (Brooks *et al.*, 2006). They are most diverse in arid and semi-arid regions, which suggests that their distributions are driven by ecological and evolutionary processes that differ from other vertebrate taxa, and these regions have been previously unrecognized as conservation priorities because other vertebrate taxa could be more efficiently protected elsewhere (Roll *et al.*, 2017).

While reptiles overall have been found to use similar habitats to mammals and birds (Cox et al. 2022), treating all reptile species as uniform may be problematic, as reptile taxonomic groups (lizard, snakes and turtles) exhibit large differences in species richness hotspots (Roll *et al.*, 2017) as well as habitat use, making them vulnerable to different anthropogenic impacts (Cox *et al.*, 2022).

Because reptiles are ectothermic, they are likely to be strongly influenced by climate warming, with some species already experiencing body temperatures above their physiological optima. This could indicate a higher vulnerability of these species to climate warming compared to species from cold environments (Diele-Viegas *et al.*, 2018). Nonetheless, temperate species are also likely to be vulnerable, assuming that their physiological adaptations for living in cold environments may hinder their ability to cope with hotter climates (Monasterio *et al.*, 2013). However, previous studies assessing climate change impacts on reptile species are biased

towards certain species and taxonomic groups as well as zoogeographic realms (Diele-Viegas *et*
al., 2020).

Here, we aimed to provide a detailed account of projected climate change impacts on global
reptile distributions and diversity, looking at species-specific changes as well as broad-scale
geographic trends across and within different taxonomic groups. We assessed changes in reptile
species richness globally, within each zoogeographic realm and across their respective climate
space. For each species, we further quantified the change in range extent, range overlap and
range distribution and again assessed differences across zoogeographic realms and taxonomic
groups. Given that we cannot model range-restricted species, we also performed a more general
assessment of species-specific changes in climatic space across both modeled and non-modeled
species.

Methods
90

Species data
91

Until recently, global reptile distribution data was unavailable, but this has changed with the
release of the global distribution database by the Global Assessment of Reptile Distributions
(GARD) initiative (Roll *et al.*, 2017) and more recently the release of the full set of IUCN
reptile range maps (IUCN, 2022). We obtained global range maps of 10,064 reptile species
from the GARD initiative (Roll *et al.*, 2017). The range maps cover lizards, snakes, turtles,
worm lizards, crocodiles and the tuatara, but in this paper, similar to (Roll *et al.*, 2017), we only
contrast snakes, turtles and paraphyletic lizards (the latter of which we subsequently refer to as
lizards for simplicity).

Range maps were gridded to a 0.5° x 0.5° grid in WGS84, to align with the available climate
data (see next section), considering any grid cell that intersected with a species' range polygon as
a presence. As range maps only provide information on a species' presence but not on a species'
absence, pseudo-absence data for each species were generated by randomly selecting grid cells
with no presence, using a distance-weighted approach, where grid cells closer to the range edge
were favored over grid cells further away (see Hof *et al.*, 2018). The number of absences was
either equal to the number of presences for species with 1000 or more presences or 1000
absences for species with less than 1000 presences, as a minimum of 1000 absences considerably
increases model performance (Barbet-Massin *et al.*, 2012). For each species we derived 10
replicate sets of pseudo-absences, to account for the variability in model accuracy because of the
random sampling of pseudo-absence data. Barbet-Massin *et al.* (2012) found that depending on
the number of pseudo-absences and model algorithm chosen, 5 to 12 replicates provide the best
model quality. Given the relative high number of pseudo-absences, 10 replicates should thus
result in a good quality of our models. For each of these 10 sets we created a separate model, but
the results were then averaged across the 10 sets.

Climate data
115

Global bias-corrected daily climate (minimum temperature, maximum temperature and
precipitation) data at a spatial resolution of 0.5° (WGS84) were obtained from the meteorological
forcing dataset 'EartH2Observe, WFDEI and ERA-Interim data Merged and Bias-corrected for
ISIMIP' (EWEMBI; Lange, 2016) for current conditions (1980 - 2009) and from the Inter-

Sectoral Impact Model Intercomparison Project phase 2b (ISIMP2b; Frieler *et al.*, 2017) for
future simulations (2036 - 2065 & 2066 - 2095). Future climate simulations were available from
four global circulation models (GCMs; GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR and
MIROC5) and for three representative concentration pathways (RCPs; RCP2.6, RCP6.0 and
RCP8.5) under the Coupled Model Intercomparison Project Phase 5 (CMIP5). Future climate
simulations strongly depend on the GCM used (Watterson, 2019). The four GCMs chosen by
ISIMIP2b capture a large range of plausible future climate projections of all the different GCMs
available (Frieler *et al.*, 2017), while the RCPs represent different emission scenarios according
to the ongoing and future trajectories of global CO_2 emissions (Van Vuuren *et al.*, 2011).

Monthly means of each climatic variable over the respective 30-year time periods, centered
around 1995, 2050 and 2080, and for each future scenario (GCM & RCP) were used to calculate
19 bioclimatic variables (see Table S1.1 in Appendix S1 in Supporting Information) using the
biovars() function of the 'dismo' package (Hijmans *et al.*, 2021) in R (R Core Team, 2021).
Bioclimatic variables represent annual trends, seasonality and extreme or limiting environmental
factors and are thus more biologically meaningful variables than temperature or precipitation
alone.

Species Distribution Models (SDMs)

Species distribution models (SDMs) are a common way of assessing species-specific responses
to climate change (Guisan & Thuiller, 2005), but are also used to assess climate change impacts
on biodiversity (Thuiller *et al.*, 2005). SDMs statistically infer a relationship between the
observed distribution of a species to the underlying environmental conditions (Elith &
Leathwick, 2009) and can then be used to project current distributions into the future (Elith *et al.*,
2010), assuming that the species maintains its climatic niche (Wiens & Graham, 2005). By doing
this for multiple species, these projections can be combined to assess future changes in species
richness (e.g. Hof *et al.*, 2018).

We fitted species distribution models using the presence/pseudo-absence data of a species as
response variable and the derived bioclimatic variables for current (1995) conditions as
explanatory variable. The number and choice of explanatory variables used strongly influences
the outcome of the SDMs (Petitpierre *et al.*, 2017). We thus performed a rigorous variable
selection approach. After pre-selecting the 10 most commonly used variables from the literature
(see Porfirio *et al.*, 2014), all potential combinations of three and four bioclimatic variables with
a low Pearson correlation ($r \leq 0.7$) were used to model 10 % of all species (N= 987), which
were randomly selected, using a Generalized Additive Model (GAM) approach (see Hof *et al.*,
2018). The model performance of the different variable combinations was tested and models for
all species were fitted using the best-performing variable combination, which was temperature
seasonality, maximum temperature of the warmest month, annual precipitation and precipitation
seasonality (see Fig. S1.1).

Projections based on SDMs vary considerably among model algorithms (Thuiller *et al.*, 2019),
therefore we fitted two different modeling algorithms with a good performance and
discrimination capacity (Meynard & Quinn, 2007; Elith *et al.*, 2010), an additive model (GAM)
and a regression tree based model (Generalized Boosted Regression Models (GBM)).

GAMs were fitted with a Bernoulli response, a logit link and thin-plate regression splines using
the 'mgcv' package (Wood, 2003, 2011) in R (R Core Team, 2021). GBMs were fitted with the
'gbm' package (Greenwell *et al.*, 2020) in R (R Core Team, 2021) and the optimal parameter
settings for learning rate (0.01 and 0.001), tree complexity (1, 2 and 3) and number of trees
(1000-10000) for each species were identified by cross-validation (Bagchi *et al.*, 2013).

Spatial autocorrelation in species distributions can bias parameter estimates and error
probabilities (Kühn, 2007). Two different methods were used to account for spatial
autocorrelation in the SDMs. Species with equal or more than 50 presences were modeled using
an ecoregion-blocking approach. Here the world was divided into 10 blocks, based on a
representative subset of the climatic space across each of the world's ecoregions (Olson *et al.*,
2001, Bagchi *et al.*, 2013). Subsequently 10 models, per species, were built leaving out one
block at a time, using the left out block for model evaluation (Bagchi *et al.*, 2013). For species
with 10 to 49 presences, we split the data into 10 datasets by repeatedly randomly selecting 70%
of the data, using the left-out 30% for model evaluation. Species occurring in less than 10 grid
cells (N = 3602, Table 1) were not modeled, as the sampling size would be too low to produce
useful results (Hernandez et al. 2006). Together with the 10 different pseudo-absence sets this
resulted in 100 models for each model algorithm and species.

The performance of the fitted SDMs was evaluated by calculating the overall AUC for each
species (the average AUC across the 10 blocks and the 10 sets of pseudo-absences). Models with
an overall AUC smaller than 0.7 were dropped (N = 166), which left us with SDMs for 6296
reptile species (see Fig. S1.2), which represents 62.6 % of the total number of available species
by GARD (Table 1). In addition to assessing the model fit of the individual models, we also
compared the observed species richness with the projected current richness per grid cell (Fig
S1.3) in order to assess the performance of all our models when looking at changes in species
richness.

The same modeling approach has been used previously to assess climate change impacts on
amphibians, birds and mammals, see Hof *et al.* (2018) and Biber *et al.* (2020). The former
provides a more detailed explanation of the modeling methodology, while the latter gives a
thorough account of the caveats and uncertainties associated with species distribution models.

Future projections

Future species distributions were projected using the future bioclimatic variables for the two
future time periods (2050, 2080), each GCM (GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR
and MIROC5), each RCP (RCP2.6, RCP6.0 and RCP8.5) and and both model algorithms (GAM
& GBM). Model results are presented as the ensemble mean, across the four GCMs and two
model algorithms (GAM & GBM) considered.

Future projections of each species were limited to the extent of their original and the neighboring
ecoregions, to avoid predictions to areas with analogue climatic conditions. Future projections
were further limited by applying a species-specific dispersal buffer. For most species considered
here species-specific dispersal distances are still unknown (Nathan *et al.*, 2012), hence we used
species-specific dispersal buffers which were based on the diameter (d) of the largest range
polygon of a species. We used three species-specific dispersal scenarios (d/4, d/8, d/16, see Fig.
S1.4) and provide a detailed comparison of these in the Supporting Information (see Appendix

S5). Here we provide the results under the medium dispersal scenario (d/8), which corresponds
to a mean dispersal distance of 2.4 km per year.

Impact analysis

The current and future probabilities of occurrence of the individual SDMs were thresholded into
binary presence-absence data using species-specific thresholds according to the true skill statistic
(MaxTSS; Allouche *et al.*, 2006; Fig. S1.5 d). Thresholded species occurrences were then used
to calculate current and future species richness, as well as richness increase, decrease, change
and relative change (%). Richness increase and decrease were identified by using the presence
information of each individual species and then summing up the number of species that newly
occur in a given grid cell (species increase) or species that disappear from the respective grid cell
(species decrease).

Summing up the thresholded species occurrences frequently overestimates species richness
(Calabrese et al., 2014), thus we also present the results using the sum of the raw non-
thresholded probabilities of occurrence of each species in the Supporting Information (see
Appendix S4).

We calculated the projected species richness for each cell of a global grid of 0.5° x 0.5°
resolution globally as well as for each zoogeographic realm, as defined by Holt *et al.* (2013), for
each time period. We then tested for significant changes in species richness over time using a
paired t-test with Holm correction. To assess how species richness and richness changes are
related to the overall change in climatic conditions, we assessed both across univariate
temperature and precipitation as well as the interaction of temperature and precipitation
conditions. To assess potential future climate effects on individual species, we quantified the
proportion of change in range extent, the proportion of range overlap and the direction and
distance in range shift. The proportion of change in range extent was calculated based on the
projected current and future range extent of a species.(Fig. S1.5 f). The proportion of range
overlap was calculated by extracting the total area of spatial range overlap between the projected
current and future range extent and dividing it by the projected current range extent (Fig. S1.5 g).
To assess the magnitude of the range shift for each species, we derived the range centroid for
both time periods and calculated the distance and direction of the projected range shift (Fig. S1.5
g).

Given that 37.4% of all reptile species for which data were available could not be modeled
(largely due to their restricted range extent, Table 1), we performed an additional analysis
considering all 10,064 species for which data were available. We used the same 4 bioclimatic
variables used for the SDMs to transform the multidimensional climate data in a two-
dimensional climate space using the first two axes of a principal component analysis (PCA).
PCAs were performed for both current and future conditions, considering the same GCMs, RCPs
and time periods as before. Explained variance of the first two PCA axes was above 75 % under
all scenarios (Fig. S1.6). For each scenario combination, we then calculated the Euclidean
distance between the two PCA axes of current and future conditions, to get a measure of climatic
change (Fig. S1.7). We then extracted the climatic distance for the gridded locations of each
species and compared the climatic distance of modeled and non-modeled (range-restricted)
species using a non-paired t-test with Holm correction.

Where no specific groups (lizards, snakes, turtles) are mentioned, we present the results for all
reptile species together, but see Appendix S2 for group-specific results. Results are presented for
the year 2080 under a medium representative concentration pathway (RCP6.0). A sensitivity
analysis with regards to the variation across years and RCPs is shown in the Supporting
Information (see Appendix S6).

Results

Spatial changes

Projected reptile richness (sum of thresholded SDM projections) based on a resolution of 0.5° x
0.5° for current conditions varied between 0 (high latitudes) and 251 in the tropics, with
particular hotspots in Brazil, Cameroon and Indonesia (Fig. 1a). Overall, reptile richness was
dominated by lizard species (N = 3695), followed by snakes (N = 2305), while turtle species
only contributed marginally to the total number of modeled species (N = 296, Table 1, Fig. 1b).

For future conditions, a large number of reptile species were projected to disappear and at the
same time a large number of new species were projected to appear in Brazil and Australia, while
other regions showed either a strong species decrease or increase (Fig. 1 c, d). The strongest
future decreases of species richness were projected east of the Caspian Sea and in South Africa
(Fig. 1 c), while strong future increases were projected in the south-west of China and in the
eastern United States (Fig. 1 d). Overall, projected richness decrease was stronger than richness
increase, which resulted in a stronger net loss in species richness from 1995 to 2080 (Fig. 1 c, d,
e). The lowest net loss in species richness was projected for Brazil, Australia and South Africa,
while the highest net gain was projected for south-west China and the western United States (Fig.
1 e). Relative change (%) was projected to be negative in particular for most of the southern
hemisphere, while the high northern latitudes showed a strong positive relative change (Fig. 1 f).

Spatial patterns in species richness changes varied strongly across the three taxa, with lizards
seeing both strong increases and decreases in Australia, snakes showing a strong decrease in
South America and turtles seeing a strong increase in eastern part of North America (Fig. S2.9 &
Fig. S2.11). All three taxa showed a net gain in species richness in northern latitudes, while
lizards showed the strongest net loss in parts of Australia, snakes in large parts of South America
and turtles in parts of South America and southern Africa (Fig. S2.10 & Fig. S2.12).

Global and zoogeographic realm changes

Summed up across all 0.5° x 0.5° grid cells globally, species richness was projected to decline
significantly ($p < 0.01$) from 1995 to 2080 globally, with a decline in mean reptile richness per
grid cell from 58.4 ± 0.24 (SE) in 1995 to 53.39 ± 0.19 in 2080 (Fig. 1b, Fig. 2 a). 8 out of 11
zoogeographic realms showed a significant decline in reptile richness by 2080 (Fig. 2 b, c, d, f, g,
h, j, k), while the Nearctic and Palearctic realm showed a significant increase (Fig. 2 e, i) and the
Sino-Japanese realm showed no significant change (Fig. 2 l).

Looking at the global averages of species richness for each 0.5° grid cell for the three reptile
groups separately, snakes had the highest mean species richness (μ_{mean} = 30.4 ± 0.15), followed
by lizard (μ_{mean} = 24.3 ± 0.09 SE) and turtle richness (μ_{mean} = 3.71 ± 0.02 SE; Fig. 1b, Fig. S2.8).
Globally, similar to the total reptile richness the individual taxonomic groups (lizards, snakes and
turtles) all showed a significant decline in species richness , while there were slight differences

across the individual realms (Fig. S2.13 – S2.15). Lizards only showed a significant increase in
richness in the Palearctic realm and no significant change in richness in the Sino-Japanese realm,
while in all other realms they showed a significant decrease (Fig. S2.13). Snake and turtle
richness increased significantly in the Nearctic and Palearctic realm. Snake richness decreased
significantly in all other realms apart from the Sino-Japanese one (Fig. S2.14), while turtle
richness significantly decreased in all other realms apart from the Saharo-Arabian and the Sino-
Japanese one (Fig. S2.15).

Biophysical changes

Reptile richness strongly differed across conditions of varying combinations of temperature and
precipitation (Fig. 3 a, b, c). For 1995 reptile richness was projected to be highest in areas with a
temperature around 28.5°C, a precipitation of about 5500 mm and when considering temperature
and precipitation together in warm, but moist regions (21°C & 3000 mm, Fig. 3 c). The climatic
conditions with the highest richness shifted to even more extreme (warmer & wetter) novel
climate conditions by 2080 (Fig. 3 a, b, d).

Looking at the species richness change across the 2-dimensional climate space, net change was
positive at the upper precipitation limits across all temperatures and the very hot and very dry
conditions and negative throughout the entire precipitation range especially for the higher
temperatures. Overall, the negative change was much stronger and more pronounced than the
positive net change (Fig. 3 e). The highest positive and negative relative change values were
clustered, both occurred at the upper precipitation limits at low and medium temperatures (Fig. 3
f). A considerable proportion of the climate space (29.5 %) was shifting towards novel climatic
conditions, for which no change in species richness could be estimated, while only few discrete
climatic conditions as well as very cold & very dry conditions (4.75 %) got lost (Fig. 3 e, f).

Species-specific range changes

The range extent of most species (N = 6021) showed a considerable decrease (μ_{mean} = -27.7 ±
0.16 SE, Fig. 4 a, c, e). Lizard species showed the strongest decline (μ_{mean} = -31.8 ± 0.22 SE) in
range extent (Fig. 4 a), followed by snakes (μ_{mean} = -22.6 ± 0.25 SE), while almost equal numbers
of turtle species showed a decline (N = 274) and an increase (n = 205) with decreases being
much more pronounced than increases (μ_{mean} = -17.5 ± 0.72 SE, Fig. 4 e). Almost half of the
modeled reptile species (N = 3029) showed a strong change in range position, demonstrated by a
relatively low range overlap (<= 60 %), which was consistent across all three groups (Fig. 4 b, d,
f).

Most of the range centroids (58 %) of all reptile species fell within the Neotropical (N = 1133),
Afrotropical (N = 1039), Oriental (N = 785) and Australian realm (N = 698). Turtle species had
50 % of their range centroids in the Nearctic (N = 58), Oriental (N = 53) and Afrotropical realm
(N = 38), while lizards and snakes reflected the overall, total reptile patterns (Fig. 5 d). Range
centroids were highly clustered within the different realms, which reflects the overall richness
hotspots, and hardly any centroids were found in the high northern latitudes (Fig. 5 d). By 2080
species centroids were projected to shift by a mean distance of 111 km ± 0.9 SE mostly towards
a southern direction. Lizards showed a shift towards all directions, with a slight trend in the
number of species towards the South (Fig. 5 a), while snakes and turtles show a more
pronounced shift of species towards the North (Fig. 5 b, c). Turtle ranges shifted by the largest
distances, followed by snakes (Fig. 5 a, b, c). The northern realms (Nearctic, Saharo-Arabian,

Paleartic und Sino-Japanese) showed a dominant shift towards the North, while the southern
realms (Neotropical, Afrotropical and Australian) showed a dominant shift towards the South.
This was also reflected in the taxon-specific range shifts for the Northern and Southern
hemisphere (Fig. S3.16 in Appendix S3). The Panamanian, Madagascan and Oriental realms also
showed a northerly shift, while the Oceanian realm showed a bi-directional shift to the
Northwest and Southeast (Fig. 5 d, Fig. S3.17). Large realms had a larger proportion of species
that shifted their range for a larger distance (Fig. 5, Fig. S3.17).

Non-modeled species

37.4 % of reptile species for which data would have been available could not be modeled using
SDMs, either due to a small sample size or a low model performance (Table 1). We found that
the species that could not be modeled showed a significantly higher ($p < 0.05$) mean climatic
distance between current and future conditions compared to the modeled species and thus
occurred in areas that are projected to experience a stronger change in climatic conditions. This
pattern was consistent across all three taxa, although the difference was less significant in turtles
($p < 0.05$) compared to the other three taxa ($p < 0.0001$; Fig. 6).

Sensitivity analysis

Looking at the sum of occurrence probabilities, we found similar spatial patterns and a similar
magnitude in change compared to the sum of thresholded occurrences (species richness) (Fig.
S4.18 - S4.20). Projected richness values and their future changes were slightly larger under a
larger dispersal ability (d/4), but overall all results were consistent across the three dispersal
scenarios considered (Fig. S5.21 - S5.26). Climate change impacts on future species richness
increased over time, with stronger effects seen for 2080 than 2050, and the strongest impacts
being observed under a high emission scenario (RCP8.5) compared to the two lower scenarios
(Fig. S6.27 - S6.38).

Discussion

354 Reptile richness was projected to decrease significantly across most parts of the world in the
355 future (Fig. 1 & 2). This effect was visible across lizards, snakes and turtles alike, although
356 regional and species-specific responses differed across the three groups (Fig. S2.8 – S2.12).

Spatial changes

Reptile richness is projected to decrease in Brazil, Australia and South Africa (Fig. 1 c, e). These
areas overlap strongly with the biotic convergence zones, areas with a high spatial concentration
of Lepidosaurians (i.e. snakes and lizards) identified by Diele-Viegas *et al.* (2020), which were
also found to cover a large number of Lepidosaurian species vulnerable to climate change (Diele-
Viegas *et al.*, 2020). Huey *et al.* (2012) further found that ectotherms sharing climate
vulnerability traits seem to be concentrated in lowland tropical forests. Combining climate-based
SDMs with land-use change information Newbold (2018) created future projections for 20938
vertebrate species and found that Brazil is strongly affected by vertebrate diversity loss due to
climate change, and together with Australia is also likely to be strongly affected by future land-
use changes, especially under a high-emission scenario (RCP8.5). Given that Brazil in particular
does not only host a high reptile richness (Roll *et al.*, 2017; Fig. 1 a), but has recently been found
to inhabit a large number of threatened reptile species (Cox *et al.*, 2022), this highlights the

responsibility of this mega-diverse country for also protecting reptile diversity. South-west China
and the western United States were projected to show a net gain in reptile richness under climate
warming (Fig. 1 e), but they also belong to those areas where most reptile species are threatened
by habitat loss from agriculture and logging or harvesting (Böhm *et al.*, 2013). These projected
losses through habitat change could potentially counteract positive effects climate warming
might have.

The strong variation of species richness changes across regions and taxa, that we found, are,
obviously, reflecting their original richness patterns. Species richness of amphibians, birds and
mammals together is a good spatial surrogate for species richness of all reptiles combined and of
snakes, but not for lizard or turtle richness (Roll *et al.*, 2017). Thus, it is not surprising that the
areas with the highest decline in overall reptile richness (see Fig. 1) strongly overlap with the
areas of highest projected changes in vertebrate species richness (amphibians, birds and
mammals) found by Hof *et al.* (2018), although global reptile richness is largely constrained by
temperature, while global richness of all other vertebrate groups is mostly constrained by the
availability of energy and water (Qian, 2010). Historical shifts in geographical ranges and
climatic niches further showed that niche shifts in endotherms are significantly faster than in
ectotherms (Rolland *et al.*, 2018).

Global and zoogeographic realm changes

Globally reptile richness was projected to decline significantly, from about 58 to 53 (9.4 %)
species on average per grid cell from 1995 to 2080 (Fig. 2 a). This projected percentage change
in average future reptile richness (9.4 %) is considerably lower than the reptile richness decline
due to climate warming predicted by Newbold (2018). However, while Newbold (2018) also use
SDMs to infer future reptile richness changes, he only used a subset of reptile species, for which
IUCN range maps were available back then. In addition, he also applied a much smaller dispersal
buffer (0.5 km per year), which might indicate that our projections provide a rather optimistic
scenario. Newbold (2018) further found that reptiles together with amphibians are
disproportionately sensitive to future human land-use. Given the synergistic effect of future
climate and land-use change on biodiversity (Brook *et al.*, 2008) as well as species populations
(Williams *et al.*, 2022), land-use change is likely to further exacerbate climate change impacts on
global reptile distribution and diversity.

Changes in reptile richness differed among zoogeographic realms, but species richness declined
significantly across most realms over both time periods (Fig. 2 b, c, d, f, g, h, j, k). Lizards,
snakes and turtles all showed similar declines in global species richness and across most realms,
but slightly differed across individual realms. This is in line with a previous study covering
various realms from tropical to temperate regions which found that 60 % of assessed
Lepidosaurian species (N = 1114) were vulnerable to changes in climate (Diele-Viegas *et al.*,
2020). Diele-Viegas *et al.* (2020) further found that the Afrotropical, Nearctic and Sino-Japanese
realms were the three realms where Lepidosaurians were most vulnerable to climatic change,
while Lepidosaurians in the Madagascan and Oceanian realm were least vulnerable. By contrast,
we found no significant decline in projected total reptile richness for the Sino-Japanese realm
(Fig. 2 l), although individual subgroups were projected to show a significant decline in species
richness from 1995 to 2050 (Fig. S2.13 – S2.15). Both the Madagascan and Oceanian realm
were further projected to significantly decrease in species richness both for all reptile species
(Fig. 2 d, g) and all three subgroups (Fig. S2.13 – S2.15). However, given that the Madagascan

realm, specifically, boasts over 90% of endemic reptile species and genera (Glaw & Vences,
2007) and that both realms are composed of island territories which are usually considered
highly vulnerable to climate change and might also be affected by future sea level rise and
erosion (Diele-Viegas *et al.*, 2020), our estimates might still underestimate potential climate
change impacts in these realms. Interestingly, turtle richness seemed to be least affected by
climate change, when looking at the different realms (Fig. S2.15), although a recent review has
deemed turtles as the vertebrate group with the highest extinction risk, being strongly affected by
habitat loss, human consumption and pet trading (Stanford *et al.*, 2020).

Biophysical changes

Reptile richness differed strongly across temperature and precipitation, with the highest richness
being observed in warm and moist conditions. Under future climate change, the climatic
conditions with high species richness were projected to shift to even more extreme (warmer &
wetter) conditions (Fig. 3). This is cause for concern given that tropical forest and desert lizards
already live in environmental conditions that are close to their thermal limits (Sinervo *et al.*,
2010), while desert and temperate lizard species have been found to be more constrained in their
thermoregulatory ability to deal with heat stress than tropical species (Anderson *et al.*, 2022).
Reptiles cannot regulate their body temperature internally, so are strongly dependent on solar
energy captured by the environment to regulate their body temperature (Huey, 1982). This might
lead to overheating when temperatures reach beyond a species' critical limit, which makes them
particularly susceptible to climatic changes (Sinervo *et al.*, 2018). However, this might be
compensated by other biological processes that help species to buffer climate change effects, i.e.
genomic and phenotypic plasticity (Rodríguez *et al.*, 2017) as well as behavioral and
physiological adaptation (Sunday *et al.*, 2014). Overall, the persistence of reptile species would
be much more affected by climate cooling than warming, but increasing droughts, which will be
a consequence of continued warming, have been suggested to pose a significant future threat to
European reptiles (Araújo *et al.*, 2006). Climate warming will likely have an additional impact
on reptiles that have temperature-dependent sex determination. Altered sex ratios will not only
result in a higher extinction risk of local populations, but, together with a reduction in nesting
sites due to habitat destruction and fragmentation, will also affect the dispersal and potential
range expansion of a species. Therefore it could also have an impact on population demography
and size unless temperature shifts in sex determination or female nest-site choice evolves in pace
with rising temperatures (Boyle *et al.*, 2016; Gibbons *et al.*, 2000).

Species-specific range changes

The range extent of most species considerably decreased, with lizard species showing the
strongest decline (Fig. 4 a). Most reptiles further showed a strong decline in range overlap, which
was consistent across all three groups (Fig. 4). This is in line with results by Warren *et al.*
(2018), who found that projected future range losses of more than 50% occur in 8 - 52% of
considered reptile species by 2100 depending on the climate scenario considered, although this
study included only a fraction of all reptile species (N=1850) and no species dispersal was
considered.

Compared to other terrestrial vertebrate groups (especially birds and mammals), reptile have
small geographic ranges, indicating also narrower niche requirements. This is likely to make
them more susceptible to future climate change (Newbold, 2018), but also to other threats such

as habitat loss or invasive species (Böhm *et al.*, 2013, Cox *et al.*, 2022). Reptiles are a
paraphyletic class with a diverse range of body forms, habitat affinities and functional roles
(Pincheira-Donoso *et al.*, 2013), likely resulting in their response to climatic and habitat changes
to be equally varied. In addition, cascading effects generated by disease, invasive species, habitat
loss and climate change might lead to declines of sympatric species and a faster deterioration of
ecosystem structure than anticipated by climate change alone (Zipkin *et al.*, 2020).

The majority of reptile species showed projected shifts towards the South, which was largely
driven by range shifts in lizards (Fig. 5). Comparing the Northern and Southern hemisphere (Fig.
S3.16), we found a clear overall poleward shift in species ranges across all three groups, which
has also been previously found in various other taxonomic groups (Chen *et al.*, 2011). Turtle
ranges were projected to shift farthest, followed by snakes (Fig. 5). The relatively short range
shift distances in lizard species are probably due to the fact that lizards have the smallest range
extent across the three groups (Roll *et al.*, 2017), which, given that our dispersal buffers are
based on range extent, also resulted in smaller dispersal distances for lizards compared to the two
other groups (Fig. S1.4). This is, however, obviously somewhat counterintuitive and in stark
contrast to generally perceived differences in mobility characteristics (e.g. movement speed)
among lizards, snakes and turtles. While it is hard to tackle this problem in a globally and
taxonomically comprehensive study as ours, it calls for more efforts in collecting data on actual,
empirically quantified dispersal distances in order to consider them in SDM exercises and large-
scale biogeographical analyses.

Non-modeled species

Range-restricted (non-modeled) species are projected to experience significantly higher shift in
climatic distance than modeled species (Fig. 6), indicating that range-restricted species are also
likely to be strongly affected by climate change. This highlights once more that sample size
restrictions of SDMs are likely to downplay climate change effects on narrow-ranging and
threatened species (Platts *et al.*, 2014). Hof *et al.* (2018) also found significant impacts of climate
and land-use changes on range-restricted vertebrate species, excluding reptiles. However, similar
to the latter study we only look at climate anomalies (Euclidean distance between current and
future climatic conditions) as a metric of climate change, while different metrics have been
found to show contrasting climate change patterns on a global scale (Garcia *et al.*, 2014). In
addition to climate change effects, habitat modification has been found to have a stronger impact
on small-ranging reptile species, as well as species with a small clutch size (Doherty *et al.*,
2020). Range-restricted reptile species are further often evolutionarily unique (Murali *et al.*,
2021) and have been found to overlap least with current conservation priority areas (Roll *et al.*,
2017, Cox *et al.*, 2022).

Sensitivity analysis

Our results were strongly dependent on the dispersal assumption, time period and emission
scenario (RCP) considered (see Appendix S5 & S6). As expected, the overall patterns and
richness changes were more pronounced in a later time period and scenarios representing higher
levels of greenhouse gas emissions, as these reflect potential futures with a higher level of
climate warming. Thuiller *et al.* (2019) have previously assessed the uncertainty originating from
dispersal, model algorithm, GCM and RCP on the future biodiversity scenario of amphibians,
birds and mammals and found that model algorithm and RCP have the strongest influence.

On the contrary, larger dispersal distances imply that reptile species are able to move larger
distances in order to track their optimal climatic niche and thus provide more optimistic potential
changes in species occurrence and richness patterns. Reptile-specific studies have either
considered no dispersal at all (Araújo *et al.*, 2006; Warren *et al.*, 2018) or a dispersal rate of 0.5
km per year (Newbold, 2018). We use species-specific dispersal buffers with an average of 2.4
km per year (Fig. S1.4). While these buffers might be optimistic, they are based on the
transparent rationale of a range-size dependent dispersal buffer, i.e. species with a small range
size have a smaller dispersal buffer than species with a large range size, which avoids the
unlikely assumption of uniform dispersal distances across species. Given that our model results,
as well as the underlying climate scenarios, are based on a 0.5° grid size (ca. 50 x 50 km), small
differences in dispersal distance do not have a strong impact on our results (see Appendix S5).
Nevertheless, these considerations again highlight the challenges to sensibly account for the
influence of dispersal in global climate impact assessments on species distributions and diversity.

The projected changes in species distributions help to investigate potential changes in global
reptile richness patterns and to highlight potential hotspots of climate change impacts. They also
allow to compare climate change vulnerability across taxonomic groups and help to identify
areas where conservation efforts might be most urgently needed (Voskamp et al. 2022).
However, SDMs are always simplifications and the resulting projections have to be interpreted
with caution.

To improve SDMs, future studies should try to consider additional factors, such as biotic
interactions (Schleuning *et al.*, 2020) and the reshuffling of species communities (Voskamp *et
al.*, in prep), which might lead to a change in competitive balance (Ockendon *et al.*, 2014),
altered predator-prey relationships (Harley, 2011) or changes in functional diversity (Stewart *et
al.*, 2022) and thus the provision of ecosystem functions and services (Pecl *et al.*, 2017). While
the above factors might improve climate change impact projections in the future, such modelling
exercises will never reflect the truth, as a species' response to climate change will be strongly
influenced by behaviour, diel rhythm, thermoregulatory potential and microclimatic conditions
(Anderson *et al.*, 2022). This complexity highlights the need for integrative approaches when
investigating species' responses to climate change (Hof, 2021).

Conclusion

Our study shows that reptiles are likely to be impacted by future climate change, globally but
also within most zoogeographic realms. These impacts are projected to have a considerable
effect on the extent and location of species' geographic ranges. Thus, to prevent large scale
declines in reptile species it is pivotal to lower CO_2 emissions in order to stop on-going climate
change but also to maintain adequate habitats of sufficient size and quality, especially of
grassland and savanna habitats (Roll *et al.*, 2017). Furthermore, it is necessary to establish new
protected areas that help to prevent the extinction of particularly vulnerable species, i.e. by
establishing high-elevation climate refugia within current species ranges (Sinervo *et al.*, 2018).

References

References

Allouche, O., Tsoar, A. & Kadmon, R. (2006) Assessing the accuracy of species distribution
models: Prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, **43**,
1223–1232.

Anderson, R. O., Meiri, S., & Chapple, D. G. (2022) The biogeography of warming tolerance
in lizards. *Journal of Biogeography*.

Araújo, M.B., Thuiller, W. & Pearson, R.G. (2006) Climate warming and the decline of
amphibians and reptiles in europe. *Journal of Biogeography*, **33**, 1712–1728.

Bagchi, R., Crosby, M., Huntley, B., Hole, D.G., Butchart, S.H., Collingham, Y., Kalra, M.,
Rajkumar, J., Rahmani, A., Pandey, M. & others (2013) Evaluating the effectiveness of
conservation site networks under climate change: Accounting for uncertainty. *Global
Change Biology*, **19**, 1236–1248.

Baisero, D., Visconti, P., Pacifici, M., Cimatti, M. & Rondinini, C. (2020) Projected global loss
of mammal habitat due to land-use and climate change. *One Earth*, **2**, 578–585.

Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences
for species distribution models: how, where and how many?. *Methods in Ecology and
Evolution*, *3*(2), 327-338.

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. (2012) Impacts of
climate change on the future of biodiversity. *Ecology Letters*, **15**, 365–377.

Biber, M.F., Voskamp, A., Niamir, A., Hickler, T. & Hof, C. (2020) A comparison of
macroecological and stacked species distribution models to predict future global terrestrial
vertebrate richness. *Journal of Biogeography*, **47**, 114–129.

Böhm, M., Collen, B., Baillie, J.E., Bowles, P., Chanson, J., Cox, N., Hammerson, G., Hoffmann,
M., Livingstone, S.R., Ram, M. & others (2013) The conservation status of the world's
reptiles. *Biological Conservation*, **157**, 372–385.

Bowler, D.E., Hof, C., Haase, P., Kröncke, I., Schweiger, O., Adrian, R., Baert, L., Bauer, H.-G.,
Blick, T., Brooker, R.W. & others (2017) Cross-realm assessment of climate change impacts
on species' abundance trends. *Nature Ecology & Evolution*, **1**, 1–7.

Boyle, M., Schwanz, L., Hone, J. & Georges, A. (2016) Dispersal and climate warming
determine range shift in model reptile populations. *Ecological Modelling*, **328**, 34–43.

Brook, B.W., Sodhi, N.S., & Bradshaw, C.J.A. (2008) Synergies among extinction drivers
under global change. *Trends in Ecology & Evolution*, **23**, 453–460.

Brooks, T. M., Mittermeier, R. A., Da Fonseca, G. A., Gerlach, J., Hoffmann, M., Lamoreux, J. F.,
& others (2006) Global biodiversity conservation priorities. S*cience*, **313**, 58-61.

Calabrese, J.M., Certain, G., Kraan, C. & Dormann, C.F. (2014) Stacking species distribution models and adjusting bias by linking them to macroecological models. *Global Ecology and Biogeography*, **23**, 99–112.

Chen, I.-C., Hill, J.K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. (2011) Rapid range shifts of species associated with high levels of climate warming. *Science*, **333**, 1024–1026.

Cox, N., Young, B. E., Bowles, P., Fernandez, M., Marin, J., Rapacciuolo, G. & others (2022) A global reptile assessment highlights shared conservation needs of tetrapods. *Nature*, **605**, 285-290.

Diele-Viegas, L.M., Figueroa, R., Vilela, B. & Rocha, C. (2020) Are reptiles toast? A worldwide evaluation of lepidosauria vulnerability to climate change. *Climatic Change*, **159**, 581–599.

Diele-Viegas, L. M., & Rocha, C. F. D. (2018) Unraveling the influences of climate change in Lepidosauria (Reptilia). *Journal of Thermal Biology*, **78**, 401-414.

Doherty, T.S., Balouch, S., Bell, K., Burns, T.J., Feldman, A., Fist, C., Garvey, T.F., Jessop, T.S., Meiri, S. & Driscoll, D.A. (2020) Reptile responses to anthropogenic habitat modification: A global meta-analysis. *Global Ecology and Biogeography*, **29**, 1265–1279.

Elith, J., Kearney, M. & Phillips, S. (2010) The art of modelling range-shifting species. *Methods in Ecology and Evolution*, **1**, 330–342.

Elith, J. & Leathwick, J.R. (2009) Species distribution models: Ecological explanation and prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*, **40**, 677–697.

Frieler, K., Lange, S., Piontek, F., Reyer, C.P., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel, K. & others (2017) Assessing the impacts of 1.5°c global warming–simulation protocol of the inter-sectoral impact model intercomparison project (ISIMIP2b). *Geoscientific Model Development*, **10**, 4321–4345.

Garcia, R.A., Cabeza, M., Rahbek, C. & Araújo, M.B. (2014) Multiple dimensions of climate change and their implications for biodiversity. *Science*, **344**, 1247579.

Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tuberville, T.D., Metts, B.S., Greene, J.L., Mills, T., Leiden, Y., Poppy, S. & others (2000) The global decline of reptiles, déjà vu amphibians: Reptile species are declining on a global scale. Six significant threats to reptile populations are habitat loss and degradation, introduced invasive species, environmental pollution, disease, unsustainable use, and global climate change. *BioScience*, **50**, 653–666.

Gibson-Reinemer, D.K. & Rahel, F.J. (2015) Inconsistent range shifts within species highlight idiosyncratic responses to climate warming. *PloS One*, **10**, e0132103.

Glaw, F. & Vences, M. (2007) *A field guide to the amphibians and reptiles of madagascar*, Vences & Glaw.

Greenwell, B., Boehmke, B., Cunningham, J. & Developers, G. (2020) *GBM: Generalized*
boosted regression models.

Guisan, A., & Thuiller, W. (2005) Predicting species distribution: offering more than simple
habitat models. *Ecology Letters*, *8*(9), 993-1009.

Harley, C.D. (2011) Climate change, keystone predation, and biodiversity loss. *Science*, **334**,
1124–1127.

Hernandez, P. A., Graham, C. H., Master, L. L., & Albert, D. L. (2006) The effect of sample size
and species characteristics on performance of different species distribution modeling
methods. *Ecography*, **29**, 773-785.

Hijmans, R.J., Phillips, S., Leathwick, J. & Elith, J. (2021) *Dismo: Species distribution modeling*.

Hof, C. (2021) Towards more integration of physiology, dispersal and land-use change to
understand the responses of species to climate change. *Journal of Experimental Biology*,
224, jeb238352.

Hof, C., Araújo, M.B., Jetz, W. & Rahbek, C. (2011) Additive threats from pathogens, climate
and land-use change for global amphibian diversity. *Nature*, **480**, 516–519.

Hof, C., Voskamp, A., Biber, M.F., Böhning-Gaese, K., Engelhardt, E.K., Niamir, A., Willis, S.G. &
Hickler, T. (2018) Bioenergy cropland expansion may offset positive effects of climate
change mitigation for global vertebrate diversity. *Proceedings of the National Academy of*
Sciences, **115**, 13294–13299.

Holt, B.G., Lessard, J.-P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D., Fabre, P.-H.,
Graham, C.H., Graves, G.R., Jønsson, K.A. & others (2013) An update of wallace's
zoogeographic regions of the world. *Science*, **339**, 74–78.

Huey, R.B. (1982) *Temperature, physiology, and the ecology of reptiles. Biology of the reptilia*,
Citeseer.

Huey, R.B., Kearney, M.R., Krockenberger, A., Holtum, J.A., Jess, M. & Williams, S.E. (2012)
Predicting organismal vulnerability to climate warming: Roles of behaviour, physiology and
adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**,
1665–1679.

IPBES (2019) Summary for policymakers of the global assessment report on biodiversity
and ecosystem services of the intergovernmental science-policy platform on biodiversity
and ecosystem services. *IPBES secretariat, Bonn, Germany*.

IPCC (2021) Climate change 2021: The physical science basis. Working group i contribution
to the sixth assessment report of the intergovernmental panel on climate change.
Cambridge University Press.

IPCC (2022) Climate change 2022: Impacts, adaptation and vulnerability. Working group II contribution to the sixth assessment report of the intergovernmental panel on climate change. *Cambridge University Press*.

IUCN (2022) The IUCN red list of threatened species. Available at https://www.iucnredlist.org/resources/spatial-data-download. *Cambridge University Press*.

Kühn, I. (2007) Incorporating spatial autocorrelation may invert observed patterns. *Diversity and Distributions*, **13**, 66–69.

Lange, S. (2016) EartH2Observe, WFDEI and ERA-interim data merged and bias-corrected for ISIMIP (EWEMBI).

Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J. & Grenouillet, G. (2020) Species better track climate warming in the oceans than on land. *Nature Ecology & Evolution*, **4**, 1044–1059.

Meynard, C.N. & Quinn, J.F. (2007) Predicting species distributions: A critical comparison of the most common statistical models using artificial species. *Journal of Biogeography*, **34**, 1455–1469.

Monasterio, C., Shoo, L. P., Salvador, A., Iraeta, P., & Díaz, J. A. (2013) High temperature constrains reproductive success in a temperate lizard: implications for distribution range limits and the impacts of climate change. *Journal of Zoology*, **291**, 136-145.

Murali, G., Gumbs, R., Meiri, S., & Roll, U. (2021) Global determinants and conservation of evolutionary and geographic rarity in land vertebrates. *Science Advances*, **7(42)**, eabe5582.

Nathan, R., Klein, E., Robledo-Arnuncio, J.J. & Revilla, E. (2012) *Dispersal kernels*, Oxford University Press Oxford, UK.

Newbold, T. (2018) Future effects of climate and land-use change on terrestrial vertebrate community diversity under different scenarios. *Proceedings of the Royal Society B*, **285**, 20180792.

Ockendon, N., Baker, D.J., Carr, J.A., White, E.C., Almond, R.E., Amano, T., Bertram, E., Bradbury, R.B., Bradley, C., Butchart, S.H. & others (2014) Mechanisms underpinning climatic impacts on natural populations: Altered species interactions are more important than direct effects. *Global Change Biology*, **20**, 2221–2229.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V., Underwood, E.C., D'amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C. & others (2001) Terrestrial ecoregions of the world: A new map of life on EarthA new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. *BioScience*, **51**, 933–938.

Pacifici, M., Foden, W.B., Visconti, P., Watson, J.E., Butchart, S.H., Kovacs, K.M., Scheffers, B.R.,
Hole, D.G., Martin, T.G., Akçakaya, H.R. & others (2015) Assessing species vulnerability to
climate change. *Nature Climate Change*, **5**, 215–224.

Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.-C., Clark, T.D.,
Colwell, R.K., Danielsen, F., Evengård, B. & others (2017) Biodiversity redistribution under
climate change: Impacts on ecosystems and human well-being. *Science*, **355**, eaai9214.

Petitpierre, B., Broennimann, O., Kueffer, C., Daehler, C., & Guisan, A. (2017) Selecting
predictors to maximize the transferability of species distribution models: Lessons from
cross-continental plant invasions. *Global Ecology and Biogeography*, **26**, 275-287.

Pincheira-Donoso, D., Bauer, A. M., Meiri, S., & Uetz, P. (2013) Global taxonomic diversity of
living reptiles. *PloS One*, **8**, e59741.

Platts, P.J., Garcia, R.A., Hof, C., Foden, W., Hansen, L.A., Rahbek, C. & Burgess, N.D. (2014)
Conservation implications of omitting narrow-ranging taxa from species distribution
models, now and in the future. *Diversity and Distributions*, **20**, 1307–1320.

Porfirio, L. L., Harris, R. M., Lefroy, E. C., Hugh, S., Gould, S. F., Lee, G. & others (2014)
Improving the use of species distribution models in conservation planning and
management under climate change. *PLoS One*, **9**, e113749.

Qian, H. (2010) Environment–richness relationships for mammals, birds, reptiles, and
amphibians at global and regional scales. *Ecological Research*, **25**, 629–637.

R Core Team (2021) *R: A language and environment for statistical computing*, R Foundation
for Statistical Computing, Vienna, Austria.

Rodríguez, A., Rusciano, T., Hamilton, R., Holmes, L., Jordan, D. & Wollenberg Valero, K.C.
(2017) Genomic and phenotypic signatures of climate adaptation in an anolis lizard.
Ecology and Evolution, **7**, 6390–6403.

Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A.M., Bernard, R., Böhm, M., Castro-
Herrera, F., Chirio, L., Collen, B. & others (2017) The global distribution of tetrapods reveals
a need for targeted reptile conservation. *Nature Ecology & Evolution*, **1**, 1677–1682.

Rolland, J., Silvestro, D., Schluter, D., Guisan, A., Broennimann, O. & Salamin, N. (2018) The
impact of endothermy on the climatic niche evolution and the distribution of vertebrate
diversity. *Nature Ecology & Evolution*, **2**, 459–464.

Scheffers, B.R., De Meester, L., Bridge, T.C., Hoffmann, A.A., Pandolfi, J.M., Corlett, R.T.,
Butchart, S.H., Pearce-Kelly, P., Kovacs, K.M., Dudgeon, D. & others (2016) The broad
footprint of climate change from genes to biomes to people. *Science*, **354**, aaf7671.

Schleuning, M., Neuschulz, E. L., Albrecht, J., Bender, I. M., Bowler, D. E., Dehling, D. M. &
others (2020) Trait-based assessments of climate-change impacts on interacting species.
Trends in Ecology & Evolution, **35**(4), 319-328.

Sinervo, B., Mendez-De-La-Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Villagrán-Santa
Cruz, M. & others (2010) Erosion of lizard diversity by climate change and altered thermal
niches. *Science*, **328**, 894-899.

Sinervo, B., Miles, D.B., Wu, Y., MÉNDEZ-DE LA CRUZ, F.R., Kirchhof, S. & Qi, Y. (2018)
Climate change, thermal niches, extinction risk and maternal-effect rescue of toad-headed
lizards, phrynocephalus, in thermal extremes of the arabian peninsula to the qinghai—
tibetan plateau. *Integrative Zoology*, **13**, 450–470.

Stanford, C. B., Iverson, J. B., Rhodin, A. G., van Dijk, P. P., Mittermeier, R. A., Kuchling, G. &
others (2020) Turtles and tortoises are in trouble. *Current Biology*, **30**, R721-R735.

Stewart, P.S., Voskamp, A., Santini, L., Biber, M.F., Hof, C., Willis, S.G. & Tobias, J.A. (2022)
Global impacts of climate change on avian functional diversity. *Ecology Letters*.

Sunday, J.M., Bates, A.E., Kearney, M.R., Colwell, R.K., Dulvy, N.K., Longino, J.T. & Huey, R.B.
(2014) Thermal-safety margins and the necessity of thermoregulatory behavior across
latitude and elevation. *Proceedings of the National Academy of Sciences*, **111**, 5610–5615.

Thuiller, W., Guéguen, M., Renaud, J., Karger, D.N. & Zimmermann, N.E. (2019) Uncertainty
in ensembles of global biodiversity scenarios. *Nature Communications*, **10**, 1–9.

Thuiller, W., Lavorel, S., Araújo, M.B., Sykes, M.T. & Prentice, I.C. (2005) Climate change
threats to plant diversity in europe. *Proceedings of the National Academy of Sciences*, **102**,
8245–8250.

Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K. & others
(2011) The representative concentration pathways: an overview. *Climatic change*, **109**, 5-
31.

Voskamp, A., Baker, D.J., Stephens, Philip A, Howard, C., Butchart, S.H. & Willis, S.G. (in prep)
Impacts of climate change on the world's terrestrial bird assemblages.

Voskamp, A., Butchart, S.H., Baker, D.J., Wilsey, C.B. & Willis, S.G. (2021) Site-based
conservation of terrestrial bird species in the caribbean and central and south america
under climate change. *Frontiers in Ecology and Evolution*, **9**, 234.

Voskamp, A., Fritz, S. A., Koecke, V., Biber, M., Nogueira-Brockmeyer, T., Bertzky, B. & others
(2022) How to resolve conflicting conservation objectives: A decision support tool for the
global selection of multi-purpose protected areas. *bioRxiv*.

Warren, R., Price, J., Graham, E., Forstenhaeusler, N. & VanDerWal, J. (2018) The projected
effect on insects, vertebrates, and plants of limiting global warming to 1.5°c rather than 2°c.
Science, **360**, 791–795.

Watterson, I. G. (2019) Indices of climate change based on patterns from CMIP5 models,
and the range of projections. *Climate Dynamics*, **52**, 2451-2466.

Wiens, J.J. & Graham, C.H. (2005) Niche conservatism: Integrating evolution, ecology, and
conservation biology. *Annual Review of Ecology, Evolution, and Systematics*, **36**, 519–539.

Williams, J.J., Freeman, R., Spooner, F. & Newbold, T. (2022) Vertebrate population trends
are influenced by interactions between land use, climatic position, habitat loss and climate
change. *Global Change Biology*, **28**, 797–815.

Wood, S.N. (2011) Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. *Journal of the Royal Statistical*
Society (B), **73**, 3–36.

Wood, S.N. (2003) Thin-plate regression splines. *Journal of the Royal Statistical Society (B)*,
65, 95–114.

Zipkin, E.F., DiRenzo, G.V., Ray, J.M., Rossman, S. & Lips, K.R. (2020) Tropical snake diversity
collapses after widespread amphibian loss. *Science*, **367**, 814–816.

Data Accessibility Statement

760 GARD range maps are available from https://doi.org/10.5061/dryad.83s7k. EWEMBI and
761 ISIMIP2b climate data are available from https://esg.pik-potsdam.de/search/isimip/?
762 product=input. The code for the species distribution models can be found at
763 **https://github.com/xxxx/xxxx**, whereas the code and data for the performed data analysis and
764 the presented figures, can be found at **https://github.com/xxxx/xxxx**.

Tables

766 **Table 1.** Number of species that were excluded from the species distribution models due to their
767 restricted range or low model performance.

Taxonomic group	Lizard	Snake	Turtle	Total
No. of species with available data	6328	3414	322	10064
No. of range-restricted species (N < 10, removed)	2536	1047	19	3602
No. of species with low model performance (AUC < 0.7, removed)	97	62	7	166
Total number of species modeled	**3695**	**2305**	**296**	**6296**
Percentage of available species modeled	58.4 %	67.5 %	91.9 %	62.6 %

Figures

Figure 1. a) Map of projected global terrestrial reptile species richness (1995), b) frequency of
species richness by taxonomic group (lizard, snake, turtle and total) with mean values
highlighted by vertical line and c) increase, d) decrease, e) net change and f) relative change (%)
in reptile species richness for all modeled reptile species (N = 6296) for the year 2080 under a
medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).
Results are presented as the ensemble mean, across the four global circulation models (GCMs)
and two model algorithms (GAM & GBM) considered. All maps are based on 0.5° x 0.5° grid
cells, which have been projected to Mollweide equal-area projection (EPSG:54009). Grey areas
are regions for which no projections are available. Note that the colour scales differ among the
individual panels.

780
Figure 2. Terrestrial reptile species richness across the globe and for each zoogeographic realm
(Afrotropical, Australian, Madagascan, Nearctic, Neotropical, Oceanian, Oriental, Palearctic,
Panamanian, Saharo-Arabian & Sino-Japanese) over time (1995, 2080) based on all modeled
reptile species (N = 6296). Results are presented as the ensemble mean, across the four global
circulation models (GCMs) and two model algorithms (GAM & GBM) considered, under a
medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).
Statistical difference between years was tested using a paired Student's t-test with Holm
correction ($p < 0.05$ = *, $p < 0.01$ = **, $p < 0.001$ = ***, $p < 0.0001$ = ****). Plots show mean
(red point & label), median (black horizontal line), 25th to 75th percentiles (box), entire range of
data (violin & data points) and density of values (width of violin). Figure 5 provides a map
outlining the different zoogeographic realms.

792
Figure 3. Univariate relationship of current (1995) and future (2080 RCP6.0) reptile species
richness with a) temperature, b) precipitation and the bivariate relationship of temperature and
precipitation with reptile species richness for c) 1995 and d) 2080 RCP6.0 and the respective e)
net richness change and f) relative richness change (%) under a medium dispersal scenario (d/8).
Heatmaps and lines show the mean and ribbons the standard deviation in variance across space,
global circulation models (GCMs) and the two model algorithms (GAM & GBM).

Figure 4. Frequency plots of the mean number of reptile species (a) lizard, c) snake, e) turtle) and their potential future change (%) in range extent (total area of occupied grid cells) and the mean number of reptile species (b) lizard, d) snake, f) turtle) per potential range overlap class (0-20, 20-40, 40-60, 60-80, 80-100). Error margins/bars show standard deviation across the four global circulation models (GCMs) and the two model algorithms (GAM & GBM) used. Both shown for 2080 under a medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).

807
Figure 5. Cumulative direction and distance of potential range centroid changes per taxonomic
group (b) lizard, c) snake, d) turtle) and d) range centroids (points on map) and the number of
species and their directional shift in range centroid position per zoogeographic realm (inset polar
plots). Results are presented as the ensemble mean, across the four global circulation models
(GCMs) and two model algorithms (GAM & GBM) considered, for the year 2080 under a
medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).
Inset bar chart shows the number of species, which have their range centroid located in the
respective realm. Numbers show the total number of species per realm, while the bars are colour
coded by the three different taxonomic groups. Please note that a considerable number of species
(n = 518) have their range centroid located outside of the zoogeographic realm boundaries and
thus were not associated with an realm and so are not considered for the polar plots in Fig. 5 d).

Figure 6. Mean climatic distance for modeled and non-modeled species, split by taxonomic group (a) lizard, b) snake, c) turtle and d) total). Statistical difference between modeled and non-modeled species was tested using a Student's t-test with Holm correction ($p < 0.05$ = *, $p < 0.01$ = **, $p < 0.001$ = ***, $p < 0.0001$ = ****). Plots show mean (red point & label), median (black horizontal line), 25th to 75th percentiles (box), entire range of data (violin & data points) and density of values (width of violin). Results are shown as the ensemble mean across the four global circulation models (GCMs), for the year 2080 under a medium representative concentration pathway (RCP6.0).