Potential future climate change effects on global reptile distribution and diversity

Matthias F. Biber[1,*], Alke Voskamp[2], Christian Hof[1]

[1]Technical University of Munich, Terrestrial Ecology Research Group, Department for Life

Science Systems, School of Life Sciences, 85354 Freising, Germany

[2]Senckenberg Biodiversity and Climate Research Centre (SBiK-F), 60325 Frankfurt, Germany

Running head: Global reptiles under climate change

Abstract

Aim: Until recently, complete information on global reptile distributions has not been widely available. Here, we provide the first comprehensive climate impact assessment for reptile distributions at a global scale.

Location: Global, excluding Antarctica

Time period: 1995, 2050, 2080

Major taxa studied: Reptiles

Methods: We performed species distribution models for 6296 reptile species and assessed potential global as well as realm-specific changes in species richness, the change in global species richness across climatic space and species-specific changes in distribution and range extent and overlap, under future climate change. To assess the future climatic impact of 3768 non-modeled species, we compared the future change in climatic conditions between both modeled and non-modeled species.

Results: Reptile richness was projected to decline significantly over time, globally but also for most zoogeographic realms, with the strongest decrease in Brazil, Australia and South Africa. Species richness was highest in warm, but moist regions, which were projected to shift further to climate extremes in the future. Extents of occurrence were projected to decline considerably in the future, with a low overlap between projected current and future ranges. Shifts in range centroids differed among realms and taxa, with a dominating global poleward shift. Non-modeled species were significantly stronger affected by climatic changes than modeled species.

Main conclusions: Reptile richness was projected to decrease significantly across most parts of the world with ongoing future climate change. This effect is visible across lizards, snakes and turtles alike and has considerable impact on species' extent of occurrence (EOO) and range distribution. Together with other anthropogenic impacts, such as habitat loss and harvesting, this

32 is cause for concern. Given the historical lack of information on global reptile distributions, this
33 calls for an re-assessment of global conservation efforts towards reptile species, with specific
34 focus on anticipated future climatic changes.

35 **KEYWORDS:** species distribution model, turtle, snake, lizard, ISIMIP, bioclim, global
36 warming, environmental niche model

Introduction

Emissions from anthropogenic activities have lead to an increase in global surface temperature of
around 1°C in the last 100 years. This has already led to changes in weather and climate
extremes in every region across the globe (IPCC, 2021). Unless emissions are vastly reduced in
the coming decades, global warming will continue and exceed 1.5°-2°C compared to pre-
industrial levels by the end of the 21st century (IPCC, 2021).

Climate change has already had adverse effects on biodiversity and ecosystem functioning and
these effects are likely to worsen as warming proceeds in future (IPBES, 2019; IPCC, 2022).
Climate change impacts on ecological processes scales from genes to entire ecosystems, can
affect organisms, populations or entire communities and vary between physiological,
morphological, phenological and distributional shifts (Bellard *et al.*, 2012; Scheffers *et al.*,
2016). Especially changes in species abundance and distribution due to climate change have
already been frequently observed (Bowler *et al.*, 2017; Lenoir *et al.*, 2020), with many species
shifting their range towards higher latitudes and elevations (Chen *et al.*, 2011). However, some
species also respond to climate change by idiosyncratic range shifts (Gibson-Reinemer & Rahel,
2015).

Species distribution models (SDMs) are a common way of assessing species-specific responses
to climate change (e.g. Engelhardt *et al.* (2020)), but also to assess climate change impacts on
biodiversity (Thuiller *et al.*, 2005). SDMs statistically infer a relationship between the observed
distribution of a species to the underlying climatic conditions (Elith & Leathwick, 2009) and can
then be used to project current distributions into the future (Elith *et al.*, 2010), assuming that the
species maintains its climatic niche (Wiens & Graham, 2005). By doing this for multiple species,
these projections can be combined to assess future changes in species richness (Hof *et al.*, 2018;
Newbold, 2018; Thuiller *et al.*, 2019).

In the past most climate change impact assessments on vertebrate biodiversity have focused on
endotherms (birds & mammals). Reptiles, though accounting for a third of global terrestrial
vertebrate diversity, have been largely ignored (Pacifici *et al.*, 2015). Previous studies that
assessed climate change impacts on reptiles species, have either only used a subset of species
(Warren *et al.*, 2018; Newbold, 2018) or have not been of global extent (Araújo *et al.*, 2006).
Until recently, this was largely due to the unavailability of global reptile distribution data, but
this has changed with the release of the global distribution database by the Global Assessment of
Reptile Distributions (GARD) initiative (Roll *et al.*, 2017) and the release of the full set of IUCN
reptile range maps (IUCN, 2022).

Global biodiversity assessments further often either consider overall effects on a single taxon
(Baisero *et al.*, 2020; Voskamp *et al.*, 2021) or compare multiple taxa (Hof *et al.*, 2018;
Newbold, 2018; Warren *et al.*, 2018; Thuiller *et al.*, 2019), but only very rarely compare
different taxonomic groups within one taxon (but see, for example Hof *et al.* (2011)). This may

be problematic as pooling all species may obscure the evolutionary and biogeographic history of
major lineages, and from a global conservation perspective ignores the fact that hotspots of total
reptile richness hardly overlap with those of lizard or turtle richness (Roll *et al.*, 2017).

Here, we tried to fill the gaps outlined above, by providing a detailed account of projected
climate change impacts on global reptile distributions and diversity, looking at species-specific
changes as well as broad-scale geographic trends across and within different taxonomic groups.
We assessed changes in reptile richness globally, within each zoogeographic realm and across
their respective climate space. For each species, we further quantified the change in extent of
occurrence (EOO), range overlap and range distribution and again assessed differences across
zoogeographic realms and taxonomic groups. Given that we cannot model range restricted
species, we also performed a more general assessment of species-specific changes in climatic
space across both modeled and non-modeled species.

Methods

Species data

We obtained global range maps of 10,064 reptile species from the Global Assessment of Reptile
Distributions (GARD, https://doi.org/10.5061/dryad.83s7k). Roll *et al.* (2017) provide a detailed
description of the methodology used for deriving the range maps. The range maps cover lizards,
snakes, turtles, worm lizards, crocodiles and the tuatara, but in this paper, similar to (Roll *et al.*,
2017), we only contrast snakes, turtles and paraphyletic lizards (the latter of which we
subsequently refer to as lizards for simplicity).

Range maps were gridded to a 0.5° x 0.5° grid in WGS84, considering any cell that intersected
with the range polygon, and pseudo-absence data for each species were generated by randomly
selecting absences using a distance-weighted approach (see Hof *et al.*, 2018). The number of
absences was either equal to the number of presences or 1000 absences for species with less than
1000 presences. For each species we derived 10 sets of pseudo-absences, which were modeled
separately.

Climate data

Global bias-corrected daily climate (minimum temperature, maximum temperature and
precipitation) data at a spatial resolution of 0.5° (WGS84) was obtained from the meteorological
forcing dataset 'EartH2Observe, WFDEI and ERA-Interim data Merged and Bias-corrected for
ISIMIP' (EWEMBI; Lange, 2016) for current conditions (1980 - 2009) and from the Inter-
Sectoral Impact Model Intercomparison Project phase 2b (ISIMP2b; Frieler *et al.*, 2017) for
future simulations (2036 - 2065 & 2066 - 2095). Future climate simulations were available from
four global circulation models (GCMs; GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR and
MIROC5) and for three representative concentration pathways (RCPs; RCP2.6, RCP6.0 and
RCP8.5). Monthly means of each climatic variable over the respective 30-year time periods,
centered around 1995, 2050 and 2080, and for each future scenario (GCM & RCP) were used to
calculate 19 bioclimatic variables using the 'dismo' package (Hijmans *et al.*, 2021) in R (R Core
Team, 2021). We used the same model selection approach as described in Hof *et al.* (2018) and
fitted our models using the best-performing combination of four explanatory variables, which
was temperature seasonality, maximum temperature of the warmest month, annual precipitation
and precipitation seasonality.

Species Distribution Models (SDMs)

116
Projections based on SDMs vary considerably among the model algorithm considered, for this
we fitted two modeling algorithms with a good performance and discrimination capacity
(Meynard & Quinn, 2007; Elith *et al.*, 2010), an additive (Generalized Additive Model (GAM))
and a regression tree based model (Generalized Boosted Regression Models (GBM)).

GAMs were fitted with a Bernoulli response, a logit link and thin-plate regression splines using
the 'mgcv' package (Wood, 2003, 2011) in R (R Core Team, 2021). GBMs were fitted with the
'gbm' package (Greenwell *et al.*, 2020) in R (R Core Team, 2021) and the optimal parameter
settings for learning rate (0.01 and 0.001), tree complexity (1, 2 and 3) and number of trees
(1000-10000) for each species were identified by cross-validation (Bagchi *et al.*, 2013).

Spatial autocorrelation in species distributions can bias parameter estimates and error
probabilities (Kühn, 2007). Two different methods were used to account for spatial
autocorrelation in the SDMs. Species with equal or more than 50 presences were modeled using
an ecoregion-blocking approach. Here the world was divided into 10 blocks, based on a
representative subset of the climatic space of each of the world's ecoregions (Olson *et al.*, 2001),
and 10 models were built leaving out one block at a time, using the left out block for model
evaluation (Bagchi *et al.*, 2013). For range-restricted species (10 - 49 presences), we split the
data into 10 datasets by repeatedly randomly selecting 70% of the data, using the left-out 30%
for model evaluation. Species occurring in less than 10 grid cells were not modeled (N = 3602,
Table 1).

The performance of the fitted SDMs was evaluated by calculating the overall AUC for each
species (the average AUC across the 10 blocks and the 10 sets of pseudo-absences). Models with
an overall AUC smaller than 0.7 were dropped (N = 166), which left us with SDMs for 6296
reptile species (see Fig. S1.1 in Appendix S1 in Supporting Information), which represents 62.6
% of the total number of available species by GARD (Table 1).

The same modeling approach has been used previously to assess climate change impacts on
birds, amphibians and reptiles, see Hof *et al.* (2018) and Biber *et al.* (2020). The former provides
a more detailed explanation of the modeling methodology, while the latter gives a thorough
account of the caveats and uncertainties associated with species distribution models.

Future projections

Future species distributions were derived by predicting the models using the future bioclimatic
variables of the two future time periods (2050, 2080) and the respective future scenario (GCM &
RCP). Future projections of each species were limited to their original and the extent of their
neighboring ecoregions, to avoid predictions to areas with analogue climatic conditions. Future
projections were further limited by applying a species-specific dispersal buffer. For most species
considered here species-specific dispersal distances are still unknown (Nathan *et al.*, 2012),
hence we used species-specific dispersal buffers, which were based on the diameter (d) of the
largest range polygon of a species. We used three species-specific dispersal scenarios (d/4, d/8,
d/16, see Fig. S1.2 in Appendix S1) and provide a detailed comparison of these in the Supporting
Information (see Appendix S4), but below provide the results under the medium dispersal
scenario d/8, with a mean dispersal distance of 2.4 km per year.

Impact analysis

The current and future probabilities of occurrence of the individual SDMs were thresholded into
binary presence-absence data using species-specific thresholds according to the true skill statistic
(MaxTSS; Allouche *et al.,* 2006). Thresholded species occurrences were then used to calculate
current and future species richness, as well as richness increase, decrease, change and relative
change (%). Richness increase and decrease was identified by using the presence information of
each individual species and then summing up the number of species that newly occur in a given
grid cell (species increase) or species that disappear from the respective grid cell (species
decreae).

As stacked thresholded data frequently overestimates species richness, we also present the results
using the stacked raw probabilities of occurrence, as suggested by Calabrese *et al.* (2014), of the
individual SDMs without thresholding where applicable, in the Supporting Information (see
Appendix S3).

We calculated the overall mean projected species richness globally and for each zoogeographic
realm, as defined by Holt *et al.* (2013), for each time period and tested for significant changes in
species richness over time using a paired t-test with Holm correction. To assess how species
richness and changes in species richness are related to the overall change in climatic conditions,
we assessed both across univariate temperature and precipitation as well as the interaction of
temperature and precipitation conditions. To assess potential future climate effects on individual
species, we quantified the proportion of change in EOO and the proportion of range overlap for
species. And to assess the geographic change between each current and future species range, we
calculated the range centroid for current and future conditions of each species and then identified
the distance and direction of change between current and future range centroids.

Given that 37.4% of reptile species for which data was available could not be modeled (largely
due to their restricted range extent, Table 1), we performed an additional analysis considering all
10,064 species for which data were available. We used the same 4 bioclimatic variables used for
the SDMs to transform the multidimensional climate data in a two-dimensional climate space
using the first two axes of a principal component analysis (PCA). PCAs were performed for both
current and future conditions, considering the same GCMs, RCPs and time periods as before
(Fig. S1.3). For each scenario combination, we then calculated the Euclidean distance between
the two PCA-axes of current and future conditions, to get a measure of climatic change (Fig.
S1.4). We then extracted the climatic distance for the gridded locations of each species and
compared the climatic distance of modeled and non-modeled species using a non-paired t-test
with Holm correction.

Where no specific groups (lizards, snakes, turtles) are mentioned, we present the results for all
reptile species together. Results are presented as the ensemble mean, across the four GCMs and
two model algorithms considered, for the year 2080 under a medium representative
concentration pathway (RCP6.0). A sensitivity analysis with regards to the variation across years
and RCPs is shown in the Supporting Information (see Appendix S4).

Results

Projected reptile richness for current conditions varied between 0 (high latitudes) and 251 in the
tropics, with particular hotspots in Brazil, Cameroon and Indonesia (Fig. 1a). Overall, reptile

richness was dominated by lizard species (N = 3695), followed by snakes (N = 2305), while
turtle species only contributed marginally to the total number of modeled species (N = 296,
Table 1). Looking at the spatial configuration of species richness across the three groups, snakes
had the highest mean species richness (μ_{mean} = 30.4 ± 0.15), followed by lizard (μ_{mean} = 24.3 ±
0.09 SE) and turtle richness (μ_{mean} = 3.71 ± 0.02 SE), while the mean total richness was 58.4 ±
0.24 (SE; Fig. 1b, Fig. S2.5, Appendix S2). A large number of reptile species was projected to
disappear and at the same time a large number of new species were projected to appear in Brazil
and Australia, while other regions showed either a strong species decrease or increase (Fig. 1
c,d). The strongest future species decreases were projected east of the Caspian Sea and in South
Africa (Fig. 1 c), while strong future increases were predicted in the south-west of China and in
the eastern United States (Fig. 1 d). Overall, species decrease was stronger than species increase,
which resulted in a stronger negative net change in species richness from 1995 to 2080 (Fig.
1c,d,e). The lowest negative net change in species richness was observed in Brazil, Australia and
South Africa, while the highest positive net change was projected for south-west China and the
western United States (Fig. 1 e). Relative change (%) was projected to be negative in particular
for most of the southern hemisphere, while the high northern latitudes showed a strong positive
relative change (Fig. 1 f).

Spatial patterns in species richness changes varied strongly across the three taxa, with lizards
seeing both strong increases and decreases in Australia, snakes showing a strong decrease in
South America and turtles seeing a strong increase in eastern North America (Fig. S2.6). All
three taxa showed a positive net change in species richness in northern latitudes (Fig. S2.7),
while lizards decreased in parts of Australia (Fig. S2.7 a,b), snakes in large parts of South
America (Fig. S2.7 c, d) and turtles showed a strong negative net change in parts of South
America and southern Africa (Fig. S2.7 f).

Globally reptile richness was projected to decline significantly (p < 0.01) from 1995 to 2080,
with a decline in mean reptile richness from 58.4 ± 0.24 (SE) in 1995 to 53.39 ± 0.19 (SE) in
2080 (Fig. 2 a). 8 out of 11 zoogeographic realms showed a significant decline in reptile richness
by 2080 (Fig. 2 b, c, d, f, g, h, j, k), while the Nearctic and Palearctic realm showed a significant
increase (Fig. 2 e, i) and the Sino-Japanese realm showed no significant change (Fig. 2 l).

The different taxonomic groups (lizards, snakes and turtles) showed changes in global species
richness that were similar to all reptiles, while there were slight differences across the individual
realms. Lizards only showed a significant increase in richness in the Palearctic realm and no
significant change in richness in the Sino-Japanese realm, while in all other realms they showed
a significant decrease (Fig. S2.8). Snake and turtle richness increased significantly in the
Nearctic and Palearctic realm. Snake richness decreased significantly in all other realms apart
from the Sino-Japanese one, while turtle richness significantly decreased in all other realms apart
from the Saharo-Arabian and the Sino-Japanese one (Fig. S2.9, S2.10).

Reptile richness strongly differed across conditions of varying combinations of temperature and
precipitation (Fig. 3 a,b,c). For 1995 reptile richness was projected to be highest in areas with a
temperature around 28.5°C, a precipitation of about 5500 mm and when considering temperature
and precipitation together in warm, but moist regions (21°C & 3000 mm, Fig. 3 c). The climatic
conditions with the highest richness shifted to even more extreme (warmer & wetter) novel
climate conditions by 2080 (Fig. 3 a,b,d).

Looking at the species richness change across the 2-dimensional climate space, net change was
positive at the upper precipitation limits across all temperatures and the very hot and very dry
conditions and negative throughout the entire precipitation range especially for the higher
temperatures. Overall the negative change was much stronger and more pronounced than the
positive net change (Fig. 3 e). The highest positive and negative relative change values were
clustered, both occurred at the upper precipitation limits at low and medium temperatures (Fig. 3
f). A considerable proportion of the climate space (29.5 %) was shifting towards novel climatic
conditions, for which no change in species richness could be estimated, while only few discrete
climatic conditions as well as very cold & very dry conditions (4.75 %) got lost (Fig. 3 e,f).

The EOO of most species (n = 6021) showed a considerable decrease (μmean = -27.7 ± 0.16 SE,
Fig. 4 a,c,e). Lizard species showed the strongest decline (μmean = -31.8 ± 0.22 SE) in EOO
(Fig. 4 a), followed by snakes (μmean = -22.6 ± 0.25 SE), while almost equal numbers of turtle
species showed a decline (n = 274) and an increase (n = 205) with decreases being much more
pronounced than increases (μmean = -17.5 ± 0.72 SE, Fig. 4 e). Almost half of the modeled
reptile species (n = 3029) showed a strong change in range position, demonstrated by a relatively
low range overlap (<= 60 %), which was consistent across all three groups (Fig. 4 b,d,f).

Most of the range centroids (58 %) of all reptile species fell within the Neotropical (n = 1133),
Afrotropical (n = 1039), Oriental (n = 785) and Australian realm (n = 698). Turtle species had 50
% of their range centroids in the Nearctic (n = 58), Oriental (n = 53) and Afrotropical realm (n =
38), while lizards and snakes reflected the overall, total reptile, patterns (Fig. 5 d). Range
centroids were highly clustered within the different realms, which reflects the overall richness
hotspots, and hardly any centroids were found in the high northern latitudes (Fig.5 d). By 2080
species centroids were projected to shift by a mean distance of 111 km ± 0.9 SE mostly towards
a southerly direction. Lizards showed a shift towards all directions, with a slight trend in the
number of species towards the South (Fig. 5 a), while snakes and turtles show a more
pronounced shift of species towards the North (Fig. 5 b, c). Turtle ranges shifted by the largest
distances, followed by snakes (Fig. 5 a-c). The northern realms (Nearctic, Saharo-Arabian,
Paleartic und Sino-Japanese) showed a dominant shift towards the North, while the southern
realms (Neotropical, Afrotropical and Australian) showed a dominant shift towards the South.
The Panamanian, Madagascan and Oriental realms also showed a northerly shift, while the
Oceanian realm showed a bi-directional shift to the Northwest and Southeast (Fig. 5 d, Fig.
S2.11). Large realms had a larger proportion of species that shifted their range for a larger
distance (Fig. S2.11).

37.4 % of reptile species, for which data would have been available, could not be modeled using
SDMs, either due to a small sample size or a low model performance (Table 1). We found that
the species that could not be modeled showed a significant higher mean climatic distance
between current and future conditions compared to the modeled species and thus occurred in
areas that experience a stronger change in climatic conditions. This pattern was consistent across
all three taxa (Fig. 6).

Looking at the sum of occurrence probabilities, we found similar spatial patternsand a similar
magnitude in change (Fig. S3.13 - S3.15). Projected richness values and their future changes
were slightly larger under a larger dispersal ability (d/4), but overall all results were consistent
across the three dispersal scenarios considered (Fig. S4.16 - S4.21). Climate change impacts on
future species richness increased over time, with stronger effects seen for 2080 than 2050, and

the strongest impacts being observed under a high emission scenario (RCP8.5) compared to the
two lower scenarios (Fig. S5.22 - S5.32).

Discussion

Reptile richness was projected to decrease significantly across most parts of the world in the
future (Fig. 1 & 2). This effect was visible across lizards, snakes and turtles alike, although
regional and species-specific responses differed across the three groups (Fig. S2.7 - S2.10).

Reptile richness is projected to decrease in Brazil, Australia and South Africa, and to increase in
south-western China and the western United Sates (Fig. 1 e). These areas overlap strongly with
the biotic convergence zones identified for the conservation of Lepidosaurians (Diele-Viegas *et*
al., 2020). Brazil in particular is not only characterized by a high reptile richness (Roll *et al.*
(2017), Fig. 1 a), but also hosts a large proportion of threatened reptile species (Böhm *et al.,*
2013). Furthermore, most protected areas within Brazil are only of low conservation status
(IUCN Categories V - VI) and the stricter ones underrepresent a large proportion of the Brazilian
biophysical environment (Baldi *et al.,* 2019). While the areas that were projected to show an
increase in reptile richness with climate warming, i.e. south-west China and the western United
States, partially overlap with the areas that have the largest proportion of reptile species, they are
also the ones most threatened by habitat loss from agriculture and logging or harvesting (Böhm
et al., 2013).

Species richness changes varied strongly across different regions and different taxa, with lizards
seeing a strong species increase and decrease in Australia, snakes seeing a strong decrease in
species richness in South America and turtles seeing a strong increase in the eastern part of the
United States (Fig. S2.7). All three taxa saw a positive net change in the northern latitudes (Fig.
S2.7), while lizards saw a decrease in parts of Australia (Fig. S2.7 a,b), snakes in large parts of
South America (Fig S.2.7 c,d) and turtles saw a strong relative net change in parts of Australia
and northern Africa (Fig S2.7 f). These differences across groups are also reflected in their
original richness patterns. Species richness of amphibians, birds and mammals together is a good
spatial surrogate for species richness of all reptiles combined and of snakes, but not for lizard or
turtle richness (Roll *et al.,* 2017). Thus, it is not surprising that the areas with the highest decline
in overall reptile richness (see Fig. 1) strongly overlap with the areas of highest projected
changes in vertebrate species richness (amphibians, birds and mammals) found by Hof *et al.*
(2018), although global reptile richness is largely constrained by temperature, while global
richness of all other vertebrate groups is mostly constrained by the availability of energy and
water (Qian, 2010). Historical shifts in geographical ranges and climatic niches further showed
that niche shifts in endotherms are significantly faster than in ectotherms (Rolland *et al.,* 2018).
Newbold (2018) further found that Brazil is strongly affected by vertebrate diversity loss due to
climate change, and together with Australia is also likely to be strongly affected by future land-
use changes, especially under a high-emission scenario (RCP8.5).

Globally reptile richness was projected to decline significantly, from about 58 to 53 (9.4 %)
species on average per grid cell from 1995 to 2080 (Fig. 2 a). This estimate is slightly lower than
the reptile richness decline predicted by Newbold (2018). One explanation for this difference
could be that Newbold (2018) only used a subset of species but they also applied a much smaller
dispersal buffer, which might indicate that our projections provide a rather optimistic scenario.
Newbold (2018) further found that reptiles together with amphibians are disproportionately

sensitive to future human land-use. Given the synergistic effect of future climate and land-use
change on biodiversity (Brook et al., 2008) as well as species population s (Williams et al.,
2022), land-use change might further exacerbate climate change impacts on global reptile
distribution and diversity.

Changes in reptile richness differed among zoogeographic realms, but species richness declined
significantly across most realms over both time periods (Fig. 2 b,c,d,f,g,h,j,k). Lizards, snakes
and turtles all showed similar declines in global species richness and across most realms, but
slightly differed across individual realms. This is in line with a previous study covering various
realms from tropical to temperate regions which found that 60 % of assessed Lepidosaurian
species (n = 1114) were vulnerable to changes in climate (Diele-Viegas et al., 2020). We found
that only the Oceanian and Madagascan realms did not show a consistent decline in species
richness, which partially corresponds with the results of Diele-Viegas et al. (2020), who found
that the Madagascan and Oceanian realm were the ones where Lepidosaurians were the least
vulnerable to climatic change. However, given that the Madagascan realm boasts over 90% of
endemic reptile species and genera (Glaw & Vences, 2007) and that both realms are completely
composed of island territories which are usually considered highly vulnerable to climate change
and might also be affected by future sea level rise and erosion (Diele-Viegas et al., 2020), our
estimates might in fact underestimate potential climate change impacts in these realms.

Reptile richness differed strongly across temperature and precipitation, with the highest richness
being observed in warm, but moist conditions. Under future climate, the climatic conditions with
high species richness were projected to shift to even more extreme (warmer & wetter) conditions
(Fig. 3). Reptiles cannot regulate their body temperature internally, so are strongly dependent on
solar energy captured by the environment to regulate their body temperature (Huey, 1982). This
might lead to overheating when temperatures reach beyond a species' critical limit, which makes
them particularly susceptible to climatic changes (Sinervo et al., 2018). However, this might be
compensated by other biological processes that help species to buffer climate change effects, i.e.
genomic and phenotypic plasticity (Rodrı́guez et al., 2017) as well as behavioral and
physiological adaptation Sunday et al. (2014). Overall, the persistence of reptile species would
be much more affected by climate cooling than warming, but increasing droughts, which will be
a consequence of continued warming, have been suggested to pose a significant future threat to
European reptiles (Araújo et al., 2006). Climate warming will likely have an additional impact
on reptiles that have temperature-dependent sex determination. Altered sex ratios will not only
result in a higher extinction risk of local populations, but, together with a reduction in nesting
sites due to habitat destruction and fragmentation, will also affect the dispersal and potential
range expansion of a species. Therefore it could also have an impact on population demography
and size unless temperature shifts in sex determination or female nest-site choice evolves in pace
with rising temperatures (Boyle et al., 2016; Gibbons et al., 2000).

The EOO of most species considerably decreased, with lizard species showing the strongest
decline (Fig. 4 a). Most reptiles further showed a strong decline in range overlap, which was
consistent across all three groups (Fig. 4). This is in line with results by Warren et al. (2018),
who found that range losses of more than 50% occur in 35% of considered reptile species,
although this study included only a fraction of all reptile species and dispersal was not
considered.

Terrestrial reptiles have narrow niche requirements and small ranges, compared to other
vertebrate groups, which makes them more susceptible to threats such as habitat loss or invasive
species (Böhm *et al.*, 2013). 21 % of reptilian species are currently threatened with extinction
(IUCN, 2022), while adequate baseline data to inform conservation actions is often missing (Roll
et al., 2017). In addition, cascading effects generated by disease, invasive species, habitat loss
and climate change might lead to declines of sympatric species and a faster deterioration of
ecosystem structure than anticipated by climate change alone (Zipkin *et al.*, 2020).

The majority of reptile species showed a shift towards the South, which was largely driven by
range shifts in lizards. Turtle ranges overall shifted farthest, followed by snakes (Fig. 5). This is
likely due to the fact that lizards have the smallest range sizes across the three groups (Roll *et al.*,
2017), which in our case also resulted in smaller dispersal distances (Fig. S1.3).

Range restricted (non-modeled) species are projected to experience significantly higher shift in
climatic distance than modeled species (Fig. 6), indicating that range-restricted species are
disproportionately affected by climate change. This highlights once more that sample size
restrictions of SDMs downplay climate change effects on narrow-ranging and threatened species
(Platts *et al.*, 2014). Hof *et al.* (2018) also found significant impacts of climate and land-use
changes on range-restricted vertebrate species, excluding reptiles. However, similar to the latter
study we only look at climate anomalies (Euclidean distance between current and future climatic
conditions) as a metric of climate change, while different metrics have been found to show
contrasting climate change patterns on a global scale (Garcia *et al.*, 2014). In addition to climate
change effects, habitat modification has been found to have a more negative effect on small-
ranging reptile species, as well as species with a small clutch size (Doherty *et al.*, 2020). While
habitat modification would be an additional factor worthwhile to consider in future impact
assessments, this would go beyond the scope of this study.

Our results were strongly dependent on the dispersal, time period and emission scenario (RCP)
considered, but the overall patterns and richness changes were consistent throughout (see
Supporting Information). Previous studies have highlighted differences among dispersal, time
period and RCP across various vertebrate taxa (Thuiller *et al.*, 2019), but also specifically for
reptiles (Araújo *et al.*, 2006; Newbold, 2018; Warren *et al.*, 2018). Reptile-specific studies have
either considered no dispersal at all (Araújo *et al.*, 2006; Warren *et al.*, 2018) or a dispersal rate
of 0.5 km per year (Newbold, 2018). We use a slightly higher biologically informed dispersal
scenario, with an average of 2-5 km per year, which might be overly optimistic. The considered
dispersal distances are species-specific and thus strongly depend on the range size of each
species. Given that our model results as well as the underlying climate scenarios are based on a
0.5° grid size, small differences in the dispersal distances considered here did not have a strong
impact on our results (see Appendix S4).

The projected changes in species distributions help to investigate potential changes in global
reptile richness patterns and to highlight hotspots of climate change impacts. They also allow to
compare climate change vulnerability across taxonomic groups. However, these projections are
far from reality and have to be interpreted with caution. Future studies should try to consider
additional factors, such as biotic interactions (Davis *et al.*, 1998) and the reshuffling of species
communities (Voskamp *et al.*, in prep), which might lead to a change in competitive balance
(Ockendon *et al.*, 2014), altered predator-prey relationships (Harley, 2011) or changes in

functional diversity (Stewart *et al.*, 2022) and thus the provision of ecosystem functions and
services (Pecl *et al.*, 2017).

Conclusion

In addition to climate and land-use changes, reptile species are threatened by habitat loss and
degradation, invasive species, environmental pollution, disease and biological resource use (e.g.,
hunting and timber harvesting) (Gibbons *et al.*, 2000, Chapple2021).

Existing protected areas, sites of biodiversity significance and global conservation schemes do
not effectively protect reptiles, particularly lizards and turtles (Roll *et al.*, 2017) and 61% of the
world's skinks do not overlap at all with protected areas (Chapple *et al.*, 2021). The climatic
space of protected areas is projected to shift considerably with future climate warming (Elsen *et*
al., 2020), though even if existing nature reserves would adequately represent reptiles now, they
might be inadequate to preserve biodiversity in the future.

Our study shows that reptiles are likely to be considerably impacted by future climate change,
globally but also within most zoogeographic realms. These impacts are projected to have a
considerable effect on species' extent of occurrence and range position. Thus, to prevent large
scale declines in reptile species it is detrimental to lower CO_2 emissions in order to stop on-going
climate change but also to maintain adequate habitats of sufficient size and quality, especially of
grassland and savanna habitats (Roll *et al.*, 2017). Furthermore it is necessary to establish new
protected areas that help to prevent the extinction of particularly vulnerable species, i.e. by
establishing high-elevation climate refugia within current species ranges (Sinervo *et al.*, 2018).

References

Allouche, O., Tsoar, A. & Kadmon, R. (2006) Assessing the accuracy of species distribution
models: Prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, **43**,
1223–1232.

Araújo, M.B., Thuiller, W. & Pearson, R.G. (2006) Climate warming and the decline of
amphibians and reptiles in europe. *Journal of Biogeography*, **33**, 1712–1728.

Bagchi, R., Crosby, M., Huntley, B., Hole, D.G., Butchart, S.H., Collingham, Y., Kalra, M.,
Rajkumar, J., Rahmani, A., Pandey, M. & others (2013) Evaluating the effectiveness of
conservation site networks under climate change: Accounting for uncertainty. *Global*
Change Biology, **19**, 1236–1248.

Baisero, D., Visconti, P., Pacifici, M., Cimatti, M. & Rondinini, C. (2020) Projected global loss
of mammal habitat due to land-use and climate change. *One Earth*, **2**, 578–585.

Baldi, G., Schauman, S., Texeira, M., Marinaro, S., Martin, O.A., Gandini, P. & Jobbágy, E.G.
(2019) Nature representation in south american protected areas: Country contrasts and
conservation priorities. *PeerJ*, **7**, e7155.

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. (2012) Impacts of
climate change on the future of biodiversity. *Ecology Letters*, **15**, 365–377.

Biber, M.F., Voskamp, A., Niamir, A., Hickler, T. & Hof, C. (2020) A comparison of
macroecological and stacked species distribution models to predict future global terrestrial
vertebrate richness. *Journal of Biogeography*, **47**, 114–129.

Böhm, M., Collen, B., Baillie, J.E., Bowles, P., Chanson, J., Cox, N., Hammerson, G., Hoffmann,
M., Livingstone, S.R., Ram, M. & others (2013) The conservation status of the world's
reptiles. *Biological conservation*, **157**, 372–385.

Bowler, D.E., Hof, C., Haase, P., Kröncke, I., Schweiger, O., Adrian, R., Baert, L., Bauer, H.-G.,
Blick, T., Brooker, R.W. & others (2017) Cross-realm assessment of climate change impacts
on species' abundance trends. *Nature ecology & evolution*, **1**, 1–7.

Boyle, M., Schwanz, L., Hone, J. & Georges, A. (2016) Dispersal and climate warming
determine range shift in model reptile populations. *Ecological Modelling*, **328**, 34–43.

Brook, B.W., Sodhi, N.S., & Bradshaw, C.J.A. (2008) Synergies among extinction drivers
under global change. *Trends in Ecology & Evolution*, *23*, 453–460.

Calabrese, J.M., Certain, G., Kraan, C. & Dormann, C.F. (2014) Stacking species distribution
models and adjusting bias by linking them to macroecological models. *Global Ecology and
Biogeography*, **23**, 99–112.

Chapple, D.G., Roll, U., Böhm, M., Aguilar, R., Amey, A.P., Austin, C.C., Baling, M., Barley, A.J.,
Bates, M.F., Bauer, A.M. & others (2021) Conservation status of the world's skinks
(scincidae): Taxonomic and geographic patterns in extinction risk. *Biological Conservation*,
257, 109101.

Chen, I.-C., Hill, J.K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. (2011) Rapid range shifts of
species associated with high levels of climate warming. *Science*, **333**, 1024–1026.

Davis, A.J., Jenkinson, L.S., Lawton, J.H., Shorrocks, B. & Wood, S. (1998) Making mistakes
when predicting shifts in species range in response to global warming. *Nature*, **391**, 783–
786.

Diele-Viegas, L.M., Figueroa, R., Vilela, B. & Rocha, C. (2020) Are reptiles toast? A worldwide
evaluation of lepidosauria vulnerability to climate change. *Climatic Change*, **159**, 581–599.

Doherty, T.S., Balouch, S., Bell, K., Burns, T.J., Feldman, A., Fist, C., Garvey, T.F., Jessop, T.S.,
Meiri, S. & Driscoll, D.A. (2020) Reptile responses to anthropogenic habitat modification: A
global meta-analysis. *Global Ecology and Biogeography*, **29**, 1265–1279.

Elith, J., Kearney, M. & Phillips, S. (2010) The art of modelling range-shifting species.
Methods in Ecology and Evolution, **1**, 330–342.

Elith, J. & Leathwick, J.R. (2009) Species distribution models: Ecological explanation and
prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*, **40**,
677–697.

Elsen, P.R., Monahan, W.B., Dougherty, E.R. & Merenlender, A.M. (2020) Keeping pace with
climate change in global terrestrial protected areas. *Science advances*, **6**, eaay0814.

Engelhardt, E.K., Neuschulz, E.L. & Hof, C. (2020) Ignoring biotic interactions overestimates
climate change effects: The potential response of the spotted nutcracker to changes in
climate and resource plants. *Journal of Biogeography*, **47**, 143–154.

Frieler, K., Lange, S., Piontek, F., Reyer, C.P., Schewe, J., Warszawski, L., Zhao, F., Chini, L.,
Denvil, S., Emanuel, K. & others (2017) Assessing the impacts of 1.5°c global warming–
simulation protocol of the inter-sectoral impact model intercomparison project (ISIMIP2b).
Geoscientific Model Development, **10**, 4321–4345.

Garcia, R.A., Cabeza, M., Rahbek, C. & Araújo, M.B. (2014) Multiple dimensions of climate
change and their implications for biodiversity. *Science*, **344**, 1247579.

Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tuberville, T.D., Metts, B.S., Greene, J.L.,
Mills, T., Leiden, Y., Poppy, S. & others (2000) The global decline of reptiles, déjà vu
amphibians: Reptile species are declining on a global scale. Six significant threats to reptile
populations are habitat loss and degradation, introduced invasive species, environmental
pollution, disease, unsustainable use, and global climate change. *BioScience*, **50**, 653–666.

Gibson-Reinemer, D.K. & Rahel, F.J. (2015) Inconsistent range shifts within species
highlight idiosyncratic responses to climate warming. *PloS One*, **10**, e0132103.

Glaw, F. & Vences, M. (2007) *A field guide to the amphibians and reptiles of madagascar*,
Vences & Glaw.

Greenwell, B., Boehmke, B., Cunningham, J. & Developers, G. (2020) *Gbm: Generalized
boosted regression models*,.

Harley, C.D. (2011) Climate change, keystone predation, and biodiversity loss. *Science*, **334**,
1124–1127.

Hijmans, R.J., Phillips, S., Leathwick, J. & Elith, J. (2021) *Dismo: Species distribution
modeling*,.

Hof, C., Araújo, M.B., Jetz, W. & Rahbek, C. (2011) Additive threats from pathogens, climate
and land-use change for global amphibian diversity. *Nature*, **480**, 516–519.

Hof, C., Voskamp, A., Biber, M.F., Böhning-Gaese, K., Engelhardt, E.K., Niamir, A., Willis, S.G. &
Hickler, T. (2018) Bioenergy cropland expansion may offset positive effects of climate
change mitigation for global vertebrate diversity. *Proceedings of the National Academy of
Sciences*, **115**, 13294–13299.

Holt, B.G., Lessard, J.-P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D., Fabre, P.-H.,
Graham, C.H., Graves, G.R., Jønsson, K.A. & others (2013) An update of wallace's
zoogeographic regions of the world. *Science*, **339**, 74–78.

Huey, R.B. (1982) *Temperature, physiology, and the ecology of reptiles. Biology of the reptilia*,
Citeseer.

IPBES (2019) Summary for policymakers of the global assessment report on biodiversity
and ecosystem services of the intergovernmental science-policy platform on biodiversity
and ecosystem services. *IPBES secretariat, Bonn, Germany*.

IPCC (2021) Climate change 2021: The physical science basis. Working group i contribution
to the sixth assessment report of the intergovernmental panel on climate change.
Cambridge University Press.

IPCC (2022) Climate change 2022: Impacts, adaptation and vulnerability. Working group II
contribution to the sixth assessment report of the intergovernmental panel on climate
change. *Cambridge University Press*.

IUCN (2022) The IUCN red list of threatened species. Available at
https://www.iucnredlist.org/resources/spatial-data-download. *Cambridge University*
Press.

Kühn, I. (2007) Incorporating spatial autocorrelation may invert observed patterns.
Diversity and Distributions, **13**, 66–69.

Lange, S. (2016) EartH2Observe, WFDEI and ERA-interim data merged and bias-corrected
for ISIMIP (EWEMBI).

Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J. & Grenouillet, G.
(2020) Species better track climate warming in the oceans than on land. *Nature Ecology &*
Evolution, **4**, 1044–1059.

Meynard, C.N. & Quinn, J.F. (2007) Predicting species distributions: A critical comparison of
the most common statistical models using artificial species. *Journal of Biogeography*, **34**,
1455–1469.

Munoz, M.M., Stimola, M.A., Algar, A.C., Conover, A., Rodriguez, A.J., Landestoy, M.A., Bakken,
G.S. & Losos, J.B. (2014) Evolutionary stasis and lability in thermal physiology in a group of
tropical lizards. *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20132433.

Nathan, R., Klein, E., Robledo-Arnuncio, J.J. & Revilla, E. (2012) *Dispersal kernels*, Oxford
University Press Oxford, UK.

Newbold, T. (2018) Future effects of climate and land-use change on terrestrial vertebrate
community diversity under different scenarios. *Proceedings of the Royal Society B*, **285**,
20180792.

Ockendon, N., Baker, D.J., Carr, J.A., White, E.C., Almond, R.E., Amano, T., Bertram, E.,
Bradbury, R.B., Bradley, C., Butchart, S.H. & others (2014) Mechanisms underpinning
climatic impacts on natural populations: Altered species interactions are more important
than direct effects. *Global Change Biology*, **20**, 2221–2229.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V., Underwood,
E.C., D'amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C. & others (2001) Terrestrial
ecoregions of the world: A new map of life on EarthA new global map of terrestrial
ecoregions provides an innovative tool for conserving biodiversity. *BioScience*, **51**, 933–
938.

Pacifici, M., Foden, W.B., Visconti, P., Watson, J.E., Butchart, S.H., Kovacs, K.M., Scheffers, B.R.,
Hole, D.G., Martin, T.G., Akçakaya, H.R. & others (2015) Assessing species vulnerability to
climate change. *Nature Climate Change*, **5**, 215–224.

Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.-C., Clark, T.D.,
Colwell, R.K., Danielsen, F., Evengård, B. & others (2017) Biodiversity redistribution under
climate change: Impacts on ecosystems and human well-being. *Science*, **355**, eaai9214.

Platts, P.J., Garcia, R.A., Hof, C., Foden, W., Hansen, L.A., Rahbek, C. & Burgess, N.D. (2014)
Conservation implications of omitting narrow-ranging taxa from species distribution
models, now and in the future. *Diversity and distributions*, **20**, 1307–1320.

Qian, H. (2010) Environment–richness relationships for mammals, birds, reptiles, and
amphibians at global and regional scales. *Ecological Research*, **25**, 629–637.

R Core Team (2021) *R: A language and environment for statistical computing*, R Foundation
for Statistical Computing, Vienna, Austria.

Rodríguez, A., Rusciano, T., Hamilton, R., Holmes, L., Jordan, D. & Wollenberg Valero, K.C.
(2017) Genomic and phenotypic signatures of climate adaptation in an anolis lizard.
Ecology and evolution, **7**, 6390–6403.

Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A.M., Bernard, R., Böhm, M., Castro-
Herrera, F., Chirio, L., Collen, B. & others (2017) The global distribution of tetrapods reveals
a need for targeted reptile conservation. *Nature Ecology & Evolution*, **1**, 1677–1682.

Rolland, J., Silvestro, D., Schluter, D., Guisan, A., Broennimann, O. & Salamin, N. (2018) The
impact of endothermy on the climatic niche evolution and the distribution of vertebrate
diversity. *Nature Ecology & Evolution*, **2**, 459–464.

Scheffers, B.R., De Meester, L., Bridge, T.C., Hoffmann, A.A., Pandolfi, J.M., Corlett, R.T.,
Butchart, S.H., Pearce-Kelly, P., Kovacs, K.M., Dudgeon, D. & others (2016) The broad
footprint of climate change from genes to biomes to people. *Science*, **354**, aaf7671.

Sinervo, B., Miles, D.B., Wu, Y., MÉNDEZ-DE LA CRUZ, F.R., Kirchhof, S. & Qi, Y. (2018)
Climate change, thermal niches, extinction risk and maternal-effect rescue of toad-headed
lizards, phrynocephalus, in thermal extremes of the arabian peninsula to the qinghai—
tibetan plateau. *Integrative zoology*, **13**, 450–470.

Stewart, P.S., Voskamp, A., Santini, L., Biber, M.F., Hof, C., Willis, S.G. & Tobias, J.A. (2022)
Global impacts of climate change on avian functional diversity. *Ecology Letters*.

Sunday, J.M., Bates, A.E., Kearney, M.R., Colwell, R.K., Dulvy, N.K., Longino, J.T. & Huey, R.B.
(2014) Thermal-safety margins and the necessity of thermoregulatory behavior across
latitude and elevation. *Proceedings of the National Academy of Sciences*, **111**, 5610–5615.

Thuiller, W., Guéguen, M., Renaud, J., Karger, D.N. & Zimmermann, N.E. (2019) Uncertainty
in ensembles of global biodiversity scenarios. *Nature Communications*, **10**, 1–9.

Thuiller, W., Lavorel, S., Araújo, M.B., Sykes, M.T. & Prentice, I.C. (2005) Climate change
threats to plant diversity in europe. *Proceedings of the National Academy of Sciences*, **102**,
8245–8250.

Tingley, R., Meiri, S. & Chapple, D.G. (2016) Addressing knowledge gaps in reptile
conservation. *Biological Conservation*, **204**, 1–5.

Voskamp, A., Baker, D.J., Stephens, Philip A, Howard, C., Butchart, S.H. & Willis, S.G. (in prep)
Impacts of climate change on the world's terrestrial bird assemblages.

Voskamp, A., Butchart, S.H., Baker, D.J., Wilsey, C.B. & Willis, S.G. (2021) Site-based
conservation of terrestrial bird species in the caribbean and central and south america
under climate change. *Frontiers in Ecology and Evolution*, **9**, 234.

Warren, R., Price, J., Graham, E., Forstenhaeusler, N. & VanDerWal, J. (2018) The projected
effect on insects, vertebrates, and plants of limiting global warming to 1.5°c rather than 2°c.
Science, **360**, 791–795.

Wiens, J.J. & Graham, C.H. (2005) Niche conservatism: Integrating evolution, ecology, and
conservation biology. *Annual Review of Ecology, Evolution, and Systematics*, **36**, 519–539.

Williams, J.J., Freeman, R., Spooner, F. & Newbold, T. (2022) Vertebrate population trends
are influenced by interactions between land use, climatic position, habitat loss and climate
change. *Global change biology*, **28**, 797–815.

Wood, S.N. (2011) Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. *Journal of the Royal Statistical
Society (B)*, **73**, 3–36.

Wood, S.N. (2003) Thin-plate regression splines. *Journal of the Royal Statistical Society (B)*,
65, 95–114.

Zipkin, E.F., DiRenzo, G.V., Ray, J.M., Rossman, S. & Lips, K.R. (2020) Tropical snake diversity
collapses after widespread amphibian loss. *Science*, **367**, 814–816.

Data Accessibility Statement

GARD range maps are available from https://doi.org/10.5061/dryad.83s7k. EWEMBI and
ISIMIP2b climate data are available from https://esg.pik-potsdam.de/search/isimip/?
product=input. The code for the species distribution models can be found at
https://github.com/xxxx/xxxx, whereas the code and data for the performed data analysis and
the presented figures, can be found at **https://github.com/xxxx/xxxx**.

Tables

Table 1. Number of species that were excluded from the species distribution models due to their
restricted range or low model performance.

Taxonomic group	Lizard	Snake	Turtle	Total
No. of species with available data	6328	3414	322	10064
No. of range restricted species (N < 10, removed)	2536	1047	19	3602
No. of species with low model performance (AUC < 0.7, removed)	97	62	7	166
Total number of species modeled	**3695**	**2305**	**296**	**6296**
Percentage of available species modeled	58.4 %	67.5 %	91.9 %	62.6 %

 Figures

 Figure 1. a) Map of projected global terrestrial reptile species richness (1995), b) frequency of
 species richness by taxonomic group (lizard, snake, turtle and total) and c) increase, d) decrease,
 e) net change and f) relative change (%) in reptile species richness under the representative
 concentration pathways RCP6.0 and dispersal scenario d/8 for the time period 2080. All maps
 are in Mollweide equal-area projection (EPSG:54009). Grey areas are regions for which no
 projections are available. Note that the colour scales differ between the individual panels.

641
Figure 2. Terrestrial reptile species richness across the globe and for each zoogeographic realm
(Afrotropical, Australian, Madagascan, Nearctic, Neotropical, Oceanian, Oriental, Palearctic,
Panamanian, Saharo-Arabian & Sino-Japanese) over time (1995, 2080) under the representative
concentration pathways RCP6.0 and the dispersal scenario d/8. Statistical difference between
years was tested using a paired Student's t-test with Holm correction ($p < 0.05$ = *, $p < 0.01$ =
, $p < 0.001$ = ***, $p < 0.0001$ = **). Plots show mean (red point & label), median (black
horizontal line), 25th to 75th percentiles (box), entire range of data (violin & data points) and
density of values (width of violin). Figure 5 provides a map outlining the different
zoogeographic realms.

651
Figure 3. Univariate relationship of current (1995) and future (2080 RCP6.0) reptile species
richness with a) temperature, b) precipitation and the bivariate relationship of temperature and
precipitation with reptile species richness for c) 1995 and d) 2080 RCP6.0 and the respective e)
net richness change and f) relative richness change (%) under the dispersal scenario d/8. Lines
show the mean and ribbons the standard deviation in variance across space, global circulation
model and algorithm.

658
Figure 4. Frequency plots of the mean number of reptile species (a) lizard, b) snake, c) turtle)
and their potential future change (%) in extent of occurrence (EOO) and the mean number of
reptile species (d) lizard, e) snake, f) turtle) per potential range overlap class (0-20, 20-40, 40-60,
60-80, 80-100). Error margins/bars show standard deviation across the different global
circulation models and model algorithms used. Both shown for 2080 under the representative
concentration pathway RCP6.0 and the dispersal scenario d/8.

665
Figure 5. Cumulative direction and distance of potential range centroid changes per taxonomic
group (b) lizard, c) snake, d) turtle) and d) range centroids (points on map) and the number of
species and their directional shift in range centroid position per zoogeographic realm (inset polar
plots) for 2080 under the representative concentration pathway RCP6.0 and the dispersal
scenario d/8. Please note that some range centroids did not fall inside the zoogeographic realm
boundaries and thus were not associated with an realm (n = 477).

672
Figure 6. Mean climatic distance for modeled and non-modeled species, split by taxonomic
group (a) lizard, b) snake, c) turtle and d) total). Statistical difference between modeled and non-
modeled species was tested using a Student's t-test with Holm correction ($p < 0.05$ = *, $p < 0.01$
= **, $p < 0.001$ = ***, $p < 0.0001$ = ****). Plots show mean (red point & label), median (black
horizontal line), 25th to 75th percentiles (box), entire range of data (violin & data points) and
density of values (width of violin). Results are shown for 2080 under the representative
concentration pathway RCP6.0.