Potential future climate change effects on global reptile distributions and diversity

Matthias F. Biber[1,*], Alke Voskamp[2], Christian Hof[1]

[1]Technical University of Munich, Terrestrial Ecology Research Group, Department for Life Science Systems, School of Life Sciences, 85354 Freising, Germany

[2]Senckenberg Biodiversity and Climate Research Centre (SBiK-F), 60325 Frankfurt, Germany

Running head: Global reptiles under climate change

Abstract

Aim: Until recently, complete information on global reptile distributions has not been widely available. Here, we provide the first comprehensive climate impact assessment for reptiles on a global scale.

Location: Global, excluding Antarctica

Time period: 1995, 2050, 2080

Major taxa studied: Reptiles

Methods: We modelled the distribution of 6,296 reptile species and assessed potential global as well as realm-specific changes in species richness, the change in global species richness across climate space, and species-specific changes in range extent, overlap and position under future climate change. To assess the future climatic impact on 3,768 range-restricted species, which could not be modelled, we compared the future change in climatic conditions between both modelled and non-modelled species.

Results: Reptile richness was projected to decline significantly over time, globally but also for most zoogeographic realms, with the greatest decrease in Brazil, Australia and South Africa. Species richness was highest in warm and moist regions, with these regions being projected to shift further towards climate extremes in the future. Range extents were projected to decline considerably in the future, with a low overlap between current and future ranges. Shifts in range centroids differed among realms and taxa, with a dominating global poleward shift. Non-modelled species were significantly stronger affected by projected climatic changes than modelled species.

Main conclusions: With ongoing future climate change, reptile richness is likely to decrease significantly across most parts of the world. This effect as well as considerable impacts on species' range extent, overlap, and position were visible across lizards, snakes and turtles alike. Together with other anthropogenic impacts, such as habitat loss and harvesting of species, this is a cause for concern. Given the historical lack of global reptile distributions, this calls for a re-assessment of global reptile conservation efforts, with a specific focus on anticipated future climate change.

36   **KEYWORDS:** bioclim, biodiversity, environmental niche model, global warming, ISIMIP,
37   lizard, snake, species distribution model, species richness, turtle

Introduction

Emissions from anthropogenic activities have led to an increase in global surface temperature of
around 1°C in the last 100 years. This has already led to changes in the weather and the
occurrence of climate extremes in every region across the globe (IPCC, 2021). Unless emissions
are vastly reduced in the coming decades, global warming will continue and exceed 1.5°-2°C
compared to pre-industrial levels by the end of the 21st century (IPCC, 2021).

Climate change has already had adverse effects on biodiversity and ecosystem functioning and
these effects are likely to worsen as warming continues in the future (IPBES, 2019; IPCC, 2022).
Climate change impacts on ecological processes at scales ranging from genes to entire
ecosystems can affect organisms, populations, or entire communities, and vary between
physiological, morphological, phenological and distributional shifts (Bellard *et al.*, 2012;
Scheffers *et al.*, 2016). In particular, changes in species abundance and distribution due to
climate change have already been frequently observed (Bowler *et al.*, 2017; Lenoir *et al.*, 2020),
with many species shifting their range towards higher latitudes and elevations (Chen *et al.*,
2011). However, some species also respond to climate change through idiosyncratic range shifts
(Gibson-Reinemer & Rahel, 2015).

In the past most climate change impact assessments on vertebrate biodiversity have focused on
endotherms (birds & mammals). Reptiles, although they account for a third of global terrestrial
vertebrate diversity, have been largely ignored (Pacifici *et al.*, 2015), and previous assessments
of climate change impacts on reptile species have either used only a subset of species (Warren *et
al.*, 2018; Newbold, 2018) or have not been of global extent (Araújo *et al.*, 2006). Moreover,
global biodiversity assessments often either consider overall effects on a single taxon (Baisero *et
al.*, 2020; Voskamp *et al.*, 2021) or compare multiple taxa (Hof *et al.*, 2018; Newbold, 2018;
Warren *et al.*, 2018; Thuiller *et al.*, 2019), but only very rarely compare different taxonomic
groups within one taxon (but see, e.g. Hof *et al.* 2011).

Reptiles have in the past also often been neglected when assessing global conservation priorities
(Brooks *et al.*, 2006). They are most diverse in arid and semi-arid regions, which suggests that
their distributions are driven by ecological and evolutionary processes that differ from other
vertebrate taxa, and these regions have been previously unrecognised as conservation priorities
because other vertebrate taxa could be more efficiently protected elsewhere (Roll *et al.*, 2017).

While reptiles as a whole have been found to use similar habitats to mammals and birds (Cox et
al. 2022), treating all reptile species as uniform may be problematic because reptile taxonomic
groups (lizard, snakes and turtles) exhibit large differences in species richness hotspots (Roll *et
al.*, 2017) and habitat use, making them vulnerable to different anthropogenic impacts (Cox *et
al.*, 2022).

As reptiles are ectothermic, they are likely to be strongly influenced by climate warming, with
some species already experiencing body temperatures above their physiological optima. This
could indicate a higher vulnerability of these species to climate warming compared to species
from cold environments (Diele-Viegas *et al.*, 2018). Nonetheless, temperate species are also
likely to be vulnerable, assuming that their physiological adaptations for living in cold

environments may hinder their ability to cope with hotter climates (Monasterio *et al.*, 2013).
However, previous studies assessing climate change impacts on reptile species are biased
towards certain species and taxonomic groups as well as zoogeographic realms (Diele-Viegas *et*
al., 2020).

The aim of our research was to provide a detailed account of projected climate change impacts
on global reptile distributions and diversity, looking at species-specific changes as well as broad-
scale geographic trends across and within different taxonomic groups. We assessed changes in
reptile species richness globally, within each zoogeographic realm and across their respective
climate space. For each species, we further quantified the change in range extent, range overlap
and range distribution and again assessed differences across zoogeographic realms and
taxonomic groups. Given that we cannot model range-restricted species, we also performed a
more general assessment of species-specific changes in climate space across both modelled and
non-modelled species.

Methods

Species data

Until recently, global reptile distribution data was unavailable, but this has changed with the
release of the global distribution database by the Global Assessment of Reptile Distributions
(GARD) initiative (Roll *et al.*, 2017) and more recently the release of the full set of IUCN reptile
range maps (IUCN, 2022). We obtained global range maps of 10,064 reptile species from the
GARD initiative (Roll *et al.*, 2017). The range maps cover lizards, snakes, turtles, worm lizards,
crocodiles and the tuatara, but in this paper, in an approach similar to Roll *et al.* (2017), we only
contrast snakes, turtles and paraphyletic lizards (for simplicity, we subsequently refer to the latter
as lizards).

Range maps were gridded to a 0.5° x 0.5° grid in WGS84, to align with the available climate
data (see next section), considering any grid cell that intersected with a species' range polygon as
a presence. As range maps only provide information on a species' presence but not on a species'
absence, pseudo-absence data for each species were generated by randomly selecting grid cells
with no presence, using a distance-weighted approach, where grid cells closer to the range edge
were favoured over grid cells further away (see Hof *et al.,* 2018). The number of absences was
either equal to the number of presences for species with 1000 or more presences or 1000
absences for species with less than 1000 presences because a minimum of 1000 absences
considerably increases model performance (Barbet-Massin *et al.,* 2012). For each species we
derived 10 replicate sets of pseudo-absences, to account for the variability in model accuracy
because of the random sampling of pseudo-absence data. Barbet-Massin *et al.* (2012) found that
depending on the number of pseudo-absences and model algorithm chosen, 5 to 12 replicates
provide the best model quality. Given the relatively high number of pseudo-absences, 10
replicates should thus result in high quality for our models. We created a separate model for each
of these 10 sets, but the results were then averaged across the 10 sets.

Climate data

Global bias-corrected daily climate (minimum temperature, maximum temperature and
precipitation) data at a spatial resolution of 0.5° (WGS84) were obtained from the meteorological

forcing dataset 'EartH2Observe, WFDEI and ERA-Interim data Merged and Bias-corrected for
ISIMIP' (EWEMBI; Lange, 2016) for current conditions (1980 - 2009) and from the Inter-
Sectoral Impact Model Intercomparison Project phase 2b (ISIMP2b; Frieler *et al.*, 2017) for
future simulations (2036 - 2065 & 2066 - 2095). Future climate simulations were available from
four global circulation models (GCMs; GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR and
MIROC5) and for three representative concentration pathways (RCPs; RCP2.6, RCP6.0 and
RCP8.5) under the Coupled Model Intercomparison Project Phase 5 (CMIP5). Future climate
simulations strongly depend on the GCM used (Watterson, 2019). The four GCMs chosen by
ISIMIP2b capture a large range of plausible future climate projections from all the different
GCMs available (Frieler *et al.*, 2017), while the RCPs represent different emission scenarios
depending on the ongoing and future trajectories of global CO_2 emissions (Van Vuuren *et al.*,
2011).

Monthly means of each climate variable over the respective 30-year time periods, centred around
1995, 2050 and 2080, and for each future scenario (GCM & RCP) were used to calculate 19
bioclimatic variables (see Table S1.1 in Appendix S1 in Supporting Information) using the
biovars() function of the 'dismo' package (Hijmans *et al.*, 2021) in R (R Core Team, 2021).
Bioclimatic variables represent annual trends, seasonality and extreme or limiting environmental
factors and are thus more biologically meaningful variables than temperature or precipitation
alone.

Species Distribution Models (SDMs)

Species distribution models (SDMs) are a common way of assessing species-specific responses
to climate change (Guisan & Thuiller, 2005), but are also used to assess climate change impacts
on biodiversity (Thuiller *et al.*, 2005). SDMs statistically infer a relationship between the
observed distribution of a species and the underlying environmental conditions (Elith &
Leathwick, 2009), and can then be used to project current distributions into the future (Elith *et*
al., 2010), assuming that the species maintains its climatic niche (Wiens & Graham, 2005). By
doing this for multiple species, these projections can be combined to assess future changes in
species richness (e.g. Hof *et al.*, 2018).

We fitted species distribution models using the presence/pseudo-absence data of a species as
response variable and the derived bioclimatic variables for current (1995) conditions as
explanatory variables. The number and choice of explanatory variables used strongly influences
the outcome of the SDMs (Petitpierre *et al.*, 2017). We thus performed a rigorous variable
selection approach. After pre-selecting the 10 most commonly used variables from the literature
(see Porfirio *et al.*, 2014), all potential combinations of three and four bioclimatic variables with
a low Pearson correlation ($r \leq 0.7$) were used to model 10 % of all species (N= 987), which
were randomly selected, using a Generalised Additive Model (GAM) approach (see Hof *et al.*,
2018). The model performance of the different variable combinations was tested and models for
all species were fitted using the best-performing variable combination, which was temperature
seasonality, maximum temperature of the warmest month, annual precipitation and precipitation
seasonality (see Fig. S1.1).

Projections based on SDMs vary considerably among model algorithms (Thuiller *et al.*, 2019),
therefore we fitted two different modelling algorithms with good performance and discrimination

capacity (Meynard & Quinn, 2007; Elith *et al.*, 2010), an additive model (GAM) and a
regression tree based model (Generalised Boosted Regression Models (GBM)).

GAMs were fitted with a Bernoulli response, a logit link and thin-plate regression splines using
the 'mgcv' package (Wood, 2003, 2011) in R (R Core Team, 2021). GBMs were fitted with the
'gbm' package (Greenwell *et al.*, 2020) in R (R Core Team, 2021) and the optimal parameter
settings for learning rate (0.01 and 0.001), tree complexity (1, 2 and 3) and number of trees
(1000-10000) for each species were identified by cross-validation (Bagchi *et al.*, 2013).

Spatial autocorrelation in species distributions can bias parameter estimates and error
probabilities (Kühn, 2007). Two different methods were used to account for spatial
autocorrelation in the SDMs. Species with equal or more than 50 presences were modelled using
an ecoregion-blocking approach. Here the world was divided into 10 blocks based on a
representative subset of the climate space across each of the world's ecoregions (Olson *et al.*,
2001, Bagchi *et al.*, 2013). Subsequently, 10 models per species were built leaving out one block
at a time, using the left-out block for model evaluation (Bagchi *et al.*, 2013). For species with 10
to 49 presences, we split the data into 10 datasets by repeatedly randomly selecting 70% of the
data, then using the left-out 30% for model evaluation. Species occurring in less than 10 grid
cells (N = 3602, Table 1) were not modelled because the sampling size would be too low to
produce meaningful results (Hernandez et al. 2006). Together with the 10 different pseudo-
absence sets this resulted in 100 models for each model algorithm and species.

The performance of the fitted SDMs was evaluated by calculating the overall AUC for each
species (the average AUC across the 10 blocks and the 10 sets of pseudo-absences). Models with
an overall AUC smaller than 0.7 were dropped (N = 166), which left us with SDMs for 6296
reptile species (see Fig. S1.2), which represents 62.6 % of the total number of available species
from GARD (Table 1). In addition to assessing the model fit of the individual models, we also
compared the observed species richness with the projected current richness per grid cell (Fig
S1.3) in order to assess the performance of all our models when looking at changes in species
richness.

The same modelling approach has been used previously to assess climate change impacts on
amphibians, birds and mammals, see Hof *et al.* (2018) and Biber *et al.* (2020). The former
provides a more detailed explanation of the modelling methodology, while the latter gives a
thorough account of the caveats and uncertainties associated with species distribution models.

Future projections

Future species distributions were projected using the future bioclimatic variables for the two
future time periods (2050, 2080), each GCM (GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR
and MIROC5), each RCP (RCP2.6, RCP6.0 and RCP8.5) and both model algorithms (GAM &
GBM). Model results are presented as the ensemble mean across the four GCMs and two model
algorithms (GAM & GBM) considered.

Future projections of each species were limited to the extent of their original and the
neighbouring ecoregions to prevent predictions of areas with analogue climatic conditions.
Future projections were further limited by applying a species-specific dispersal buffer. For most
species considered in this paper, species-specific dispersal distances are still unknown (Nathan *et*
al., 2012), hence we used species-specific dispersal buffers which were based on the diameter (d)

of the largest range polygon of a species. We used three species-specific dispersal scenarios (d/4,
d/8, d/16, see Fig. S1.4) and provide a detailed comparison of these in the Supporting
Information (see Appendix S5). Here we provide results under a medium dispersal scenario
(d/8), which corresponds to a mean dispersal distance of 2.4 km per year.

Impact analysis

The current and future probabilities of occurrence of the individual SDMs were thresholded into
binary presence-absence data using species-specific thresholds according to the true skill statistic
(MaxTSS; Allouche *et al.,* 2006; Fig. S1.5 d). Thresholded species occurrences were then used
to calculate current and future species richness, as well as richness increase, decrease, change
and relative change (%). Richness increase and decrease were identified by using the presence
information of each individual species and then summing up the number of species that newly
occur in a given grid cell (species increase) or species that disappear from the respective grid cell
(species decrease).

Summing up the thresholded species occurrences frequently overestimates species richness
(Calabrese et al., 2014), thus we also present the results using the sum of the raw non-
thresholded probabilities of occurrence of each species in the Supporting Information (see
Appendix S4).

We calculated the projected species richness for each cell of a global grid of 0.5° x 0.5°
resolution globally as well as for each zoogeographic realm, as defined by Holt *et al.* (2013), for
each time period. We then tested for significant changes in species richness over time using a
paired t-test with Holm correction. To assess how species richness and richness changes are
related to the overall change in climatic conditions, we assessed both against univariate
temperature and precipitation as well as the interaction of temperature and precipitation
conditions. To assess potential future climate effects on individual species, we quantified the
percentage of change in range extent, the percentage of range overlap and the direction and
distance in range shift. The percentage of change in range extent was calculated based on the
projected current and future range extent of a species (Fig. S1.5 f). The percentage of range
overlap was calculated by extracting the total area of spatial range overlap between the projected
current and future range extent and dividing it by the projected current range extent (Fig. S1.5 g).
To assess the magnitude of the range shift for each species, we derived the range centroid for
both time periods and calculated the distance and direction of the projected range shift (Fig. S1.5
g).

Given that 37.4% of all reptile species for which data were available could not be modelled
(largely due to their restricted range extent, Table 1), we performed an additional analysis
considering all 10,064 species for which data were available. We used the same 4 bioclimatic
variables used for the SDMs to transform the multidimensional climate data to a two-
dimensional climate space using the first two axes of a principal component analysis (PCA).
PCAs were performed for both current and future conditions, taking into consideration the same
GCMs, RCPs, and time periods as previously. The explained variance of the first two PCA axes
was above 75 % under all scenarios (Fig. S1.6). For each scenario combination, we then
calculated the Euclidean distance between the two PCA axes of current and future conditions, to
obtain a measure of climatic change (Fig. S1.7). We then extracted the climatic distance for the

gridded locations of each species and compared the climatic distance of modelled and non-
modelled (range-restricted) species using a non-paired t-test with Holm correction.

Where no specific groups (lizards, snakes, turtles) are mentioned, we present the results for all
reptile species together. Nevertheless, Appendix S2 presents group-specific results. Results are
presented for the year 2080 under a medium representative concentration pathway (RCP6.0). A
sensitivity analysis with regards to the variation across years and RCPs is shown in the
Supporting Information (see Appendix S6).

Results

Spatial changes

Projected reptile richness (sum of thresholded SDM projections) based on a resolution of 0.5° x
0.5° for current conditions varied between 0 at high latitudes and 251 in the tropics, with
particular hotspots in Brazil, Cameroon and Indonesia (Fig. 1a). Overall, reptile richness was
dominated by lizard species (N = 3695), followed by snakes (N = 2305), while turtle species
only contributed marginally to the total number of modelled species (N = 296, Table 1, Fig. 1b).

For future conditions, a large number of reptile species were projected to disappear and at the
same time a large number of new species were projected to appear in Brazil and Australia, while
other regions showed either a strong species decrease or increase (Fig. 1 c, d). The greatest future
decreases of species richness were projected east of the Caspian Sea and in South Africa (Fig. 1
c), while strong future increases were projected in the south-west of China and in the eastern
United States (Fig. 1 d). Overall, the projected richness decrease was greater than the richness
increase, which resulted in a greater net loss in species richness from 1995 to 2080 (Fig. 1 c, d,
e). The lowest net loss in species richness was projected for Brazil, Australia and South Africa,
while the highest net gain was projected for south-west China and the western United States (Fig.
1 e). Relative change (%) was projected to be negative - in particular for most of the southern
hemisphere, while the high northern latitudes showed a strong positive relative change (Fig. 1 f).

Spatial patterns in species richness changes varied greatly across the three taxa, with lizards
seeing both strong increases and decreases in Australia, snakes showing a strong decrease in
South America and turtles seeing a strong increase in the eastern part of North America (Fig.
S2.9 & Fig. S2.11). All three taxa showed a net gain in species richness in northern latitudes,
while lizards showed the greatest net loss in parts of Australia, snakes in large parts of South
America and turtles in parts of South America and southern Africa (Fig. S2.10 & Fig. S2.12).

Global and zoogeographic realm changes

Summed up across all 0.5° x 0.5° grid cells globally, species richness was projected to decline
significantly ($p < 0.01$) from 1995 to 2080 globally, with a decline in mean reptile richness per
grid cell from 58.4 ± 0.24 (SE) in 1995 to 53.39 ± 0.19 in 2080 (Fig. 1b, Fig. 2 a). 8 out of 11
zoogeographic realms showed a significant decline in reptile richness by 2080 (Fig. 2 b, c, d, f, g,
h, j, k), while the Nearctic and Palearctic realms showed a significant increase (Fig. 2 e, i) and
the Sino-Japanese realm showed no significant change (Fig. 2 l).

Looking at the global averages of species richness for each 0.5° grid cell for the three reptile
groups separately, snakes had the highest mean species richness ($\mu_{mean} = 30.4 \pm 0.15$), followed

by lizard (μ_{mean} = 24.3 ± 0.09 SE) and turtle richness (μ_{mean} = 3.71 ± 0.02 SE; Fig. 1b, Fig. S2.8).
Globally, similar to the total reptile richness, the individual taxonomic groups (lizards, snakes
and turtles) all showed a significant decline in species richness, while there were slight
differences across the individual realms (Fig. S2.13 – S2.15). Lizards only showed a significant
increase in richness in the Palearctic realm and no significant change in richness in the Sino-
Japanese realm, while in all other realms they showed a significant decrease (Fig. S2.13). Snake
and turtle richness increased significantly in the Nearctic and Palearctic realms. Snake richness
decreased significantly in all other realms apart from the Sino-Japanese one (Fig. S2.14), while
turtle richness significantly decreased in all other realms apart from the Saharo-Arabian and the
Sino-Japanese one (Fig. S2.15).

Biophysical changes

Reptile richness varied greatly across conditions with varying combinations of temperature and
precipitation (Fig. 3 a, b, c). For 1995, reptile richness was projected to be highest in areas with a
temperature of about 28.5°C, a precipitation of about 5500 mm and when considering
temperature and precipitation together in warm and moist regions (21°C & 3000 mm, Fig. 3 c).
The climatic conditions with the highest richness shifted to even more extreme (warmer &
wetter) novel climate conditions by 2080 (Fig. 3 a, b, d).

Looking at the species richness change across the 2-dimensional climate space, net change was
positive at the upper precipitation limits across all temperatures and the very hot and very dry
conditions and negative throughout the entire precipitation range especially for the higher
temperatures. Overall, the negative change was much greater and more pronounced than the
positive net change (Fig. 3 e). The highest positive and negative relative change values were
clustered, both occurred at the upper precipitation limits at low and medium temperatures (Fig. 3
f). A considerable percentage of the climate space (29.5 %) was shifting towards novel climatic
conditions, for which no change in species richness could be estimated, while only few discrete
climatic conditions as well as very cold & very dry conditions (4.75 %) got lost (Fig. 3 e, f).

Species-specific range changes

The range extent of most species (N = 6021) showed a considerable decrease (μ_{mean} = -27.7 ±
0.16 SE, Fig. 4 a, c, e). Lizard species showed the greatest decline (μ_{mean} = -31.8 ± 0.22 SE) in
range extent (Fig. 4 a), followed by snakes (μ_{mean} = -22.6 ± 0.25 SE), while almost equal numbers
of turtle species showed a decline (N = 274) and an increase (n = 205) with decreases being
much more pronounced than increases (μ_{mean} = -17.5 ± 0.72 SE, Fig. 4 e). Almost half of the
modelled reptile species (N = 3029) showed a strong change in range position, demonstrated by a
relatively low range overlap (<= 60 %), which was consistent across all three groups (Fig. 4 b, d,
f).

Most of the range centroids (58 %) of all reptile species fell within the Neotropical (N = 1133),
Afrotropical (N = 1039), Oriental (N = 785) and Australian realm (N = 698). Turtle species had
50 % of their range centroids in the Nearctic (N = 58), Oriental (N = 53) and Afrotropical realm
(N = 38), while lizards and snakes reflected the overall, total reptile patterns (Fig. 5 d). Range
centroids were highly clustered within the different realms, which reflects the overall richness
hotspots, and hardly any centroids were found in the high northern latitudes (Fig. 5 d). By 2080
species centroids were projected to shift by a mean distance of 111 km ± 0.9 SE primarily
towards the South. Lizards showed a shift in all directions, with a slightly greater number of

species exhibiting a shift towards the South (Fig. 5 a), while snakes and turtles showed a more
pronounced shift of species towards the North (Fig. 5 b, c). Turtle ranges shifted by the largest
distances, followed by snakes (Fig. 5 a, b, c). The northern realms (Nearctic, Saharo-Arabian,
Palearctic und Sino-Japanese) showed a dominant shift towards the North, while the southern
realms (Neotropical, Afrotropical and Australian) showed a dominant shift towards the South.
This was also reflected in the taxon-specific range shifts for the Northern and Southern
hemisphere (Fig. S3.16 in Appendix S3). The Panamanian, Madagascan and Oriental realms also
showed a northerly shift, while the Oceanian realm showed a bi-directional shift to the
Northwest and Southeast (Fig. 5 d, Fig. S3.17). Large realms had a greater percentage of species
that shifted their range over a greater distance (Fig. 5, Fig. S3.17).

Non-modelled species

37.4 % of reptile species for which data would have been available could not be modelled using
SDMs, either due to a small sample size or a low model performance (Table 1). We found that
the species that could not be modelled showed a significantly greater ($p < 0.05$) mean climatic
distance between current and future conditions than did the modelled species and thus occurred
in areas that are projected to experience a greater change in climatic conditions. This pattern was
consistent across all three taxa

Sensitivity analysis

Looking at the sum of occurrence probabilities, we found similar spatial patterns and a similar
magnitude of change compared to the sum of thresholded occurrences (species richness) (Fig.
S4.18 - S4.20). Projected richness values and their future changes were slightly higher assuming
a larger dispersal ability (d/4), but on the whole all results were consistent across the three
dispersal scenarios considered (Fig. S5.21 - S5.26). Climate change impacts on future species
richness increased over time, with greater effects seen for 2080 than 2050, and the greatest
impacts being observed under a high emission scenario (RCP8.5) compared to the two lower
scenarios (Fig. S6.27 - S6.38).

Discussion

Reptile richness was projected to decrease significantly across most parts of the world in the
future (Fig. 1 & 2). This effect was apparent for lizards, snakes and turtles alike, although
regional and species-specific responses differed across the three groups (Fig. S2.8 – S2.12).

Spatial changes

Reptile richness is projected to decrease in Brazil, Australia and South Africa (Fig. 1 c, e). These
areas overlap significantly with the biotic convergence zones, areas with a high spatial
concentration of Lepidosaurians (i.e. snakes and lizards) identified by Diele-Viegas *et al.* (2020),
which were also found to cover a large number of the Lepidosaurian species vulnerable to
climate change (Diele-Viegas *et al.*, 2020). Huey *et al.* (2012) further found that ectotherms
sharing climate vulnerability traits seem to be concentrated in lowland tropical forests.
Combining climate-based SDMs with land-use change information, Newbold (2018) created
future projections for 20938 vertebrate species and found that Brazil is strongly affected by
vertebrate diversity loss due to climate change, and, together with Australia, is also likely to be
strongly affected by future land-use changes, especially under a high-emission scenario

(RCP8.5). Given that Brazil in particular does not only host a high reptile richness (Roll *et al.,*
2017; Fig. 1 a), but has recently been found to host a large number of threatened reptile species
(Cox *et al.*, 2022), this highlights the responsibility of this mega-diverse country for protecting
reptile diversity. South-western China and the western United States were projected to show a
net gain in reptile richness under climate warming (Fig. 1 e), but they also belong to those areas
where most reptile species are threatened by habitat loss from agriculture and logging or the
harvesting of species (Böhm *et al.*, 2013). These projected losses due to habitat change could
potentially counteract any positive effects climate warming might have.

The high variation in species richness changes across regions and taxa that we found obviously
reflects their original richness patterns. Species richness of amphibians, birds and mammals
together is a good spatial surrogate for species richness of all reptiles combined and of snakes,
but is not a good surrogate for lizard or turtle richness (Roll *et al.*, 2017). Thus, it is not
surprising that the areas with the highest decline in overall reptile richness (see Fig. 1) overlap
significantly with the areas of highest projected changes in vertebrate species richness
(amphibians, birds and mammals) found by Hof *et al.* (2018), although global reptile richness is
largely constrained by temperature, while global richness of all other vertebrate groups is
primarily constrained by the availability of energy and water (Qian, 2010). Further, historical
shifts in geographic ranges and climatic niches have further demonstrated that niche shifts in
endotherms are significantly faster than in ectotherms (Rolland *et al.*, 2018).

Global and zoogeographic realm changes

Globally reptile richness was projected to decline significantly, from an average of about 58 to
about 53 (9.4 %) species per grid cell from 1995 to 2080 (Fig. 2 a). This projected percentage
change in average future reptile richness (9.4 %) is considerably lower than the reptile richness
decline due to climate warming predicted by Newbold (2018). However, while Newbold (2018)
also used SDMs to infer future reptile richness changes, he used only a subset of reptile species,
for which IUCN range maps were available at that time. In addition, he also applied a much
smaller dispersal buffer (0.5 km per year), which might indicate that our projections provide a
rather optimistic scenario. Newbold (2018) also found that reptiles, together with amphibians,
are disproportionately sensitive to future human land-use. Given the synergistic effect of future
climate and land-use changes on biodiversity (Brook *et al.*, 2008) as well as species populations
(Williams *et al.*, 2022), land-use change is likely to further exacerbate climate change impacts on
global reptile distribution and diversity.

Changes in reptile richness differed among zoogeographic realms, but species richness declined
significantly across most realms over both time periods (Fig. 2 b, c, d, f, g, h, j, k). Lizards,
snakes and turtles all showed similar declines in species richness globally and across most
realms, but differed slightly across individual realms. This is in line with a previous study
covering various realms from tropical to temperate regions which found that 60 % of assessed
Lepidosaurian species (N = 1114) were vulnerable to changes in climate (Diele-Viegas *et al.*,
2020). Diele-Viegas *et al.* (2020) further found that the Afrotropical, Nearctic and Sino-Japanese
realms were the three realms where Lepidosaurians were most vulnerable to climatic change,
while Lepidosaurians in the Madagascan and Oceanian realm were least vulnerable. By contrast,
we found no significant decline in projected total reptile richness for the Sino-Japanese realm
(Fig. 2 l), although individual subgroups were projected to show a significant decline in species
richness from 1995 to 2050 (Fig. S2.13 – S2.15). Both the Madagascan and Oceanian realm were

further projected to significantly decrease in species richness including for all reptile species
(Fig. 2 d, g) and all three subgroups (Fig. S2.13 – S2.15). However, given that the Madagascan
realm, specifically, boasts over 90% of endemic reptile species and genera (Glaw & Vences,
2007) and that both realms are composed of island territories which are usually considered
highly vulnerable to climate change and might also be affected by future sea level rise and
erosion (Diele-Viegas *et al.*, 2020), our estimates might still underestimate potential climate
change impacts in these realms. Interestingly, turtle richness seemed to be least affected by
climate change, when looking at the different realms (Fig. S2.15), although a recent review has
deemed turtles as the vertebrate group with the highest extinction risk, as it is strongly affected
by habitat loss, human consumption and pet trading (Stanford *et al.*, 2020).

Biophysical changes

Reptile richness differed significantly with temperature and precipitation, with the highest
richness being observed in warm and moist conditions. Under future climate change scenarios,
the climatic conditions with high species richness were projected to shift to even more extreme
(warmer & wetter) conditions (Fig. 3). This gives cause for concern given that tropical forest and
desert lizards already live in environmental conditions that are close to their thermal limits
(Sinervo *et al.*, 2010), while desert and temperate lizard species have been found to be less able
to regulate their temperature in order to deal with heat stress than tropical species (Anderson *et*
al., 2022). Reptiles cannot regulate their body temperature internally, so are strongly dependent
on using solar energy captured by the environment to regulate their body temperature (Huey,
1982). This might lead to overheating when temperatures go beyond a species' critical limit,
which makes them particularly susceptible to climatic changes (Sinervo *et al.*, 2018). However,
this might be compensated by other biological processes that help species to buffer climate
change effects, i.e. genomic and phenotypic plasticity (Rodríguez *et al.*, 2017) as well as
behavioural and physiological adaptation (Sunday *et al.*, 2014). Overall, the persistence of reptile
species would be much more affected by climate cooling than warming, but it has been
suggested that increasing droughts, which will be a consequence of continued warming, pose a
significant future threat to European reptiles (Araújo *et al.*, 2006). It is likely that climate
warming will have an additional impact on reptiles that have temperature-dependent sex
determination. Altered sex ratios will not only result in a higher extinction risk for local
populations, but, together with a reduction in nesting sites due to habitat destruction and
fragmentation, will also affect the dispersal and potential range expansion of a species. Therefore
it could also have an impact on population demography and size unless temperature shifts in sex
determination or female nest-site choice evolves in pace with rising temperatures (Boyle *et al.*,
2016; Gibbons *et al.*, 2000).

Species-specific range changes

The range extent of most species were projected to considerably decrease, with lizard species
showing the greatest decline (Fig. 4 a). Further, most reptiles also showed a strong decline in
range overlap, which was consistent across all three groups (Fig. 4). This is in line with results
by Warren *et al.* (2018), who found that projected future range losses of more than 50% occur in
8 - 52% of considered reptile species by 2100 depending on the climate scenario considered,
although this study included only a fraction of all reptile species (N=1850) and no species
dispersal was considered.

Compared to other terrestrial vertebrate groups (especially birds and mammals), reptiles have small geographic ranges, which also indicates narrower niche requirements. This is not only likely to make them more susceptible to future climate change (Newbold, 2018), but also to other threats such as habitat loss or invasive species (Böhm *et al.*, 2013, Cox *et al.*, 2022). Reptiles are a paraphyletic class with a diverse range of body forms, habitat affinities and functional roles (Pincheira-Donoso *et al.*, 2013), which will probably result in their response to climatic and habitat changes being equally varied. In addition, cascading effects generated by disease, invasive species, habitat loss and climate change might lead to declines of sympatric species and a faster deterioration of ecosystem structure than anticipated for climate change alone (Zipkin *et al.*, 2020).

The majority of reptile species showed projected shifts towards the South, which was largely driven by range shifts in lizards (Fig. 5). Comparing the Northern and Southern hemispheres (Fig. S3.16), we found a clear overall poleward shift in species ranges across all three groups, which has also been found previously in various other taxonomic groups (Chen *et al.*, 2011). Turtle ranges were projected to shift farthest, followed by snakes (Fig. 5). The relatively short range shift distances in lizard species are probably due to the fact that lizards have the smallest range extents across the three groups (Roll *et al.*, 2017), which, given that our dispersal buffers are based on range extent, also resulted in smaller dispersal distances for lizards compared to the two other groups (Fig. S1.4). This is, however, obviously somewhat counter-intuitive and in stark contrast to generally perceived differences in mobility characteristics (e.g. movement speed) among lizards, snakes and turtles. While it is hard to tackle this problem in a globally and taxonomically comprehensive study like ours, it underlines the need for more efforts in collecting data on actual, empirically quantified dispersal distances in order to consider them in SDM exercises and large-scale biogeographical analyses.

Non-modelled species

Range-restricted (non-modelled) species were projected to experience significantly higher shift in climatic distance than modelled species (Fig. 6), indicating that range-restricted species are also likely to be strongly affected by climate change. This highlights once more that sample size restrictions of SDMs are likely to downplay climate change effects on narrow-ranging and threatened species (Platts *et al.*, 2014). Hof *et al.* (2018) also found significant impacts of climate and land-use changes on range-restricted vertebrate species, excluding reptiles. However, similar to the latter study, we only looked at climate anomalies (Euclidean distance between current and future climatic conditions) as a metric of climate change, while different metrics have been found to indicate contrasting climate change patterns on a global scale (Garcia *et al.*, 2014). In addition to climate change effects, habitat modification has been found to have a greater impact on range-restricted reptile species, as well as species with a small clutch size (Doherty *et al.*, 2020). Range-restricted reptile species are further often evolutionarily unique (Murali *et al.*, 2021) and have been found to overlap least with current conservation priority areas (Roll *et al.*, 2017, Cox *et al.*, 2022).

Sensitivity analysis

Our results were strongly dependent on the dispersal assumption, time period and emission scenario (RCP) considered (see Appendices S5 & S6). As expected, the overall patterns and richness changes were more pronounced in a later time period and scenarios representing higher

levels of greenhouse gas emissions, as these reflect potential futures with a higher level of
climate warming. Thuiller *et al.* (2019) have previously assessed the uncertainty originating from
dispersal, model algorithm, GCM and RCP on the future biodiversity scenario of amphibians,
birds and mammals and found that model algorithm and RCP have the greatest influence.

On the contrary, greater dispersal distances imply that reptile species are able to move greater
distances in order to track their optimal climatic niche and thus provide more optimistic potential
changes in species occurrence and richness patterns. Reptile-specific studies have either
considered no dispersal at all (Araújo *et al.*, 2006; Warren *et al.*, 2018) or a dispersal rate of 0.5
km per year (Newbold, 2018). We use species-specific dispersal buffers with an average of 2.4
km per year (Fig. S1.4). While these buffers might be optimistic, they are based on the
transparent rationale of a range-size dependent dispersal buffer, i.e. species with a small range
size have a smaller dispersal buffer than species with a large range size, which avoids the
unlikely assumption of uniform dispersal distances across species. Given that our model results,
as well as the underlying climate scenarios, are based on a 0.5° grid size (about 50 x 50 km),
small differences in dispersal distance do not have a strong impact on our results (see Appendix
S5). Nevertheless, these considerations again highlight the challenges in sensibly accounting for
the influence of dispersal in global climate impact assessments on species distributions and
diversity.

The projected changes in species distributions help to investigate potential changes in global
reptile richness patterns and in highlighting potential hotspots of climate change impacts. They
also allow comparison of climate change vulnerability across taxonomic groups and help to
identify areas where conservation efforts might be most urgently needed (Voskamp et al. 2022).
However, SDMs are always simplifications and the resulting projections need to be interpreted
with caution.

To improve SDMs, future studies should try to consider additional factors, such as biotic
interactions (Schleuning *et al.*, 2020) and the reshuffling of species communities (Voskamp *et*
al., in prep), which might lead to a change in competitive balance (Ockendon *et al.*, 2014),
altered predator-prey relationships (Harley, 2011) or changes in functional diversity (Stewart *et*
al., 2022) and thus the provision of ecosystem functions and services (Pecl *et al.*, 2017). While
the above factors might improve climate change impact projections in the future, such modelling
exercises will never reflect the truth, as a species' response to climate change will be strongly
influenced by behaviour, diel rhythm, thermoregulatory potential and microclimatic conditions
(Anderson *et al.*, 2022). This complexity highlights the need for integrative approaches when
investigating species' responses to climate change (Hof, 2021).

Conclusion

Our study shows that reptiles are not only likely to be impacted by future climate change,
globally but also within most zoogeographic realms. These impacts are projected to have a
considerable effect on the extent and location of species' geographic ranges. Thus, to prevent
large scale declines in reptile species, not only is it of key importance to lower CO_2 emissions in
order to stop on-going climate change, but also to maintain adequate habitats of sufficient size
and quality, especially grassland and savanna habitats (Roll *et al.*, 2017). Furthermore, it is
necessary to establish new protected areas that will help to prevent the extinction of particularly

vulnerable species, i.e. by establishing high-elevation climate refugia within current species

ranges (Sinervo *et al.*, 2018).

Data Accessibility Statement

GARD range maps are available from https://doi.org/10.5061/dryad.83s7k. EWEMBI and ISIMIP2b climate data are available from https://data.isimip.org/10.5880/pik.2019.004 and https://data.isimip.org/search/query/ISIMIP2b%20Input/tree/ISIMIP2b/InputData/climate/ atmosphere/. The code for creating the species distribution models can be found at https://github.com/christianhof/BioScen1.5_SDM, whereas the code and data for the performed data analysis and the presented figures, can be found on Dryad (doi: 10.5061/dryad.rn8pk0pgb; Biber et al. 2023).

References

Allouche, O., Tsoar, A. & Kadmon, R. (2006) Assessing the accuracy of species distribution models: Prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, **43**, 1223–1232.

Anderson, R. O., Meiri, S., & Chapple, D. G. (2022) The biogeography of warming tolerance in lizards. *Journal of Biogeography*.

Araújo, M.B., Thuiller, W. & Pearson, R.G. (2006) Climate warming and the decline of amphibians and reptiles in Europe. *Journal of Biogeography*, **33**, 1712–1728.

Bagchi, R., Crosby, M., Huntley, B., Hole, D.G., Butchart, S.H., Collingham, Y., Kalra, M., Rajkumar, J., Rahmani, A., Pandey, M. & others (2013) Evaluating the effectiveness of conservation site networks under climate change: Accounting for uncertainty. *Global Change Biology*, **19**, 1236–1248.

Baisero, D., Visconti, P., Pacifici, M., Cimatti, M. & Rondinini, C. (2020) Projected global loss of mammal habitat due to land-use and climate change. *One Earth*, **2**, 578–585.

Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: how, where and how many?. *Methods in Ecology and Evolution*, *3*(2), 327-338.

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. (2012) Impacts of climate change on the future of biodiversity. *Ecology Letters*, **15**, 365–377.

Biber, M.F., Voskamp, A., Niamir, A., Hickler, T. & Hof, C. (2020) A comparison of macroecological and stacked species distribution models to predict future global terrestrial vertebrate richness. *Journal of Biogeography*, **47**, 114–129.

Biber, M.F., Voskamp, A., & Hof, C. (2023) Data from: Potential future climate change effects on global reptile distribution and diversity. Dryad. https://doi.org/10.5061/dryad.rn8pk0pgb.

Böhm, M., Collen, B., Baillie, J.E., Bowles, P., Chanson, J., Cox, N., Hammerson, G., Hoffmann, M., Livingstone, S.R., Ram, M. & others (2013) The conservation status of the world's reptiles. *Biological Conservation*, **157**, 372–385.

Bowler, D.E., Hof, C., Haase, P., Kröncke, I., Schweiger, O., Adrian, R., Baert, L., Bauer, H.-G., Blick, T., Brooker, R.W. & others (2017) Cross-realm assessment of climate change impacts on species' abundance trends. *Nature Ecology & Evolution*, **1**, 1–7.

Boyle, M., Schwanz, L., Hone, J. & Georges, A. (2016) Dispersal and climate warming determine range shift in model reptile populations. *Ecological Modelling*, **328**, 34–43.

Brook, B.W., Sodhi, N.S., & Bradshaw, C.J.A. (2008) Synergies among extinction drivers under global change. *Trends in Ecology & Evolution*, *23*, 453–460.

Brooks, T. M., Mittermeier, R. A., Da Fonseca, G. A., Gerlach, J., Hoffmann, M., Lamoreux, J. F., & others (2006) Global biodiversity conservation priorities. S*cience*, **313**, 58-61.

Calabrese, J.M., Certain, G., Kraan, C. & Dormann, C.F. (2014) Stacking species distribution
models and adjusting bias by linking them to macroecological models. *Global Ecology and*
Biogeography, **23**, 99–112.

Chen, I.-C., Hill, J.K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. (2011) Rapid range shifts of
species associated with high levels of climate warming. *Science*, **333**, 1024–1026.

Cox, N., Young, B. E., Bowles, P., Fernandez, M., Marin, J., Rapacciuolo, G. & others (2022) A
global reptile assessment highlights shared conservation needs of tetrapods. *Nature*, **605**, 285-
290.

Diele-Viegas, L.M., Figueroa, R., Vilela, B. & Rocha, C. (2020) Are reptiles toast? A worldwide
evaluation of lepidosauria vulnerability to climate change. *Climatic Change*, **159**, 581–599.

Diele-Viegas, L. M., & Rocha, C. F. D. (2018) Unraveling the influences of climate change in
Lepidosauria (Reptilia). *Journal of Thermal Biology*, **78**, 401-414.

Doherty, T.S., Balouch, S., Bell, K., Burns, T.J., Feldman, A., Fist, C., Garvey, T.F., Jessop,
T.S., Meiri, S. & Driscoll, D.A. (2020) Reptile responses to anthropogenic habitat modification:
A global meta-analysis. *Global Ecology and Biogeography*, **29**, 1265–1279.

Elith, J., Kearney, M. & Phillips, S. (2010) The art of modelling range-shifting species. *Methods*
in Ecology and Evolution, **1**, 330–342.

Elith, J. & Leathwick, J.R. (2009) Species distribution models: Ecological explanation and
prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*, **40**,
677–697.

Frieler, K., Lange, S., Piontek, F., Reyer, C.P., Schewe, J., Warszawski, L., Zhao, F., Chini, L.,
Denvil, S., Emanuel, K. & others (2017) Assessing the impacts of 1.5°c global warming–
simulation protocol of the inter-sectoral impact model intercomparison project (ISIMIP2b).
Geoscientific Model Development, **10**, 4321–4345.

Garcia, R.A., Cabeza, M., Rahbek, C. & Araújo, M.B. (2014) Multiple dimensions of climate
change and their implications for biodiversity. *Science*, **344**, 1247579.

Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tuberville, T.D., Metts, B.S., Greene,
J.L., Mills, T., Leiden, Y., Poppy, S. & others (2000) The global decline of reptiles, déjà vu
amphibians: Reptile species are declining on a global scale. Six significant threats to reptile
populations are habitat loss and degradation, introduced invasive species, environmental
pollution, disease, unsustainable use, and global climate change. *BioScience*, **50**, 653–666.

Gibson-Reinemer, D.K. & Rahel, F.J. (2015) Inconsistent range shifts within species highlight
idiosyncratic responses to climate warming. *PloS One*, **10**, e0132103.

Glaw, F. & Vences, M. (2007) *A field guide to the amphibians and reptiles of Madagascar*,
Vences & Glaw.

Greenwell, B., Boehmke, B., Cunningham, J. & Developers, G. (2020) *GBM: Generalized*
boosted regression models.

Guisan, A., & Thuiller, W. (2005) Predicting species distribution: offering more than simple
habitat models. *Ecology Letters*, *8*(9), 993-1009.

Harley, C.D. (2011) Climate change, keystone predation, and biodiversity loss. *Science*, **334**,
1124–1127.

Hernandez, P. A., Graham, C. H., Master, L. L., & Albert, D. L. (2006) The effect of sample size
and species characteristics on performance of different species distribution modelling methods.
Ecography, **29**, 773-785.

Hijmans, R.J., Phillips, S., Leathwick, J. & Elith, J. (2021) *Dismo: Species distribution*
modelling.

Hof, C. (2021) Towards more integration of physiology, dispersal and land-use change to
understand the responses of species to climate change. *Journal of Experimental Biology*, **224**,
jeb238352.

Hof, C., Araújo, M.B., Jetz, W. & Rahbek, C. (2011) Additive threats from pathogens, climate
and land-use change for global amphibian diversity. *Nature*, **480**, 516–519.

Hof, C., Voskamp, A., Biber, M.F., Böhning-Gaese, K., Engelhardt, E.K., Niamir, A., Willis,
S.G. & Hickler, T. (2018) Bioenergy cropland expansion may offset positive effects of climate
change mitigation for global vertebrate diversity. *Proceedings of the National Academy of*
Sciences, **115**, 13294–13299.

Holt, B.G., Lessard, J.-P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D., Fabre, P.-
H., Graham, C.H., Graves, G.R., Jønsson, K.A. & others (2013) An update of Wallace's
zoogeographic regions of the world. *Science*, **339**, 74–78.

Huey, R.B. (1982) *Temperature, physiology, and the ecology of reptiles*. *Biology of the reptilia*,
Citeseer.

Huey, R.B., Kearney, M.R., Krockenberger, A., Holtum, J.A., Jess, M. & Williams, S.E. (2012)
Predicting organismal vulnerability to climate warming: Roles of behaviour, physiology and
adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**, 1665–
1679.

IPBES (2019) Summary for policymakers of the global assessment report on biodiversity and
ecosystem services of the intergovernmental science-policy platform on biodiversity and
ecosystem services. *IPBES secretariat, Bonn, Germany*.

IPCC (2021) Climate change 2021: The physical science basis. Working group i contribution to
the sixth assessment report of the intergovernmental panel on climate change. *Cambridge*
University Press.

IPCC (2022) Climate change 2022: Impacts, adaptation and vulnerability. Working group II
contribution to the sixth assessment report of the intergovernmental panel on climate change.
Cambridge University Press.

IUCN (2022) The IUCN red list of threatened species. Available at https://www.iucnredlist.org/resources/spatial-data-download. *Cambridge University Press*.

Kühn, I. (2007) Incorporating spatial autocorrelation may invert observed patterns. *Diversity and Distributions*, **13**, 66–69.

Lange, S. (2016) EartH2Observe, WFDEI and ERA-interim data merged and bias-corrected for ISIMIP (EWEMBI).

Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J. & Grenouillet, G. (2020) Species better track climate warming in the oceans than on land. *Nature Ecology & Evolution*, **4**, 1044–1059.

Meynard, C.N. & Quinn, J.F. (2007) Predicting species distributions: A critical comparison of the most common statistical models using artificial species. *Journal of Biogeography*, **34**, 1455–1469.

Monasterio, C., Shoo, L. P., Salvador, A., Iraeta, P., & Díaz, J. A. (2013) High temperature constrains reproductive success in a temperate lizard: implications for distribution range limits and the impacts of climate change. *Journal of Zoology*, **291**, 136-145.

Murali, G., Gumbs, R., Meiri, S., & Roll, U. (2021) Global determinants and conservation of evolutionary and geographic rarity in land vertebrates. *Science Advances*, **7(42)**, eabe5582.

Nathan, R., Klein, E., Robledo-Arnuncio, J.J. & Revilla, E. (2012) *Dispersal kernels*, Oxford University Press Oxford, UK.

Newbold, T. (2018) Future effects of climate and land-use change on terrestrial vertebrate community diversity under different scenarios. *Proceedings of the Royal Society B*, **285**, 20180792.

Ockendon, N., Baker, D.J., Carr, J.A., White, E.C., Almond, R.E., Amano, T., Bertram, E., Bradbury, R.B., Bradley, C., Butchart, S.H. & others (2014) Mechanisms underpinning climatic impacts on natural populations: Altered species interactions are more important than direct effects. *Global Change Biology*, **20**, 2221–2229.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V., Underwood, E.C., D'amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C. & others (2001) Terrestrial ecoregions of the world: A new map of life on Earth. *BioScience*, **51**, 933–938.

Pacifici, M., Foden, W.B., Visconti, P., Watson, J.E., Butchart, S.H., Kovacs, K.M., Scheffers, B.R., Hole, D.G., Martin, T.G., Akçakaya, H.R. & others (2015) Assessing species vulnerability to climate change. *Nature Climate Change*, **5**, 215–224.

Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.-C., Clark, T.D., Colwell, R.K., Danielsen, F., Evengård, B. & others (2017) Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, **355**, eaai9214.

Petitpierre, B., Broennimann, O., Kueffer, C., Daehler, C., & Guisan, A. (2017) Selecting
predictors to maximize the transferability of species distribution models: Lessons from cross-
continental plant invasions. *Global Ecology and Biogeography*, **26**, 275-287.

Pincheira-Donoso, D., Bauer, A. M., Meiri, S., & Uetz, P. (2013) Global taxonomic diversity of
living reptiles. *PloS One*, **8**, e59741.

Platts, P.J., Garcia, R.A., Hof, C., Foden, W., Hansen, L.A., Rahbek, C. & Burgess, N.D. (2014)
Conservation implications of omitting narrow-ranging taxa from species distribution models,
now and in the future. *Diversity and Distributions*, **20**, 1307–1320.

Porfirio, L. L., Harris, R. M., Lefroy, E. C., Hugh, S., Gould, S. F., Lee, G. & others (2014)
Improving the use of species distribution models in conservation planning and management
under climate change. *PLoS One*, **9**, e113749.

Qian, H. (2010) Environment–richness relationships for mammals, birds, reptiles, and
amphibians at global and regional scales. *Ecological Research*, **25**, 629–637.

R Core Team (2021) *R: A language and environment for statistical computing*, R Foundation for
Statistical Computing, Vienna, Austria.

Rodríguez, A., Rusciano, T., Hamilton, R., Holmes, L., Jordan, D. & Wollenberg Valero, K.C.
(2017) Genomic and phenotypic signatures of climate adaptation in an anolis lizard. *Ecology and*
Evolution, **7**, 6390–6403.

Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A.M., Bernard, R., Böhm, M., Castro-
Herrera, F., Chirio, L., Collen, B. & others (2017) The global distribution of tetrapods reveals a
need for targeted reptile conservation. *Nature Ecology & Evolution*, **1**, 1677–1682.

Rolland, J., Silvestro, D., Schluter, D., Guisan, A., Broennimann, O. & Salamin, N. (2018) The
impact of endothermy on the climatic niche evolution and the distribution of vertebrate diversity.
Nature Ecology & Evolution, **2**, 459–464.

Scheffers, B.R., De Meester, L., Bridge, T.C., Hoffmann, A.A., Pandolfi, J.M., Corlett, R.T.,
Butchart, S.H., Pearce-Kelly, P., Kovacs, K.M., Dudgeon, D. & others (2016) The broad
footprint of climate change from genes to biomes to people. *Science*, **354**, aaf7671.

Schleuning, M., Neuschulz, E. L., Albrecht, J., Bender, I. M., Bowler, D. E., Dehling, D. M. &
others (2020) Trait-based assessments of climate-change impacts on interacting species. *Trends*
in Ecology & Evolution, **35**(4), 319-328.

Sinervo, B., Mendez-De-La-Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Villagrán-Santa
Cruz, M. & others (2010) Erosion of lizard diversity by climate change and altered thermal
niches. *Science*, **328**, 894-899.

Sinervo, B., Miles, D.B., Wu, Y., MÉNDEZ-DE LA CRUZ, F.R., Kirchhof, S. & Qi, Y. (2018)
Climate change, thermal niches, extinction risk and maternal-effect rescue of toad-headed
lizards, phrynocephalus, in thermal extremes of the Arabian peninsula to the Qinghai—Tibetan
plateau. *Integrative Zoology*, **13**, 450–470.

Stanford, C. B., Iverson, J. B., Rhodin, A. G., van Dijk, P. P., Mittermeier, R. A., Kuchling, G. &
others (2020) Turtles and tortoises are in trouble. *Current Biology*, **30**, R721-R735.

Stewart, P.S., Voskamp, A., Santini, L., Biber, M.F., Hof, C., Willis, S.G. & Tobias, J.A. (2022)
Global impacts of climate change on avian functional diversity. *Ecology Letters*.

Sunday, J.M., Bates, A.E., Kearney, M.R., Colwell, R.K., Dulvy, N.K., Longino, J.T. & Huey,
R.B. (2014) Thermal-safety margins and the necessity of thermoregulatory behavior across
latitude and elevation. *Proceedings of the National Academy of Sciences*, **111**, 5610–5615.

Thuiller, W., Guéguen, M., Renaud, J., Karger, D.N. & Zimmermann, N.E. (2019) Uncertainty
in ensembles of global biodiversity scenarios. *Nature Communications*, **10**, 1–9.

Thuiller, W., Lavorel, S., Araújo, M.B., Sykes, M.T. & Prentice, I.C. (2005) Climate change
threats to plant diversity in europe. *Proceedings of the National Academy of Sciences*, **102**,
8245–8250.

Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K. & others
(2011) The representative concentration pathways: an overview. *Climatic change*, **109**, 5-31.

Voskamp, A., Baker, D.J., Stephens, Philip A, Howard, C., Butchart, S.H. & Willis, S.G. (in
prep) Impacts of climate change on the world's terrestrial bird assemblages.

Voskamp, A., Butchart, S.H., Baker, D.J., Wilsey, C.B. & Willis, S.G. (2021) Site-based
conservation of terrestrial bird species in the Caribbean and Central and South America under
climate change. *Frontiers in Ecology and Evolution*, **9**, 234.

Voskamp, A., Fritz, S. A., Koecke, V., Biber, M., Nogueira-Brockmeyer, T., Bertzky, B. &
others (2022) How to resolve conflicting conservation objectives: A decision support tool for the
global selection of multi-purpose protected areas. *bioRxiv*.

Warren, R., Price, J., Graham, E., Forstenhaeusler, N. & VanDerWal, J. (2018) The projected
effect on insects, vertebrates, and plants of limiting global warming to 1.5°c rather than 2°c.
Science, **360**, 791–795.

Watterson, I. G. (2019) Indices of climate change based on patterns from CMIP5 models, and the
range of projections. *Climate Dynamics*, **52**, 2451-2466.

Wiens, J.J. & Graham, C.H. (2005) Niche conservatism: Integrating evolution, ecology, and
conservation biology. *Annual Review of Ecology, Evolution, and Systematics*, **36**, 519–539.

Williams, J.J., Freeman, R., Spooner, F. & Newbold, T. (2022) Vertebrate population trends are
influenced by interactions between land use, climatic position, habitat loss and climate change.
Global Change Biology, **28**, 797–815.

Wood, S.N. (2011) Fast stable restricted maximum likelihood and marginal likelihood estimation
of semiparametric generalized linear models. *Journal of the Royal Statistical Society (B)*, **73**, 3–
36.

Wood, S.N. (2003) Thin-plate regression splines. *Journal of the Royal Statistical Society (B)*, **65**,
95–114.

Zipkin, E.F., DiRenzo, G.V., Ray, J.M., Rossman, S. & Lips, K.R. (2020) Tropical snake
diversity collapses after widespread amphibian loss. *Science*, **367**, 814–816.

Tables

Table 1. Number of species that were excluded from the species distribution models due to their
restricted range or low model performance.

Taxonomic group	Lizard	Snake	Turtle	Total
No. of species with available data	6328	3414	322	10064
No. of range-restricted species (N < 10, removed)	2536	1047	19	3602
No. of species with low model performance (AUC < 0.7, removed)	97	62	7	166
Total number of species modelled	**3695**	**2305**	**296**	**6296**
Percentage of available species modelled	58.4 %	67.5 %	91.9 %	62.6 %

Figures

Figure 1. a) Map of projected global terrestrial reptile species richness (1995), b) frequency of species richness by taxonomic group (lizard, snake, turtle and total) with mean values highlighted by vertical line and c) increase, d) decrease, e) net change and f) relative change (%) in reptile species richness for all modelled reptile species (N = 6296) for the year 2080 under a medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8). Results are presented as the ensemble mean, across the four global circulation models (GCMs) and two model algorithms (GAM & GBM) considered. All maps are based on 0.5° x 0.5° grid cells, which have been projected to Mollweide equal-area projection (EPSG:54009). Grey areas are regions for which no projections are available. Note that the colour scales differ among the individual panels.

786
Figure 2. Terrestrial reptile species richness across the globe and for each zoogeographic realm
(Afrotropical, Australian, Madagascan, Nearctic, Neotropical, Oceanian, Oriental, Palearctic,
Panamanian, Saharo-Arabian & Sino-Japanese) over time (1995, 2080) based on all modelled
reptile species (N = 6296). Results are presented as the ensemble mean across the four global
circulation models (GCMs) and two model algorithms (GAM & GBM) considered, under a
medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).
The statistical difference between years was tested using a paired Student's t-test with Holm
correction ($p < 0.05$ = *, $p < 0.01$ = **, $p < 0.001$ = ***, $p < 0.0001$ = ****). Plots show mean
(red point & label), median (black horizontal line), 25th to 75th percentiles (box), entire range of
data (violin & data points) and density of values (width of violin). Figure 5 provides a map
outlining the different zoogeographic realms.

798
Figure 3. Univariate relationship of current (1995) and future (2080 RCP6.0) reptile species
richness with a) temperature, b) precipitation; and the bivariate relationship of temperature and
precipitation with reptile species richness for c) 1995 and d) 2080 RCP6.0; and the respective e)
net richness change and f) relative richness change (%) under a medium dispersal scenario (d/8).
Heat maps and lines show the mean and ribbons the standard deviation in variance across space,
global circulation models (GCMs) and the two model algorithms (GAM & GBM).

Figure 4. Frequency plots of the mean number of reptile species (a) lizard, c) snake, e) turtle) and their potential future change (%) in range extent (total area of occupied grid cells) and the mean number of reptile species (b) lizard, d) snake, f) turtle) per potential range overlap class (0-20, 20-40, 40-60, 60-80, 80-100). Error margins/bars indicate the standard deviation across the four global circulation models (GCMs) and the two model algorithms (GAM & GBM) used. Both shown for 2080 under a medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).

Figure 5. Cumulative direction and distance of potential range centroid changes per taxonomic
group (a) lizard, b) snake, c) turtle) and d) range centroids (points on map) and the number of
species and their directional shift in range centroid position per zoogeographic realm (inset polar
plots). Results are presented as the ensemble mean across the four global circulation models
(GCMs) and two model algorithms (GAM & GBM) considered, for the year 2080 under a
medium representative concentration pathway (RCP6.0) and a medium dispersal scenario (d/8).
The inset bar chart shows the number of species that have their range centroid located in the
respective realm. Numbers show the total number of species per realm, while the bars are colour
coded by the three different taxonomic groups. Please note that a considerable number of species
(n = 518) have their range centroid located outside of the zoogeographic realm boundaries and
thus were not associated with any realm and so are not considered for the polar plots in Fig. 5 d).

825
Figure 6. Mean climatic distance for modelled and non-modelled species, split by taxonomic
group (a) lizard, b) snake, c) turtle and d) total). The statistical difference between modelled and
non-modelled species was tested using a Student's t-test with Holm correction ($p < 0.05$ = *, $p <$
0.01 = **, $p < 0.001$ = ***, $p < 0.0001$ = ****). Plots show mean (red point & label), median
(black horizontal line), 25th to 75th percentiles (box), entire range of data (violin & data points)
and density of values (width of violin). The results are shown as the ensemble mean across the
four global circulation models (GCMs) for the year 2080 under a medium representative
concentration pathway (RCP6.0).

Supplementary Material

Appendix S1 Methodological details

Appendix S2 Taxon-specific results

Appendix S3 Additional range change results

Appendix S4 Summed probability results

Appendix S5 Variation across dispersal distances

Appendix S6 Variation across RCPs and years