Population genomics provide insights into the global genetic structure of *Colletotrichum*

***graminicola,* the causal agent of maize anthracnose**

Flávia Rogério[1], Riccardo Baroncelli[1,2], Francisco Borja Cuevas-Fernández[1], Sioly Becerra[1],

JoAnne Crouch[3], Wagner Bettiol[4], M. Andrea Azcárate-Peril[5], Martha Malapi-Wight[6],

Veronique Ortega[7], Javier Betran[8], Albert Tenuta[9] José S. Dambolena[10], Paul D. Esker[11],

Pedro Revilla[12], Tamra A. Jackson-Ziems[13], Jürg Hiltbrunner[14], Gary Munkvold[15], Ivica

Buhiniček[16], José L. Vicente-Villardón[17], Serenella A. Sukno[1*], Michael R. Thon[1*]

[1]Instituto de Investigación en Agrobiotecnología (CIALE), Departamento de Microbiología y Genética,

Universidad de Salamanca, Villamayor, 37185 Salamanca, Spain. [2] Department of Agricultural and Food

Sciences (DISTAL), University of Bologna, Viale Fanin 44, 40126 Bologna, Italy. [3]United States Department of

Agriculture, Foreign Disease and Weed Science Unit, 1301 Ditto Avenue, Fort Detrick MD, 21702 USA. [4]

Embrapa Environment, Jaguariúna, SP, Brazil. [5]Center for Gastrointestinal Biology and Disease, Division of

Gastroenterology and Hepatology, and UNC Microbiome Core, Department of Medicine, School of Medicine,

University of North Carolina, Chapel Hill, NC 27599, USA. [6]USDA-Animal and Plant Health Inspection

Services, Biotechnology Regulatory Services, Riverdale, MD 20737, USA. [7] Syngenta Seeds La Grangette,

32220 Lombez France. [8] Bayer Crop Science/Monsanto SAS 82170, Monbequi, France. [9] Ontario Ministry of

Agriculture, Food, and Rural Affairs, University of Guelph-Ridgetown, Ridgetown, Ontario, Canada. [10]

Facultad de Ciencias Exactas Físicas y Naturales, Universidad Nacional de Córdoba, IMBIV-CONICET-ICTA

Av Vélez Sarsfield 1611, Ciudad Universitaria, Córdoba, Argentina. [11]Department of Plant Pathology and

Environmental Microbiology, Pennsylvania State University, State College, PA, 16801, United States. [12]Misión

Biológica de Galicia, Spanish National Research Council (CSIC), PO Box 2836080, Pontevedra, Spain.

[13]Department of Plant Pathology, University of Nebraska–Lincoln, 68583-0722, United States. [14]Federal

Department of Economic Affairs, Agroscope Reckenholzstrasse 191, 8046 Zurich, Switzerland. [15]Department of

Plant Pathology and Microbiology, Iowa State University, Ames, IA 5001, United States. [16] Bc Institute for

Breeding and Production of Field Crops, 10370 Dugo Selo, Croatia. [17] Statistics Department University of

Salamanca, Salamanca, Spain.

* Co-senior authors

***Corresponding authors:** Serenella A. Sukno, Institute for Agribiotechnology Research (CIALE), Departamento de Microbiología
y Genética, Universidad de Salamanca, Villamayor, 37185 Salamanca, Spain. E-mail: ssukno@usal.es https://orcid.org/0000-0003-3248-
6490; Michael R. Thon, Institute for Agribiotechnology Research (CIALE), Departamento de Microbiología y Genética, Universidad de
Salamanca, Villamayor, 37185 Salamanca, Spain. E-mail: mthon@usal.es https://orcid.org/0000-0002-7225-7003

Abstract

Background: *Colletotrichum graminicola*, the causal agent of maize anthracnose, is an

important crop disease worldwide. Understanding the genetic diversity and mechanisms

underlying genetic variation in pathogen populations is crucial to the development of

effective control strategies. The genus *Colletotrichum* is largely recognized as asexual, but

several species have been reported to have a sexual cycle. Here, we employed a population

genomics approach to investigate the genetic diversity and reproductive biology of *C.*

graminicola isolates infecting maize. We sequenced 108 isolates of *C. graminicola* collected

in 14 countries using restriction site-associated DNA sequencing (RAD-Seq) and whole-

genome sequencing (WGS). **Results:** Clustering analyses based on single-nucleotide

polymorphisms showed populational differentiation at a global scale, with three genetic

groups delimited by continental origin, compatible with short-dispersal of the pathogen, and

geographic subdivision. Distinct levels of genetic diversity were observed between these

clades, suggesting different evolutionary histories. Intra and inter-continental migration was

predicted between Europe and South America, likely associated with the movement of

contaminated germplasm. Low clonality and evidence of genetic recombination were

detected from the analysis of linkage disequilibrium and the pairwise homoplasy index (PHI)

test for clonality. We show evidence that even if rare (possibly due to losses of sex and

meiosis-associated genes) *C. graminicola* can undergo sexual recombination based on lab

assays and genomic analyses. **Conclusions:** Our results support hypotheses of intra and

intercontinental pathogen migration and genetic recombination with great impact on *C.*

graminicola population structure.

Keywords: Fungal plant pathogen, population genetics, phylogeography, phytopathology,

recombination, migration

63   **Introduction**

Crop diseases are one of the biggest concerns in global agriculture responsible for

major losses in food production [1,2]. Fungi are primary causal agents of plant diseases and

can cause catastrophic damage [3,4]. Disease outbreaks caused by fungal plant pathogens

during recent decades have increased and are recognized as emergent threats to food security

worldwide [5–8]. Human activity tends to intensify fungal disease dispersal by promoting

pathogen movement far from their native location [9,10]. The modification of natural

environments caused by modern agriculture provides new opportunities for natural selection

[10]. The employ of crops with a limited genetic diversity increases the risks of global

disease spread since pathogen genotypes can rapidly spread through genetically uniform host

populations [11,12]. Additionally, ongoing climate changes can shift plant-pathogen

distributions on a global scale since temperature changes may affect diseases development,

hence acting in the spread and survival of pathogens and their establishment in hitherto

unsuitable regions [8,13,14].

Maize (*Zea mays*) is the second most extensively cultivated cereal crop globally, with

a harvested area of almost 200 million hectares [15]. Together with rice and wheat, maize

supplies nearly half daily calories in human food [14, 16]. Although many diseases

negatively impact maize production, anthracnose is one of the most important diseases, it is

so difficult to control that requires to use of integrated practices [17–20]. Anthracnose disease

is caused by *Colletotrichum graminicola*, an ascomycete fungus that can infect most plant

tissues, though the stalk rot and seedling blight can bring more significant economic damage

[21–23]. Taxonomically, *C. graminicola* belongs to the Graminicola species complex, a well-

defined monophyletic clade encompassing 16 *Colletotrichum* species pathogenic to the

Poaceae family [23–25].

Prior to 1960, maize anthracnose was a minor leaf spot disease in the USA, and the

causal agent *C. graminicola* was primarily studied for its taxonomic significance [22,26].

However, this view changed during the following decades when devastating outbreaks caused

unexpected losses across north-central and eastern regions of the USA [19,22]. Afterwards,

the disease became increasingly common, and in some seasons extremely destructive

throughout the eastern USA [27–29]. The occurrence of these outbreaks are thought to be

associated with the use of hybrids particularly susceptible to anthracnose, more favorable

weather conditions, and the emergence of more aggressive strains of the pathogen with rapid

spread through the maize belt [30–32]. Nowadays, maize anthracnose is a well-established

endemic disease in the USA, and better managed mainly through the use of resistant

cultivars, but it still can cause important damage in some regions of the country [33–37].

Given the history of the disease, it may have the potential to become much more significant

in the future.

Outside of the USA, *C. graminicola* is globally distributed and is of major importance

in the Americas, and other regions of the world [38]. In North America, Canada reports

variable disease incidence, but anthracnose is still the cause of significant concern [36,39,40].

In addition to temperate areas, maize anthracnose is also reported in tropical and subtropical

regions, including Brazil and Argentina, where warm and humid conditions contribute to

augment the disease [41–48]. Since the first report of anthracnose in Europe, (Italy in 1852)

[49], several countries have reported the occurrence of the disease, including Germany,

France, and ex Czechoslovakia [50–52], and more recently Croatia, Portugal, Switzerland,

Bosnia and Herzegovina [53–57]. Recently, maize anthracnose also was reported in Asia

[58], indicating that the pathogen is increasing its geographic range.

Pathogens widely disseminated over continents indicate an intense dispersal of their

propagules, either from aerial dispersal, or via human-mediated movement of infected plant

material, and/or seeds [11,59–61]. The nature of fungal propagules, i.e., their small and

numerous spores, favor long-distance dispersal resulting in the spread of pathogens across

and between continents [11]). Although many pathogens are cosmopolitan in distribution,

their populations often can differ in some aspects, for instance, virulence, resistance to

chemical controls, and genetic diversity [62]. The migration of plant pathogens can have

epidemiological and evolutionary implications since the movement of individuals from one

population to another may result in gene flow when migrants contribute their genes to the

gene pool of another population [63]. *Colletotrichum graminicola* is seed-transmitted, also

known as a seed-born pathogen [64–66], therefore, contaminated seeds serve as inoculum

sources, and promote pathogen movement. Furthermore, though the transport of seeds and

infected plant material, fungi can naturally be spread via the dispersal of both sexual and

asexual spores [11,59,67]. In addition to the dispersal process, the reproductive biology of a

species is crucial to understanding the evolutionary forces that shape the genetic diversity of

pathogen populations over time and space [68]. Asexual reproduction predominates in the

most of plant-pathogenic fungi, but many species undergo sexual cycles, and many others

employ both reproductive strategies [63]. Sexual reproduction produces genetic

recombination by the independent assortment of chromosomes, increasing genotype diversity

and generating a random association between alleles at different loci [69,70]. Yet, alternative

mechanisms, such as parasexual reproduction, and cryptic sexual cycle can also create

genetic recombination in fungal species [71,72]. While sexual reproduction may be costly (in

terms of fitness) and disrupt favorable gene combinations, experimental evidence has

suggested that sex in fungi may increase the rate of adaptation to new environments [73,74].

Recombination can create particular genotypes through the combination of more fit alleles,

which can accelerate the evolution of resistance to multiple fungicides and overtake R genes

of resistant cultivars [69,75,76].

The genetic control underlying sexual development in the genus *Colletotrichum* is not

well known and differs from the typical bipolar mating system of ascomycetes [77–79]. All

species studied so far contain just one of the two mating types alleles typical of ascomycetes

(*MAT1-1/2*) [23], which would seemingly limit their ability to mate. However, for at least

some *Colletotrichum* species, including *C. graminicola* sexual reproduction has been

demonstrated. Although the teleomorph of *C. graminicola* has been characterized under

laboratory conditions [78,79], the asexual state is the one commonly found in the field

conditions, which plays a significant role in the lifecycle. This species is reported to be a

homothallic species [78], i.e., possesses both sets of genes and thus can complete the sexual

cycle independently [80]. However, some strains are self-sterile and cross-fertile [79,81],

which means that some strains are capable of reproducing sexually when cultured alone,

while other isolates require a mating partner to produce sexual offspring.

Genetic recombination is reported as an important factor in the ecology and evolution

of many *Colletotrichum* species [82–87]. However, only a few population studies have

characterized the genetic diversity of grass-infecting *Colletotrichum* species, such as *C.*

sublineola and, *C. cereale* [85–89], and little is known about *C. graminicola* population

diversity. Studies using single marker genes and limited sample size showed high genetic

variability on *C. graminicola* isolates [87,90]. Therefore, in this study, we investigated the

global population structure of *C. graminicola* to get insights into the reproductive biology of

this species. Our population genomic analyses using single-nucleotide polymorphisms

(SNPs) revealed population subdivision into three genetic clades with isolates grouped

according to their country of origin, with distinct levels of genetic diversity between them,

suggesting different evolutionary histories. We found intra and intercontinental migration,

mainly between Europe and South America, likely associated with the movement of

contaminated seeds. We discussed the possible evolutionary origins of these genetic groups

and the impact of recombination on the genetic structure as well the potential of maize

anthracnose becoming a much more significant disease in the future.

Materials And Methods

Sampling and isolation of *C. graminicola*

Fresh collections of *C. graminicola* samples were obtained from symptomatic maize stems provided by an international network of maize researchers and producers, allowed sampling from nine different countries where anthracnose disease is prevalent (Fig.1). Eleven *C. graminicola* isolates originally collected from five countries (Germany, Japan, Netherlands, Nigeria, and Zimbabwe) [91,92] were obtained from public culture collections (Supplementary Table S1) and also included in the collection. Symptomatic stems were dissected, surface disinfected, and plated in culture medium to fungal isolation, as previously described [21,57]. Fungal structures (hyphal tips, acervuli) emerging from maize tissue were transferred to potato dextrose agar (PDA) medium allowed to grow for 7 days at 23°C then single-spore purified. Monosporic cultures were stored at -80°C from suspension conidia in 7.5% skim milk (Merck, Darmstadt Germany) mixed on silica gel (S/0730/53 Fisher) [93]. Isolates were verified as *C. graminicola* by inspection of colony morphology, spore size and shape, and sequencing of the internal transcribed spacer (*ITS*) region of rDNA using the primers ITS4 and ITS5 [94], following PCR amplification conditions as described by Rech et al. [85]. The amplified ITS fragments were purified using NucleoSpin gel and PCR Clean-up kits (Macherey Nagel, Düren, Germany) and sequenced using Sanger technology by University of Salamanca (Spain). Specie identity was confirmed by BLAST searches performed against in GenBank (NCBI). All materials used in this study were used in accordance with the rights and obligations of the recipient as described in the International Treaty on Plant Genetic Resources for Food and Agriculture adopted by the FAO on November 3, 2001.

Figure1. Worldwide *Colletotrichum graminicola* sampling.

DNA extraction, library preparation, and sequencing

Mycelia from monosporic cultures were incubated on an orbital shaker in potato

dextrose broth (PDB) (Difco, Becton, Dickinson and Company, Franklin Lakes, USA) for

three days at 25°C and 150 rpm under continuous light [54]. Genomic DNA was extracted

using DNeasy Plant Mini Kits (Qiagen Inc., Valencia, California, USA) and quantified using

a Qubit 3.0 fluorimeter (Invitrogen, Waltham, USA). DNA for *C. graminicola* isolates used

for whole genome sequencing (WGS) was extracted using the cetyltrimethylammonium-

bromide (CTAB) protocol adapted from Baek and Kernerley [95].

New DNA sequence datasets were generated from 101 *C. graminicola* isolates using

either RAD-seq or WGS; sequences for an additional seven isolates were also included in the

analyses [92] (Supplementary Table S1, S2). A total of 88 isolates were genotyped with

RAD-seq by Floragenex Inc. (Beaverton, Oregon, USA). Briefly, RAD-seq libraries with

sample-specific barcode sequences were constructed from DNA digested with restriction

enzyme PstI, then single-end sequenced (1 x 100 bp) on one lane using TrueSeq PE150 kit

(Illumina, Inc., San Diego, CA) on an Illumina HiSeq 2000 instrument. WGS data was

obtained from eight *C. graminicola* isolates (CBS252.59, F-64330-2, F-64330-7, F-40330-1,

SW-8046-2, CRO-I-35, CR-10370-50, SW-8046-2) from libraries prepared using the Nextera

XT DNA kit (Illumina, Inc, San Diego, CA, and sequenced on Illumina HiSeq4000 (2x150

bp). WGS data was obtained from six *C. graminicola* isolates (ARG-2700-2, CA-CHAT-1,

CA-N8H, CRO-I-41, P-7565-072-1, SW-8046-1) from paired-end libraries prepared using

the Illumina TruSeq DNA Nano kit and sequenced on an Illumina MiSeq (2x300 bp).

Sequence quality was checked on FASTQC v0.11.7 (Babraham Bioinformatics) and low-

quality reads and adaptors were trimmed with TRIMMOMATIC v0.3.8 [96]. *De novo* assemblies

were performed with SPADES v3.11.1 [97]. Low coverage scaffolds were identified and

removed; scaffolds from the mitochondrial genome and rRNA clusters were identified and

masked. The assembly completeness was assessed based on the sordariomyceta_odb9 lineage

dataset using BUSCO v.3 [98].

Read mapping and SNP calling

The RAD-Seq reads were inspected with FASTQC v0.11.7 (Babraham Bioinformatics,

Cambridge, UK) to confirm quality and were demultiplexed and quality filtered with the

process_radtags program in STACKS v.2.10 [99]. The demultiplexed reads were trimmed using

TRIMMOMATIC v.0.3.8 [96] using the parameters ('LEADING:15 TRAILING:15

SLIDINGWINDOW:4:15 MINLEN:36') and aligned to the *C. graminicola* reference genome

M1.001 (NCBI accession number SAMN02953757) using BWA v.0.7.8 [100]. Genetic

variants were identified with FREEBAYES v.0.9.21 [102]. The WGS reads were quality

trimmed with TRIMMOMATIC v.0.3.8 and aligned to the reference genome with SPEEDSEQ

v.0.1.2 [101], which uses BWA-MEM for read mapping and FREEBAYES for variant

identification. The vcf files were merged and filtered using VCFTOOLS [103] to remove

nonbiallelic SNPs do not present in all isolates, sites with quality scores lower than 30, and

minor allele frequency (MAF) lower than 3%.

Population structure and genetic diversity

We defined a population as a group of individuals of the same species that occupy that

share the same area and colonize the same type of host simultaneously, then sharing a

common evolutionary history [63]. Analysis of molecular variance (AMOVA) was

performed with the hypothesis of no genetic differentiation using the package POPPR v2.8.6

[106]. Weir and Cockerham's genetic differentiation index (F_{ST}) was calculated with the R

package HIERFSTAT using the WC84 method [107]. We used three clustering methods to

examine the C. *graminicola* population structure. First, we used a principal component

analysis (PCA), with *a priori* geographical knowledge from sampling (by continent). This

analysis was conducted in the R environment with the package ADEGENET v2.0.1 [108] using

the DUDI.PCA function. Second, we performed a discriminate analysis of principal

components (DAPC) with R package ADEGENET v2.0.1 [108]. The FIND.CLUSTER function

was used to identify the most likely grouping in the data based on the Bayesian information

criterion (BIC) calculated for 1 to 10 clusters. Third, to confirm the pattern of population

subdivision inferred by the PCA and DAPC we used the fast maximum-likelihood genetic

clustering method SNAPCLUST [109]. The function SNAPCLUST.CHOOSEK was applied as a

prior step to determine the optimal number of clusters (K), based on the Bayesian information

criterion (BIC). Additionally, we implemented a neighbor network using the software

SPLITSTREE V4 [110] to visualize phylogenetic signals, indicated by the presence of

reticulations in the network, from a distance matrix with default parameters.

The number of multilocus genotypes (MLGs) was calculated using the R package

POPPR v2.8.6 [106]. We used the MGL.FILTER function to determine the true number of MLGs

from a genetic distance obtained by DISS.DIST function using the threshold determined in the

cutoff_predictor tool. To visualize relationships among genotypes a minimum spanning

network (MSN) was generated using the *msn* function in the R package POPPR. Genetic

diversity was estimated by comparison of the number of SNP variations by counting the

number of segregating sites between individuals, using the FIND.VARIATIONS tool in

GENEIOUS [111].

Linkage disequilibrium

The coefficient of linkage disequilibrium (r^2) [113] was calculated using VCFTOOLS

[103]. The rate of decay in linkage disequilibrium with physical distance was calculated for

all pairs of SNPs less than 100 kb apart using the option --hap-r2 and excluding sites with

minor allele frequencies below 5%. We also used the pairwise homoplasy index (PHI) test

implemented in SPLITSTREE V4 [110] to test the null hypothesis of clonality.

Investigation of putative sex-related genes

To identify orthologues of sex-related genes in the *C. graminicola* genomes, BLASTp

searches (e-value cutoff 1e-3) were performed using for the query a set of 96 gene with a role

in sexual reproduction in fungal model systems [114,115]. All nucleotide sequences were

checked manually to confirm no interruption in the translation frame. For this analysis, we

used the 12 *C. graminicola* assemblies generated in this study (see Supplementary Table S2

for the summary assemblies statistics), plus seven *C. graminicola* genome assemblies used by

Rech et al. [92] and genome assemblies from nine other fungal species: *Colletotrichum*

fiorinae ([116]; Genome accession: JARH00000000.1), *Colletotrichum fructicola* ([117];

Genome accession: ANPB00000000.1), *Colletotrichum gloeosporioides* ([118]; Genome

accession: AMYD00000000.1), *Colletotrichum higginsianum* ([119]; Genome accession:

LTAN00000000.1), *Colletotrichum nymphaeae* ([120]; Genome accession:

JEMN00000000.1)*, Colletotrichum salicis* ([120]; Genome accession: JFFI00000000.1),

Colletotrichum simmondsii ([120]; Genome accession: JFBX00000000.1), *Colletotrichum*

sublineola ([121]; Genome accession: JMSE00000000.1), *Colletotrichum orbiculare* ([117];

Genome accession: AMCV00000000.1), *Verticillium dahliae* ([122]; Genome accession:

ABJE00000000.1),

Pathogenicity assays

Pathogenicity characterization was performed by *in vivo* leaf blight assays using a

derivative of the susceptible maize inbred line Mo940 [123]. Plants were cultured in a

greenhouse for two weeks (V3 developmental stage), and inoculation via *C. graminicola*

spore suspensions were performed according to Vargas et al. [124]. Virulence was quantified

by measuring the area of necrotic lesions on leaves 4 days after inoculation. Symptomatic

leaves were excised from the plants, scanned with a flat-bed scanner, and quantified using the

imaging processing software Paint.NET v.4.2.16 (dotPDN LCC) [54], with three biological

replicates per isolate. For the comparison of differences in virulence among isolates, Analysis

of Variance (ANOVA) with Bonferroni and Sidak corrections for pairwise comparisons was

performed. Seventeen isolates were assessed, divided into batches because we could not

manage the whole set simultaneously in the laboratory. Given the possibility of a batch effect,

one of the strains (M1.001) was analyzed in all the batches to estimate the possible systematic

differences among them. Once all the isolates were analyzed, we compared the isolate

repeated in the different batches using ANOVA and obtained highly significant differences (p

< 0.0001), indicating a clear batch effect. Therefore, to eliminate the systematic differences

among batches, we standardized each strain using the mean and the standard deviation of the

control of its batch.

Results

Population genetic analyses

A total of 5811 biallelic variants were detected among 108 *C. graminicola* isolates

collected from 14 countries, which were used for population genetic analyses (Fig.1;

Supplementary Table S1). Analysis of SNP density showed an average of 1.4 SNPs per 10 kb

across contigs. Information about reads and coverage per individual are listed in the

supplementary table S1.

Clustering methods revealed population subdivision, supporting the existence of three

genetic groups, with isolates grouped by their continent of origin corresponding to the

isolates from South America, Europe, and North America, henceforth referred to as Brazilian

(BR), European (EU), and North American (NAM) clades (Fig. 2). The PCA and DAPC

grouped individuals according to their geographic origin (Fig. 2a, Supplementary Fig. S1).

Nevertheless, the isolates LAR318, CBS113173, and MAFF511343, from Zimbabwe,

Nigeria, and Japan respectively, obtained from culture collections did not group with the

main three genetic clusters. We do not discard the possibility of their geographic origins may

not be correctly reported in the collections. The SNAPCLUST function based on the BIC

criterion supported the clustering pattern by showing the optimum value of $K = 3$

clusters (Supplementary Fig. S2). The neighbor-net network generated by SPLITSTREE

identified three groups in the data (Fig. 3). Clusters in the network were connected by

extensive reticulated branches (looping) and long branches among samples, indicating a large

evolutionary distance separating individuals. AMOVA performed between *C. graminicola*

clusters showed that 65.5% of the total genetic variance was within clusters and 34.5% was

among them. Pairwise F_{ST} values between clusters were significant and revealed strong

differentiation, ranging from 0.26 to 0.42 (Supplementary Table S3).

Among the 108 isolates, we detected 76 MLGs unique and 15 MLGs shared between

at least two isolates. Most MLGs included isolates from the EU clade (13 MLGs in the total)

and shared between isolates from the same clade, and country (Supplementary Table S4).

However, one MLG encompassed isolates from France and Switzerland (F-40300-3 and SW-

8046-3), and another with isolates from Argentina and Croatia (ARG-2700-2 and CRO-I-35).

Two isolates from culture collections (M1.001 and LAR318) from USA and Nigeria were

also shared. Such genotypes shared were almost identical genetically, with few SNPs

between samples (Supplementary Table S4). The minimum spanning network (MSN) showed

a similar relationship among the aforementioned genotypes. The European cluster displayed

an inner position with MLGs closer genetically, while Brazilian and North American clusters

were placed on the edges of the network, likely due to their separate evolutionary history.

(a)   (b)

345   **Figure 2.** Population subdivision of *Colletotrichum graminicola*. (A) Principal components

346   analysis (PCA). Isolates are colored according to their continent of origin. (B) Minimum

347   spanning network (MSN) showing the multilocus genotypes (MLGs) based on Nei's genetic

348   distance. The distance between nodes represents the genetic distance between MLGs.

350   To estimate genetic diversity, we computed the number of segregating sites within

351   genetic groups. Genetic diversity was twice as high in NAM compared to BR and EU clades

352   (Table 1). Based on all the SNPs from the full dataset, NAM showed almost all sites

353   segregating when considering all isolates belonging to this clade. In contrast, isolates from

354   the EU clade showed fewer segregating SNPs and consequently lower genetic diversity.

Table 1. Single nucleotide polymorphism (SNP) variation within *Colletotrichum graminicola*

clusters.

Cluster	SNPs within clusters
BR	2484
EU	2077
NAM	5264

*Standard deviation in parentheses.

Figure 3. Neighbor-Net network showing relationships between *Colletotrichum graminicola*

isolates.

Footprints of recombination

We assessed the genomic impact of recombination by analyzing patterns of linkage

disequilibrium. For this, we computed LD on the genetic clusters separately to avoid

combining evolutionarily distinct populations, which can lead to an overestimate of the

linkage equilibrium [69]. For all genetic clusters, the LD decayed quickly with physical

distance, reaching 50% of its maximum value in less than 1kb, suggesting a random

association of alleles at different loci, likely caused by recombination events. We also used

the pairwise homoplasy index (PHI) test implemented in SPLITSTREE to test the null

hypothesis of no recombination, which rejected the null hypothesis for all three genetic

populations ($p < 0.001$).

To test the hypothesis that *C. graminicola* sex and meiosis-associated genes are being

maintained, we explore the presence of 96 genes previously identified as having roles in

sexual reproduction in Ascomycetes (Supplementary Table S5). We observed the presence of

94 genes out of the 96 sex-related genes in three *C. graminicola* genomes (M1.001,

LARS318 and CBS113173), while all the other strains were missing three or more genes. A

summary of the 15 genes showing the highest variability in presence/absence in the *C.*

graminicola genomes is shown in Table 2.

Figure 4. Linkage disequilibrium (LD) decay plots of three genetic populations of

Colletotrichum graminicola (European, Brazilian, and North America). LD was calculated

for all pairs of SNPs less than 100 kb distance apart.

Table 2. *Colletotrichum graminicola* orthologs of *Fusarium graminearum, Neurospora*

crassa, Podospora anserina, Saccharomyces cerevisiae, and *Schizosaccharomyces pombe*

sex-related genes. Fifteen genes showing the highest variability in terms of present/absence in

the *C. graminicola* genomes. Genes absent are marked with an x.

Gene[1]	Gene annotation / putative function	MI.00I	DMSZ63127	LARS318	CBS11517/5	NRRL47511	NRRL13649	CKO-I-35	F40500-I	CKI057/0-50	F64550-I	SW8046-2	CBS252.59	F64550-2	SW8046-I	P7565-07212	CA-CHA1-I	CKO-I-41	CA-N8H	ARG-2/00-2
GLRG_06331T0	Meiotic recombination protein REC4							x	x	x	x			x	x			x	x	
GLRG_03069T0	Splicing factor 3B subunit 4						x	x	x	x	x	x	x							
GLRG_06080T0	Strand exchange protein RAD55p					x	x	x	x	x	x	x	x	x	x	x		x		x
GLRG_04547T0	Cell cycle checkpoint protein RAD17													x						
GLRG_06289T0	Histone H2A.Z							x		x			x							
GLRG_10150T0	Casein kinase I							x	x	x		x	x	x			x			
GLRG_02257T0	Nucleotide excision repair protein RAD23									x										
GLRG_02375T0	DNA topoisomerase							x	x	x	x	x	x	x						

Gene	Description																		
GLRG_02369T0	Ku80 protein							x	x	x	x		x						
GLRG_02331T0	Chromosome segregation protein SUDA							x											
GLRG_00702T0	WD repeat-containing protein slp1				x														
GLRG_00302T0	Histone-lysine N-methyltransferase								x										
GLRG_03812T0	Peptide pheromone maturation							x											
GLRG_00900T0	Histone deacetylase HOS2	x																	
GLRG_09043T0	G protein-coupled receptor gpcr1							x	x	x		x							

[1]Genes from *C. graminicola* locus tag from reference genome strain M1.001 (O'Connell et al. 2012).

Pathogenicity

Seventeen isolates representative of the subclades identified in the phylogeny inference and showing diversity of colony morphology and pigmentation were chosen for pathogenic characterization (Supplementary Fig. S3). Considering the continental origin, the following isolates were chosen, America: M1.001, A-52339-1, ARG-2700-5, BR-85925-1, CA-NOP-2CO, CA-CHAT-1, and JAB2; Africa: CBS113173, LARS318; Europe: CBS252.59, CR-42230-1, CRO-I-41, F-64330-2, P-7565-072-3, SI-9253-1, and SW-8046-2; Asia: MAFF511343. All isolates, except two - CBS252.59 and MAFF511343, were pathogenic on the maize cultivar evaluated producing necrotic lesions at the inoculation points by 4 days post inoculation (dpi) (Fig. 5). Isolates CBS252.59 and MAFF511343 also failed to produce necrotic lesions at 6 dpi (not shown). A high variation in virulence among the isolates was found compared to the strain M1.001. We compared all the strains to the virulence of M1.001 using a classical analysis of variance $p < 0.0001$). Isolates ARG-2700-5, P-7565-072-3, SI-9253-1, CA-CHAT-1, and CR-42230-1 had significantly higher virulence

levels, while isolate JAB2 was significantly less virulent. Population structure was not

correlated with virulence.

Figure 5. Pathogenic characterization of *Colletotrichum graminicola* isolates. (a) Graphic

representation of pairwise comparisons of 14 isolates in order of virulence (relative to strain

M1.001 used as control strain) in ascending order. The isolates CBS252.59 and

MAFF511343 were non-pathogenic to the maize cultivar evaluated and were not included in

the figure. Isolates marked with an asterisk are significantly different from isolate M1.001.

(b) Necrotic lesions on maize leaves at 4 days after inoculation with spore suspensions of the

isolates.

Discussion

We employed a population genomics approach to understand the genetic diversity of

C. graminicola isolates infecting maize sampled from fourteen countries. Using SNP markers

generated through RAD-seq and WGS, we investigated the distribution of genetic variation

considering our samples as a global population. We found populational differentiation at a

worldwide scale, with three genetic groups delimited by continental origin, compatible with

short dispersal of the pathogen, and geographic subdivision. However, we observed migration

intra and intercontinental migration, likely associated with the movement of contaminated

seeds. The genetic clusters, corresponding to Brazilian, European, and North American

clades, showed distinct levels of genetic diversity that could reflect different evolutionary

histories. Evidence of genetic recombination was detected from analysis of linkage

disequilibrium and the PHI test for clonality. Investigation of sex-related genes revealed gene

losses in most isolates. We show evidence that even if rare (possibly due to losses of sex and

meiosis-associated genes) *C. graminicola* can undergo sexual recombination based on lab

assays and genomic analyses. We present the possible evolutionary origins of these genetic

groups and the impact of recombination on genetic structure and the potential of maize

anthracnose becoming a much more significant disease in the future.

Clustering analyses based on single-nucleotide polymorphisms showed strong

population differentiation, grouping of the isolates according to their country of origin. The

European clade showed less genetic variation, a higher number of repeated MLGs, and lower

pairwise differentiation among clusters, suggesting a larger clonal component as compared to

the other clades. Interestingly, all genotypes shared were almost identical genetically,

suggesting that gene flow can be dependent on the clonal isolates generated by asexual

reproduction. Most MLGs were shared between isolates from the same cluster and country,

however, one MLG encompassed isolates from France and Switzerland (more than 1000 km

apart) and another one with isolates from Argentina and Croatia. In our cluster analyses, the

European clade encompassed all isolates from Argentina. Together, these results indicate

intra and intercontinental migration of the pathogen. We argue a likely European origin of the

isolates from Argentina due to the movement of maize germplasm into South America. It is

common for maize breeders from North America and Europe to use South America

(including Argentina) fields for advancing breeding programs during the northern hemisphere

winter. As Argentina is a maize exporter, breeding programs can represent a possible source

of pathogen importation and consequently a source of pathogen introductions into new areas,

for instance in Europe or elsewhere. This result is particularly important from the

epidemiological point of view, with direct implications on disease management since

migration can cause the movement of more virulent and/or fungicide resistant genotypes.

The predominant form of dissemination of *C. graminicola* is thought to be by

contaminated seeds [64–66]. Fungi can naturally be spread via the dispersal of both sexual

and asexual spores, caused mainly by the transport of infected plant material [11,59,67].

Dispersal of asexual conidia of *C. graminicola* is strongly dependent on water splash and

blowing raindrops, responsible for short-distance dispersal [22]. Although airborne dispersal

is not typically documented, ascospore dispersal is associated with long-distance

dissemination since such spores can be ejected up and dispersed by air currents reaching

longer distances [125,126]. If sexual reproduction occurs, even sporadically in this species, it

can contribute to long distance dissemination, leading to a broader pathogen distribution.

The North American clade encompassed isolates from the United States, Canada, and

one strain from Africa obtained from a culture collection. This cluster is more genetically

diverse and seems to have a separate evolutionary history. Although the first report in the

literature of a *Colletotrichum* species infecting maize came from Europe (Italy, 1852) we

argue that the North American cluster may be the more ancient lineage (or ancestral lineage)

since maize anthracnose was described for the first time in 1914 in the United States [127].

Considering that Mesoamerica is the maize center of origin, most probably Mexico,

[128,129] and the center of origin of pathogenic species usually correspond to the center of

origin of their host [130], given the proximity to the USA, this lineage could be the founder

of the outbreak during the 1970s and may have served as the source for introductions in other

continents. The highest genetic diversity found in this lineage also corroborates this

hypothesis, since the center of origin of a pathogen is likely to consist of populations with

more genetic variability than recently founded populations [131]. *C. graminicola* may have

evolved together with the host in Central America, and due to the expansion of maize

production in the USA green belt the pathogen had an opportunity to spread across the vast

maize growing areas. Alternatively, our sampling, which consisted of a large range of years

(1975 to 2016), might have overestimated the genetic diversity of this group. A more

contemporary sampling can reveal the current genetic variation of populations from North

America.

In South America, isolates from Argentina and Brazil were grouped into separate

clades, suggesting independent introductions. In Brazil, the first disease report was in 1965 in

the São Paulo state (Campina's city) [132]. In the country, maize is planted in two distinct

growing seasons, the first crop planted in the summer and the second crop during the winter.

This practice contributes to increasing primary inoculum in the area. A significant reduction

in maize yield in tested hybrids was reported by Cota et al. [43] which was associated, among

other factors, with a higher pathogen aggressiveness and genetic differentiation in

populations during the crop cycle. This could indicate changes in the population size and

their genetic composition that may have contributed to a higher disease virulence in the

country.

We found molecular evidence of genetic recombination for the three *C. graminicola*

populations detected in this study. The rapid decay of linkage disequilibrium and extensive

reticulations in the neighbor network is consistent with a possible history of recombination.

Furthermore, the PHI test for clonality detected recombination within each population ($P =$

0.0). These results corroborate a previous study that found evidence of recombination using

the PHI test at the chromosome level [92]. Similar findings of low and fast LDecay are

reported in other fungal plant pathogens which suggests the occurrence of genetic

mechanisms responsible for creating such patterns in genomes [82,133,134]. In addition, the

signature of repeat-induced point mutation (RIP), a genome defense mechanism that occurs

during sexual reproduction, [135,136] was detected in *C. graminicola* genomes [137,138].

RIP mutations may indicate an ancestral state of sexual reproduction or that meiosis

cryptically occurs in nature [71,139].

Despite *C. graminicola* being reported as a homothallic species, strains may exhibit

homothallism and heterothallism, but this phenomenon is not yet fully understood [22].

Investigation of fertility through crosses of the strains M1.001 and M5.001 produced an

offspring exclusively heterothallic, which led Vaillancourt et al. [79] to hypothesize that

unbalanced heterothallism could be involved in the mating system. Our investigation of sex-

related genes revealed gene losses in most isolates and in those species known to be self-

fertile such as *C. salicis,* and in those known to be self-sterile and cross-fertile like *C. fiorinae*

[120]. Interestingly the only two genes missing in all *C. graminicola* genomes analyzed were

ppg1 and *pre2,* which form a pheromone receptor pair. The importance of this signaling pair

varies among homothallic ascomycetes. However, none of the deletion mutants obtained in

Sordaria macrospora, *Aspergillus nidulans* and *Gibberella zeae* completely lost the

capability to sexually cross [115].

Considering this evidence and the high completeness of the assemblies we conclude

that at least three of the *C. graminicola* isolates maintained all the genes required to undergo

sexual reproduction. This finding also suggests that some lineages have lost the ability for

sexual reproduction or that not all sex-related genes reported in other species are required in

Colletotrichum spp. [22]. Keeping in mind the presence of homothallic and heterothallic

strains in this species, the lack of genes required to sexually cross in one of the strains could

mean failure in sexual reproduction. In this scenario, sexual reproduction would become rare

(but not inconceivable) due to the low chance of two strains having the full set of genes

encounter themselves and mating.

The reproductive biology of the genus *Colletotrichum* is mixed, and the sexual state

of several species is unknown although several species are known to have the capacity for

sexual reproduction. The sexual capability of *C. graminicola* is well-documented [78,140].

We demonstrated experimentally the sexual capability of *C. graminicola* isolates mating in

lab conditions, as described by Vaillancourt et al. [79] (see Supplementary Fig. S4 for

details), although the teleomorph has not yet been found in field conditions. Although the

absence of the sexual state in nature could cause less recombination, population genetic

studies show that several *Colletotrichum* species known as exclusively asexual, exhibit

evidence of genetic recombination [83,141,142]. Some studies have described alternative

mechanisms, such as the parasexual cycle as responsible for creating genetic recombination

in *Colletotrichum* species [79,84,143–145].

Species with the potential for genetic exchange are more likely to overcome control

measures employed since recombination can result in new combinations of alleles that

increase diversity and speed up adaptation at the field scale [63,146]. This study gathered

evidence that *C. graminicola* populations may have been affected by genetic recombination,

but we also showed evidence that even if rare (possibly due to frequent losses of sex and

meiosis-associated genes) *C. graminicola* can undergo sexual recombination. Fine-scale

investigations at the genome level can be employed to understand the real impact of genetic

recombination on genetic diversity and adaptation, for instance, if genetic exchange enhances

virulence factors (also called fitness-related traits) in these populations, as suggested for other

fungal species [147–151]. That can be particularly important in the agricultural pathosystem

given that crop pathogens are constantly facing a homogeneous environment that imposes

strong directional selection that leads to the evolution of more virulent and drug-resistant

pathogen populations [130,152].

We did not find a correlation between virulence and population structure; however,

we observed a substantial difference in the spectrum of virulence. Several isolates were more

aggressive than the strain M1.001 which was originally collected from North America in

1972 during the US. outbreak [153]. This finding is particularly important because this strain

represents the genetic diversity existent in the outbreak and strains more aggressive are

currently present in nature, which may be a risk to the development of new, even more,

destructive outbreaks. The pathogenicity assay was performed in one maize line (Mo940)

notably susceptible to anthracnose. The use of distinct maize genotypes can result in different

responses to fungal infection, as already reported in previous studies, and it should be

considered in further pathogenic characterizations [42,154,155]. Another significant aspect is

that resistance to maize anthracnose is inherited quantitatively, i.e., host resistance is

controlled by several genes with additive genetic effects [156–158]. This fact may correlate

with the range of virulence found in this study. The three genetic clades of *C. graminicola*

detected may have distinct evolutionary origins, hence encompassing a diverse combination

of virulence genes, which could cause different defense responses from the host. We also

highlighted that the large variability in virulence should be considered for the screening of

resistant maize varieties. Population studies can support breeding programs by investigations

of intraspecific variability of *C. graminicola* populations since the effectiveness of genetic

resistance depends on knowledge of the genetic structure of pathogens.

Conclusions

Our study represents the first investigation of the population structure of *C. graminicola*

infecting maize. The three main clades (North America, Brazil, and Europe) have different

levels of genetic diversity suggesting that each has a unique evolutionary history. We also

found evidence of recent migration of isolates between Europe and Argentina, possibly due to

the importation of infected plant material. Although the sexual state of *C. graminicola* has

only been reported in lab conditions, we show strong evidence that genetic recombination,

whether by sexual reproduction, or by alternative mechanisms (such as the parasexual cycle)

plays an important role in *C. graminicola* population structure. In contrast to the traditional

view of *C. graminicola* being mainly asexual, we conclude that genetic recombination is

frequent. Fine-scale investigations at the genome level can be employed to understand the

real impact of recombination and gene flow on the genetic diversity and adaptation of this

species, with a direct effect on control measures and breeding programs aiming at resistance.

Finally, we warn of the danger that maize anthracnose has the potential to become much

more significant in the future, mainly due to highly aggressive isolates currently present in

maize fields and the geographic range expansion of the pathogen, in addition to increased

susceptibility of agroecosystems accentuated by ongoing climate changes.

Additional Files

Supplementary Fig. S1. A) Bayesian information criteria (BIC) indicating the most probable

number of genetic groups by discriminant analysis of principal components analysis (DAPC).

B) Scatterplot from DAPC of *Colletotrichum graminicola* isolates assigned with *a priori*

geographical information (by continent).

Supplementary Fig. S2. Bayesian information criteria (BIC) indicating the most probable

number of genetic groups using the SNAPCLUST function.

Supplementary Fig. S3. Colony morphology of *Colletotrichum graminicola* showing

phenotypic diversity among isolates. The colonies were cultivated on PDA medium, at 23ºC

under continuous light for 6 days.

Supplementary Fig. S4. Sexual structures of *Colletotrichum graminicola* produced by

crossing from strains M1.001 (CB130836) and M5.001 (CBS 130839). The cross-inoculation

experiment was performed under laboratory conditions as described by Vaillancourt and

Hanau 1991. (a) Agar plugs containing strains growing on autoclaved leaves of maize. (b)

Mature perithecia, picture taken with Olympus SZX9 stereomicroscope outfitted with an

Olympus DP70 camera. c) Ascus and ascospores. d) Ascospores.

Supplementary Table S1. Isolates of *Colletotrichum graminicola* used in this study.

Supplementary Table S2. Summary statistics of *Colletotrichum graminicola* genomes

generated in this study.

Supplementary Table S3. Measure of pairwise genetic differentiation between genetic

clusters of *Colletotrichum graminicola.*

Supplementary Table S4. Multilocus genotypes (MLGs) among *Colletotrichum*

graminicola isolates.

Supplementary Table S5. *Colletotrichum graminicola* orthologs of *Neurospora crassa,*

Saccharomyces cerevisiae, Podospora anserina, Fusarium graminearum, and

Schizosaccharomyces pombe sex-related genes.

Data Availability

The vcf file generated in this study is provided as Additional file 1. The genome assemblies

were submitted to GenBank, and their accession numbers are in Supplementary Table S2.

Abbreviations

ANOVA: Analysis of Variance; BIC: Bayesian information criterion; BLAST: Basic Local

Alignment Search Tool; BR: Brazilian clade; CTAB: cetyltrimethylammonium-bromide; EU:

European clade; FST: Genetic differentiation index; GTR: general Time-Reversible model of

nucleotide substitution; ITS: internal transcribed spacer; MAF: minor allele frequency;

MLGs: multilocus genotypes; NAM: North American clade; MSN: minimum spanning

network; NCBI: National Center for Biotechnology Information; PDA: potato dextrose agar;

PDB: potato dextrose broth; PCA: principal component analysis; PHI: pairwise homoplasy

index; RAD-Seq: restriction site-associated DNA sequencing; pairwise homoplasy index;

SNPs: single-nucleotide polymorphisms; WGS: whole-genome sequencing.

Competing Interests

The authors declare that they have no competing interests.

Funding

This research was supported by grants AGL2015-66362-R and RTI2018-093611-B-I00 from

the Ministerio de Ciencia e Innovación (MICINN) of Spain AEI/10.13039/501100011033

and the grant SA165U13 from the Junta de Castilla y Léon. F.R. was supported by the EU

NextGeneration project EU/PRTR (FJC2020-043351-I). S.A.S. was supported by

POSDOC2019_USAL48 from the University of Salamanca. R.B. was supported by the

postdoctoral program of USAL (Programme II). S.B. was supported by a fellowship program

from the regional government of Castilla y León and Ph.D. fellowship to F.B.C.F. BES-2016-

078373 from the Ministerio de Educación of Spain. W.B was supported by a productivity

fellowship from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq

307855/2019-8). Genomes sequencing were funded by the UNC Microbiome Core, which is

funded in part by the Center for Gastrointestinal Biology and Disease (CGIBD P30

DK034987) and the UNC Nutrition Obesity Research Center (NORC P30 DK056350).

P.D.E. was partially supported by the USDA National Institute of Food and Federal

Appropriations under Project PEN04660 and Accession Number 1016474.

Competing Interest

663   The authors declare that they have no competing interests.

Authors' Contributions

S.A.S. and M.R.T conceived the study and designed the project; S.A.S performed fungal

isolation and DNA extraction; M.R.T., F.R., R.B., S.B. J.L.V.V., S.A.S, and F.B.C.F.

performed analyses; M.R.T., F.R., R.B, and S.A.S. wrote the manuscript; F.B.C.F. and S.A.S.

performed the pathogenic characterization; M.A.A.P., M.M.W. and J.A.C. performed genome

sequencing; I.B., W.B., V.O., J.B., A.T., J.S.D., P.D.E., P.R., T.J.Z., J.H., G.M. collected the

samples. All authors reviewed the manuscript.

Acknowledgments

674   The authors thank Maria Elena Bayón Sánchez for her assistance in the isolation of samples

675   and pathogenicity tests, Lisa J. Vaillancourt, and the Genetic Resources Center (NARO) for

676   providing some isolates.

References

1. Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. The global

burden of pathogens and pests on major food crops. *Nat Ecol Evol*. Nature Publishing Group;

2019; doi: 10.1038/s41559-018-0793-y.

2. Fones HN, Bebber DP, Chaloner TM, Kay WT, Steinberg G, Gurr SJ. Threats to global

food security from emerging fungal and oomycete crop pathogens. *Nat Food*. Springer US;

2020; doi: 10.1038/s43016-020-0075-0.

3. Strange RN, Scott PR. Plant Disease: A Threat to Global Food Security. *Annu Rev*

Phytopathol. 2005; doi: 10.1146/annurev.phyto.43.113004.133839.

4. Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, Stukenbrock EH, et al.. Threats posed

by the fungal kingdom to humans, wildlife, and agriculture. *MBio*. 2020; doi:

10.1128/mBio.00449-20.

5. Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, et al..

Emerging fungal threats to animal, plant and ecosystem health. *Nature*. Nature Publishing

Group; 2012; doi: 10.1038/nature10947.

6. Santini A, Ghelardini L, De Pace C, Desprez-Loustau ML, Capretti P, Chandelier A, et al..

Biogeographical patterns and determinants of invasion by forest pathogens in Europe. *New*

Phytol. 2013; doi: 10.1111/j.1469-8137.2012.04364.x.

7. Fones HN, Gurr SJ. NOXious gases and the unpredictability of emerging plant pathogens

under climate change. *BMC Biol*. BMC Biology; 2017; doi: 10.1186/s12915-017-0376-4.

8. Chaloner TM, Gurr SJ, Bebber DP. Plant pathogen infection risk tracks global crop yields

under climate change. *Nat Clim Chang*. Springer US; 2021; doi: 10.1038/s41558-021-01104-

8.

9. Chapman D, Purse B V., Roy HE, Bullock JM. Global trade networks determine the

distribution of invasive non-native species. *Glob Ecol Biogeogr*. 2017; doi:

10.1111/geb.12599.

10. Sikes BA, Bufford JL, Hulme PE, Cooper JA, Johnston PR, Duncan RP. Import volumes

and biosecurity interventions shape the arrival rate of fungal pathogens. *PLoS Biol*. 2018; doi:

10.1371/journal.pbio.2006025.

11. Brown JKM, Hovmøll MS. Aerial dispersal of pathogens on the global and continental

scales and its impact on plant disease. *Science (80-)*. 2002; doi: 10.1126/science.1072678.

12. Bebber DP. Range-Expanding Pests and Pathogens in a Warming World. *Annu Rev*

Phytopathol. 2015; doi: 10.1146/annurev-phyto-080614-120207.

13. Saunders DGO. Will yield gains be lost to disease? *Nat Clim Chang*. 2021; doi:

10.1038/s41558-021-01113-7.

14. Bebber DP, Ramotowski MAT, Gurr SJ. Crop pests and pathogens move polewards in a

warming world. *Nat Clim Chang*. Nature Publishing Group; 2013; doi:

10.1038/nclimate1990.

15. FAOSTAT. Food and Agriculture Organization Corporate Statistical Database. Accessed

10 Jan 2022.

16. Shiferaw B, Prasanna BM, Hellin J, Bänziger M. Crops that feed the world 6. Past

successes and future challenges to the role played by maize in global food security. *Food*

Secur. 2011; doi: 10.1007/s12571-011-0140-5.

17. Wu F, Guclu H. Global Maize Trade and Food Security: Implications from a Social

Network Model. *Risk Anal*. 2013; doi: 10.1111/risa.12064.

18. Nicoli A, Zambolim L, Costa RV da, Guimarães LJM, Lanza FE, Silva DD da, et al..

Identification of sources of resistance to anthracnose stalk rot in maize. *Ciência Rural*. 2016;

doi: 10.1590/0103-8478cr20151052.

19. Bergstrom GC, Nicholson RL. The biology of corn anthracnose: Knowledge to exploit for

improved management. *Plant Dis*. 1999; doi: 10.1094/PDIS.1999.83.7.596.

20. Munlvold G. Managing Anthracnose Stalk Rot: A moving Target. *Proc 56th Annu corn*

sorghum Res Conf. p. 196–206.

21. Sukno SA, García VM, Shaw BD, Thon MR. Root infection and systemic colonization of

maize by *Colletotrichum graminicola. Appl Environ Microbiol*. 2008; doi:

10.1128/AEM.01165-07.

22. Crouch JA, Beirn L.. Anthracnose of cereals and grasses. *Fungal Divers*. 39:19–442009;

23. Talhinhas P, Baroncelli R. *Colletotrichum* species and complexes: geographic

distribution, host range and conservation status. *Fungal Divers*. Springer Netherlands; 2021;

doi: 10.1007/s13225-021-00491-9.

24. Bhunjun CS, Phukhamsakda C, Jayawardena RS, Jeewon R, Promputtha I, Hyde KD.

Investigating species boundaries in *Colletotrichum. Fungal Divers*. Springer Netherlands;

2021; doi: 10.1007/s13225-021-00471-z.

25. Jayawardena RS, Bhunjun CS, Hyde KD, Gentekaki E, Itthayakorn P. *Colletotrichum*:

lifestyles, biology, morphospecies, species complexes and accepted species. *Mycosphere*.

2021; doi: 10.5943/mycosphere/12/1/7.

26. Ullstrup AJ, Troyer AF. A lethal leaf spot of maize. *Phytopathology*. 57:1282–31967;

27. Byrnes KJ, Carroll RB. Fungi Causing Stalk Rot of Conventional-Tillage and No-Tillage

Corn in Delaware. *Plant Dis*. 1986; doi: 10.1094/pd-70-238.

28. Hooker AL, White DG. Prevalence of Corn Stalk Rot Fungi Illinois. *Plant Dis Report*.

60:1032–41976;

29. Lipps PE. Survival of *Colletotrichum graminicola* in infested corn residues in Ohio. *Plant*

Dis. 67:102–41983;

30. Poneleit CG, Politis DJ, Wheeler H. Resistance to Corn Anthracnose. *Crop Sci*. 12:875–

61972;

31. Warren HL, Nicholson RL, Ullstrup AJ, Sharvelle EG. Observations of *Colletotrichum*

graminicola on sweet corn in Indiana. *Plant Dis Report*. 57:143–41973;

32. Leonard KJ, Thompson DL. Effects of Temperature and Host Maturity on Lesion

Development of *Colletotrichum graminicola* on Corn. Phytopathology.

33. Jirak-Peterson JC, Esker PD. Tillage, crop rotation, and hybrid effects on residue and

corn anthracnose occurrence in Wisconsin. *Plant Dis*. 2011; doi: 10.1094/PDIS-11-10-0837.

34. Farber C, Bennett JS, Dou T, Abugalyon Y, Humpal D, Sanchez L, et al.. Raman-Based

Diagnostics of Stalk Rot Disease of Maize Caused by *Colletotrichum graminicola*. *Front*

Plant Sci. 2021; doi: 10.3389/fpls.2021.722898.

35. Gorman Z, Christensen SA, Yan Y, He Y, Borrego E, Kolomiets M V.. Green leaf

volatiles and jasmonic acid enhance susceptibility to anthracnose diseases caused by

Colletotrichum graminicola in maize. *Mol Plant Pathol*. 2020; doi: 10.1111/mpp.12924.

36. Mueller DS, Wise KA, Sisson AJ, Allen TW, Bergstrom GC, Bosley DB, et al.. Corn

yield loss estimates due to diseases in the United States and Ontario, Canada from 2012 to

2015. *Plant Heal Prog*. 2016; doi: 10.1094/PHP-RS-16-0030.

37. Jackson-Ziems TA, Rees JM, Harveson RM. Common Stalk Rot Diseases of Corn. Bull.

Univ. Nebraska. p. 8.

38. : Centre for Agriculture and Bioscience International (CABI). https://cabi.org/. Accessed

13 Jan 2022.

39. Hein T: Anthracnose in corn: the eastern Canadian snapshot. TopCrop Manag.

https://www.topcropmanager.com/anthracnose-in-corn-the-eastern-canadian-snapshot-4393/

(2009).

40. Mueller DS, Wise KA, Sisson AJ, Allen TW, Bergstrom GC, Bissonnette KM, et al..

Corn yield loss estimates due to diseases in the United States and Ontario, Canada, from 2016

to 2019. *Plant Heal Prog*. 2020; doi: 10.1094/PHP-05-20-0038-RS.

41. Wheeler H, Politis DJ, Poneleit CG. Pathogenicity, Host Range, and Distribution of

Colletotrichum graminicola on Corn. *Phytopathology*. 64:293–61974;

42. Costa R V., Cota L V., Silva DD, Parreira DF, Casela CR, Landau EC, et al.. Races of

Colletotrichum graminicola pathogenic to maize in Brazil. *Crop Prot*. Elsevier Ltd; 2014;

doi: 10.1016/j.cropro.2013.10.005.

43. Cota L V., da Costa R V., Silva DD, Casela CR, Parreira DF. Quantification of Yield

Losses Due to Anthracnose Stalk Rot on Corn in Brazilian Conditions. *J Phytopathol*. 2012;

doi: 10.1111/jph.12008.

44. Costa RV, Silva DD, Cota LV, Parreira DF, Ferreira AS, Casela CR. Incidência de

Colletotrichum graminicola em colmos de genótipos de milho. *Summa Phytopathol*. 2010;

doi: 10.1590/s0100-54052010000200003.

45. Coêlho RMS, Silva HP, Brunelli KR, Camargo LEA. Controle genético da antracnose

foliar em milho. *Fitopatol Bras*. 2001; doi: 10.1590/s0100-41582001000300010.

46. De Rossi RL, Giménez Pecci MP, Guerra FA, Plaza MC, Brücher E, Guerra GD, et al..

Enfermedades Del Maíz De Siembra Tardía Causadas Por Hongos. *El mismo maíz, un nuevo*

desafío Compend Prim Congr maíz tardío. p. 1–14.

47. Eyhérabide G.. Bases para el Manejo del Cultivo de Maíz. Ed. INTA Pergam. y Minist.

Agric. Ganad.

48. Gatica SM, Gally M, Carmona M, Ramos AM, Ferreyra LI. Diferenciación genética de

aislamientos de *Colletotrichum graminicola* de la región pampeana mediante marcadores

ISSR. III Congr. Argentino Fitopatol.

49. Tiffany L, Gilman J. Species of *Colletotrichum* from Legumes. *Mycologia*. 46:52–

751954.

50. Böning K, Wallner F. Fußkrankheit und andere Schädigungen an Mais durch

Colletotrichum graminicolum (Ces.) Wilson. *Phytopathol Zeitschrift*. 9:99–1101936;

51. Messiaen CM, Lapon R, Molot P. Root necroses, stalle rots, and parasitic lodging of

maize. *Annu Epiphyte*. 10:441–741960.

52. Sychrova E, Zakopal J, Kopa K. Symptome und Erreger der Fusskrankheiten des

Getreides. *Ustred Vyzk Ust Rostl Vyroby, Ved Pr*. :31–6 1974.

53. Palaversic B, Jukic M, Buhinicek I, Vragolovic A, Kozic Z. Breeding maize for resistance

to stalk anthracnose. *Maydica*. 54:229–322009.

54. Sanz-Martín JM, Pacheco-Arjona JR, Bello-Rico V, Vargas WA, Monod M, Díaz-

Mínguez JM, et al.. A highly conserved metalloprotease effector enhances virulence in the

maize anthracnose fungus *Colletotrichum graminicola*. *Mol Plant Pathol*. 2016; doi:

10.1111/mpp.12347.

55. Sukno SA, Sanz-Martín JM, González-Fuente M, Hiltbrunner J, Thon MR. First report of

anthracnose stalk rot of maize caused by *Colletotrichum graminicola* in Switzerland. *Plant*

Dis. 2014; doi: https://doi.org/10.1094/PDIS-09-13-0912-PDN.

56. Winter W, Menzi M. Inbred maize lines: infection and symptoms of important fungal

diseases. *Landwirtschaft Schweiz*. 4:599–6061991.

57. Cuevas-Fernández FB, Robledo-Briones AM, Baroncelli R, Trkulja V, Thon MR,

Buhinicek I, et al.. First report of *Colletotrichum graminicola* causing maize anthracnose in

Bosnia and Herzegovina. *Plant Dis*. 2019; doi: 10.1094/PDIS-06-19-1224-PDN.

58. Duan CX, Guo C, Yang ZH, Sun SL, Zhu ZD, Wang XM. First report of anthracnose leaf

blight of maize caused by *Colletotrichum graminicola* in China. *Plant Dis*. 2019; doi:

10.1094/PDIS-12-18-2140-PDN.

59. Gladieux P, Feurtey A, Hood ME, Snirc A, Clavel J, Dutech C, et al.. The population

biology of fungal invasions. *Mol Ecol*. 2015; doi: 10.1111/mec.13028.

60. Desprez-Loustau ML, Robin C, Buée M, Courtecuisse R, Garbaye J, Suffert F, et al.. The

fungal dimension of biological invasions. *Trends Ecol Evol*. 2007; doi:

10.1016/j.tree.2007.04.005.

61. Gottwald T, Luo W, Posny D, Riley T, Louws F. A probabilistic census-travel model to

predict introduction sites of exotic plant, animal and human pathogens. *Philos Trans R Soc B*.

2019; doi: 10.1098/rstb.2018.0260.

62. Barrett LG, Thrall PH, Burdon JJ, Linde CC. Life history determines genetic structure

and evolutionary potential of host-parasite interactions. *Trends Ecol Evol*. 2008; doi:

10.1016/j.tree.2008.06.017.

63. Milgroom MG. Population biology of plant pathogens: genetics, ecology, and evolution.

Am. Phytopathol. Soc. St. Paul, MN.

64. Tahir R, Ahmad U. SEED-BORNE FUNGAL DISEASES OF MAIZE (Zea mays L.): A

REVIEW Fazal. *J Agroteknologi dan Perkeb*. 4:43–602021;

65. Warren HL. Kernel Infection, Seedling Blight, and Wilt of Maize Caused by

Colletotrichum graminicola. *Phytopathology*. 1975; doi: 10.1094/phyto-65-620.

66. Somda I, Sanou J, Sanon P. Seed Born Infection of Farmer-Saved Maize Seeds by

Pathogenic Fungi and Their Transmission to Seedlings. *Plant Pathol J*. 7:98–1032008;

67. Aylor DE. Spread of plant disease on a continental scale: Role of aerial dispersal of

pathogens. *Ecology*. 2003; doi: 10.1890/01-0619.

68. Milgroom MG, Fry WE. Contributions of Population Genetics to Plant Disease

Epidemiology and Management. *Adv Bot Res*. 1997; doi: 10.1016/S0065-2296(08)60069-5.

69. Milgroom MG. Recombination and the Multilocus Structure of Fungal Populations. *Annu*

Rev Phytopathol. 34:457–771996;

70. Taylor JW, Hann-Soden C, Branco S, Sylvain I, Ellison CE. Clonal reproduction in fungi.

Proc Natl Acad Sci U S A. 2015; doi: 10.1073/pnas.1503159112.

71. da Silva LL, Moreno HLA, Correia HLN, Santana MF, De Queiroz MV. *Colletotrichum*:

species complexes, lifestyle, and peculiarities of some sources of genetic variability. *Appl*

Enviromental Microbiol. 2020; doi: 10.1007/s00253-020-10363-y.

72. Nieuwenhuis BPS, James TY. The frequency of sex in fungi. *Philos Trans R Soc B Biol*

Sci. 2016; doi: 10.1098/rstb.2015.0540.

73. Goddard MR, Godfray HCJ, Austin Bur. Sex increases the efficacy of natural selection in

experimental yeast populations. *Nature*. 2005; doi: 10.1029/2001jb001613.

74. de Vienne DM, Giraud T, Gouyon PH. Lineage Selection and the Maintenance of Sex.

PLoS One. 2013; doi: 10.1371/journal.pone.0066906.

75. Posada D, Crandall KA, Holmes EC. Recombination in Evolutionary Genomics. *Annu*

Rev Genet. 2002; doi: 10.1146/annurev.genet.36.040202.111115.

76. Croll D, Lendenmann MH, Stewart E, McDonald BA. The impact of recombination

hotspots on genome evolution of a fungal plant pathogen. *Genetics*. 2015; doi:

863 10.1534/genetics.115.180968.

864 77. Crouch J, O'Connell R, Gan P, Buiate E, Torres MF, Beirn L, et al.. The Genomics of

865 *Colletotrichum*. In: R. D, A. L-P, Kole C., editors. *Genomics Plant-Associated Fungi*

866 *Monocot Pathog*.

867 78. Politis DJ. The Identity and Perfect State of *Colletotrichum graminicola*. *Mycol Soc Am*.

868 67:56–621975;

869 79. Vaillancourt L, Wang J, Hanau R, Rollins J, Du M. Genetic Analysis of Cross Fertility

870 between Two Self-Sterile Strains of *Glomerella graminicola*. *Mycologia*. 2000; doi:

871 10.2307/3761501.

872 80. Billiard S, López-Villavicencio M, Hood ME, Giraud T. Sex, outcrossing and mating

873 types: Unsolved questions in fungi and beyond. *J Evol Biol*. 2012; doi: 10.1111/j.1420-

874 9101.2012.02495.x.

875 81. Wilson AM, Lelwala R V., Taylor PWJ, Wingfield MJ, Wingfield BD. Unique patterns

876 of mating pheromone presence and absence could result in the ambiguous sexual behaviors of

877 *Colletotrichum* species. *G3 Genes, Genomes, Genet*. 2021; doi: 10.1093/g3journal/jkab187.

878 82. Rogério F, Van Oosterhout C, Ciampi☐Guillardi M, Correr FH, Hosaka GH, Cros☐Arteil

879 S, et al.. Means, motive, and opportunity for biological invasions: genetic introgression in a

880 fungal pathogen. *Mol Ecol*. 2022; doi: 10.1111/mec.16366.

881 83. Diao Y, Zhang C, Xu J, Lin D, Liu L, Mtung'e OG, et al.. Genetic differentiation and

882 recombination among geographic populations of the fungal pathogen *Colletotrichum*

883 *truncatum* from chili peppers in China. *Evol Appl*. 2015; doi: 10.1111/eva.12233.

884 84. Souza-Paccola EA, Fávaro LCL, Casela CR, Paccola-Meirelles LD. Genetic

885 Recombination in *Colletotrichum sublineolum*. *J Phytopathol*. 2003; doi: 10.1046/j.1439-

0434.2003.00727.x.

85. Gale LR. A Population Genetic Approach to Variation in *Colletotrichum graminicola*, the

Causal Agent of Sorghum Anthracnose. *Sorghum Millets Dis*.

86. Crouch JA, Clarke BB, Hillman BI. Unraveling evolutionary relationships among the

divergent lineages of *Colletotrichum* causing anthracnose disease in turfgrass and corn.

Phytopathology. 2006; doi: 10.1094/PHYTO-96-0046.

87. Crouch JA, Tredway LP, Clarke BB, Hillman BI. Phylogenetic and population genetic

divergence correspond with habitat for the pathogen *Colletotrichum cereale* and allied taxa

across diverse grass communities. *Mol Ecol*. 2009; doi: 10.1111/j.1365-294X.2008.04008.x.

88. Rosewich UL, Pettway RE, McDonald BA, Duncan RR, Frederiksen RA. Genetic

structure and temporal dynamics of a *Colletotrichum graminicola* population in a sorghum

disease nursery. *Phytopathology*. 1998; doi: 10.1094/PHYTO.1998.88.10.1087.

89. Xavier K V., Mizubuti ESG, Queiroz M V., Chopra S, Vaillancourt L. Genotypic and

Pathogenic Diversity of *Colletotrichum sublineola* Isolates from Sorghum (*Sorghum bicolor*)

and Johnsongrass (*S. halepense*) in the Southeastern United States. *Phytopathology*. 2018;

doi: https://doi.org/10.1094/PDIS-04-18-0562-RE.

90. Guthrie PAI, Magill CW, Frederiksen RA, Odvody GN. Random Amplified Polymorphic

DNA Markers: A System for Identifying and Differentiating Isolates of *Colletotrichum*

graminicola. Phytopathology.

91. Du M, Schardl CL, Nuckles EM, Vaillancourt LJ. Using mating-type gene sequences for

improved phylogenetic resolution of *Colletotrichum* species complexes. *Mycologia*. 97:641–

582005;

92. Rech GE, Sanz-Martín JM, Anisimova M, Sukno SA, Thon MR. Natural selection on

coding and noncoding DNA sequences is associated with virulence genes in a plant

pathogenic fungus. *Genome Biol Evol*. 2014; doi: 10.1093/gbe/evu192.

93. Tuite JF. Plant Pathological Methods: Fungi and Bacteria. Burgess Publ. Co.

Minneapolis⬜: Burgess Publishing Company;

94. White TJ, Bruns T, Lee S, Taylor J. Amplification and Direct Sequencing of Fungal

Ribosomal Rna Genes for Phylogenetics. *Innis MA, Glefand JJ, White TJ PCR Protoc a Guid*

to methods Appl Acad Press San Diego. 1990; doi: 10.1016/b978-0-12-372180-8.50042-1.

95. Baek JM, Kenerley CM. The arg2 gene of *Trichoderma virens*: Cloning and development

of a homologous transformation system. *Fungal Genet Biol*. 1998; doi:

10.1006/fgbi.1997.1025.

96. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence

data. *Bioinformatics*. 2014; doi: 10.1093/bioinformatics/btu170.

97. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.. SPAdes:

A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J*

Comput Biol. 19:455–772012;

98. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:

Assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. 2015; doi: 10.1093/bioinformatics/btv351.

99. Rochette NC, Rivera-Colón AG, Catchen JM. Stacks 2: Analytical methods for paired-

end sequencing improve RADseq-based population genomics. *Mol Ecol*. 2019; doi:

10.1111/mec.15253.

100. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*. 2009; doi: 10.1093/bioinformatics/btp324.

101. Chiang C, Layer RM, Faust GG, Lindberg MR, Rose DB, Garrison EP, et al.. SpeedSeq:

Ultra-fast personal genome analysis and interpretation. *Nat Methods*. 2015; doi:

10.1038/nmeth.3505.

102. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.

arXiv:12073907 [q-bio]. :1 2012;

103. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al.. The variant

call format and VCFtools. *Bioinformatics*. 2011; doi: 10.1093/bioinformatics/btr330.

104. Ortiz M.. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for

phylogenetic analysis. *DOI:105281/zenodo2540861*. 2019;

105. Katoh K, Misawa K, Kuma KI, Miyata T. MAFFT: A novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Res*. 2002; doi:

10.1093/nar/gkf436.

106. Kamvar ZN, Brooks JC, Grünwald NJ. Novel R tools for analysis of genome-wide

population genetic data with emphasis on clonality. *Front Genet*. 2015; doi:

10.3389/fgene.2015.00208.

107. Goudet J. HIERFSTAT, a package for R to compute and test hierarchical F-statistics.

Mol Ecol Notes. 2005; doi: 10.1111/j.1471-8286.2004.00828.x.

108. Jombart T, Devillard S, Balloux F. Discriminant analysis of principal components: a

new method for the analysis of genetically structured populations. *BMC Genet*. 2010; doi:

10.1371/journal.pcbi.1000455.

109. Beugin MP, Gayet T, Pontier D, Devillard S, Jombart T. A fast likelihood solution to the

genetic clustering problem. *Methods Ecol Evol*. 2018; doi: 10.1111/2041-210X.12968.

110. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. *Mol*

Biol Evol. 2006; doi: 10.1093/molbev/msj030.

111. Primer Geneious. https://www.geneious.com. Accessed June 21, 2020.

113. Hill WG, Robertson A. Linkage disequilibrium in finite populations. *Theor Appl Genet*.

1968; doi: 10.1007/bf01245622.

114. Short DPG, Gurung S, Hu X, Inderbitzin P, Subbarao K V.. Maintenance of sex-related

genes and the co- Occurrence of both mating types in *Verticillium dahliae*. *PLoS One*. 2014;

doi: 10.1371/journal.pone.0112145.

115. Liang X, Yao L, Hao X, Li B, Kong Y, Yuyi Lin a MC, et al.. Molecular Dissection of

Perithecial Mating Line Development in *Colletotrichum fructicola*, a Species with a

Nontypical Mating System Featuring Plus-to-Minus Switch and Plus-Minus- Mediated

Sexual Enhancement. *Appl Env Microbiol*. 2021; doi: https:// doi.org/10.1128/AEM.00474-

21.

116. Baroncelli R, Sreenivasaprasad S, Sukno SA, Thon MR, Holub E. Draft genome

sequence of *Colletotrichum acutatum* sensu lato (*Colletotrichum fioriniae*). *Genome*

Announc. 2014; doi: 10.1128/genomeA.00112-14.

117. Gan P, Ikeda K, Irieda H, Narusaka M, O'Connell RJ, Narusaka Y, et al.. Comparative

genomic and transcriptomic analyses reveal the hemibiotrophic stage shift of *Colletotrichum*

fungi. *New Phytol*. 2013; doi: 10.1111/nph.12085.

118. Alkan N, Friedlander G, Ment D, Prusky D, Fluhr R. Simultaneous transcriptome

analysis of *Colletotrichum gloeosporioides* and tomato fruit pathosystem reveals novel fungal

pathogenicity and fruit defense strategies. *New Phytol*. 2015; doi: 10.1111/nph.13087.

119. Zampounis A, Pigné S, Dallery J-F, Wittenberg AHJ, Zhou S, Schwartz DC, et al..

Genome Sequence and Annotation of *Colletotrichum higginsianum*, a Causal Agent of

Crucifer Anthracnose Disease. *Genome Announc*. 2016; doi: 10.1128/genomeA.00821-16.

120. Baroncelli R, Amby DB, Zapparata A, Sarrocco S, Vannacci G, Le Floch G, et al.. Gene

family expansions and contractions are associated with host range in plant pathogens of the

genus *Colletotrichum. BMC Genomics*. BMC Genomics; 2016; doi: 10.1186/s12864-016-

2917-6.

121. Baroncelli R, Sanz-Martin JM, Rech GE, Sukno SA, Thon MR. Draft Genome Sequence

of *Colletotrichum sublineola,* a Destructive Pathogen of Cultivated Sorghum. *Genome*

Announc. 2014; doi: 10.1128/genomeA.00540-14.

122. Klosterman SJ, Subbarao K V, Kang S, Veronese P, Gold SE, Thomma BPHJ, et al..

Comparative genomics yields insights into niche adaptation of plant vascular wilt pathogens.

PLoS Pathog. 2011; doi: 10.1371/journal.ppat.1002137.

123. Warren HL, Nicholson RL, Turner M.. Field reaction of corn inbreds to *Colletotrichum*

graminicola. *Plant Dis Report*. 59:767–91975;

124. Vargas WA, Sanz Martín JM, Rech GE, Rivera LP, Benito EP, Díaz-Mínguez JM, et al..

Plant defense mechanisms are activated during biotrophic and necrotrophic development of

Colletotricum graminicola in maize. *Plant Physiol*. 2012; doi: 10.1104/pp.111.190397.

125. Alaniz S, Cuozzo V, Martínez V, Stadnik MJ, Mondino P. Ascospore infection and

Colletotrichum species causing *Glomerella* leaf spot of apple in Uruguay. *Plant Pathol J*.

2019; doi: 10.5423/PPJ.OA.07.2018.0145.

126. Taylor J, Jacobson D, Fisher M. The Evolution of Asexual Fungi: Reproduction,

specification and classification. *Annu Rev Phytopathol*. 37:197–2461999;

127. Wilson G.. The identity of the anthracnose of grasses in the United States.

Phytopathology. 4Wilson, G:106–121914;

128. Sawers RJH, Sanchez Leon NL. Origins of maize: A further paradox resolved. *Front*

Genet. 2011; doi: 10.3389/fgene.2011.00053.

129. Matsuoka Y, Vigouroux Y, Goodman MM, Sanchez JG, Buckler E, Doebley J. A single

domestication for maize shown by multilocus microsatellite genotyping. *Proc Natl Acad Sci*

U S A. 2002; doi: 10.1073/pnas.052125199.

130. Stukenbrock EH, McDonald BA. The Origins of Plant Pathogens in Agro-Ecosystems.

Annu Rev Phytopathol. 2008; doi: 10.1146/annurev.phyto.010708.154114.

131. Templeton AR, Routman E, Phillips CA. Separating Population Structure from

Population History: A Cladistic Analysis of the Geographical Distribution of Mitochondrial

DNA Haplotypes in the Tiger Salamander, *Ambystoma tigrinurn*. *Genetics*. 140:767–821995;

132. Silveira AP, Figueiredo MF, Cruz BP. Ocorrência de antracnose do milho no Estado de

São Paulo. *Biol*. 31:192–41965;

133. Talas F, McDonald BA. Genome-wide analysis of *Fusarium graminearum* field

populations reveals hotspots of recombination. *BMC Genomics*. BMC Genomics; 2015; doi:

10.1186/s12864-015-2166-0.

134. Derbyshire M, Denton-Giles M, Hane JK, Chang S, Mousavi-Derazmahalleh M,

Raffaele S, et al.. Selective sweeps in populations of the broad host range plant pathogenic

fungus *Sclerotinia sclerotiorum*. *bioRxiv*. 2018; doi: 10.1101/352930.

135. Cambareri EB, Singer MJ, Selker EU. Recurrence of repeat-induced point mutation

(RIP) in *Neurospora crassa*. *Genetics*. 1991; doi: 10.1093/genetics/127.4.699.

136. van Wyk S, Wingfield BD, De Vos L, van der Merwe NA, Steenkamp ET. Genome-

Wide Analyses of Repeat-Induced Point Mutations in the Ascomycota. *Front Microbiol*.

2021; doi: 10.3389/fmicb.2020.622368.

137. Braga RM, Santana MF, da Costa RV, Brommonschenkel SH, de Araújo EF, de Queiroz

MV. Transposable elements belonging to the Tc1-Mariner superfamily are heavily mutated in

Colletotrichum graminicola. *Mycologia*. 2014; doi: 10.3852/13-262.

138. Clutterbuck AJ. Genomic evidence of repeat-induced point mutation (RIP) in

filamentous ascomycetes. *Fungal Genet Biol*. Elsevier Inc.; 2011; doi:

10.1016/j.fgb.2010.09.002.

139. Crouch JA, Glasheen BM, Giunta MA, Clarke BB, Hillman BI. The evolution of

transposon repeat-induced point mutation in the genome of *Colletotrichum cereale*:

Reconciling sex, recombination and homoplasy in an "asexual" pathogen. *Fungal Genet Biol*.

2008; doi: 10.1016/j.fgb.2007.08.004.

140. Vaillancourt LJ, Hanau RM. Genetic and morphological comparisons of *Glomerella*

(*Colletotrichum*) isolates from maize and from sorghum. *Exp Mycol*. 1992; doi:

10.1016/0147-5975(92)90030-U.

141. Ciampi-Guillardi M, Baldauf C, Souza AP, Silva-Junior GJ, Amorim L. Recent

Introduction and Recombination in *Colletotrichum acutatum* Populations Associated with

Citrus Postbloom Fruit Drop Epidemics in São Paulo, Brazil. *Phytopathology*. 2014; doi:

10.1094/phyto-06-13-0165-r.

142. Rogério F, Oosterhout C van, Ciampi-Guillardi M, Correr FH, Hosaka GK, Cros-Arteil

S, et al.. Means, motive, and opportunity for biological invasions : genetic introgression in a

fungal pathogen. *bioRxiv*. 2021:2021.09.25.4612172021;

143. Roca MG, Davide LC, Mendes-Costa MC, Wheals A. Conidial anastomosis tubes in

Colletotrichum. *Fungal Genet Biol*. 2003; doi: 10.1016/S1087-1845(03)00088-4.

144. Rosada LJ, Franco CCS, Sant'anna JR, Kaneshima EN, Gonçalves-Vidigal MC, Castro-

Prado MAA. Parasexuality in Race 65 *Colletotrichum lindemuthianum* Isolates. *J Eukaryot.*

2010; doi: 10.1111/j.1550-7408.2010.00486.x.

145. Nordzieke DE, Sanken A, Antelo L, Raschke A, Deising HB, Pöggeler S. Specialized

infection strategies of falcate and oval conidia of *Colletotrichum graminicola Fungal Genet*

Biol. Elsevier; 2019; doi: 10.1016/j.fgb.2019.103276.

146. Mcdonald BA. How can research on pathogen population biology suggest disease

management strategies? The example of barley scald (*Rhynchosporium commune*). *Plant*

Pathol. 2016; doi: 10.1111/ppa.12415.

147. Hessenauer P, Feau N, Gill U, Schwessinger B, Brar GS, Hamelin RC. Evolution and

Adaptation of Forest and Crop Pathogens in the Anthropocene. *Phytopathology*. 2020; doi:

10.1094/phyto-08-20-0358-fi.

148. Stukenbrock EH. The Role of Hybridization in the Evolution and Emergence of New

Fungal Plant Pathogens. *Phytopathology*. 2016; doi: 10.1094/phyto-08-15-0184-rvw.

149. Sun Y, Corcoran P, Menkis A, Whittle CA, Andersson SGE, Johannesson H. Large-

scale introgression shapes the evolution of the mating-type chromosomes of the filamentous

ascomycete *Neurospora tetrasperma*. *PLoS Genet*. 2012; doi: 10.1371/journal.pgen.1002820.

150. Gladieux P, Ropars J, Badouin H, Branca A, Aguileta G, De Vienne DM, et al.. Fungal

evolutionary genomics provides insight into the mechanisms of adaptive divergence in

eukaryotes. *Mol Ecol*. 2014; doi: 10.1111/mec.12631.

151. Ahmed S, Tourvieille D, Labrouhe D, Delmotte F. Emerging virulence arising from

hybridisation facilitated by multiple introductions of the sunflower downy mildew pathogen

Plasmopara halstedii. *Fungal Genet Biol*. Elsevier Inc.; 2012; doi:

10.1016/j.fgb.2012.06.012.

152. Hartmann FE, Mcdonald BA, Croll D. Genome-wide evidence for divergent selection

between populations of a major agricultural pathogen. *Mol Ecol*. 2018; doi:

10.1111/mec.14711.

153. O'Connell RJ, Thon MR, Hacquard S, Amyotte SG, Kleemann J, Torres MF, et al..

Lifestyle transitions in plant pathogenic *Colletotrichum* fungi deciphered by genome and

transcriptome analyses. *Nat Genet*. Nature Publishing Group; 2012; doi: 10.1038/ng.2372.

154. Nicholson RL. Criteria for Evaluation of Resistance to Maize Anthracnose.

Phytopathology.

155. Weihmann F, Eisermann I, Becher R, Krijger JJ, Hübner K, Deising HB, et al..

Correspondence between symptom development of *Colletotrichum graminicola* and fungal

biomass, quantified by a newly developed qPCR assay, depends on the maize variety. *BMC*

Microbiol. 2016; doi: 10.1186/s12866-016-0709-4.

156. Lim SM. Estimates of Heterosis and Combining Ability for Resistance of Maize to

Colletotrichum graminicola. *Phytopathology*. 1978; doi: 10.1094/phyto-68-1336.

157. Matiello RR, Brunelli KR, Lopes MTG, Morello RMSC, Silva HP da, Camargo LEA.

Inheritance of resistance to anthracnose stalk rot (*Colletotrichum graminicola*) in tropical

maize inbred lines. *Crop Breed Appl Biotechnol*. 2012; doi: 10.1590/s1984-

70332012000300003.

158. Carson ML, Hooker AL. Inheritance of Resistance to Stalk Rot of Corn Caused by

Colletotrichum graminicola. *Phytopathology*. 71:1190–61981.

1090