Allele-specific **knockouts reveal a role for** *apontic-like* **in the**

evolutionary loss of larval pigmentation in the domesticated

silkworm, *Bombyx mori*

Kenta Tomihara[1*], Peter Andolfatto[2], Takashi Kiuchi[1*]

[1]Graduate School of Agricultural and Life Sciences, The University of Tokyo, Yayoi,

Bunkyo-ku, Tokyo 113-8657, Japan

[2]Department of Biological Sciences, Columbia University, New York, NY 10026, USA

*Corresponding Author:

Kenta Tomihara: tomihara@g.ecc.u-tokyo.ac.jp

Takashi Kiuchi: kiuchi@g.ecc.u-tokyo.ac.jp

Running title: Genome editing in *Bombyx mandarina*

Abstract

The domesticated silkworm, *Bombyx mori*, and its wild progenitor, *B. mandarina*, are extensively studied as a model case of the evolutionary process of domestication. A conspicuous difference between these species is the dramatic reduction in pigmentation in both larval and adult *B. mori*. Here we evaluate the efficiency of CRISPR/Cas9-targeted knockouts of pigment-related genes as a tool to understand their potential contributions to domestication-associated pigmentation loss in *B. mori*. To demonstrate the efficacy of targeted knockouts in *B. mandarina*, we generated a homozygous CRISPR/Cas9-targeted knockout of *yellow-y*. In *yellow-y* knockout mutants, black body color became lighter throughout the larval, pupal and adult stages, confirming a role for this gene in pigment formation. Further, we performed *allele-specific* CRISPR/Cas9-targeted knockouts of the pigment-related transcription factor, *apontic-like* (*apt-like*) in *B. mori* × *B. mandarina* F_1 hybrid individuals. Knockout of the *B. mandarina* allele of *apt-like* in F_1 embryos results in depigmented patches on the dorsal integument of larvae, whereas corresponding knockouts of the *B. mori* allele consistently exhibit normal F_1 larval pigmentation. These results demonstrate a contribution of *apt-like* to the evolution of reduced pigmentation in *B. mori*. Together, our results demonstrate the feasibility of CRISPR/Cas9-targeted knockouts as a tool for understanding the genetic basis of traits associated with *B. mori* domestication.

Key words: *Bombyx mandarina*, domestication, pigmentation, CRISPR/Cas9, reciprocal hemizygosity test.

Brief abstract

Bombyx mori and its wild progenitor are an important model for the study of phenotypic evolution associated with domestication. As proof-of-principle, we used CRISPR/Cas9 to generate targeted knockouts of two pigmentation-related genes. By generating a homozygous knockout of *yellow-y* in *B. mandarina*, we confirmed this gene's role in pigment formation. Further, by generating *allele-specific* knockouts of *apontic-like* (*apt-like*) in *B. mori* × *B. mandarina* F_1 hybrids, we establish that evolution of *apt-like* contributed to reduced pigmentation during *B. mori* domestication.

Graphical TOC/Abstract

Bombyx mandarina F_1 (*B. mori* × *B. mandarina*)

yellow-y knockout

B. mandarina specific *apt-like* knockout

Introduction

The silkworm, *Bombyx mori*, was domesticated over 5000 years ago from its wild progenitor species, *B. mandarina*. Under long-term artificial selection, *B. mori* acquired various characteristics suitable for sericulture. For example, the weight of the cocoon shell of *B. mori* is much higher than that of *B. mandarina* (Ômura 1950, Li *et al.* 2017, Fang *et al.* 2020). In addition, *B. mori* moths lost their flying ability due to the degeneration of their flight muscles and a reduction in wing stiffness (Lu *et al.* 2020). Among the most conspicuous domestication-associated traits is a marked reduction in pigmentation in *B. mori* larvae and adults relative to *B. mandarina* (Figure 1A). Curiously, depigmentation is a major trait contributing to the so-called "domestication syndrome" observed in a variety of domesticated animals, but for reasons that are not well-understood (Wilkins *et al.* 2014).

The genetic basis of pigmentation loss associated with *B. mori* domestication is not yet known. Previous studies reported that melanin synthesis pathway genes *tyrosine hydroxylase* (*TH*) and *aspartate decarboxylase* (*ADC*, also known as *black*) were potentially targets of selection during silkworm domestication (Yu *et al.* 2011, Xiang *et al.* 2018). Pigmentation patterning genes are also likely to be associated with the body color differences between *B. mori* and *B. mandarina*. The wild-type *B. mori* larvae are largely white but exhibit melanic spots (*i.e.* eye spots, crescent spots and star spots, Figure 1A). *B. mandarina* larvae, on the other hand, are substantially darker, and exhibit extensive dorsal pigment patterning that includes banding and spots (Figure 1A).

In *B. mori*, distinct alleles of the genetic locus, *p*, encode at least 15 different

larval markings such as spots, stripes, and banding (Yoda *et al.* 2014). The gene

underlying allelic variation at *p*, *apontic-like* (*apt-like*), encodes a transcription factor

that is likely to regulate the expression of melanin synthesis pathway genes such as

yellow-y, *ebony*, *TH*, *Dopa decarboxylase* (*DDC*) and *laccase 2* (Futahashi *et al.* 2008,

Yoda *et al.* 2014) (Figure 1B). While *apt-like* has been implicated in pigmentation

differences among *B. mori* strains, it's potential role in depigmentation of *B. mori* during

domestication is not known. A hybrid strain (semiconsomic T02), in which chromosome

2 of *B. mandarina* has been substituted into the genomic background of *B. mori*, exhibits

a phenotype similar to that of the *B. mori* allele *moricaud* (p^M) (Fujii *et al.* 2021). Since

apt-like resides on chromosome 2, dorsal pigmentation patterning on the larvae of *B.*

mandarina that is absent in *B. mori* was hypothesized to be controlled by the expression

of *apt-like* (Yoda *et al.* 2014, Fujii *et al.* 2021), although direct evidence is lacking.

The genetic analysis of domestication-associated loss of pigmentation in *B.*

mori has been challenging because deficiencies in candidate genes can result in lethality.

For example, in the fruit fly, *Drosophila melanogaster*, *TH*-deficient (*pale*) mutants die

at embryonic stage (Neckameyer and White 1993). In *B. mori*, *TH*-deficient mutants

(*sch lethal*, sch^l) and RNAi-mediated knockdowns of *TH* are both lethal at embryonic

stage (Liu *et al.* 2010). In addition, *apt* mutants in *D. melanogaster* die at embryonic

stage (Eulenberg and Schuh 1997, Gellon *et al.* 1997), and RNAi-mediated knockdown

of *apt-like* in *B. mori* embryos results in death before hatching (Yoda *et al.* 2014).

The issue of lethality is likely to continue to impede the genetic analysis of

pigmentation loss and other domestication-associated traits in *B. mori*. Recently, several

candidate genes associated with domestication in *B. mori* have been identified using

quantitative trait locus (QTL) mapping, including silk production (Li *et al.* 2017, Fang

et al. 2020), larval climbing ability (Wang, Lin, *et al.* 2020), and mimicry (Wang, Lin,

et al. 2020). In addition, a recent population genetic analysis identified 300 candidate

genes as targets of recent selection in *B. mori*, some of which are likely to be associated

with silk production and voltinism (Xiang *et al.* 2018). Despite these efforts, reverse

genetic approaches such as targeted gene knockouts and editing (Takasu *et al.* 2010,

2013, Wang *et al.* 2013), gene silencing (Quan *et al.* 2002), or transgenesis (Tamura *et*

al. 2000) will likely be required to fully understand the function of these candidate genes

and their potential contribution to domestication-associated traits in *B. mori*. Notably,

the functions of domestication candidate genes have been tested by reverse genetics

approaches in *B. mori* (Xiang *et al.* 2018) but these approaches, to our knowledge, have

not yet been implemented in *B. mandarina*.

Here we use CRISPR/Cas9-targeted knockouts of two candidate pigmentation

genes in two distinct contexts. First, we demonstrate the feasibility of CRISPR/Cas9-

targeted knockouts in *B. mandarina* by generating a homozygous *yellow-y* knockout

strain. Next, to circumvent the lethal effects of knocking out a second candidate gene,

apt-like, we use *allele-specific* CRISPR/Cas9-targeted knockouts in *B. mori* × *B.*

mandarina F_1 hybrids. The latter experiment also comprises a test (the "reciprocal

hemizygosity test", Steinmetz *et al.* 2002, Stern 2014) of the contribution of *apt-like*

evolution to the domestication-associated loss of pigmentation in *B. mori*.

Results and discussion

A CRISPR/Cas9-targeted knockout of *yellow-y* in *B. mandarina*.

To demonstrate the feasibility of CRISPR/Cas9-targeted knockouts in *B.*

mandarina, we focused on known pigmentation-related genes. *TH* is among the most

compelling candidate genes in the melanin synthesis pathway that seems likely to

underlie domestication-associated loss of pigmentation in *B. mori* (Yu *et al.* 2011, Xiang

et al. 2018). However, *TH* knockout mutants are predicted to be lethal (Neckameyer and

White 1993, Liu *et al.* 2010), rendering them difficult to study. Thus, we decided to

target *yellow-y* (Figure 1B), a melanin synthesis gene that is downstream of *TH* and

functions in melanin synthesis in wide range of insects including *B. mori* and other

Lepidoptera and is predicted to be non-essential (Futahashi *et al.* 2008, Zhang *et al.* 2017,

Chen *et al.* 2018, Matsuoka and Monteiro 2018, Liu *et al.* 2020, Wang, Huang, *et al.*

2020, Han *et al.* 2021, Shirai *et al.* 2021). After confirming the coding sequence

annotation (CDS) of the *B. mandarina yellow-y* gene, we designed a unique CRISPR-

RNA (crRNA) target site in exon 2 (Figure 2A, Table S1).

In *B. mori*, CRISPR/Cas9-targeted genome editing requires microinjection into

non-diapausing eggs (Kanda and Tamura 1994). We obtained non-diapausing eggs by

rearing *B. mandarina* larvae under 16 h-light/8 h-dark conditions (Kobayashi 1990). We

then injected a mixture of crRNA, trans-activating crRNA (tracrRNA), and Cas9 into

336 *B. mandarina* embryos. Among 16 hatched larvae (G_0 generation), nine grew to

adult moths (Table 1). We crossed six G_0 adults with wild-type moths and obtained

generation 1 (G_1) eggs. Using a heteroduplex mobility assay on the PCR products from

G_1 embryos, we confirmed that mutations were introduced at the target site in five of the

six G_1 broods (Table 1, Figure S1), showing that CRISPR/Cas9-induced mutations of

yellow-y were heritable. We then crossed G_1 siblings with each other and obtained

yellow-y homozygous knockout individuals carrying a five-nucleotide deletion followed

by a single-nucleotide substitution (Figure 2B), which results in a frame-shift and

premature stop codon. We designated this mutant allele *yellow-y$^{\Delta 5}$* and used it for further

analyses.

B. mandarina yellow-y$^{\Delta 5}$ homozygotes hatched normally and their development

was comparable to that of wild-type individuals (Figure 3). In homozygous *yellow-y$^{\Delta 5}$*

neonate larvae, the larval integument and the head capsule are reddish brown instead of

the normal black (Figure 3A) and in final instar larvae, spots and dorsal pigmentation

patterns are lighter than that of the wild type (Figure 3B). Later in development, the

pupal integument of homozygous *yellow-y$^{\Delta 5}$* mutants exhibits reddish color instead of

the normal black (Figure 3C), and the body and wing spot markings of *yellow-y$^{\Delta 5}$* adult

moths are lighter than that of wild-type (Figure 3D). Further, the phenotypes of

heterozygous *+/yellow-y$^{\Delta 5}$* individuals were comparable to that of wild type (data not

shown). Together, these observations suggest that, as observed in *B. mori* (Futahashi *et*

al. 2008), *yellow-y* is a non-essential gene contributing to melanin pigment synthesis in

B. mandarina and loss-of-function is recessive.

Comparative data from other species suggests that *yellow-y* functions

differently among lepidopteran species. For example, while *yellow-y* loss-of-function is

also found to be recessive in several other lepidopteran species (Liu *et al.* 2020, Wang,

Huang, *et al.* 2020, Han *et al.* 2021, Shirai *et al.* 2021), it is dominant in the black

cutworm, *Agrotis ipsilon* (Chen *et al.* 2018)*.* Further, unlike *Bombyx*, *yellow-y*

knockouts in *Agrotis* are susceptible to dehydration (Chen *et al.* 2018) and mutants in

Spodoptera exhibit defects in body development, copulation, oviposition, and

hatchability (Liu *et al.* 2020, Han *et al.* 2021, Shirai *et al.* 2021). These observations

suggest that although the function of *yellow-y* in melanin synthesis is conserved,

additional *yellow-y* functions might be diverged among lepidopteran species. *yellow*

genes are a rapidly evolving gene family, and loss or duplication of some *yellow* genes

have been observed in Lepidoptera (Chen *et al.* 2018, Liu *et al.* 2020, Han *et al.* 2021,

Shirai *et al.* 2021). It is possible that functions of *yellow-y* outside of melanin synthesis

in *Bombyx* are compensated for by other *yellow* paralogs. In Lepidoptera, the function

of most *yellow* genes remained unknown except for *yellow-y* (Futahashi *et al.* 2008,

Zhang *et al.* 2017, Chen *et al.* 2018, Matsuoka and Monteiro 2018, Liu *et al.* 2020, Wang,

Huang, *et al.* 2020, Han *et al.* 2021, Shirai *et al.* 2021), *yellow-e* (Ito *et al.* 2010), *yellow-*

d (Zhang *et al.* 2017) and *yellow-h2/3* (Zhang *et al.* 2017). Further studies of *yellow*

genes including *yellow-y* in diverse insect species are required to understand the

functions and evolution of *yellow* paralogs.

While our results demonstrate the feasibility of genome editing in *B. mandarina*,

these experiments are challenging due to low post-injection hatchability rates (4.8% in

our experiment). A recent study showed that the hatchability of *B. mandarina* eggs is

generally lower than *B. mori* eggs, even under normal conditions (Zhu *et al.* 2019). To

compare the hatchability of *B. mori* and *B. mandarina* embryos after injection, we

injected three commonly used buffers (see Methods) and distilled water into embryos of

both species and compared hatching rates. We found that the post-injection hatchability

of *B. mandarina* embryos (0–8.3 %) is substantially lower than that of *B. mori* (22.9–

62.5 %) in all experimental conditions (Table 2), suggesting that *B. mandarina* embryos

are more sensitive to injection.

***Allele-specific* knockouts of *apt-like* in interspecific F$_1$ hybrids.**

Knockouts of some pigmentation pathway genes, such as *apt-like* and *TH*, are

predicted to be lethal (Neckameyer and White 1993, Eulenberg and Schuh 1997, Gellon

et al. 1997, Liu *et al.* 2010, Yoda *et al.* 2014). Considering this obstacle to the study of

essential genes, together with the observation (above) of reduced hatchability of injected

B. mandarina embryos, we instead opted for the alternative strategy of injecting *B. mori*

× *B. mandarina* F$_1$ hybrids. Specifically, we conducted an *allele-specific* CRISPR/Cas9-

targeted knockout of *apt-like* in F$_1$ embryos (from crosses between *B. mori* females and

B. mandarina males). We reasoned that, because the components of these F$_1$ eggs is

derived from the *B. mori* mother, these embryos should have post-injection hatchability

similar to that of *B. mori*. We designed species-specific crRNA targeting *apt-like*, for

which the targeted protospacer adjacent motif (PAM) sequence is only present in either

B. mori- or *B. mandarina*-derived sequence (Methods, Figure 4, Table S1). For each

targeted allele, we injected a mixture of crRNA, tracrRNA and Cas9 into 48 F_1 embryos.

To confirm that mutations were specifically introduced into the targeted allele, we

extracted genomic DNA from adult legs and PCR-amplified the target sites. We then

carried out heteroduplex mobility assays using a microchip electrophoresis system (Ota

et al. 2013, Ansai *et al.* 2014), which show that various mutations were introduced into

the *apt-like* target sequence in both of the *allele-specific* knockout series (Figure S2).

The PCR products obtained from two representative individuals of each *allele-specific*

knockout series were then cloned and sequenced (Figure 4). The sequences of cloned

PCR products confirmed that various mutations were introduced into the target sites in

both knockout series, some of which cause frameshifts and associated premature stop

codons. All mutations detected in this experiment were specifically introduced only into

the targeted allele.

In addition to spots, normal F_1 larvae have a dark body with darker greyish

brown banding covering wide range of the dorsal surface, similar to the pattern for *B.*

mandarina (Figure 1A, Figure S3). As predicted, post-injection hatchability of F_1

embryos was high whether targeting the *B. mori* (60%) or the *B. mandarina* (66%)

allele. In the *apt-like* knockout series targeting the *B. mori* allele, all of the 26 larvae that

survived to the fifth instar stage exhibited normal body color (Table 3, Figure 5). In

contrast, for the *apt-like* knockout series targeting the *B. mandarina* allele, 24 of the 29

larvae that survived to the fifth instar stage exhibited white patches on dorsal

pigmentation pattern banding that varied in size (Table 3, Figure 5, Figure S4). This

result establishes a role for *apt-like* in body pigment formation and that the *B.*

mandarina-derived *apt-like* allele is dominant with respect to this trait. Notably,

however, the pigmentation of F_1 pupae and adult stages of the knockout series targeting

the *B. mori* or the *B. mandarina* allele both exhibited normal body color (data not shown).

Additionally, larval spots, which are also predicted to be controlled by *apt-like* (Yoda *et*

al. 2014), were also not affected the F_1 series targeting either the *B. mori* or the *B.*

mandarina allele (Figure 5, Figure S4). Since both wild-type ($+^p$) *B. mori* and *B.*

mandarina larvae exhibit spots (Figure 1A), we conclude that the *B. mori* or the *B.*

mandarina-derived *apt-like* alleles are both sufficient to direct the formation of larval

spots.

Our results have implications beyond merely confirming a role for *apt-like* in

B. mandarina larval pigmentation. The *allele-specific* knockouts of *apt-like* in F_1 hybrids

allows us to compare the phenotype of genetically identical hybrids that differ only at

the target locus (a framework called the "reciprocal hemizygosity test", Steinmetz *et al.*

2002, Stern 2014). As such, we can attribute the loss of pigmentation in the series

targeting the *B. mandarina* allele to evolution at the *apt-like* gene in *B. mori,* rather than

exclusively at a *trans*-acting factor. The Apt-like proteins of *B. mori* (p50T) and *B.*

mandarina (Sakado) differ by only one amino acid substitution: Alanine to Valine at

residue 188 (see DDBJ accession numbers LC706749 and LC706750). However, this

substitution is not observed in *B. mandarina* collected at different locations (see NCBI

accession numbers SRR6111377, SRR6111379, SRR6111381 and SRR6111382),

suggesting this is not a fixed amino acid difference between species. This implies that

evolution of an *apt-like cis*-regulatory element contributes to the observed phenotypic

difference between species.

Conclusion

Here we demonstrate the utility of CRISPR/cas9 genome editing in *B.*

mandarina and *B. mori* × *B. mandarina* F_1 hybrids to the study the function and

evolution of domestication-associated candidate genes. Focusing on two pigmentation-

related genes, we show that *apt-like* plays a role in larval body pigmentation patterning

(Figure 5), whereas *yellow-y* plays a more general role in pigmentation that is not pattern

dependent or specific to developmental life-stage (Figure 3B). These results are

consistent with the proposed roles of Apt-like as a transcription factor responsible for

larval color patterning, and Yellow-y as an enzyme in the melanin synthesis pathway

under control of Apt-like (Futahashi *et al.* 2008, Yoda *et al.* 2014). Further, using the

framework of the reciprocal hemizygosity test, we show that *apt-like* has evolved in *B.*

mori in a way that has specifically reduced larval body pigmentation, without affecting

the formation of larval spots or adult body pigmentation.

Despite being a powerful tool to study gene function and evolution, the *allele-*

specific knockout approach has several limitations. First, it requires sequence variants

distinguishing the parents of the F_1 hybrid that result in a PAM-site that is only present

in one of two species, limiting the potential to design *allele-specific* targets. However,

this limitation may be overcome by using Cas proteins that recognize different PAM-

sites (Leenay and Beisel 2017). For example, while the *Streptococcus pyogenes* Cas9

(*Sp*Cas9) protein has been used here, the Cas12a, which recognizes a distinct PAM

sequence, has also been implemented in *B. mori* (Dong *et al.* 2020). In addition, recent

studies have reported that *Sp*Cas9 can be modified to recognize alternative PAM

sequences (Kleinstiver *et al.* 2015, 2016). A second limitation is that, given the

mosacism of knockouts in G_0 individuals, one cannot exclude the possibility of false-

negative phenotyping results. The cleavage efficiency of the CRISPR/Cas9 system is

affected by several features, such as the sequences of PAM-distal and PAM-proximal

regions of the guide RNA, the genomic context of the targeted DNA, as well as GC-

content and secondary structure of the guide RNA (Liu *et al.* 2016). To minimize this

problem, one can screen a large number of G_0 individuals and confirm that mutations

were introduced with high efficiency (as in Figure 4 and Figure S2).

Despite these limitations, our results highlight several advantages of *allele-*

specific knockouts in the F_1 over knockouts in *B. mandarina*. First, our results show that

B. mori (female) × *B. mandarina* (male) F_1 embryos are substantially more tolerant to

injection compared to *B. mandarina*. Second, *allele-specific* knockouts in the F_1 permit

the study of essential genes (such as *apt-like*) at which knockouts are expected to be

homozygous lethal and recessive with respect to the phenotype. Finally, *allele-specific*

CRISPR/Cas9-targeted knockouts in F_1 hybrids have the added utility of identifying loci

that have diverged in function between *B. mori* and *B. mandarina*, and contributing to

domestication-related traits in *B. mori* using the framework of the reciprocal

hemizygosity test. Thus, our study showcases the multifaceted utility of *allele-specific*

knockouts in F_1 hybrids in the study of gene function and evolution.

Experimental procedures

Insects

The *B. mori* strain p50T, a single-paired descendant of individuals of strain p50

(a   derivative   from   Daizo;

https://shigen.nig.ac.jp/silkwormbase/ViewStrainDetail.do?name=p50), is maintained

at our laboratory. The *B. mandarina* strain, Sakado, was originally collected in Sakado-

city, Saitama, Japan, in 1982. Since then, it has been maintained at our laboratory by

sib-mating. All larvae were reared on fresh mulberry leaves or artificial diet (SilkMate

PS, NOSAN) under continuous 12 h-light/12 h-dark conditions at 25 °C with the

exceptions described below. For injections in to *B. mandarina* embryos, we obtained

non-diapausing eggs by rearing *B. mandarina* larvae under continuous long-day

conditions (16 h-light/8 h-dark) at 25 °C (Kobayashi 1990). We then collected eggs in

crosses between emerged adults. To generate *B. mori* × *B. mandarina* F_1 hybrid

embryos for injection, we first incubated *B. mori* eggs at 15 ºC under continuous

darkness (Kogure 1933). The hatched larvae were then reared under continuous 16 h-

light/8 h-dark condition at 25 °C and females were crossed to *B. mandarina* males.

Confirmation of the *yellow-y* and *apt-like* coding sequences in *B. mandarina*.

Total RNA was extracted from the integument of fourth instar *B. mandarina*

larvae using TRIzol (Thermo Fisher Scientific). Complementary DNA (cDNA) was

reverse transcribed from total RNA using TaKaRa RNA PCR Kit (TaKaRa). Reverse

transcriptase-PCR was performed using KOD One polymerase (TOYOBO). PCR

products were cloned into pGEM-T Easy Vector (Promega) and Sanger-sequenced

using the FASMAC sequencing service (Kanagawa, Japan).

Knockout of *yellow-y* in *B. mandarina*

An unique crRNA target sequence in the *B. mori* genome was selected using

CRISPRdirect (https://crispr.dbcls.jp) (Table S1) (Naito *et al.* 2015). The uniqueness

of the target sequence in the *B. mandarina* genome was then confirmed by performing

blastn at SilkBase (http://silkbase.ab.a.u-tokyo.ac.jp). A mixture of crRNA, tracrRNA

and Cas9 Nuclease protein NLS (600 ng/µL; NIPPON GENE) in injection buffer (100

mM KOAc, 2 mM Mg(OAc)$_2$, 30 mM HEPES-KOH; pH 7.4) was injected into each

embryo within 3 h after oviposition (Yamaguchi *et al.* 2011).

The injected embryos (G_0 generation) were incubated at 25 °C in a humidified

Petri dish until hatching. Adult G_0 moths were crossed with wild-type *B. mandarina*,

and G_1 eggs were obtained. To detect heritable CRISPR/Cas9-induced mutations, ten

G_1 eggs were collected into one tube, and genomic DNA was prepared using the

HotSHOT method (Truett *et al.* 2000). The region containing the target site of *yellow-y*

crRNA was PCR-amplified using KOD One polymerase (TOYOBO). Mutations at the

target site were detected by heteroduplex mobility assay using the MultiNA microchip

electrophoresis system (SHIMAZU) with the DNA-500 reagent kit (Ota *et al.* 2013,

Ansai *et al.* 2014).

Adult G_1 moths from broods with heritable CRISPR/Cas9-induced mutations were

crossed with each other to obtain homozygous knockout mutants (G_2). To confirm

CRISPR/Cas9-induced mutations, we prepared genomic DNA, PCR-amplified the

target region and detected mutations as described above using G_2 adult moths. To

determine the precise nature of insertions, deletions and substitutions, PCR products

obtained from G_2 individuals were directly Sanger-sequenced using the FASMAC

sequencing service (Kanagawa, Japan).

Comparison of post-injection hatchability.

Three commonly used buffers (injection buffer 1 (Yamaguchi *et al.* 2011),

injection buffer 2 (Tamura *et al.* 2000), PBS buffer) and distilled water into embryos of

B. mori or *B. mandarina* within 3 h after oviposition. The injected embryos were

incubated at 25 °C in a humidified Petri dish until hatching.

***Allele-specific* gene knockouts in F_1 hybrids.**

A PAM sequence is necessary for target recognition and following DNA

cleavage in CRISPR/Cas system (Hsu *et al.* 2013, Anders *et al.* 2014), and *Sp*Cas9,

which we used in this study, recognizes 5'-NGG-3' as PAM. We specifically targeted

*Sp*Cas9 PAM-sites that differed in sequence between *B. mori* and *B. mandarina* (Figure

4, Table S1), allowing *allele-specific* DNA cleavage and gene knockout (Courtney *et*

al. 2015, Christie *et al.* 2017). A mixture of crRNA, tracrRNA and Cas9 was injected

to F_1 embryos as described above. To evaluate CRISPR/Cas9 cleavage efficiency, we

extracted genomic DNA from G_0 adult legs and PCR-amplified the target region as

described above. PCR products were cloned into pGEM-T Easy Vector and Sanger-

sequenced using an ABI3130xl genetic analyzer (Applied Biosystems).

Acknowledgements

We are grateful to Katsuya Satta and Hisashi Tobita for assistance with the

experiments. We are also grateful to Professor Susumu Katsuma for helpful discussions.

We thank the Institute for Sustainable Agro-ecosystem Services, The University of

Tokyo, for facilitating the mulberry cultivation and the Biotron Facility at the

University of Tokyo for rearing the silkworms. This work was supported by JSPS

KAKENHI grant number JP20J22954 to KT and JP20H02997 to TK.

Data Availability

366   Full-length coding sequences of *B. mori yellow-y*, *B. mandarina yellow-y*, *B.

367 mori apt-like* and *B. mandarina apt-like* are available on the DDBJ under accession

368 numbers of LC706747, LC706748, LC706749 and LC706750, respectively.

Author contributions

371   KT, PA and TK designed the study. KT performed most of the experiments. KT

372 wrote the manuscript with intellectual input from TK and PA. All authors edited and

373 approved the final version of the manuscript and agree to be accountable for all aspects

374 of the work.

References

References

Anders, C., Niewoehner, O., Duerst, A., and Jinek, M., 2014. Structural basis of

PAM-dependent target DNA recognition by the Cas9 endonuclease. *Nature*, 513

(7519), 569–573.

Ansai, S., Inohaya, K., Yoshiura, Y., Schartl, M., Uemura, N., Takahashi, R., and

Kinoshita, M., 2014. Design, evaluation, and screening methods for efficient

targeted mutagenesis with transcription activator-like effector nucleases in

medaka. *Development, Growth & Differentiation*, 56 (1), 98–107.

Chen, X., Cao, Y., Zhan, S., Zhang, Y., Tan, A., and Huang, Y., 2018. Identification

of *yellow* gene family in *Agrotis ipsilon* and functional analysis of *Aiyellow-y* by

CRISPR/Cas9. *Insect Biochemistry and Molecular Biology*, 94, 1–9.

Christie, K.A., Courtney, D.G., DeDionisio, L.A., Shern, C.C., De Majumdar, S.,

Mairs, L.C., Nesbit, M.A., and Moore, C.B.T., 2017. Towards personalised

allele-specific CRISPR gene editing to treat autosomal dominant disorders.

Scientific Reports, 7 (1), 16174.

Courtney, D.G., Moore, J.E., Atkinson, S.D., Maurizi, E., Allen, E.H.A., Pedrioli,

D.M.L., McLean, W.H.I., Nesbit, M.A., and Moore, C.B.T., 2015. CRISPR/Cas9

DNA cleavage at SNP-derived PAM enables both *in vitro* and *in vivo KRT12*

mutation-specific targeting. *Gene Therapy*, 23 (1), 108–112.

Dong, Z., Qin, Q., Hu, Z., Zhang, X., Miao, J., Huang, L., Chen, P., Lu, C., and Pan,

M., 2020. CRISPR/Cas12a mediated genome editing enhances *Bombyx mori*

resistance to BmNPV. *Frontiers in Bioengineering and Biotechnology*, 8, 841.

Eulenberg, K.G. and Schuh, R., 1997. The tracheae defective gene encodes a bZIP

protein that controls tracheal cell movement during *Drosophila* embryogenesis.

The EMBO Journal, 16 (23), 7156–7165.

Fang, S., Zhou, Q., Yu, Q., and Zhang, Z., 2020. Genetic and genomic analysis for

cocoon yield traits in silkworm. *Scientific Reports*, 10 (1), 5682.

Fujii, T., Kiuchi, T., Daimon, T., Ito, K., Katsuma, S., Shimada, T., Yamamoto, K.,

and Banno, Y., 2021. Development of interspecific semiconsomic strains

between the domesticated silkworm, *Bombyx mori* and the wild silkworm, *B.*

mandarina. *Journal of Insect Biotechnology and Sericology*, 90 (2), 33–40.

Futahashi, R., Sato, J., Meng, Y., Okamoto, S., Daimon, T., Yamamoto, K., Suetsugu,

Y., Narukawa, J., Takahashi, H., Banno, Y., Katsuma, S., Shimada, T., Mita, K.,

and Fujiwara, H., 2008. *yellow* and *ebony* are the responsible genes for the larval

color mutants of the silkworm *Bombyx mori*. *Genetics*, 180 (4), 1995–2005.

Gellon, G., Harding, K.W., McGinnis, N., Martin, M.M., and McGinnis, W., 1997. A

genetic screen for modifiers of deformed homeotic function identifies novel

genes required for head development. *Development*, 124 (17), 3321–3331.

Han, W., Tang, F., Zhong, Y., Zhang, J., and Liu, Z., 2021. Identification of *yellow*

gene family and functional analysis of *Spodoptera frugiperda yellow-y* by

CRISPR/Cas9. *Pesticide Biochemistry and Physiology*, 178, 104937.

Hsu, P.D., Scott, D.A., Weinstein, J.A., Ran, F.A., Konermann, S., Agarwala, V., Li,

Y., Fine, E.J., Wu, X., Shalem, O., Cradick, T.J., Marraffini, L.A., Bao, G., and

Zhang, F., 2013. DNA targeting specificity of RNA-guided Cas9 nucleases.

Nature Biotechnology, 31 (9), 827–832.

Ito, K., Katsuma, S., Yamamoto, K., Kadono-Okuda, K., Mita, K., and Shimada, T.,

2010. Yellow-e determines the color pattern of larval head and tail spots of the

silkworm *Bombyx mori*. *Journal of Biological Chemistry*, 285 (8), 5624–5629.

Kanda, T. and Tamura, T., 1994. Microinjection method for DNA in early embryos of

the silkworm, *Bombyx mori*, using air-pressure. *Bulletin of the National Institute*

of Sericultural and Entomological Science, 2, 31-46 (in Japanese with English

summary).

Kleinstiver, B.P., Pattanayak, V., Prew, M.S., Tsai, S.Q., Nguyen, N.T., Zheng, Z.,

and Joung, J.K., 2016. High-fidelity CRISPR–Cas9 nucleases with no detectable

genome-wide off-target effects. *Nature*, 529 (7587), 490–495.

Kleinstiver, B.P., Prew, M.S., Tsai, S.Q., Topkar, V. V., Nguyen, N.T., Zheng, Z.,

Gonzales, A.P.W., Li, Z., Peterson, R.T., Yeh, J.-R.J., Aryee, M.J., and Joung,

J.K., 2015. Engineered CRISPR-Cas9 nucleases with altered PAM specificities.

Nature, 523 (7561), 481–485.

Kobayashi, J., 1990. Effects of photoperiod on the induction of egg diapause of

tropical races of the domestic silkworm, *Bombyx mori*, and the wild silkworm, *B.*

mandarina. *Japan Agricultural Research Quarterly*, 23 (3), 202–205.

Kogure, M., 1933. The influence of light and temperature on certain characters of the

silkworm, *Bombyx mori*. *Journal of the Faculty of Agriculture, Kyushu*

University, 4 (1), 1–93.

Leenay, R.T. and Beisel, C.L., 2017. Deciphering, communicating, and engineering

the CRISPR PAM. *Journal of Molecular Biology*, 429 (2), 177–191.

Li, C., Tong, X., Zuo, W., Luan, Y., Gao, R., Han, M., Xiong, G., Gai, T., Hu, H.,

Dai, F., and Lu, C., 2017. QTL analysis of cocoon shell weight identifies

BmRPL18 associated with silk protein synthesis in silkworm by pooling

sequencing. *Scientific Reports*, 7 (1), 17985.

Liu, C., Yamamoto, K., Cheng, T.C., Kadono-Okuda, K., Narukawa, J., Liu, S.P.,

Han, Y., Futahashi, R., Kidokoro, K., Noda, H., Kobayashi, I., Tamura, T.,

Ohnuma, A., Banno, Y., Dai, F.Y., Xiang, Z.H., Goldsmith, M.R., Mita, K., and

Xia, Q.Y., 2010. Repression of *tyrosine hydroxylase* is responsible for the *sex-*

linked chocolate mutation of the silkworm, *Bombyx mori. Proceedings of the*

National Academy of Sciences of the United States of America, 107 (29), 12980–

12985.

Liu, X., Han, W., Ze, L., Peng, Y., Yang, Y., Zhang, J., Yan, Q., and Dong, S., 2020.

Clustered regularly interspaced short palindromic repeats/CRISPR-associated

protein 9 mediated knockout reveals functions of the *yellow-y* gene in

Spodoptera litura. Frontiers in Physiology, 11, 1643.

Liu, X., Homma, A., Sayadi, J., Yang, S., Ohashi, J., and Takumi, T., 2016. Sequence

features associated with the cleavage efficiency of CRISPR/Cas9 system.

Scientific Reports, 6 (1), 19675.

Lu, K., Liang, S., Han, M., Wu, C., Song, J., Li, C., Wu, S., He, S., Ren, J., Hu, H.,

Shen, J., Tong, X., and Dai, F., 2020. Flight muscle and wing mechanical

properties are involved in flightlessness of the domestic silkmoth, *Bombyx mori*.

Insects, 11 (4), 220.

Matsuoka, Y. and Monteiro, A., 2018. Melanin pathway genes regulate color and

morphology of butterfly wing scales. *Cell reports*, 24 (1), 56–65.

Naito, Y., Hino, K., Bono, H., and Ui-Tei, K., 2015. CRISPRdirect: software for

designing CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics*,

31 (7), 1120–1123.

Neckameyer, W.S. and White, K., 1993. *Drosophila* Tyrosine hydroxylase is encoded

by the *pale* locus. *Journal of Neurogenetics*, 8 (4), 189–199.

Ômura, S., 1950. Researches on the behavior and ecological characteristics of the

wild silkworm *Bombyx mandarina*. *Bulletin of the Imperial Sericultural*

Experiment Station, 13 (3), 79-130 (in japanese with English summary).

Ota, S., Hisano, Y., Muraki, M., Hoshijima, K., Dahlem, T.J., Grunwald, D.J., Okada,

Y., and Kawahara, A., 2013. Efficient identification of TALEN-mediated

genome modifications using heteroduplex mobility assays. *Genes to Cells*, 18

(6), 450–458.

Quan, G.-X., Kim, I., Kômoto, N., Sezutsu, H., Ote, M., Shimada, T., Kanda, T.,

Mita, K., Kobayashi, M., and Tamura, T., 2002. Characterization of the

kynurenine 3-monooxygenase gene corresponding to the *white egg 1* mutant in

the silkworm *Bombyx mori*. *Molecular Genetics and Genomics*, 267 (1), 1–9.

Shirai, Y., Ohde, T., and Daimon, T., 2021. Functional conservation and

diversification of *yellow-y* in lepidopteran insects. *Insect Biochemistry and*

Molecular Biology, 128, 103515.

Steinmetz, L.M., Sinha, H., Richards, D.R., Spiegelman, J.I., Oefner, P.J., McCusker,

J.H., and Davis, R.W., 2002. Dissecting the architecture of a quantitative trait

locus in yeast. *Nature*, 416 (6878), 326–330.

Stern, D.L., 2014. Identification of loci that cause phenotypic variation in diverse

species with the reciprocal hemizygosity test. *Trends in Genetics*, 30 (12), 547–

554.

Takasu, Y., Kobayashi, I., Beumer, K., Uchino, K., Sezutsu, H., Sajwan, S., Carroll,

D., Tamura, T., and Zurovec, M., 2010. Targeted mutagenesis in the silkworm

Bombyx mori using zinc finger nuclease mRNA injection. *Insect Biochemistry*

and Molecular Biology, 40 (10), 759–765.

Takasu, Y., Sajwan, S., Daimon, T., Osanai-Futahashi, M., Uchino, K., Sezutsu, H.,

Tamura, T., and Zurovec, M., 2013. Efficient TALEN construction for *Bombyx*

mori gene targeting. *PLoS ONE*, 8 (9), e73458.

Tamura, T., Thibert, C., Royer, C., Kanda, T., Eappen, A., Kamba, M., Kômoto, N.,

Thomas, J.-L., Mauchamp, B., Chavancy, G., Shirk, P., Fraser, M., Prudhomme,

J.-C., and Couble, P., 2000. Germline transformation of the silkworm *Bombyx*

mori L. using a *piggyBac* transposon-derived vector. *Nature Biotechnology*, 18

(1), 81–84.

Truett, G.E., Heeger, P.S., Mynatt, R.L., Truett, A.A., Walker, J.A., and Warman,

M.L., 2000. Preparation of PCR-quality mouse genomic DNA with hot sodium

hydroxide and tris (HotSHOT). *Biotechniques*, 29 (1), 52–54.

Wang, M., Lin, Y., Zhou, S., Cui, Y., Feng, Q., Yan, W., and Xiang, H., 2020.

Genetic mapping of climbing and mimicry: two behavioral traits degraded

during silkworm domestication. *Frontiers in Genetics*, 11, 1568.

Wang, Y., Huang, Y., Xu, X., Liu, Z., Li, J., Zhan, X., Yang, G., You, M., and You,

S., 2020. CRISPR/Cas9-based functional analysis of *yellow* gene in the

diamondback moth, *Plutella xylostella*. *Insect Science*, 28 (5), 1504–1509.

Wang, Y., Li, Z., Xu, J., Zeng, B., Ling, L., You, L., Chen, Y., Huang, Y., and Tan,

A., 2013. The CRISPR/Cas System mediates efficient genome engineering in

Bombyx mori. *Cell Research*, 23 (12), 1414–1416.

Wilkins, A.S., Wrangham, R.W., and Tecumseh Fitch, W., 2014. The "domestication

syndrome" in mammals: a unified explanation based on neural crest cell

behavior and genetics. *Genetics*, 197 (3), 795–808.

Xiang, H., Liu, X., Li, M., Zhu, Y., Wang, L., Cui, Y., Liu, L., Fang, G., Qian, H.,

Xu, A., Wang, W., and Zhan, S., 2018. The evolutionary road from wild moth to

domestic silkworm. *Nature Ecology & Evolution*, 2 (8), 1268–1279.

Yamaguchi, J., Mizoguchi, T., and Fujiwara, H., 2011. siRNAs induce efficient RNAi

response in *Bombyx mori* embryos. *PLoS ONE*, 6 (9), e25469.

Yoda, S., Yamaguchi, J., Mita, K., Yamamoto, K., Banno, Y., Ando, T., Daimon, T.,

and Fujiwara, H., 2014. The transcription factor Apontic-like controls diverse

colouration pattern in caterpillars. *Nature Communications*, 5 (1), 4936.

Yu, H.-S., Shen, Y.-H., Yuan, G.-X., Hu, Y.-G., Xu, H.-E., Xiang, Z.-H., and Zhang,

Z., 2011. Evidence of selection at melanin synthesis pathway loci during

silkworm domestication. *Molecular Biology and Evolution*, 28 (6), 1785–1799.

Zhang, L., Martin, A., Perry, M.W., van der Burg, K.R.L., Matsuoka, Y., Monteiro,

A., and Reed, R.D., 2017. Genetic basis of melanin pigmentation in butterfly

wings. *Genetics*, 205 (4), 1537–1550.

Zhu, Y.N., Wang, L.Z., Li, C.C., Cui, Y., Wang, M., Lin, Y.J., Zhao, R.P., Wang, W.,

and Xiang, H., 2019. Artificial selection on storage protein 1 possibly

contributes to increase of hatchability during silkworm domestication. *PLOS*

Genetics, 15 (1), e1007616.

Figures and Tables

Table 1. Efficiency of CRISPR/Cas9-targeted knockout in *B. mandarina* targeting

yellow-y.

# eggs injected	# hatched larvae	# adults	# G_1 broods	# G_1 broods carrying mutant alleles
336	16	9	6	5

546   **Table 2.** The hatchability of *B. mori* and *B. mandarina* embryos injected with an

547   injection buffer 1 (Yamaguchi *et al.* 2011), injection buffer 2 (Tamura *et al.* 2000), PBS

548   buffer and distilled water.

Treatment	Composition	*B. mori*		*B. mandarina*	
		# injected eggs	# hatched larvae	# injected eggs	# hatched larvae
Injection buffer 1	100 mM KOAc, 2 mM Mg(OAc)$_2$, 30 mM HEPES-KOH; pH 7.4	48	11	48	0
Injection buffer 2	0.5 mM phosphate buffer (pH 7.0), 5 mM KCl	48	30	48	4
PBS buffer	137 mM NaCl, 2.7 mM KCl, 10 mM Na$_2$HPO$_4$, 1.76mM KH$_2$PO$_4$; pH 7.4	48	23	48	1
Distilled water	–	48	17	48	2

550   **Table 3.** Efficiency of *allele-specific* knockouts of *apt-like* in F_1 hybrids.

Targeted allele	# eggs injected	# hatched larvae	# 5th instar larvae	# mosaic 5th instar larvae	# adults (male/female)
B. mori	48	29	26	0	26 (16/10)
B. mandarina	48	32	29	24	29 (18/11)

Figure 1. (A) Larvae (top) and adult males (bottom) of *B. mori* (left) and *B. mandarina* (right). Black arrows indicate larval spot markings (eye spots, crescent spots and star spots) and red dotted lines delimit the larval dorsal pigment patterning ("banding"). The names of spots follow the nomenclature of Yoda *et al.* (2014). (B) The proposed melanin biosynthesis pathway in *Bombyx* larvae largely adapted from Futahashi *et al.* (2008) and Yoda *et al.* (2014). Orange dashed arrows indicate presumed regulation of genes by the transcription factor *apt-like*.

A

yellow-y

ATG

ACTA**CCC**<u>GGACCAATTCTCGAAGCAGC</u>AGGC

TGA

B

```
            Trp Asn Tyr Pro Asp Gln Phe Ser Lys Gln Gln Ala Leu Arg Thr Gly
Wild type   TGG AAC TAC CCG GAC CAG TTC TCG AAG CAG CAG GCT CTC AGG ACT GGT
yellow-yΔ5   TGG AAC TAC CC        AA TTC TCG AAG CAG CAG GCT CTC AGG ACT GGT
```

Figure 2. The nature of the lesion in the *yellow-y* knockout (*yellow-y*$^{\Delta 5}$) in *B. mandarina*. (A) Gene structure of *yellow-y* and the selected crRNA target site. The target sequence is underlined and the protospacer adjacent motif (PAM) site is shown in bold letters. (B) Alignment of the *yellow-y* gene sequences surrounding the crRNA target site from wild type, and G$_2$ *B. mandarina* individuals that are homozygous for a 5-base pair deletion and associated single nucleotide mutation. The target sequence is underlined, and the PAM site (CCN) is indicated in bold letters.

570 **Figure 3.** The phenotypes of wild-type and *yellow-y* knockout mutants of *B. mandarina*.

571 Representative (A) neonate larvae, (B) final-instar larvae, (C) male pupae and (D) adult

572 male moths of wild-type (left or top) and *yellow-y*$^{\Delta 5}$ (right or bottom) *B. mandarina*.

573 Arrows indicate wing spot markings in adult moths. Scale bars: 1 mm (A) or 1 cm (B-

574 D).

Figure 4. Mutations introduced by *allele-specific apt-like* mosaic knockout (mKO) in

F₁ hybrids. Annotation and inset alignments of the *apt-like* gene in both species with

selected crRNA target sites whose PAM sequences are only present in (A) *B. mori* or

(B) *B. mandarina*. Two *allele-specific* knockout individuals were sequenced for each

target. The target sequences are underlined, and the PAM sequences are shown in bold

letters. *B. mori* specific SNPs and *B. mandarina* specific single nucleotide variants are

highlighted with green and yellow shadings, respectively. Mutations introduced by

CRISPR/Cas9 system are highlighted with red shading. Numbers on the right edge

indicate the numbers of the clones identified among all cloned and sequenced PCR

products.

588   **Figure 5.** Representative fifth instar larvae of normal F_1 (left), *allele-specific apt-like*

589   mKOs targeting the *B. mori* allele (middle) and the *B. mandarina* allele (right). Red

590   arrows indicate ectopic white (depigmented) regions. Scale bars: 1 cm.

592 **Table S1** PCR primers and RNA sequences used in this study.

Primer Name	Sequence	Usage
PCR primers		
yellow-y_cloning_F02	AAGCTGCCGCGAACATAG	RT-PCR and CDS cloning
yellow-y_cloning_R02	TGCAATAGAAAGCCTCACATTT	RT-PCR and CDS cloning
apt-like_cloning_F	GAAAATAAGAGCCCACGCACT	RT-PCR and CDS cloning
apt-like_cloning_R	ACTTGAATCCGCTTTTGACG	RT-PCR and CDS cloning
yellow-y_mid_F	CGTCTTTGGGTATTAGACGTTGG	CDS cloning
yellow_y_check_F	GTACATACCTGAGCGCCACCT	Genomic PCR for mutation detection and sequencing in CRISPR/Cas9-targeted gene knockout
yellow_y_check_R	CAGCAACGATAAAGCTCCAAG	Genomic PCR for mutation detection and sequencing in CRISPR/Cas9-targeted gene knockout
Bmand_apt-like_F03	GAGGTTTTGTGTGGCGAGATG	Genomic PCR for mutation detection and sequencing in CRISPR/Cas9-targeted gene knockout
Bmand_apt-like_R02	AGGAAGTTCTGCTGGAGTCTG	Genomic PCR for mutation detection and sequencing in CRISPR/Cas9-targeted gene knockout
Bmori_apt-like_F01	CTGGACGATCACATAAAGCAC	Genomic PCR for mutation detection and sequencing in CRISPR/Cas9-targeted gene knockout
Bmori_apt-like_R01	TCTCAACTCAATGAGGAACAGC	Genomic PCR for mutation detection and sequencing in CRISPR/Cas9-targeted gene knockout
RNAs		
yellow-y_crRNA_1	GCUGCUUCGAGAAUUGGUCCGU UUUAGAGCUAUGCUGUUUUG	CRISPR/Cas9-targeted gene knockout
Bmori_apt-like_specific_crRNA	GAAGUACAGCUUCUCCUCUUGU UUUAGAGCUAUGCUGUUUUG	CRISPR/Cas9-targeted gene knockout
Bmand_apt-like_specific_crRNA	UGUCGUUCGUUGUCGAUAGUGU UUUAGAGCUAUGCUGUUUUG	CRISPR/Cas9-targeted gene knockout
tracrRNA	AAACAGCAUAGCAAGUUAAAAU AAGGCUAGUCCGUUAUCAACUU GAAAAAGUGGCACCGAGUCGGU GCUUUUUUU	CRISPR/Cas9-targeted gene knockout

Figure S1. Detection of mutations at the *yellow-y* crRNA target site using a heteroduplex

mobility shift assay. Two intervals were prepared for each G_1 brood. Ten eggs were

collected into one tube, and genomic DNA was prepared. The region containing the

target site of *yellow-y* crRNA was PCR-amplified. Multiple heteroduplex bands caused

by insertion/deletion mismatches were observed indicating the presence of DNA lesions

(insertions/deletions and/or associated nucleotide variants).

Figure S2. Detection of mutations at *apt-like* crRNA target sites using a heteroduplex

mobility shift assay. The region containing the target site of (A) *B. mori* specific *apt-*

like crRNA and (B) *B. mandarina* specific *apt-like* crRNA was PCR-amplified using

DNA prepared from G_0 adults' legs. Multiple heteroduplex bands caused by

insertion/deletion mismatches and associated nucleotide variants were observed in G_0

mosaics.

Figure S3. A representative (A) fifth instar larva and (B) adult male moth of normal F_1

(*B mori* female × *B. mandarina* male). Black arrows indicate larval spot markings and

red dotted lines outline the banding. Normal F_1 larvae have a dark body with darker

greyish brown banding covering wide range of the dorsal surface similar to *B.*

mandarina. The names of spots were referred to Yoda *et al.* (2014). Scale bars: 1 cm.

Normal F₁

B. mandarina
specific
apt-like mKO

Figure S4. Fifth instar larvae of normal F$_1$ and *B. mandarina* specific *apt-like* mKO.

Red arrows indicate ectopic white (depigmented) regions. Crescent and star spots were

not depigmented in all larvae. Scale bars: 1 cm.