A zebrafish embryo screen utilizing gastrulation for identification of anti-metastasis drugs

Joji Nakayama[1,2,3,§], Hideki Makinoshima[1,4] , and Zhiyuan Gong[3]

[1]Tsuruoka Metabolomics Laboratory, National Cancer Center, Tsuruoka, Japan

[2]Shonai Regional Industry Promotion Center, Tsuruoka, Japan.

[3]Department of Biological Science, National University of Singapore, Singapore

[4]Division of Translational Research, Exploratory Oncology Research, and Clinical Trial Center,

National Cancer Center, Kashiwa, Japan.

[§]Corresponding author:

Joji Nakayama

Tsuruoka Metabolomics Laboratory, National Cancer Center, Mizukami 246-2, Kakuganji,

Tsuruoka, Yamagata, Japan 975-0052.

E-mail: zmetastasis@gmail.com

ORCID: 0000-0003-1077-140X

Tel: +81-235-64-0980

Keywords: Metastasis, gastrulation, EMT, Phenotyping screening, zebrafish

Total words excluding references and figure legends: 2354

Number of figures: 4

Number of tables: 1

Abstract

Few models exist that allow for rapid and effective screening of anti-metastasis drugs. Here, we

present a phenotype-based chemical screen utilizing gastrulation of zebrafish embryos for

identification of anti-metastasis drugs. Based on the evidence that metastasis proceeds through

utilizing the molecular mechanisms of gastrulation, we hypothesize that chemicals which

interrupt zebrafish gastrulation might suppress metastasis of cancer cells. Thus, we developed a

drug screening protocol which uses epiboly, the first morphogenetic movement in gastrulation, as

a marker. The screen only needs zebrafish embryos and enables hundreds of chemicals to be

tested in five hours through observing epiboly progression of a test chemical-treated embryos. In

the screen, embryos at the two-cell stage are firstly corrected and then developed to the sphere

stage. The embryos are treated with a test chemical and incubated in the presence of the chemical

until vehicle-treated embryos develop to 90% epiboly stage. Finally, positive 'hit' chemicals that

interrupt epiboly progression are selected through comparing epiboly progression of the

chemical-treated embryos with that of vehicle-treated embryos under a stereoscopic microscope.

Previous study subjected 1,280 FDA-approved drugs to the screen and identified Adrenosterone

and Pizotifen as epiboly-interrupting drugs. These drugs were validated to suppress metastasis of

breast cancer cells in mice models of metastasis. Furthermore, 11β–Hydroxysteroid

Dehydrogenase 1 (HSD11β1) and serotonin receptor 2C (HTR2C), which are primary target of

Adrenosterone and Pizotifen respectively, promotes metastasis through induction of epithelial-

mesenchymal transition (EMT). That indicates the screen could be diverted to a chemical genetic

screening platform for identification of metastasis-promoting genes.

Introduction

Cancer research using zebrafish as a model has attracted attention because this model offers many unique advantages that are not readily provided by other animal models. Futhermore, the zebrafish system has been increasingly recognized as a platform for chemical screening because it provides the advantage of high-throughput screening in an in vivo vertebrate setting with physiologic relevance to humans [1-5].

Metastasis is responsible for approximately 90% of cancer-associated mortality. It proceeds through multiple steps: invasion, intravasation, survival in the circulatory system, extravasation, colonization, and metastatic tumor formation in secondary organs with angiogenesis [6-8]. Dissemination of cancer cells is an initial step of metastasis and its molecular mechanism involves local breakdown of basement membrane, loss of cell polarity, and induction of EMT [9,10]. These cellular and biological phenomena are also observed during vertebrate gastrulation in that evolutionarily conserved morphogenetic movements of epiboly, internalization, convergence, and extension cooperate to generate germ layers and sculpt the body plan [11]. In zebrafish, the first morphogenetic movement, epiboly, is initiated at approximately four hours post-fertilization (hpf) to move cells from the animal pole to eventually engulf the entire yolk cell by 10 hpf. These movements are governed by the molecular mechanisms that are induced by temporally and spatially regulated gene expression, and these mechanisms and changes in gene expression are partially observed in metastatic progression [12].

Development of the protocol

Metastasis proceeds through utilizing the molecular mechanisms of gastrulation. At least fifty common genes were shown to be involved in both gastrulation and metastasis progression (Table 1) [13-17]. The fifty genes are expressed in Xenopus or zebrafish embryos, and genetic inhibition of each of the fifty genes in these embryos interferes with gastrulation progression. Conversely, the same fifty genes are ectopically expressed in metastatic cancer cells and confer metastatic

properties on cancer cells, and genetic inhibition of each of the fifty genes suppresses metastasis

progression. These evidences led us to hypothesize that chemicals which interfere with zebrafish

gastrulation might suppress metastasis progression of cancer cells. Based on the hypothesis, we

developed a drug screening protocol which uses epiboly, the first morphogenetic movement in

gastrulation, as a marker. This screen measures the suppressor effect of each of test chemicals

through observing epiboly progression of the chemical-treated embryos (Fig. 1 and Fig. 2).

Applications of the method

This screen enables hundreds of chemicals to be tested in five hours. Our study subjected 1280

FDA-approved drugs to this screen and identified Adrenosterone and Pizotifen as epiboly-

interfering drugs. These drugs were further validated to suppress metastasis of breast cancer cells

in mouse models of metastasis (Fig. 3) [18,19]. This screen can also measure suppressor effect of

crude drugs. We subjected 120 herbal medicines to this screen and identified cinnamon bark

extract as an epiboly-interfering drug. Cinnamon bark extract was validated to suppress

metastatic dissemination of breast cancer cells in zebrafish xenograft model [20]. Moreover, this

screen can be diverted to a chemical genetic screening platform for identification of metastasis-

promoting genes. HSD11β1 and HTR2C, which are respectively primary targets of

Adrenosterone and Pizotifen, induce EMT and promote metastasis of breast cancer cells (Fig. 4)

[18,19].

Comparison with other methods

Current mouse models of metastasis are too expensive and time-consuming to use for rapid and

high-throughput screening [21,22]. Also in vitro model of metastatic dissemination such as a Boyden

chamber assay can test a limited number of chemicals in one assay and needs huge time and

effort in analyzing the results [23]. In contrast, our screen only needs zebrafish embryos and enables

hundreds of chemicals to be tested in five hours through observing epiboly progression of a test

chemical-treated embryos. Furthermore, out of the 78 chemicals which interrupt epiboly

progression of zebrafish embryos, 20 of the chemicals were validated to suppress cell motility

and invasion of highly metastatic human cancer cells without affecting cell viability in a Boyden

chamber assay. Among the 20 chemicals, Adrenosterone and Pizotifen were validated to suppress

metastasis of breast cancer cells in mice models of metastasis [18,19]. A disadvantage of this screen

is that zebrafish have orthologues to 86% of 1318 human drug targets [27]. Therefore, 75% of the

chemicals which interrupt epiboly progression of zebrafish embryos, fail to suppress cell motility

and invasion of highly metastatic human cancer cells in a Boyden chamber assay [19].

Experimental design

This screen measures suppressor effect of each of chemicals based on epiboly progression of

zebrafish embryos. Niclosamid or DMSO is used as positive or negative control, respectively.

Epiboly progression of each of chemical treated embryos is compared with that of DMSO-treated

embryos. Firstly, embryos at the two-cell stage are firstly corrected and then developed to the

sphere stage. The embryos are treated with a test chemical and incubated in the presence of the

chemical until vehicle-treated embryos develop to 90% epiboly stage. Finally, positive 'hit'

chemicals that interrupt epiboly progression are selected through comparing epiboly progression

of the chemical-treated embryos with that of vehicle-treated embryos under a stereoscopic

microscope [19].

Limitations

There is a limitation in delivering chemicals to zebrafish embryos. Zebrafish embryos are

surrounded by the acellular chorion, which is known to be about 1.5–2.5 μm thick and to consist

of three layers pierced by pore canals. The pore allows passage of water, ions, and chemicals. A

study reported molecules which are larger than 3-4 KDa fail to pass through the chorion.

Therefore, this screen may not be able to measure suppressor effect of the molecules which are

larger than 3-4 KDa [28].

Materials

REAGENTS

• Wild-type zebrafish strain

• E3 medium (5.0 mM NaCl, 0.17 mM KCl, 0.33 mM $MgSO_4$)

• FDA, EMA, and other agencies-approved chemical libraries were purchased from

Prestwick Chemical (Illkirch, France).

• Niclosamid

• DMSO

EQUIPMENT

• 24-well flat bottom plastic plates (Corning)

• Stereomicroscope (MZ75, Leica)

• Incubator (Thermo)

Procedure

Zebrafish mating setup (Day 0) _Timing 10 mins

1. On the night before collecting embryos, arrange male and female zebrafish in pairs

separated by a divider

CRITICAL STEP

Young adult zebrafish should be used for the crossing. Qualities of zebrafish embryos

affect screening efficiency.

Embryo collection, and distribution (Day 1) _Timing 10-30 min

2. Remove the divider to allow the fish to spawn.

3. To obtain zebrafish embryos of the same development stage, zebrafish were crossed for

10 mins. If more than twenty chemicals were tested, the crossing were conducted three

times at three different time points (Group A_8:30, Group B_9:00 and Group

C_9:30AM).

4. After 10 mins, set back divider to prevent zebrafish from spawning.

CRITICAL STEP

This screen measures suppressor effect of each chemical on progression of epiboly in live

zebrafish embryos. Therefore, epiboly proceeds during measuring the effect under a

stereoscopic microscope. If more than 20 chemicals are tested, screening should be

divided into more than two sessions and each of the sessions start at different time point.

For example, if 60 chemicals are screened, zebrafish should be crossed at three different

time points over 30 mins apart. To do that, 30 mins for measuring the effects would be

ensured.

5. Collect the embryos and remove dead embryos

6. Incubate the embryos at 27 ˚C for twenty mins

7. Collect embryos at the two-cell stage under stereoscopic microscope.

8. Array approximately twenty embryos into each well of a 24-well plate

9. Remove E3 medium from each well including the embryos by using a pipet

10. Add 900 µl of E3 medium to the well.

Embryo development to the sphere stage_Timing 4 hours

11. Incubate the embryos at 27 ˚C until the embryos develop to the sphere stage

CRITICAL STEP

The temperature of E3 medium affects the rate of development of zebrafish embryos.

Higher temperature accelerates the rate; conversely, lower temperature slows the rate [29].

176   Therefore, non-uniform temperature between E3 medium of each well of 24-well plate

177   containing zebrafish embryos would cause false positive.

Addition of chemicals_Timing 30 mins

12. At 30 mins before adding test chemicals to embryos, prepare 10-fold concentration of

each of the chemicals in E3 medium

13. Add 100 µl of 10-fold concentration of the medium to 900 µl of E3 medium containing

zebrafish embryos when the embryos develop to the sphere stage.

14. For example, for 60 test chemicals to be screened, they are divided into three groups.

a. First 20 test chemicals plus Niclosamid as positive control, and DMSO as negative

control are added into group A when embryos from group A develop to sphere stage.

b. Second 20 test chemicals plus Niclosamid, and DMSO were added into group B when

embryos from group B develop to sphere stage.

c. Last 20 test chemicals plus Niclosamid, and DMSO were added into group C when

embryos from group C develop to sphere stage

Development of DMSO-treated embryos to 90% epiboly stage_Timing 5 hours

15. After the addition of test chemicals, the embryos are incubated at 27 ˚C for approximately

five hours.

CRITICAL STEP

The temperature of E3 medium affects the rate of development of zebrafish embryos. Non-

uniform temperature between E3 medium of each well of 24-well plate containing zebrafish

embryos would cause false positive.

Measuring the inhibition effects of each of chemicals_Timing 30 mins

16. Comparing epiboly progression of each of chemicals-treated embryos from group A with that of DMSO-treated embryos from group A under the stereoscopic microscope when DMSO-treated embryos from group A develop to 90% epiboly stage.

17. After approximately 30 min from observing embryos in group A, measuring the inhibition effects of each of chemicals on epiboly progression of embryos in group B and when DMSO-treated embryos from group B develop to 90% epiboly stage.

18. After approximately 30 mis from observing embryos in group B, measuring the inhibition effects of each of chemicals on epiboly progression of embryos in group C and when DMSO-treated embryos in group C develop to 90% epiboly stage.

CRITICAL STEP

Epiboly proceeds during comparing epiboly progression of each of chemicals-treated embryos with that of DMSO-treated embryos under the stereoscopic microscope. Therefore, measuring the effect should be done in 30 mins.

Timing

Step 1, Zebrafish mating setup: overnight

Step 2-5, Crossing zebrafish: 10 mins

Step 6, Develop the embryos to the two-cell stage

Step 7, Collect two cell stage embryos: 20 mins

Step 8-10, Array 20 embryos into each well of 24-well plate: 30 mins

Step 11, Develop the embryos to sphere stage: 4 hours

Step 12-14, Prepare chemical drugs in E3 medium and add the medium into each well of 24-well plate: 30 mins

Step 15, Develop the embryos to 90% epiboly stage: 5 hours

Step 16-18, measuring epiboly progression: 30 mins

Troubleshooting

Qualities of zebrafish embryos affect screening efficiency. Low qualities of embryos show high frequencies of abnormal embryos with asymmetric cell cleavage, and development of the embryos arrest at early cleavage stages [30]. If a screen used low qualities of zebrafish embryos, it would generate false 'hit' chemicals since suppressor effect of a test chemical is measured by observing epiboly progression of the chemical-treated embryos. If the number of zebrafish embryos showing morphological abnormalities correlate with final concentration of a test chemical, the abnormalities may result from an effect of the test chemical on the embryos.

Anticipated results

Suppressor effects of a tested chemical on epiboly progression of zebrafish embryos are significantly affected by final concentration of the chemical. Previous study subjected 1,280 FDA-approved drugs to the screen and showed 6% (78/1280) of the tested drugs affected epiboly progression of the embryos when the embryos were treated with 10 μM. Out of the 78 epiboly-interrupting drugs, 25% of the drugs succeed to suppress cell motility and invasion of highly metastatic human cancer cells in a Boyden chamber assay. In contrast, epiboly progression was affected more severely when the embryos were treated at 50 μM. 10.3% (132/1280) of the tested drugs affected epiboly progression of the embryos, but 85 % (112/132) of the epiboly-interrupting drugs failed to suppress cell motility and invasion of highly metastatic human cancer cells in a Boyden chamber assay [19].

Author contributions statements

J.N. created a concept of this screen, designed research, conducted experiments, analyzed data, wrote an original draft, supervised this project; H.M. and Z.G. administered this project, acquired funding. All authors reviewed and edited the draft.

ORCID for corresponding author

ORCID: Joji Nakayama, 0000-0003-1077-140X

Acknowledgments

We thank Dr. Herrick (Albany Medical College) for providing pCMV-h5TH2C-VSV with us.

Illustrations of Fig. 1 were drawn by Ami Inoue (Kyoto University of the Arts). This study was

funded by grants from National Medical Research Council of Singapore (R-154000547511) and

Ministry of Education of Singapore (R-154000A23112 and R154000B88112) to ZG.

Experiments using mice were supported in part by research funds from the Yamagata prefectural

government and the City of Tsuruoka.

Competing interests

J.N., H.M., and Z.G. declare no conflict of interest.

Fig. 1

Fig. 1_Graphic schematic of a phenotype-based chemical screen using zebrafish embryos

Pairs of adult zebrafish are crossed and their embryos at the two-cell stage are collected and

arrayed into individual wells of 24-well plate. Chemicals are added into each well when the

embryos develop to the sphere stage. Epiboly progression of each of chemicals-treated embryos

are compared with that of DMSO-treated embryos under a stereoscopic microscope when

DMSO-treated embryos develop to 90% epiboly stage.

Fig. 2

| Niclosamide | Hexachlorophene | Pizotifen | Ipriflavone | DMSO |

| Tolcapone | Troglitazone | Vinpocetine | Zuclopenthxiol | Mebendazole |

| Severe delayed | Delayed | Slighly delayed | Non effect |

Fig. 2_Representative samples of the embryos that were treated with indicated chemicals.

Indicated chemicals were added when the embryos develop to the sphere stage. Embryos were

treated with 10 μM. Niclosamide-treated embryos serve as positive control and DMSO-treated

embryos serve as negative control. Epiboly progression of each of chemicals-treated embryos are

compared with that of DMSO-treated embryos under a stereoscopic microscope when DMSO-

treated embryos develop to 90% epiboly stage.

Fig. 3

Fig. 3_Pizotifen, one of epiboly-interrupting drugs, suppressed metastatic progression

of breast cancer cells in vitro and vivo

a, Effect of pizotifen on cell motility and invasion of MBA-MB-231, MDA-MB-435, and

PC9 cells. Either vehicle or pizotifen treated the cells were subjected to Boyden chamber

assays. Fetal bovine serum (1% v/v) was used as the chemoattractant in both assays. Each

experiment was performed at least twice. **b**, Representative images of primary tumors on day

10 post-injection (top panels) and metastatic burden on day 70 post-injection (bottom panels)

taken using an IVIS Imaging System. **c**, Number of metastatic nodules in the lung of either

vehicle- or pizotifen-treated mice.

Fig. 4

Fig. 4_ HTR2C, a primary target of Pizotifen, induces epithelial-to-mesenchymal

transition (EMT)-mediated metastatic dissemination of human cancer cells

a, Immunofluorescence staining of E-cadherin and vimentin expressions in the MCF7 cells.

b, Expression of E-cadherin and vimentin, and HTR2C were examined by western blotting in

the MCF7 and HaCaT cells; GAPDH loading control is shown (bottom). **c**, Representative

images of dissemination patterns of MCF7 cells expressing either the control vector (top left)

or HTR2C (lower left) in a zebrafish xenotransplantation model. White arrowheads indicate

disseminated MCF7 cells. The mean frequencies of the fish showing head, trunk, or end-tail

dissemination (right). Each value is indicated as the mean ± SEM of two independent

experiments. Statistical analysis was determined by Student's t test.

308 **Table 1. A list of the common fifty genes that are involved between gastrulation and**

309 **metastasis progression**

310

Genes	Gastrulation Defects	Ref	Effects in Metastasis	Ref
BMP	Convergence and extension	[31]	EMT	[32]
WNT	Convergence and extension	[33]	Migration and Invasion	[34]
FGF	Convergence and extension	[35]	Invision	[36]
EGF	Epiboly	[37]	Migration	[38]
PDGF	Convergence and extension	[39]	EMT	[40]
CXCL12	Migration of endodermal cells	[41]	Migration and Invasion	[42]
CXCR4	Migration of endodermal cells	[41]	Migration and Invasion	[42]
PIK3CA	Convergence and extension	[43]	Migration and Invasion	[44]
YES	Epiboly	[45]	Migration	[46]
FYN	Epiboly	[47]	Migration and Invasion	[48]
MAPK1	Epiboly	[49]	Migration	[50]
SHP2	Convergence and extension	[51]	Migration	[52]
SNAI1	Convergence and extension	[53]	EMT	[54]
SNAI2	Mesoderm & Neural crest formation	[55]	EMT	[56]
TWIST1	Mesoderm formation	[57]	EMT	[58]
TBXT	Convergence and extension	[33]	EMT	[59]
ZEB1	Epiboly	[60]	EMT	[61]
GSC	Mesodermal patterning	[62]	EMT	[63]
FOXC2	Unclear, defects in gastrulation	[64]	EMT	[65]
STAT3	Convergence and extension	[66]	Migration	[67]
POU5F1	Epiboly	[68]	EMT	[69]
EZH2	Unclear, defects in gastrulation	[70]	Invasion	[71]
EHMT2	Defects in Neurogenesis	[72]	Migration and Invasion	[73]
BMI1	Defects in skelton formation	[74]	EMT	[75]
RHOA	Convergence and extension	[76]	Migration and Invasion	[77]
CDC42	Convergence and extension	[78]	Migration and Invasion	[79]
RAC1	Convergence and extension	[80]	Migration and Invasion	[81]
ROCK2	Convergence and extension	[82]	Migration and Invasion	[83]
PAR1	Convergence and extension	[84]	Migration	[85]
PRKCI	Convergence and extension	[84]	EMT	[86]
CAP1	Convergence and extension	[87]	Migration	[88]
EZR	Epiboly	[89]	Migration	[90]
EPCAM	Epiboly	[91]	Migration and Invasion	[92]
ITGB1 / ITA5	Mesodermal Migration	[93]	Migration and Invasion	[94]
FN1	Convergence and extension	[95]	Invasion	[96]
HAS2	Dorsal migration of lateral cells	[97]	Invasion	[98]
MMP14	Convergence and extension	[99]	Invasion	[100]
COX1	Epiboly	[101]	Invasion	[102]
PTGES	Convergence and extension	[103]	Invasion	[104]
SLC39A6	Aterior migration	[105]	EMT	[106]
GNA12 / 13	Convergence and extension	[72]	Migration and Invasion	[107]
OGT	Epiboly	[108]	Migration and Invasion	[109]
CCN1	Cell Movement	[110]	Migration and Invasion	[111]
TRPM7	Convergence and extension	[112]	Migration	[113]
MAPKAPK2	Epiboly	[114]	Migration	[115]
B4GALT1	Convergence and extension	[116]	Invasion	[117]
IER2	Convergence and extension	[118]	Migration	[119]
TIP1	Convergence and extension	[120]	Migration and Invasion	[121]
PAK5	Convergence and extension	[122]	Migration	[123]
MARCKS	Convergence and extension	[124]	Migration and Invasion	[125]

311
312

References

1   MacRae, C. A. & Peterson, R. T. Zebrafish as tools for drug discovery. *Nature
REVIEWS* **14**, 721-731 (2015).
2   Zon, L. I. & Peterson, R. T. In vivo drug discovery in the zebrafish *Nat Rev Drug
Discov* **4**, 35-44 (2005).
3   Nakayama., J., Lu., J.-W., Makinoshima., H. & Gong., Z. A Novel Zebrafish Model of
Metastasis Identifies the HSD11β1 Inhibitor Adrenosterone as a Suppressor of
Epithelial-Mesenchymal Transition and Metastatic Dissemination. *Mol Cancer Res*
18, 477-487. (2020).
4   Nakayama, J. & Makinoshima, H. Zebrafish-Based Screening Models for the
Identification of Anti-Metastatic Drugs. *Molecules* **25**, 2407 (2020).
5   Nakayama, J. & Gong, Z. Transgenic zebrafish for modeling hepatocellular carcinoma.
MedComm **1**, 146-156. (2020).
6   Nguyen., D. X., Bos., P. D. & Massagué., J. Metastasis: from dissemination to organ-
specific colonization. *Nat Rev Cancer* **9**, 274-284 (2009).
7   Welch, D. R. & Hurst, D. R. Defining the Hallmarks of Metastasis *Cancer research* **79**,
3011-3027. (2019).
8   Chaffer, C. L. & Weinberg, R. A. A perspective on cancer cell metastasis *Science* **331**,
1559-1564 (2011).
9   Lu, W. & Kang, Y. Epithelial-Mesenchymal Plasticit in Cancer Progression and
Metastasis. *Developmental Cell* **49**, 361-374 (2019).
10   Tsai, J. H. & Yang, J. Epithelial–mesenchymal plasticity in carcinoma metastasis.
Genes Dev **27**, 2192-2206 (2013).
11   Solnica-Krezel, L. Conserved patterns of cell movements during vertebrate
gastrulation *Curr Biol* **15**, R213-228. , doi:10.1016/j.cub.2005.03.016. (2005).
12   White, R. J. *et al.* A high-resolution mRNA expression time course of embryonic
development in zebrafish. *eLife* **6** (2017).
13   Yang., J., Parker, A. & Geert, B. Guidelines and definitions for research on epithelial-
mesenchymal transition. *Nat Rev Mol Cell Biol* **21**, 341-352 (2020).
14   Nieto, M. A., Huang, R. Y.-J., Jackson, R. A. & Thiery, J. P. EMT: 2016 *Cell* **166**, 21-45
(2016).
15   Thiery, J. P., Acloque, H., Huang, R. Y. J. & Nieto, M. A. Epithelial-mesenchymal
transitions in development and disease *Cell* **139**, 871-890 (2009).
16   Yang, J. & Weinberg, R. A. Epithelial-mesenchymal transition: at the crossroads of
development and tumor metastasis *Dev Cell* **14**, 818-829 (2008).
17   Dongre, A. & Weinberg, R. A. New insights into the mechanisms of epithelial-
mesenchymal transition and implications for cancer. *Nat Rev Mol Cell Biol* 20, 69-84
(2019).
18   Nakayama, J. *et al.* HSD11β1 promotes EMT-mediated breast cancer metastasis.
bioRxiv **doi: https://doi.org/10.1101/2021.09.27.461934** (2021).
19   Nakayama, J. *et al.* A zebrafish embryo screen utilizing gastrulation identifies the
HTR2C inhibitor pizotifen as a suppressor of EMT-mediated metastasis. *Elife* **10**,
e70151 (2021).
20   Nakayama, J., Konno, Y., Maruyama, A., Tomita, M. & Makinoshima, H. Cinnamon
bark extract suppresses metastatic dissemination of cancer cells through inhibition
of glycolytic metabolism *J Nat Med* (2022).

21 Gómez-Cuadrado, L., Tracey, N., Ma, R., Qian, B. & Brunton, V. G. Mouse models of
metastasis: progress and prospects *Dis Model Mech* **10**, 1061-1074. (2017).
22 Hoffman, R. M. Orthotopic metastatic (MetaMouse) models for discovery and
development of novel chemotherapy *Methods Mol Med* **111**, 297-322 (2005).
23 Chen, H.-C. Boyden chamber assay *Methods Mol Biol* **294**, 15-22 (2005).
24 Sack, U., Walther, W. & Scudiero, D. Novel effect of antihelminthic Niclosamide on
S100A4-mediated metastatic progression in colon cancer. *J Natl Cancer Inst* **103**,
1018-1036 (2011).
25 Itoh, K. *et al.* An essential part for Rho-associated kinase in the transcellular invasion
of tumor cells *Nat Med. 1999 Feb;5(2):* **2**, 221-225 (1999).
26 Marshall, J.-C. A. *et al.* Effect of inhibition of the lysophosphatidic acid receptor 1 on
metastasis and metastatic dormancy in breast cancer *J Natl Cancer Inst* **104**, 1306-
1319 (2012).
27 Gunnarsson, L., Jauhiainen, A., Kristiansson, E., Nerman, O. & Larsson, D. G. J.
Evolutionary conservation of human drug targets in organisms used for
environmental risk assessments. *Environmental Science & Technology* **42**, 5807–5813
(2008).
28 Pelka, K. E., Henn, K., Keck, A., Sapel, B. & Braunbeck, T. Size does matter –
Determination of the critical molecular size for the uptake of chemicals across the
chorion of zebrafish (Danio rerio) embryos. *Aquatic Toxicology* **185**, 1-10 (2017).
29 Urushibata, H. *et al.* Control of Developmental Speed in Zebrafish Embryos Using
Different Incubation Temperatures *Zebrafish* **18**, 316-325 (2021).
30 Yilmaz, O. *et al.* Scrambled eggs: Proteomic portraits and novel biomarkers of egg
quality in zebrafish (Danio rerio). *PLoS ONE* **12**, e0188084 (2017).
31 Kondo, M. Bone morphogenetic proteins in the early development of zebrafish. *FEBS
J* **274**, 2960-2967, doi:10.1111/j.1742-4658.2007.05838.x (2007).
32 Katsuno, Y. *et al.* Bone morphogenetic protein signaling enhances invasion and bone
metastasis of breast cancer cells through Smad pathway. *Oncogene* **27**, 6322-6333,
doi:10.1038/onc.2008.232 (2008).
33 Tada, M. & Smith, J. C. Xwnt11 is a target of Xenopus Brachyury: regulation of
gastrulation movements via Dishevelled, but not through the canonical Wnt
pathway. *Development* **127**, 2227-2238 (2000).
34 Vincan, E. & Barker, N. The upstream components of the Wnt signalling pathway in
the dynamic EMT and MET associated with colorectal cancer progression. *Clin Exp
Metastasis* **25**, 657-663, doi:10.1007/s10585-008-9156-4 (2008).
35 Yang, X., Dormann, D., Munsterberg, A. E. & Weijer, C. J. Cell movement patterns
during gastrulation in the chick are controlled by positive and negative chemotaxis
mediated by FGF4 and FGF8. *Dev Cell* **3**, 425-437 (2002).
36 Nomura, S. *et al.* FGF10/FGFR2 signal induces cell migration and invasion in
pancreatic cancer. *Br J Cancer* **99**, 305-313, doi:10.1038/sj.bjc.6604473 (2008).
37 Song, S. *et al.* Pou5f1-dependent EGF expression controls E-cadherin endocytosis,
cell adhesion, and zebrafish epiboly movements. *Dev Cell* **24**, 486-501,
doi:10.1016/j.devcel.2013.01.016 (2013).
38 Lu, Z., Jiang, G., Blume-Jensen, P. & Hunter, T. Epidermal growth factor-induced
tumor cell invasion and metastasis initiated by dephosphorylation and
downregulation of focal adhesion kinase. *Mol Cell Biol* **21**, 4016-4031,
doi:10.1128/MCB.21.12.4016-4031.2001 (2001).

39 Damm, E. W. & Winklbauer, R. PDGF-A controls mesoderm cell orientation and radial
intercalation during Xenopus gastrulation. *Development* **138**, 565-575,
doi:10.1242/dev.056903 (2011).
40 Jechlinger, M. *et al.* Autocrine PDGFR signaling promotes mammary cancer
metastasis. *J Clin Invest* **116**, 1561-1570, doi:10.1172/JCI24652 (2006).
41 Mizoguchi, T., Verkade, H., Heath, J. K., Kuroiwa, A. & Kikuchi, Y. Sdf1/Cxcr4 signaling
controls the dorsal migration of endodermal cells during zebrafish gastrulation.
Development **135**, 2521-2529, doi:10.1242/dev.020107 (2008).
42 Shen, H. B., Gu, Z. Q., Jian, K. & Qi, J. CXCR4-mediated Stat3 activation is essential for
CXCL12-induced cell invasion in bladder cancer. *Tumour Biol* **34**, 1839-1845,
doi:10.1007/s13277-013-0725-z (2013).
43 Montero, J.-A., Kilian, B., Chan, J., Bayliss, P. E. & Heisenberg, C.-P. Phosphoinositide
3-Kinase Is Required for Process Outgrowth and Cell Polarization of Gastrulating
Mesendodermal Cells. *Current Biology* **13**, 1279-1289, doi:10.1016/s0960-
9822(03)00505-0 (2003).
44 Wander, S. A. *et al.* PI3K/mTOR inhibition can impair tumor invasion and metastasis
in vivo despite a lack of antiproliferative action in vitro: implications for targeted
therapy. *Breast Cancer Res Treat* **138**, 369-381, doi:10.1007/s10549-012-2389-6
(2013).
45 Tsai, W. B., Zhang, X., Sharma, D., Wu, W. & Kinsey, W. H. Role of Yes kinase during
early zebrafish development. *Dev Biol* **277**, 129-141,
doi:10.1016/j.ydbio.2004.08.052 (2005).
46 Barraclough, J., Hodgkinson, C., Hogg, A., Dive, C. & Welman, A. Increases in c-Yes
Expression Level and Activity Promote Motility But Not Proliferation of Human
Colorectal Carcinoma Cells. *Neoplasia* **9**, 745-IN732, doi:10.1593/neo.07442 (2007).
47 Sharma, D., Holets, L., Zhang, X. & Kinsey, W. H. Role of Fyn kinase in signaling
associated with epiboly during zebrafish development. *Dev Biol* **285**, 462-476,
doi:10.1016/j.ydbio.2005.07.018 (2005).
48 Yadav, V. & Denning, M. F. Fyn is induced by Ras/PI3K/Akt signaling and is required
for enhanced invasion/migration. *Mol Carcinog* **50**, 346-352, doi:10.1002/mc.20716
(2011).
49 Krens, S. F. *et al.* Distinct functions for ERK1 and ERK2 in cell migration processes
during zebrafish gastrulation. *Dev Biol* **319**, 370-383,
doi:10.1016/j.ydbio.2008.04.032 (2008).
50 Radtke, S., Milanovic, M. & Rossé, C. ERK2 but not ERK1 mediates HGF-induced
motility in non-small cell lung carcinoma cell lines. *J Cell Sci.* **126**, 2381-2391 (2013).
51 Jopling, C., van Geemen, D. & den Hertog, J. Shp2 knockdown and Noonan/LEOPARD
mutant Shp2-induced gastrulation defects. *PLoS Genet* **3**, e225,
doi:10.1371/journal.pgen.0030225 (2007).
52 Wang, F. M. *et al.* SHP-2 promoting migration and metastasis of MCF-7 with loss of
E-cadherin, dephosphorylation of FAK and secretion of MMP-9 induced by IL-1beta
in vivo and in vitro. *Breast Cancer Res Treat* **89**, 5-14, doi:10.1007/s10549-004-1002-
z (2005).
53 Ip, Y. T. & Gridley, T. Cell movements during gastrulation: snail dependent and
independent pathways. *Curr Opin Genet Dev* **12**, 423-429 (2002).

54 Batlle, E., Sancho, E., Francí, C., Domínguez, D. & Monfar, M. The transcription factor
snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nat Cell*
Biol **2**, 84-89 (2000).
55 Shi, J., Severson, C., Yang, J., Wedlich, D. & Klymkowsky, M. W. Snail2 controls
mesodermal BMP/Wnt induction of neural crest. *Development* **138**, 3135-3145,
doi:10.1242/dev.064394 (2011).
56 Medici, D., Hay, E. D. & B.R., O. Snail and Slug Promote Epithelial-Mesenchymal
Transition through -Catenin–T-Cell Factor-4-dependent Expression of Transforming
Growth Factor-3. *Molecular Biology of the Cell* **19**, 4875–4887, doi:10.1091/mbc.E08
(2008).
57 Castanon, I. & Baylies, M. K. A Twist in fate: evolutionary comparison of Twist
structure and function. *Gene* **287**, 11-22 (2002).
58 Yang, J. *et al.* Twist, a master regulator of morphogenesis, plays an essential role in
tumor metastasis. *Cell* **117**, 927-939, doi:10.1016/j.cell.2004.06.006 (2004).
59 Fernando, R. I. *et al.* The T-box transcription factor Brachyury promotes epithelial-
mesenchymal transition in human tumor cells. *J Clin Invest* **120**, 533-544,
doi:10.1172/JCI38379 (2010).
60 Vannier, C., Mock, K., Brabletz, T. & Driever, W. Zeb1 regulates E-cadherin and
Epcam (epithelial cell adhesion molecule) expression to control cell behavior in early
zebrafish development. *J Biol Chem* **288**, 18643-18659,
doi:10.1074/jbc.M113.467787 (2013).
61 Spaderna, S. *et al.* The transcriptional repressor ZEB1 promotes metastasis and loss
of cell polarity in cancer. *Cancer Res* **68**, 537-544, doi:10.1158/0008-5472.CAN-07-
5682 (2008).
62 Sander, V., Reversade, B. & De Robertis, E. M. The opposing homeobox genes
Goosecoid and Vent1/2 self-regulate Xenopus patterning. *EMBO J* **26**, 2955-2965,
doi:10.1038/sj.emboj.7601705 (2007).
63 Hartwell, K. A. *et al.* The Spemann organizer gene, Goosecoid, promotes tumor
metastasis. *Proc Natl Acad Sci U S A* **103**, 18969-18974,
doi:10.1073/pnas.0608636103 (2006).
64 Wilm, B., James, R. G., Schultheiss, T. M. & Hogan, B. L. The forkhead genes, Foxc1
and Foxc2, regulate paraxial versus intermediate mesoderm cell fate. *Dev Biol* **271**,
176-189, doi:10.1016/j.ydbio.2004.03.034 (2004).
65 Mani, S. A. *et al.* Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and
is associated with aggressive basal-like breast cancers. *Proc Natl Acad Sci U S A* **104**,
10069-10074, doi:10.1073/pnas.0703900104 (2007).
66 Miyagi, C. *et al.* STAT3 noncell-autonomously controls planar cell polarity during
zebrafish convergence and extension. *J Cell Biol* **166**, 975-981,
doi:10.1083/jcb.200403110 (2004).
67 Abdulghani, J. *et al.* Stat3 promotes metastatic progression of prostate cancer. *Am J*
Pathol **172**, 1717-1728, doi:10.2353/ajpath.2008.071054 (2008).
68 Lachnit, M., Kur, E. & Driever, W. Alterations of the cytoskeleton in all three
embryonic lineages contribute to the epiboly defect of Pou5f1/Oct4 deficient MZspg
zebrafish embryos. *Dev Biol* **315**, 1-17, doi:10.1016/j.ydbio.2007.10.008 (2008).
69 Dai, X. *et al.* OCT4 regulates epithelial-mesenchymal transition and its knockdown
inhibits colorectal cancer cell migration and invasion. *Oncol Rep* **29**, 155-160,
doi:10.3892/or.2012.2086 (2013).

70 O'Carroll, D. *et al.* The polycomb-group gene Ezh2 is required for early mouse
development. *Mol Cell Biol* **21**, 4330-4336, doi:10.1128/MCB.21.13.4330-4336.2001
(2001).
71 Ren, G. *et al.* Polycomb protein EZH2 regulates tumor invasion via the transcriptional
repression of the metastasis suppressor RKIP in breast and prostate cancer. *Cancer*
Res **72**, 3091-3104, doi:10.1158/0008-5472.CAN-11-3546 (2012).
72 Lin, F. *et al.* Essential roles of G{alpha}12/13 signaling in distinct cell behaviors
driving zebrafish convergence and extension gastrulation movements. *J Cell Biol* **169**,
777-787, doi:10.1083/jcb.200501104 (2005).
73 Chen, M. W. *et al.* H3K9 histone methyltransferase G9a promotes lung cancer
invasion and metastasis by silencing the cell adhesion molecule Ep-CAM. *Cancer Res*
70, 7830-7840, doi:10.1158/0008-5472.CAN-10-0833 (2010).
74 van der Lugt, N. M. *et al.* Posterior transformation, neurological abnormalities, and
severe hematopoietic defects in mice with a targeted deletion of the bmi-1 proto-
oncogene. *Genes Dev* **8**, 757-769 (1994).
75 Guo, B. H. *et al.* Bmi-1 promotes invasion and metastasis, and its elevated expression
is correlated with an advanced stage of breast cancer. *Mol Cancer* **10**, 10,
doi:10.1186/1476-4598-10-10 (2011).
76 Zhu, S., Liu, L., Korzh, V., Gong, Z. & Low, B. C. RhoA acts downstream of Wnt5 and
Wnt11 to regulate convergence and extension movements by involving effectors
Rho kinase and Diaphanous: use of zebrafish as an in vivo model for GTPase
signaling. *Cell Signal* **18**, 359-372, doi:10.1016/j.cellsig.2005.05.019 (2006).
77 Yoshioka, K., Nakamori, S. & Itoh, K. Overexpression of small GTP-binding protein
RhoA promotes invasion of tumor cells. *Cancer Res* **59**, 2004-2010 (1999).
78 Choi, S. C. & Han, J. K. Xenopus Cdc42 regulates convergent extension movements
during gastrulation through Wnt/Ca2+ signaling pathway. *Dev Biol* **244**, 342-357,
doi:10.1006/dbio.2002.0602 (2002).
79 Reymond, N. *et al.* Cdc42 promotes transendothelial migration of cancer cells
through beta1 integrin. *J Cell Biol* **199**, 653-668, doi:10.1083/jcb.201205169 (2012).
80 Habas, R., Dawid, I. B. & He, X. Coactivation of Rac and Rho by Wnt/Frizzled signaling
is required for vertebrate gastrulation. . *Genes Dev* **17**, 295-309., doi:10.1101/
(2003).
81 Vega, F. M. & Ridley, A. J. Rho GTPases in cancer cell biology. *FEBS Lett* **582**, 2093-
2101, doi:10.1016/j.febslet.2008.04.039 (2008).
82 Marlow, F., Topczewski, J., Sepich, D. & Solnica-Krezel, L. Zebrafish Rho kinase 2 acts
downstream of Wnt11 to mediate cell polarity and effective convergence and
extension movements. *Curr Biol* **12**, 876-884 (2002).
83 Itoh, K. *et al.* An essential part for Rho-associated kinase in the transcellular invasion
of tumor cells. *Nat Med* **5**, 221-225, doi:10.1038/5587 (1999).
84 Kusakabe, M. & Nishida, E. The polarity-inducing kinase Par-1 controls Xenopus
gastrulation in cooperation with 14-3-3 and aPKC. *EMBO J* **23**, 4190-4201,
doi:10.1038/sj.emboj.7600381 (2004).
85 Shi, X., Gangadharan, B., Brass, L. F., Ruf, W. & Mueller, B. M. Protease-activated
receptors (PAR1 and PAR2) contribute to tumor cell motility and metastasis. *Mol*
Cancer Res **2**, 395-402 (2004).
86 Gunaratne, A., Thai, B. L. & Di Guglielmo, G. M. Atypical protein kinase C
phosphorylates Par6 and facilitates transforming growth factor beta-induced

epithelial-to-mesenchymal transition. *Mol Cell Biol* **33**, 874-886,
doi:10.1128/MCB.00837-12 (2013).

87   Seifert, K., Ibrahim, H., Stodtmeister, T., Winklbauer, R. & Niessen, C. M. An
adhesion-independent, aPKC-dependent function for cadherins in morphogenetic
movements. *J Cell Sci* **122**, 2514-2523, doi:10.1242/jcs.042796 (2009).

88   Yamazaki, K. *et al.* Adenylate cyclase-associated protein 1 overexpressed in
pancreatic cancers is involved in cancer cell motility. *Lab Invest* **89**, 425-432,
doi:10.1038/labinvest.2009.5 (2009).

89   Link, V. *et al.* Identification of regulators of germ layer morphogenesis using
proteomics in zebrafish. *J Cell Sci* **119**, 2073-2083, doi:10.1242/jcs.02928 (2006).

90   Khanna, C. *et al.* The membrane-cytoskeleton linker ezrin is necessary for
osteosarcoma metastasis. *Nat Med* **10**, 182-186, doi:10.1038/nm982 (2004).

91   Slanchev, K. *et al.* The epithelial cell adhesion molecule EpCAM is required for
epithelial morphogenesis and integrity during zebrafish epiboly and skin
development. *PLoS Genet* **5**, e1000563, doi:10.1371/journal.pgen.1000563 (2009).

92   Ni, J. *et al.* Role of the EpCAM (CD326) in prostate cancer metastasis and
progression. *Cancer Metastasis Rev* **31**, 779-791, doi:10.1007/s10555-012-9389-1
(2012).

93   Skalski, M., Alfandari, D. & Darribere, T. A key function for alphav containing
integrins in mesodermal cell migration during Pleurodeles waltl gastrulation. *Dev Biol*
195, 158-173, doi:10.1006/dbio.1997.8838 (1998).

94   Felding-Habermann, B. Integrin adhesion receptors in tumor metastasis. *Clin Exp*
Metastasis **20**, 203-213 (2003).

95   Marsden, M. & DeSimone, D. W. Integrin-ECM Interactions Regulate Cadherin-
Dependent Cell Adhesion and Are Required for Convergent Extension in Xenopus.
Current Biology **13**, 1182-1191, doi:10.1016/s0960-9822(03)00433-0 (2003).

96   Malik, G. *et al.* Plasma fibronectin promotes lung metastasis by contributions to
fibrin clots and tumor cell invasion. *Cancer Res* **70**, 4327-4334, doi:10.1158/0008-
5472.CAN-09-3312 (2010).

97   Bakkers, J. *et al.* Has2 is required upstream of Rac1 to govern dorsal migration of
lateral cells during zebrafish gastrulation. *Development* **131**, 525-537,
doi:10.1242/dev.00954 (2004).

98   Kim, H. R. *et al.* Hyaluronan facilitates invasion of colon carcinoma cells in vitro via
interaction with CD44. *Cancer Res* **64**, 4569-4576, doi:10.1158/0008-5472.CAN-04-
0202 (2004).

99   Coyle, R. C., Latimer, A. & Jessen, J. R. Membrane-type 1 matrix metalloproteinase
regulates cell migration during zebrafish gastrulation: evidence for an interaction
with non-canonical Wnt signaling. *Exp Cell Res* **314**, 2150-2162,
doi:10.1016/j.yexcr.2008.03.010 (2008).

100   Perentes, J. Y. *et al.* Cancer cell-associated MT1-MMP promotes blood vessel
invasion and distant metastasis in triple-negative mammary tumors. *Cancer Res* **71**,
4527-4538, doi:10.1158/0008-5472.CAN-10-4376 (2011).

101   Cha, Y. I. *et al.* Cyclooxygenase-1-derived PGE2 promotes cell motility via the G-
protein-coupled EP4 receptor during vertebrate gastrulation. *Genes Dev* **20**, 77-86,
doi:10.1101/gad.1374506 (2006).

102 Kundu, N. & Fulton, A. M. Selective cyclooxygenase (COX)-1 or COX-2 inhibitors
control metastatic disease in a murine model of breast cancer. *Cancer Res* **62**, 2343-
2346 (2002).

103 Speirs, C. K. *et al.* Prostaglandin Gbetagamma signaling stimulates gastrulation
movements by limiting cell adhesion through Snai1a stabilization. *Development* **137**,
1327-1337, doi:10.1242/dev.045971 (2010).

104 Wang, D. & Dubois, R. N. Prostaglandins and cancer. *Gut* **55**, 115-122,
doi:10.1136/gut.2004.047100 (2006).

105 Yamashita, S. *et al.* Zinc transporter LIVI controls epithelial-mesenchymal transition
in zebrafish gastrula organizer. *Nature* **429**, 298-302, doi:10.1038/nature02545
(2004).

106 Lue, H. W. *et al.* LIV-1 promotes prostate cancer epithelial-to-mesenchymal
transition and metastasis through HB-EGF shedding and EGFR-mediated ERK
signaling. *PLoS One* **6**, e27720, doi:10.1371/journal.pone.0027720 (2011).

107 Yagi, H. *et al.* A synthetic biology approach reveals a CXCR4-G13-Rho signaling axis
driving transendothelial migration of metastatic breast cancer cells. *Sci Signal* **4**,
ra60, doi:10.1126/scisignal.2002221 (2011).

108 Webster, D. M. *et al.* O-GlcNAc modifications regulate cell survival and epiboly
during zebrafish development. *BMC Dev Biol* **9**, 28, doi:10.1186/1471-213X-9-28
(2009).

109 Lynch, T. P. *et al.* Critical role of O-Linked beta-N-acetylglucosamine transferase in
prostate cancer invasion, angiogenesis, and metastasis. *J Biol Chem* **287**, 11070-
11081, doi:10.1074/jbc.M111.302547 (2012).

110 Latinkic, B. V. Xenopus Cyr61 regulates gastrulation movements and modulates Wnt
signalling. *Development* **130**, 2429-2441, doi:10.1242/dev.00449 (2003).

111 Lin, J. *et al.* A novel anti-Cyr61 antibody inhibits breast cancer growth and metastasis
in vivo. *Cancer Immunol Immunother* **61**, 677-687, doi:10.1007/s00262-011-1135-y
(2012).

112 Liu, W. *et al.* TRPM7 regulates gastrulation during vertebrate embryogenesis. *Dev
Biol* **350**, 348-357, doi:10.1016/j.ydbio.2010.11.034 (2011).

113 Middelbeek, J. *et al.* TRPM7 is required for breast tumor cell metastasis. *Cancer Res*
72, 4250-4261, doi:10.1158/0008-5472.CAN-11-3863 (2012).

114 Holloway, B. A. *et al.* A novel role for MAPKAPK2 in morphogenesis during zebrafish
development. *PLoS Genet* **5**, e1000413, doi:10.1371/journal.pgen.1000413 (2009).

115 Kumar, B. *et al.* p38 mitogen-activated protein kinase-driven MAPKAPK2 regulates
invasion of bladder cancer by modulation of MMP-2 and MMP-9 activity. *Cancer Res*
70, 832-841, doi:10.1158/0008-5472.CAN-09-2918 (2010).

116 Machingo, Q. J., Fritz, A. & Shur, B. D. A beta1,4-galactosyltransferase is required for
convergent extension movements in zebrafish. *Dev Biol* **297**, 471-482,
doi:10.1016/j.ydbio.2006.05.024 (2006).

117 Zhu, X. *et al.* Elevated beta1,4-galactosyltransferase I in highly metastatic human
lung cancer cells. Identification of E1AF as important transcription activator. *J Biol
Chem* **280**, 12503-12516, doi:10.1074/jbc.M413631200 (2005).

118 Hong, S. K., Tanegashima, K. & Dawid, I. B. XIer2 is required for convergent extension
movements during Xenopus development. *Int J Dev Biol* **55**, 917-921,
doi:10.1387/ijdb.113288sh (2011).

119 Neeb, A. *et al.* The immediate early gene Ier2 promotes tumor cell motility and
metastasis, and predicts poor survival of colorectal cancer patients. *Oncogene* **31**,
3796-3806, doi:10.1038/onc.2011.535 (2012).
120 Besser, J., Leito, J. T., van der Meer, D. L. & Bagowski, C. P. Tip-1 induces filopodia
growth and is important for gastrulation movements during zebrafish development.
Dev Growth Differ **49**, 205-214, doi:10.1111/j.1440-169X.2007.00921.x (2007).
121 Han, M., Wang, H., Zhang, H. T. & Han, Z. The PDZ protein TIP-1 facilitates cell
migration and pulmonary metastasis of human invasive breast cancer cells in
athymic mice. *Biochem Biophys Res Commun* **422**, 139-145,
doi:10.1016/j.bbrc.2012.04.123 (2012).
122 Faure, S. *et al.* Xenopus p21-activated kinase 5 regulates blastomeres' adhesive
properties during convergent extension movements. *Dev Biol* **277**, 472-492,
doi:10.1016/j.ydbio.2004.10.005 (2005).
123 Gong, W. *et al.* P21-activated kinase 5 is overexpressed during colorectal cancer
progression and regulates colorectal carcinoma cell adhesion and migration. *Int J*
Cancer **125**, 548-555, doi:10.1002/ijc.24428 (2009).
124 Iioka, H., Ueno, N. & Kinoshita, N. Essential role of MARCKS in cortical actin dynamics
during gastrulation movements. *J Cell Biol* **164**, 169-174, doi:10.1083/jcb.200310027
(2004).
125 Rombouts, K. *et al.* Myristoylated Alanine-Rich protein Kinase C Substrate (MARCKS)
expression modulates the metastatic phenotype in human and murine colon
carcinoma in vitro and in vivo. *Cancer Lett* **333**, 244-252,
doi:10.1016/j.canlet.2013.01.040 (2013).
659