A zebrafish embryo screen utilizing gastrulation for identification of anti-metastasis drugs

Joji Nakayama[1,2,3,§], Hideki Makinoshima[1,4] , and Zhiyuan Gong[3]

[1]Tsuruoka Metabolomics Laboratory, National Cancer Center, Tsuruoka, Japan

[2]Shonai Regional Industry Promotion Center, Tsuruoka, Japan.

[3]Department of Biological Science, National University of Singapore, Singapore

[4]Division of Translational Research, Exploratory Oncology Research and Clinical Trial Center,

National Cancer Center, Kashiwa, Japan.

[§]Corresponding authors:

Joji Nakayama

Tsuruoka Metabolomics Laboratory, National Cancer Center, Mizukami 246-2, Kakuganji,

Tsuruoka, Yamagata, Japan 975-0052.

E-mail: zmetastasis@gmail.com

ORCID: 0000-0003-1077-140X

Tel: +81-235-64-0980

Keywords: Metastasis, gastrulation, EMT, Phenotyping screening, zebrafish

Total words excluding references and figure legends: 2158

Number of figures: 4

Number of tables: 1

Abstract

Few models exist that allow for rapid and effective screening of anti-metastasis drugs. Here, we present a phenotype-based chemical screen utilizing gastrulation of zebrafish embryos for identification of anti-metastasis drugs. Based on our hypothesis that small molecules which interrupt zebrafish gastrulation might suppress metastasis progression of cancer cells, we designed a drug screening concept which uses epiboly, the first morphogenetic movement in gastrulation, as a marker. The screen only needs zebrafish embryos and enables 100 chemicals to be tested in five hours through just observing epiboly progression of a test chemical-treated embryos. In the screen, embryos at two-cell stage are firstly corrected and then developed to sphere stage. The embryos are treated with a test chemical and incubated in the presence of the chemical until vehicle-treated embryos develop to 90% epiboly stage. Finally, positive 'hit' chemicals that interrupt epiboly progression are selected through comparing epiboly progression of a test chemical-treated embryos with that of vehicle-treated embryos under a stereoscopic microscope. Previous study subjected 1280 FDA-approved drugs into the screen and identified Adrenosterone and Pizotifen as epiboly-interrupting drugs. Mouse models of metastasis validated that the drugs could suppress metastasis of breast cancer cells. Furthermore, the screen can be diverted to a chemical genetic screening platform for identification of metastasis-promoting genes. Hydroxysteroid (11-Beta) Dehydrogenase 1 (HSD11b1) and serotonin receptor 2C (HTR2C), which are primary target of Adrenosterone and Pizotifen respectively, induces epithelial-mesenchymal transition (EMT) and promote metastasis of breast cancer cells.

Introduction

Cancer research using zebrafish as a model has attracted attention because this model offers many unique advantages that are not readily provided by other animal models. Moreover, the zebrafish system has been increasingly recognized as a platform for chemical screening because it provides the advantage of high-throughput screening in an in vivo vertebrate setting with physiologic relevance to humans [1-4].

Metastasis is responsible for approximately 90% of cancer-associated mortality. It proceeds through multiple steps: invasion, intravasation, survival in the circulatory system, extravasation, colonization, and metastatic tumor formation in secondary organs with angiogenesis [5-7]. Dissemination of cancer cells is an initial step of metastasis and its molecular mechanism involves local breakdown of basement membrane, loss of cell polarity, and induction of EMT [8,9]. These cellular, biological phenomena are also observed during vertebrate gastrulation in that evolutionarily conserved morphogenetic movements of epiboly, internalization, convergence and extension progress [10]. In zebrafish, the first morphogenetic movement, epiboly, is initiated at approximately four hours post fertilization (hpf) to move cells from the animal pole to eventually engulf the entire yolk cell by 10 hpf. These movements are governed by the molecular mechanisms that are induced by temporally and spatially regulated gene expression, and these mechanisms and changes in gene expression are partially observed in metastatic progression.

Development of the protocol

At least fifty common genes were shown to be involved in both gastrulation and metastasis progression (Table 1) [11-15]. The fifty genes are expressed in xenopus or zebrafish embryos, and genetic inhibition of each of the fifty genes in these embryos interferes with gastrulation progression. Conversely, the same fifty genes are ectopically expressed in metastatic cancer cells and confer metastatic properties on cancer cells, and genetic inhibition of each of the same fifty

genes suppresses metastasis progression. This evidence led us to hypothesize that small

molecules which interfere with zebrafish gastrulation might suppress metastasis progression of

cancer cells. Based on the hypothesis, we designed a drug screening concept which uses epiboly,

the first morphogenetic movement in gastrulation, as a marker. This screen measures suppressor

effect of each of chemicals based on epiboly progression of zebrafish embryos (Fig. 1 and 2).

Applications of the method

This screen enables 100 chemicals to be tested in five hours and succeeds to identified two FDA-

approved drugs and one crude drug have suppressor effects on metastasis. Adrenosterone and

Pizotifen, which were the identified drugs, were validated to suppress metastasis of breast cancer

cells in mouse models of metastasis (Fig. 3) [16,17]. Cinnamon bark extract, which was the

identified crude drug, was validated to suppress metastatic dissemination of breast cancer cells in

zebrafish xenograft model [18]. Furthermore, this screen can be diverted to a chemical genetic

screening platform for identification of metastasis-promoting genes. Hydroxysteroid (11-Beta)

Dehydrogenase 1 (HSD11b1) and serotonin receptor 2C (HTR2C), which are primary target of

Adrenosterone and Pizotifen respectively, induces epithelial-mesenchymal transition (EMT) and

promote metastasis of breast cancer cells (Fig. 4) [16,17].

Comparison with other methods

Current mouse models of metastasis are too expensive and time consuming to use for rapid and

high-throughput screening [19,20]. Also in vitro model of metastatic dissemination such as a Boyden

chamber assay can test a limited number of chemicals in one assay and needs huge times and

efforts in analyzing the results[21]. Therefore, the discovery of anti-metastatic drugs is based

essentially on accumulated knowledge of the molecular mechanisms of metastasis: identifying

therapeutic targets for inhibiting metastasis, evaluating the effect of candidate chemicals and the

markers which are used to measure a suppressor effect [22-24]. In contrast, our screen only needs

zebrafish embryos and enables 100 chemicals to be tested in five hours through just observing

epiboly progression of a test chemical-treated embryos. Furthermore, out of the 78 chemicals

which interrupt epiboly progression of zebrafish embryos, 20 of the chemicals were validated to

suppress cell motility and invasion of highly metastatic human cancer cells without affecting cell

viability in a Boyden chamber assay. Disadvantages of this screen is that 75% of the chemicals

which interrupt epiboly progression of zebrafish embryos, fail to suppress cell motility and

invasion of highly metastatic human cancer cells in a Boyden chamber assay [17].

Experimental design

This screen measures suppressor effect of each of chemicals based on epiboly progression of

zebrafish embryos. Niclosamid or DMSO is used as positive or negative control, respectively.

Epiboly progression of each of chemical treated embryos is compared with that of DMSO-treated

embryos. Firstly, embryos at two cell stage are corrected and approximately twenty embryos are

arranged into each well of 24-well plate. Each of chemicals are added when the embryos develop

to sphere stage. Suppressor effects of each of chemicals on the epiboly progression are measured

under stereoscopic microscope when DMSO-treated embryos develop to 90% epiboly stage [17].

Limitations

There is a limitation in delivering chemicals to zebrafish embryos. Zebrafish embryos are

surrounded by the acellular chorion, which is known to be about 1.5–2.5 μm thick and to consist

of three layers pierced by pore canals. The pore allows passage of water, ions, and chemicals. A

study reported molecules which are larger than 3-4KDa fail to pass through the chorion.

Therefore, this screen can not measure suppressor effect of the molecules which are larger than 3-

4KDa [25].

Materials

REAGENTS

• Wild-type AB zebrafish strain

• E3 medium (5.0 mM NaCl, 0.17 mM KCl, 0.33 mM $MgSO_4$)

• FDA, EMA, and other agencies-approved chemical libraries were purchased from

Prestwick Chemical (Illkirch, France).

• Niclosamid

• DMSO

EQUIPMENT

• 24-well flat bottom plastic plates (Corning)

• Stereomicroscope (MZ75, Leica)

• Incubator (Thermo)

Procedure

Zebrafish mating setup (Day 0) _Timing 10 mins

1. On the night before collecting embryos, arrange male and female zebrafish in pairs

separated by divider

CRITICAL STEP

Young adult zebrafish should be used for the crossing. Quality of zebrafish embryos

affect screening efficiency.

Embryo collection, and distribution (Day 1) _Timing 10-30 min

2. Remove the divider to allow the fish to spawn.

3. To obtain same development stage of zebrafish embryos, zebrafish were crossed for 10

mins. If more than twenty chemicals were tested, the crossing were conducted three times

at three different time points (Group A_8:30, Group B_9:00 and Group C_9:30AM).

4. After 10mins, set back divider to prevent zebrafish from spawning.

CRITICAL STEP

This screen measures suppressor effect of each of drugs by progression of epiboly in live

zebrafish embryos. Therefore, epiboly proceeds during measuring the effect under

stereoscopic microscope. If more than 20 chemicals were tested, screening should be

divided into more than two sessions and each of the sessions start at different time point.

For example, if screen test 60 chemicals, zebrafish should be crossed at three different

time points over 30 mins apart. To do that, 30 mins for measuring the effects would be

ensured.

5. Collect the embryos and remove dead embryos

6. Incubate the embryos at 27 ˚C for twenty mins and then collect embryos at two cell stage

under stereoscopic microscope.

7. Array approximately twenty embryos into each well of 24 well plate

8. Remove E3 medium from each well including the embryos by pipet

9. Add 900 µl of E3 medium water to the well.

Develop embryos to sphere stage_Timing 4 hours

10. Incubate the embryos at 27 ˚C until the embryos develop to sphere stage

Addition of chemicals_Timing 30 mins

11. Before 30 mins from adding chemicals to embryos, prepare E3 medium containing

chemicals. Prepare 100 µl of 10-fold concentration of chemicals-containing E3 medium

containing desired final concentration

12. Add 100 µl of the solution into each well when the embryos develop to sphere stage.

13. First 22 chemicals, Niclosamid as positive control, and DMSO as negative control are

added into group A when embryos from group A develop to sphere stage.

14. Second 22 chemicals, Niclosamid, and DMSO were added into group B when embryos from group B develop to sphere stage.

15. Last 22 chemicals, Niclosamid, and DMSO were added into group C when embryos from group C develop to sphere stage

Develop DMSO-treated embryos to 90% epiboly stage_Timing 5 hours

16. After the addition of chemicals, the embryos were incubated at 27 ˚C for approximately five hours

Measuring the inhibition effects of each of chemicals_Timing 30 mins

17. Comparing epiboly progression of each of chemicals-treated embryos from group A with that of DMSO-treated embryos from group A under stereoscopic microscope when DMSO-treated embryos from group A develop to 90% epiboly stage.

18. After approximately 30 mis from observing embryos from group A, measuring the inhibition effects of each of chemicals on epiboly progression of embryos from group B and when DMSO-treated embryos from group B develop to 90% epiboly stage.

19. After approximately 30 mis from observing embryos from group B, measuring the inhibition effects of each of chemicals on epiboly progression of embryos from group C and when DMSO-treated embryos from group C develop to 90% epiboly stage.

Timing

Step 1, Zebrafish mating setup: overnight

Step 2-4, Crossing zebrafish: 10 mins

Step 5+6, Collect two cell stage embryos: 20 mins

Step 7-9, Array 20 embryos into each well of 24-well plate: 30 mins

Step 10, Develop the embryos to sphere stage: 4 hours

Step 11-15, Prepare chemical drugs and add the solution into each well of 24-well plate: 30 mins

Step 16, Develop the embryos to 90% epiboly stage: 5 hours

Step 17-19, measuring epiboly progression: 30 mins

Troubleshooting

Quality of zebrafish embryos affect screening efficiency. A study which investigates

development of 12,919 fertilized embryos from zebrafish AB line reveals that by three hours post

fertilization, 290 (2.3%) or 2,055 (16.3%) of the fertilized eggs die or show morphological

abnormalities, respectively. Low quality of zebrafish embryos would generate false 'hit'

chemicals since our screen measures suppressor effect of a test chemical through just observing

epiboly progression of the chemical-treated embryos. If the number of zebrafish embryos

showing morphological abnormalities correlate with final concentration of a test chemical, the

abnormalities may result from an effect of the test chemical on the embryos.

Anticipated results

Suppressor effects of tested chemicals on epiboly progression of zebrafish embryos are

significantly affected by final concentration of the chemicals. Previous study showed 0.9%

(12/1280) of the drugs induced severe or complete arrest of embryonic cell movement when

embryos were treated with 10 μM. 5.2% (66/1280) of the drugs induced either delayed epiboly or

interrupted epiboly of the embryos. 93.3% (1194/1280) of drugs have no effect on epiboly

progression of the embryos. 0.6% (8/1280) of drugs induced toxic lethality. Out of the 78

epiboly-interrupting drugs, 25% of the chemicals succeed to suppress cell motility and invasion

of highly metastatic human cancer cells in a Boyden chamber assay. In contrast, epiboly

progression was affected more severely when embryos were treated with 50 μM. However,

almost of the chemicals which interrupt epiboly progression of zebrafish embryos when embryos

were treated with 50 μM, failed to suppress cell motility and invasion of highly metastatic human

cancer cells in a Boyden chamber assay [17].

Author contributions statements

J.N. created a concept of this screen, designed research, conducted experiments, analyzed data,

wrote and edited original draft, supervised this project; H.M. and Z.G. administered this project,

acquired funding.

ORCID for corresponding authors

Joj Nakayama: ORCID: 0000-0003-1077-140X

Acknowledgments

We thank Dr. Herrick (Albany Medical College) for providing pCMV-h5TH2C-VSV with us.

This study was funded by grants from National Medical Research Council of Singapore (R-

154000547511) and Ministry of Education of Singapore (R-154000A23112) to ZG. Experiments

using mice were supported in part by research funds from the Yamagata prefectural government

and the City of Tsuruoka.

Competing interests

J.N., Z.G., and H.M. declare no conflict of interest.

Fig. 1

Fig. 1_Graphic schematic of a phenotype-based chemical screen using zebrafish embryos

Pairs of adult zebrafish are crossed and their embryos at two-cell stage are collected and arrayed

into individual wells of 24-well plate. Chemicals are added into each well when the embryos

develop to sphere stage. Epiboly progression of each of chemicals-treated embryos are compared

with that of DMSO-treated embryos under a stereoscopic microscope when DMSO-treated

embryos develop to 90% epiboly stage.

Fig. 2

Niclosamide	Hexachlorophene	Pizotifen	Ipriflavone	DMSO
Tolcapone	Troglitazone	Vinpocetine	Zuclopenthxiol	Mebendazole
Severe delayed	Delayed		Slighly delayed	Non effect

Fig. 2_Representative samples of the embryos that were treated with indicated chemicals.

Indicated chemicals were added when the embryos develop to the sphere stage. Embryos were

treated with 10 μM. Niclosamide-treated embryos serve as positive control and DMSO-treated

embryos serve as negative control. Epiboly progression of each of chemicals-treated embryos are

compared with that of DMSO-treated embryos under a stereoscopic microscope when DMSO-

treated embryos develop to 90% epiboly stage.

Fig. 3

Vehicle-treated mice   **Pizotifen-treated mice**

Fig. 3_Pizotifen, one of epiboly-interrupting drugs, suppressed metastatic progression

of breast cancer cells in vitro and vivo

a, Effect of pizotifen on cell motility and invasion of MBA-MB-231, MDA-MB-435, and

PC9 cells. Either vehicle or pizotifen treated the cells were subjected to Boyden chamber

assays. Fetal bovine serum (1% v/v) was used as the chemoattractant in both assays. Each

experiment was performed at least twice. b, Representative images of primary tumors on day

10 post injection (top panels) and metastatic burden on day 70 post injection (bottom panels)

taken using an IVIS Imaging System. c, Number of metastatic nodules in the lung of either

vehicle- or pizotifen-treated mice.

Fig. 4

Fig. 4_ HTR2C, a primary target of Pizotifen, induces epithelial-to-mesenchymal

transition (EMT)-mediated metastatic dissemination of human cancer cells

a, Immunofluorescence staining of E-cadherin and vimentin expressions in the MCF7 cells.

b, Expression of E-cadherin and vimentin, and HTR2C was examined by western blotting in

the MCF7 and HaCaT cells; GAPDH loading control is shown (bottom). c, Representative

images of dissemination patterns of MCF7 cells expressing either the control vector (top left)

or HTR2C (lower left) in a zebrafish xenotransplantation model. White arrow heads indicate

disseminated MCF7 cells. The mean frequencies of the fish showing head, trunk, or end-tail

dissemination tabulated (right). Each value is indicated as the mean ± SEM of two

independent experiments. Statistical analysis was determined by Student's t test.

Table 1. A list of the common fifty genes that are involved between gastrulation and

metastasis progression

Genes	Gastrulation Defects	Ref	Effects in Metastasis	Ref
BMP	Convergence and extension	26	EMT	27
WNT	Convergence and extension	28	Migration and Invasion	29
FGF	Convergence and extension	30	Invision	31
EGF	Epiboly	32	Migration	33
PDGF	Convergence and extension	34	EMT	35
CXCL12	Migration of endodermal cells	36	Migration and Invasion	37
CXCR4	Migration of endodermal cells	36	Migration and Invasion	37
PIK3CA	Convergence and extension	38	Migration and Invasion	39
YES	Epiboly	40	Migration	41
FYN	Epiboly	42	Migration and Invasion	43
MAPK1	Epiboly	44	Migration	45
SHP2	Convergence and extension	46	Migration	47
SNAI1	Convergence and extension	48	EMT	49
SNAI2	Mesoderm & Neural crest formation	50	EMT	51
TWIST1	Mesoderm formation	52	EMT	53
TBXT	Convergence and extension	28	EMT	54
ZEB1	Epiboly	55	EMT	56
GSC	Mesodermal patterning	57	EMT	58
FOXC2	Unclear, defects in gastrulation	59	EMT	60
STAT3	Convergence and extension	61	Migration	62
POU5F1	Epiboly	63	EMT	64
EZH2	Unclear, defects in gastrulation	65	Invasion	66
EHMT2	Defects in Neurogenesis	67	Migration and Invasion	68
BMI1	Defects in skelton formation	69	EMT	70
RHOA	Convergence and extension	71	Migration and Invasion	72
CDC42	Convergence and extension	73	Migration and Invasion	74
RAC1	Convergence and extension	75	Migration and Invasion	76
ROCK2	Convergence and extension	77	Migration and Invasion	78
PAR1	Convergence and extension	79	Migration	80
PRKCI	Convergence and extension	79	EMT	81
CAP1	Convergence and extension	82	Migration	83
EZR	Epiboly	84	Migration	85
EPCAM	Epiboly	86	Migration and Invasion	87
ITGB1 / ITA5	Mesodermal Migration	88	Migration and Invasion	89
FN1	Convergence and extension	90	Invasion	91
HAS2	Dorsal migration of lateral cells	92	Invasion	93
MMP14	Convergence and extension	94	Invasion	95
COX1	Epiboly	96	Invasion	97
PTGES	Convergence and extension	98	Invasion	99
SLC39A6	Aterior migration	100	EMT	101
GNA12 / 13	Convergence and extension	67	Migration and Invasion	102
OGT	Epiboly	103	Migration and Invasion	104
CCN1	Cell Movement	105	Migration and Invasion	106
TRPM7	Convergence and extension	107	Migration	108
MAPKAPK2	Epiboly	109	Migration	110
B4GALT1	Convergence and extension	111	Invasion	112
IER2	Convergence and extension	113	Migration	114
TIP1	Convergence and extension	115	Migration and Invasion	116
PAK5	Convergence and extension	117	Migration	118
MARCKS	Convergence and extension	119	Migration and Invasion	120

References

1   Zon, L. I. & Peterson, R. T. In vivo drug discovery in the zebrafish *Nat Rev Drug*
Discov **4**, 35-44 (2005).
2   Nakayama., J., Lu., J.-W., Makinoshima., H. & Gong., Z. A Novel Zebrafish Model of
Metastasis Identifies the HSD11β1 Inhibitor Adrenosterone as a Suppressor of
Epithelial-Mesenchymal Transition and Metastatic Dissemination. *Mol Cancer Res*
18, 477-487. (2020).
3   Nakayama, J. & Makinoshima, H. Zebrafish-Based Screening Models for the
Identification of Anti-Metastatic Drugs. *Molecules* **25**, 2407 (2020).
4   Nakayama, J. & Gong, Z. Transgenic zebrafish for modeling hepatocellular carcinoma.
MedComm **1**, 146-156. (2020).
5   Nguyen., D. X., Bos., P. D. & Massagué., J. Metastasis: from dissemination to organ-
specific colonization. *Nat Rev Cancer* **9**, 274-284 (2009).
6   Welch, D. R. & Hurst, D. R. Defining the Hallmarks of Metastasis *Cancer research* **79**,
3011-3027. (2019).
7   Chaffer, C. L. & Weinberg, R. A. A perspective on cancer cell metastasis *Science* **331**,
1559-1564 (2011).
8   Lu, W. & Kang, Y. Epithelial-Mesenchymal Plasticit in Cancer Progression and
Metastasis. *Developmental Cell* **49**, 361-374 (2019).
9   Tsai, J. H. & Yang, J. Epithelial–mesenchymal plasticity in carcinoma metastasis.
Genes Dev **27**, 2192-2206 (2013).
10   Solnica-Krezel, L. Conserved patterns of cell movements during vertebrate
gastrulation *Curr Biol* **15**, R213-228. , doi:10.1016/j.cub.2005.03.016. (2005).
11   Yang., J., Parker, A. & Geert, B. Guidelines and definitions for research on epithelial-
mesenchymal transition. *Nat Rev Mol Cell Biol* **21**, 341-352 (2020).
12   Nieto, M. A., Huang, R. Y.-J., Jackson, R. A. & Thiery, J. P. EMT: 2016 *Cell* **166**, 21-45
(2016).
13   Thiery, J. P., Acloque, H., Huang, R. Y. J. & Nieto, M. A. Epithelial-mesenchymal
transitions in development and disease *Cell* **139**, 871-890 (2009).
14   Yang, J. & Weinberg, R. A. Epithelial-mesenchymal transition: at the crossroads of
development and tumor metastasis *Dev Cell* **14**, 818-829 (2008).
15   Dongre, A. & Weinberg, R. A. **New insights into the mechanisms of epithelial-**
mesenchymal transition and implications for cancer. *Nat Rev Mol Cell Biol* **20**, 69-84
(2019).
16   Nakayama, J. *et al.* HSD11β1 promotes EMT-mediated breast cancer metastasis.
bioRxiv **doi: https://doi.org/10.1101/2021.09.27.461934** (2021).
17   Nakayama, J. *et al.* A zebrafish embryo screen utilizing gastrulation identifies the
HTR2C inhibitor pizotifen as a suppressor of EMT-mediated metastasis. *Elife* **10**,
e70151 (2021).
18   Nakayama, J., Konno, Y., Maruyama, A., Tomita, M. & Makinoshima, H. Cinnamon
bark extract suppresses metastatic dissemination of cancer cells through inhibition
of glycolytic metabolism *J Nat Med* (2022).
19   Gómez-Cuadrado, L., Tracey, N., Ma, R., Qian, B. & Brunton, V. G. Mouse models of
metastasis: progress and prospects *Dis Model Mech* **10**, 1061-1074. (2017).
20   Hoffman, R. M. Orthotopic metastatic (MetaMouse) models for discovery and
development of novel chemotherapy *Methods Mol Med* **111**, 297-322 (2005).

21 Chen, H.-C. Boyden chamber assay *Methods Mol Biol* **294**, 15-22 (2005).

22 Sack, U., Walther, W. & Scudiero, D. Novel effect of antihelminthic Niclosamide on
S100A4-mediated metastatic progression in colon cancer. *J Natl Cancer Inst* **103**,
1018-1036 (2011).

23 Itoh, K. *et al.* An essential part for Rho-associated kinase in the transcellular invasion
of tumor cells *Nat Med*

. 1999 Feb;5(2): **2**, 221-225 (1999).

24 Marshall, J.-C. A. *et al.* Effect of inhibition of the lysophosphatidic acid receptor 1 on
metastasis and metastatic dormancy in breast cancer *J Natl Cancer Inst* **104**, 1306-
1319 (2012).

25 Pelka, K. E., Henn, K., Keck, A., Sapel, B. & Braunbeck, T. Size does matter –
Determination of the critical molecular size for the uptake of chemicals across the
chorion of zebrafish (Danio rerio) embryos. *Aquatic Toxicology* **185**, 1-10 (2017).

26 Kondo, M. Bone morphogenetic proteins in the early development of zebrafish. *FEBS
J* **274**, 2960-2967, doi:10.1111/j.1742-4658.2007.05838.x (2007).

27 Katsuno, Y. *et al.* Bone morphogenetic protein signaling enhances invasion and bone
metastasis of breast cancer cells through Smad pathway. *Oncogene* **27**, 6322-6333,
doi:10.1038/onc.2008.232 (2008).

28 Tada, M. & Smith, J. C. Xwnt11 is a target of Xenopus Brachyury: regulation of
gastrulation movements via Dishevelled, but not through the canonical Wnt
pathway. *Development* **127**, 2227-2238 (2000).

29 Vincan, E. & Barker, N. The upstream components of the Wnt signalling pathway in
the dynamic EMT and MET associated with colorectal cancer progression. *Clin Exp
Metastasis* **25**, 657-663, doi:10.1007/s10585-008-9156-4 (2008).

30 Yang, X., Dormann, D., Munsterberg, A. E. & Weijer, C. J. Cell movement patterns
during gastrulation in the chick are controlled by positive and negative chemotaxis
mediated by FGF4 and FGF8. *Dev Cell* **3**, 425-437 (2002).

31 Nomura, S. *et al.* FGF10/FGFR2 signal induces cell migration and invasion in
pancreatic cancer. *Br J Cancer* **99**, 305-313, doi:10.1038/sj.bjc.6604473 (2008).

32 Song, S. *et al.* Pou5f1-dependent EGF expression controls E-cadherin endocytosis,
cell adhesion, and zebrafish epiboly movements. *Dev Cell* **24**, 486-501,
doi:10.1016/j.devcel.2013.01.016 (2013).

33 Lu, Z., Jiang, G., Blume-Jensen, P. & Hunter, T. Epidermal growth factor-induced
tumor cell invasion and metastasis initiated by dephosphorylation and
downregulation of focal adhesion kinase. *Mol Cell Biol* **21**, 4016-4031,
doi:10.1128/MCB.21.12.4016-4031.2001 (2001).

34 Damm, E. W. & Winklbauer, R. PDGF-A controls mesoderm cell orientation and radial
intercalation during Xenopus gastrulation. *Development* **138**, 565-575,
doi:10.1242/dev.056903 (2011).

35 Jechlinger, M. *et al.* Autocrine PDGFR signaling promotes mammary cancer
metastasis. *J Clin Invest* **116**, 1561-1570, doi:10.1172/JCI24652 (2006).

36 Mizoguchi, T., Verkade, H., Heath, J. K., Kuroiwa, A. & Kikuchi, Y. Sdf1/Cxcr4 signaling
controls the dorsal migration of endodermal cells during zebrafish gastrulation.
Development **135**, 2521-2529, doi:10.1242/dev.020107 (2008).

37 Shen, H. B., Gu, Z. Q., Jian, K. & Qi, J. CXCR4-mediated Stat3 activation is essential for
CXCL12-induced cell invasion in bladder cancer. *Tumour Biol* **34**, 1839-1845,
doi:10.1007/s13277-013-0725-z (2013).
38 Montero, J.-A., Kilian, B., Chan, J., Bayliss, P. E. & Heisenberg, C.-P. Phosphoinositide
3-Kinase Is Required for Process Outgrowth and Cell Polarization of Gastrulating
Mesendodermal Cells. *Current Biology* **13**, 1279-1289, doi:10.1016/s0960-
9822(03)00505-0 (2003).
39 Wander, S. A. *et al.* PI3K/mTOR inhibition can impair tumor invasion and metastasis
in vivo despite a lack of antiproliferative action in vitro: implications for targeted
therapy. *Breast Cancer Res Treat* **138**, 369-381, doi:10.1007/s10549-012-2389-6
(2013).
40 Tsai, W. B., Zhang, X., Sharma, D., Wu, W. & Kinsey, W. H. Role of Yes kinase during
early zebrafish development. *Dev Biol* **277**, 129-141,
doi:10.1016/j.ydbio.2004.08.052 (2005).
41 Barraclough, J., Hodgkinson, C., Hogg, A., Dive, C. & Welman, A. Increases in c-Yes
Expression Level and Activity Promote Motility But Not Proliferation of Human
Colorectal Carcinoma Cells. *Neoplasia* **9**, 745-IN732, doi:10.1593/neo.07442 (2007).
42 Sharma, D., Holets, L., Zhang, X. & Kinsey, W. H. Role of Fyn kinase in signaling
associated with epiboly during zebrafish development. *Dev Biol* **285**, 462-476,
doi:10.1016/j.ydbio.2005.07.018 (2005).
43 Yadav, V. & Denning, M. F. Fyn is induced by Ras/PI3K/Akt signaling and is required
for enhanced invasion/migration. *Mol Carcinog* **50**, 346-352, doi:10.1002/mc.20716
(2011).
44 Krens, S. F. *et al.* Distinct functions for ERK1 and ERK2 in cell migration processes
during zebrafish gastrulation. *Dev Biol* **319**, 370-383,
doi:10.1016/j.ydbio.2008.04.032 (2008).
45 Radtke, S., Milanovic, M. & Rossé, C. ERK2 but not ERK1 mediates HGF-induced
motility in non-small cell lung carcinoma cell lines. *J Cell Sci.* **126**, 2381-2391 (2013).
46 Jopling, C., van Geemen, D. & den Hertog, J. Shp2 knockdown and Noonan/LEOPARD
mutant Shp2-induced gastrulation defects. *PLoS Genet* **3**, e225,
doi:10.1371/journal.pgen.0030225 (2007).
47 Wang, F. M. *et al.* SHP-2 promoting migration and metastasis of MCF-7 with loss of
E-cadherin, dephosphorylation of FAK and secretion of MMP-9 induced by IL-1beta
in vivo and in vitro. *Breast Cancer Res Treat* **89**, 5-14, doi:10.1007/s10549-004-1002-
z (2005).
48 Ip, Y. T. & Gridley, T. Cell movements during gastrulation: snail dependent and
independent pathways. *Curr Opin Genet Dev* **12**, 423-429 (2002).
49 Batlle, E., Sancho, E., Francí, C., Domínguez, D. & Monfar, M. The transcription factor
snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nat Cell
Biol* **2**, 84-89 (2000).
50 Shi, J., Severson, C., Yang, J., Wedlich, D. & Klymkowsky, M. W. Snail2 controls
mesodermal BMP/Wnt induction of neural crest. *Development* **138**, 3135-3145,
doi:10.1242/dev.064394 (2011).
51 Medici, D., Hay, E. D. & B.R., O. Snail and Slug Promote Epithelial-Mesenchymal
Transition through -Catenin–T-Cell Factor-4-dependent Expression of Transforming
Growth Factor- 3. *Molecular Biology of the Cell* **19**, 4875–4887,
doi:10.1091/mbc.E08 (2008).

52 Castanon, I. & Baylies, M. K. A Twist in fate: evolutionary comparison of Twist
structure and function. *Gene* **287**, 11-22 (2002).
53 Yang, J. *et al.* Twist, a master regulator of morphogenesis, plays an essential role in
tumor metastasis. *Cell* **117**, 927-939, doi:10.1016/j.cell.2004.06.006 (2004).
54 Fernando, R. I. *et al.* The T-box transcription factor Brachyury promotes epithelial-
mesenchymal transition in human tumor cells. *J Clin Invest* **120**, 533-544,
doi:10.1172/JCI38379 (2010).
55 Vannier, C., Mock, K., Brabletz, T. & Driever, W. Zeb1 regulates E-cadherin and
Epcam (epithelial cell adhesion molecule) expression to control cell behavior in early
zebrafish development. *J Biol Chem* **288**, 18643-18659,
doi:10.1074/jbc.M113.467787 (2013).
56 Spaderna, S. *et al.* The transcriptional repressor ZEB1 promotes metastasis and loss
of cell polarity in cancer. *Cancer Res* **68**, 537-544, doi:10.1158/0008-5472.CAN-07-
5682 (2008).
57 Sander, V., Reversade, B. & De Robertis, E. M. The opposing homeobox genes
Goosecoid and Vent1/2 self-regulate Xenopus patterning. *EMBO J* **26**, 2955-2965,
doi:10.1038/sj.emboj.7601705 (2007).
58 Hartwell, K. A. *et al.* The Spemann organizer gene, Goosecoid, promotes tumor
metastasis. *Proc Natl Acad Sci U S A* **103**, 18969-18974,
doi:10.1073/pnas.0608636103 (2006).
59 Wilm, B., James, R. G., Schultheiss, T. M. & Hogan, B. L. The forkhead genes, Foxc1
and Foxc2, regulate paraxial versus intermediate mesoderm cell fate. *Dev Biol* **271**,
176-189, doi:10.1016/j.ydbio.2004.03.034 (2004).
60 Mani, S. A. *et al.* Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and
is associated with aggressive basal-like breast cancers. *Proc Natl Acad Sci U S A* **104**,
10069-10074, doi:10.1073/pnas.0703900104 (2007).
61 Miyagi, C. *et al.* STAT3 noncell-autonomously controls planar cell polarity during
zebrafish convergence and extension. *J Cell Biol* **166**, 975-981,
doi:10.1083/jcb.200403110 (2004).
62 Abdulghani, J. *et al.* Stat3 promotes metastatic progression of prostate cancer. *Am J
Pathol* **172**, 1717-1728, doi:10.2353/ajpath.2008.071054 (2008).
63 Lachnit, M., Kur, E. & Driever, W. Alterations of the cytoskeleton in all three
embryonic lineages contribute to the epiboly defect of Pou5f1/Oct4 deficient MZspg
zebrafish embryos. *Dev Biol* **315**, 1-17, doi:10.1016/j.ydbio.2007.10.008 (2008).
64 Dai, X. *et al.* OCT4 regulates epithelial-mesenchymal transition and its knockdown
inhibits colorectal cancer cell migration and invasion. *Oncol Rep* **29**, 155-160,
doi:10.3892/or.2012.2086 (2013).
65 O'Carroll, D. *et al.* The polycomb-group gene Ezh2 is required for early mouse
development. *Mol Cell Biol* **21**, 4330-4336, doi:10.1128/MCB.21.13.4330-4336.2001
(2001).
66 Ren, G. *et al.* Polycomb protein EZH2 regulates tumor invasion via the transcriptional
repression of the metastasis suppressor RKIP in breast and prostate cancer. *Cancer
Res* **72**, 3091-3104, doi:10.1158/0008-5472.CAN-11-3546 (2012).
67 Lin, F. *et al.* Essential roles of G{alpha}12/13 signaling in distinct cell behaviors
driving zebrafish convergence and extension gastrulation movements. *J Cell Biol* **169**,
777-787, doi:10.1083/jcb.200501104 (2005).

68 Chen, M. W. *et al.* H3K9 histone methyltransferase G9a promotes lung cancer
invasion and metastasis by silencing the cell adhesion molecule Ep-CAM. *Cancer Res*
70, 7830-7840, doi:10.1158/0008-5472.CAN-10-0833 (2010).

69 van der Lugt, N. M. *et al.* Posterior transformation, neurological abnormalities, and
severe hematopoietic defects in mice with a targeted deletion of the bmi-1 proto-
oncogene. *Genes Dev* **8**, 757-769 (1994).

70 Guo, B. H. *et al.* Bmi-1 promotes invasion and metastasis, and its elevated expression
is correlated with an advanced stage of breast cancer. *Mol Cancer* **10**, 10,
doi:10.1186/1476-4598-10-10 (2011).

71 Zhu, S., Liu, L., Korzh, V., Gong, Z. & Low, B. C. RhoA acts downstream of Wnt5 and
Wnt11 to regulate convergence and extension movements by involving effectors
Rho kinase and Diaphanous: use of zebrafish as an in vivo model for GTPase
signaling. *Cell Signal* **18**, 359-372, doi:10.1016/j.cellsig.2005.05.019 (2006).

72 Yoshioka, K., Nakamori, S. & Itoh, K. Overexpression of small GTP-binding protein
RhoA promotes invasion of tumor cells. *Cancer Res* **59**, 2004-2010 (1999).

73 Choi, S. C. & Han, J. K. Xenopus Cdc42 regulates convergent extension movements
during gastrulation through Wnt/Ca2+ signaling pathway. *Dev Biol* **244**, 342-357,
doi:10.1006/dbio.2002.0602 (2002).

74 Reymond, N. *et al.* Cdc42 promotes transendothelial migration of cancer cells
through beta1 integrin. *J Cell Biol* **199**, 653-668, doi:10.1083/jcb.201205169 (2012).

75 Habas, R., Dawid, I. B. & He, X. Coactivation of Rac and Rho by Wnt/Frizzled signaling
is required for vertebrate gastrulation. . *Genes Dev* **17**, 295-309., doi:10.1101/
(2003).

76 Vega, F. M. & Ridley, A. J. Rho GTPases in cancer cell biology. *FEBS Lett* **582**, 2093-
2101, doi:10.1016/j.febslet.2008.04.039 (2008).

77 Marlow, F., Topczewski, J., Sepich, D. & Solnica-Krezel, L. Zebrafish Rho kinase 2 acts
downstream of Wnt11 to mediate cell polarity and effective convergence and
extension movements. *Curr Biol* **12**, 876-884 (2002).

78 Itoh, K. *et al.* An essential part for Rho-associated kinase in the transcellular invasion
of tumor cells. *Nat Med* **5**, 221-225, doi:10.1038/5587 (1999).

79 Kusakabe, M. & Nishida, E. The polarity-inducing kinase Par-1 controls Xenopus
gastrulation in cooperation with 14-3-3 and aPKC. *EMBO J* **23**, 4190-4201,
doi:10.1038/sj.emboj.7600381 (2004).

80 Shi, X., Gangadharan, B., Brass, L. F., Ruf, W. & Mueller, B. M. Protease-activated
receptors (PAR1 and PAR2) contribute to tumor cell motility and metastasis. *Mol
Cancer Res* **2**, 395-402 (2004).

81 Gunaratne, A., Thai, B. L. & Di Guglielmo, G. M. Atypical protein kinase C
phosphorylates Par6 and facilitates transforming growth factor beta-induced
epithelial-to-mesenchymal transition. *Mol Cell Biol* **33**, 874-886,
doi:10.1128/MCB.00837-12 (2013).

82 Seifert, K., Ibrahim, H., Stodtmeister, T., Winklbauer, R. & Niessen, C. M. An
adhesion-independent, aPKC-dependent function for cadherins in morphogenetic
movements. *J Cell Sci* **122**, 2514-2523, doi:10.1242/jcs.042796 (2009).

83 Yamazaki, K. *et al.* Adenylate cyclase-associated protein 1 overexpressed in
pancreatic cancers is involved in cancer cell motility. *Lab Invest* **89**, 425-432,
doi:10.1038/labinvest.2009.5 (2009).

84 Link, V. *et al.* Identification of regulators of germ layer morphogenesis using
proteomics in zebrafish. *J Cell Sci* **119**, 2073-2083, doi:10.1242/jcs.02928 (2006).
85 Khanna, C. *et al.* The membrane-cytoskeleton linker ezrin is necessary for
osteosarcoma metastasis. *Nat Med* **10**, 182-186, doi:10.1038/nm982 (2004).
86 Slanchev, K. *et al.* The epithelial cell adhesion molecule EpCAM is required for
epithelial morphogenesis and integrity during zebrafish epiboly and skin
development. *PLoS Genet* **5**, e1000563, doi:10.1371/journal.pgen.1000563 (2009).
87 Ni, J. *et al.* Role of the EpCAM (CD326) in prostate cancer metastasis and
progression. *Cancer Metastasis Rev* **31**, 779-791, doi:10.1007/s10555-012-9389-1
(2012).
88 Skalski, M., Alfandari, D. & Darribere, T. A key function for alphav containing
integrins in mesodermal cell migration during Pleurodeles waltl gastrulation. *Dev Biol*
195, 158-173, doi:10.1006/dbio.1997.8838 (1998).
89 Felding-Habermann, B. Integrin adhesion receptors in tumor metastasis. *Clin Exp*
Metastasis **20**, 203-213 (2003).
90 Marsden, M. & DeSimone, D. W. Integrin-ECM Interactions Regulate Cadherin-
Dependent Cell Adhesion and Are Required for Convergent Extension in Xenopus.
Current Biology **13**, 1182-1191, doi:10.1016/s0960-9822(03)00433-0 (2003).
91 Malik, G. *et al.* Plasma fibronectin promotes lung metastasis by contributions to
fibrin clots and tumor cell invasion. *Cancer Res* **70**, 4327-4334, doi:10.1158/0008-
5472.CAN-09-3312 (2010).
92 Bakkers, J. *et al.* Has2 is required upstream of Rac1 to govern dorsal migration of
lateral cells during zebrafish gastrulation. *Development* **131**, 525-537,
doi:10.1242/dev.00954 (2004).
93 Kim, H. R. *et al.* Hyaluronan facilitates invasion of colon carcinoma cells in vitro via
interaction with CD44. *Cancer Res* **64**, 4569-4576, doi:10.1158/0008-5472.CAN-04-
0202 (2004).
94 Coyle, R. C., Latimer, A. & Jessen, J. R. Membrane-type 1 matrix metalloproteinase
regulates cell migration during zebrafish gastrulation: evidence for an interaction
with non-canonical Wnt signaling. *Exp Cell Res* **314**, 2150-2162,
doi:10.1016/j.yexcr.2008.03.010 (2008).
95 Perentes, J. Y. *et al.* Cancer cell-associated MT1-MMP promotes blood vessel
invasion and distant metastasis in triple-negative mammary tumors. *Cancer Res* **71**,
4527-4538, doi:10.1158/0008-5472.CAN-10-4376 (2011).
96 Cha, Y. I. *et al.* Cyclooxygenase-1-derived PGE2 promotes cell motility via the G-
protein-coupled EP4 receptor during vertebrate gastrulation. *Genes Dev* **20**, 77-86,
doi:10.1101/gad.1374506 (2006).
97 Kundu, N. & Fulton, A. M. Selective cyclooxygenase (COX)-1 or COX-2 inhibitors
control metastatic disease in a murine model of breast cancer. *Cancer Res* **62**, 2343-
2346 (2002).
98 Speirs, C. K. *et al.* Prostaglandin Gbetagamma signaling stimulates gastrulation
movements by limiting cell adhesion through Snai1a stabilization. *Development* **137**,
1327-1337, doi:10.1242/dev.045971 (2010).
99 Wang, D. & Dubois, R. N. Prostaglandins and cancer. *Gut* **55**, 115-122,
doi:10.1136/gut.2004.047100 (2006).

565	100	Yamashita, S. *et al.* Zinc transporter LIVI controls epithelial-mesenchymal transition
566		in zebrafish gastrula organizer. *Nature* **429**, 298-302, doi:10.1038/nature02545
567		(2004).
568	101	Lue, H. W. *et al.* LIV-1 promotes prostate cancer epithelial-to-mesenchymal
569		transition and metastasis through HB-EGF shedding and EGFR-mediated ERK
570		signaling. *PLoS One* **6**, e27720, doi:10.1371/journal.pone.0027720 (2011).
571	102	Yagi, H. *et al.* A synthetic biology approach reveals a CXCR4-G13-Rho signaling axis
572		driving transendothelial migration of metastatic breast cancer cells. *Sci Signal* **4**,
573		ra60, doi:10.1126/scisignal.2002221 (2011).
574	103	Webster, D. M. *et al.* O-GlcNAc modifications regulate cell survival and epiboly
575		during zebrafish development. *BMC Dev Biol* **9**, 28, doi:10.1186/1471-213X-9-28
576		(2009).
577	104	Lynch, T. P. *et al.* Critical role of O-Linked beta-N-acetylglucosamine transferase in
578		prostate cancer invasion, angiogenesis, and metastasis. *J Biol Chem* **287**, 11070-
579		11081, doi:10.1074/jbc.M111.302547 (2012).
580	105	Latinkic, B. V. Xenopus Cyr61 regulates gastrulation movements and modulates Wnt
581		signalling. *Development* **130**, 2429-2441, doi:10.1242/dev.00449 (2003).
582	106	Lin, J. *et al.* A novel anti-Cyr61 antibody inhibits breast cancer growth and metastasis
583		in vivo. *Cancer Immunol Immunother* **61**, 677-687, doi:10.1007/s00262-011-1135-y
584		(2012).
585	107	Liu, W. *et al.* TRPM7 regulates gastrulation during vertebrate embryogenesis. *Dev*
586		*Biol* **350**, 348-357, doi:10.1016/j.ydbio.2010.11.034 (2011).
587	108	Middelbeek, J. *et al.* TRPM7 is required for breast tumor cell metastasis. *Cancer Res*
588		**72**, 4250-4261, doi:10.1158/0008-5472.CAN-11-3863 (2012).
589	109	Holloway, B. A. *et al.* A novel role for MAPKAPK2 in morphogenesis during zebrafish
590		development. *PLoS Genet* **5**, e1000413, doi:10.1371/journal.pgen.1000413 (2009).
591	110	Kumar, B. *et al.* p38 mitogen-activated protein kinase-driven MAPKAPK2 regulates
592		invasion of bladder cancer by modulation of MMP-2 and MMP-9 activity. *Cancer Res*
593		**70**, 832-841, doi:10.1158/0008-5472.CAN-09-2918 (2010).
594	111	Machingo, Q. J., Fritz, A. & Shur, B. D. A beta1,4-galactosyltransferase is required for
595		convergent extension movements in zebrafish. *Dev Biol* **297**, 471-482,
596		doi:10.1016/j.ydbio.2006.05.024 (2006).
597	112	Zhu, X. *et al.* Elevated beta1,4-galactosyltransferase I in highly metastatic human
598		lung cancer cells. Identification of E1AF as important transcription activator. *J Biol*
599		*Chem* **280**, 12503-12516, doi:10.1074/jbc.M413631200 (2005).
600	113	Hong, S. K., Tanegashima, K. & Dawid, I. B. XIer2 is required for convergent extension
601		movements during Xenopus development. *Int J Dev Biol* **55**, 917-921,
602		doi:10.1387/ijdb.113288sh (2011).
603	114	Neeb, A. *et al.* The immediate early gene Ier2 promotes tumor cell motility and
604		metastasis, and predicts poor survival of colorectal cancer patients. *Oncogene* **31**,
605		3796-3806, doi:10.1038/onc.2011.535 (2012).
606	115	Besser, J., Leito, J. T., van der Meer, D. L. & Bagowski, C. P. Tip-1 induces filopodia
607		growth and is important for gastrulation movements during zebrafish development.
608		*Dev Growth Differ* **49**, 205-214, doi:10.1111/j.1440-169X.2007.00921.x (2007).
609	116	Han, M., Wang, H., Zhang, H. T. & Han, Z. The PDZ protein TIP-1 facilitates cell
610		migration and pulmonary metastasis of human invasive breast cancer cells in

athymic mice. *Biochem Biophys Res Commun* **422**, 139-145,
doi:10.1016/j.bbrc.2012.04.123 (2012).
117 Faure, S. *et al.* Xenopus p21-activated kinase 5 regulates blastomeres' adhesive
properties during convergent extension movements. *Dev Biol* **277**, 472-492,
doi:10.1016/j.ydbio.2004.10.005 (2005).
118 Gong, W. *et al.* P21-activated kinase 5 is overexpressed during colorectal cancer
progression and regulates colorectal carcinoma cell adhesion and migration. *Int J*
Cancer **125**, 548-555, doi:10.1002/ijc.24428 (2009).
119 Iioka, H., Ueno, N. & Kinoshita, N. Essential role of MARCKS in cortical actin dynamics
during gastrulation movements. *J Cell Biol* **164**, 169-174, doi:10.1083/jcb.200310027
(2004).
120 Rombouts, K. *et al.* Myristoylated Alanine-Rich protein Kinase C Substrate (MARCKS)
expression modulates the metastatic phenotype in human and murine colon
carcinoma in vitro and in vivo. *Cancer Lett* **333**, 244-252,
doi:10.1016/j.canlet.2013.01.040 (2013).
626