SARS-CoV-2 Spike N-Terminal Domain modulates TMPRSS2-dependent viral

entry and fusogenicity

Bo Meng[1,2,*], Rawlings Datir[1,2], Jinwook Choi[3], CITIID-NIHR BioResource COVID-19 Collaboration, John

Bradley[2,4], Kenneth GC Smith[1,2], Joo Hyeon Lee[3,5], Ravindra K. Gupta[1,2,6,*]

[1] Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), Cambridge, UK.

[2] Department of Medicine, University of Cambridge, Cambridge, UK.

[3] Wellcome-MRC Cambridge Stem Cell Institute, Cambridge, UK.

[4] NIHR Bioresource, Cambridge, UK.

[5] Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge,

UK.

[6] Africa Health Research Institute, Durban, South Africa

*Correspondence to :

Bo Meng bm432@cam.ac.uk or Ravindra K. Gupta rkg20@cam.ac.uk

Key words: SARS-CoV-2, NTD, fusogenicity, TMPRSS2, Organoid, Delta, Omicron

Abstract

Over 20 mutations have been identified in the N-Terminal Domain (NTD) of SARS-CoV-2 spike and yet few of them are fully characterised. Here we first examined the contribution of the NTD to infection and cell-cell fusion by constructing different VOC-based chimeric spikes bearing B.1617 lineage (Delta and Kappa variants) NTDs and generating spike pseudotyped lentivirus (PV). We found the Delta NTD on a Kappa or WT background increased spike S1/S2 cleavage efficiency and virus entry, specifically in Calu-3 lung cells and airway organoids, through use of TMPRSS2. Delta was previously shown to have fast cell-cell fusion kinetics and increased fusogenicity that could be conferred to WT and Kappa variant spikes by transfer of the Delta NTD. Moving to contemporary variants, we found that BA.2 had higher entry efficiency in a range of cell types as compared to BA.1. BA.2 showed higher fusogenic activity than BA.1, but the BA.2 NTD could not confer higher fusion to BA.1 spike. There was low efficiency of TMPRSS2 usage by both BA.1 and BA.2, and chimeras of Omicron BA.1 and BA.2 spikes with a Delta NTD did not result in more efficient use of TMRPSS2 or cell-cell fusogenicity. We conclude that the NTD allosterically modulates S1/S2 cleavage and spike-mediated functions such as entry and cell-cell fusion in a spike context dependent manner, and allosteric interactions may be lost when combining regions from more distantly related spike proteins. These data may explain the lack of dominant SARS-CoV-2 inter-variant recombinants bearing breakpoints within spike.

Introduction

The SARS-CoV-2 furin cleavage site (also known as a polybasic cleavage site), cleaved in producer
cells by a ubiquitously expressed protease furin (Hoffmann et al., 2020a), is believed to be one of the
main reasons behind the success of SARS-CoV2 worldwide (Johnson et al., 2021; Peacock et al.,
2021). Upon virus release the trimeric spike engages with angiotensin converting enzyme 2 (ACE2) of
the target cells to initiate virus entry (Jackson et al., 2022; Peng et al., 2021). Depending on the
abundance of the cofactor TMPRSS2 and the status of spike cleavage (Hoffmann et al., 2020b; Ou et
al., 2021a, 2021b; Park et al., 2016; Whittaker et al., 2021), the virus either enters through the
TMPRSS2 mediated route by fusing at the plasma membrane or via late endosomes through a
secondary cleavage preceding the concealed fusion peptide at S2' (Figure 1A). The latter does not
require a pre-cleaved spike at S1/S2, as cathepsin can mediate cleavage of both S1/S2 and S2'.

Kappa (B.1.617.1) and Delta (B.1.617.2) SARS-CoV-2 variants were first detected in India in late 2020
(Dhar et al., 2021; Ferreira et al., 2021). Although Kappa, the earliest detected B.1.617 variant in
India, displayed greater escape to vaccine-elicited antibody responses (McCallum et al., 2021a),
Delta surpassed Kappa to become the dominant strain in India and worldwide by mid-2021
(Mlcochova et al., 2021). The S gene encodes 8 and 9 non-synonymous mutations in Kappa and Delta,
respectively and four of which are shared by both (Figure 1A). The NTD is more variable between the
two (3 and 4 mutations relative to Wu-1 in Kappa and Delta, respectively). In contrast, the RBD, each
bearing two mutations, is more constrained most likely due to its obligatory role in engaging with
ACE2. Diverse mutations in the NTD are not unique to Kappa or Delta and have been documented in
Alpha and many other VOCs (McCarthy et al., 2021). Additionally, both Kappa and Delta share two
conserved mutations at D614G and P681R and a unique mutation at S2; Q1071H for Kappa and
D950N for Delta.

We previously showed that both Kappa and Delta spikes exhibit highly efficient cleavage of S1/S2
over 614G WT (Mlcochova et al., 2021). Intriguingly Delta appears to be superior to Kappa in
entering Calu3 cells and organoids expressing endogenous level of ACE2 and TMPRSS2. Receptor
binding is modestly increased in Delta but lower than the preceding strains of VOCs, for example
Alpha (Collier et al., 2021a; Liu et al., 2022; Ulrich et al., 2022), indicating enhanced receptor binding
cannot solely explain the higher transmissibility of Delta (Barton et al., 2021; McCallum et al., 2021a;
Ramanathan et al., 2021; Saville et al., 2022; Supasa et al., 2021). Moreover, cryo-EM structures of
Delta and Kappa trimeric spike show that both RBD and S2 adopt a remarkably similar geometry

suggesting those two regions are less likely to be accountable for the increased entry of Delta over

Kappa (Zhang et al., 2021a). *In vitro* studies using replication competent virus isolates showed that

Delta has fast replication kinetics in Calu3, human airway epithelium cells and in airway organoids

(Mlcochova et al., 2021). However, the underlying molecular mechanism for the high transmissibility

of Delta over Kappa in the real world is elusive.

Our published data also showed that the RBD on its own did not confer higher infectivity to Kappa

(Ferreira et al., 2021), suggesting the NTD may be responsible for the increased infectivity. The NTD

interacts with cofactors L-SIGN and DC-SIGN at the cell surface (Lempp et al., 2021); blockade of

these proteins can effectively neutralise the virus in ACE2 non-overexpressing cells, suggesting NTD

and RBD may work cooperatively. The cooperativity of NTD and the RBD is additionally supported by

the identification of infectivity enhancing antibodies specifically targeting the NTD domain (Li et al.,

2021; Liu et al., 2021), and the observation that binding of the 4A8 monoclonal antibody in the NTD

modulates the RBD into an up position (Chi et al., 2020; Díaz-Salinas et al., 2022). Interestingly such

antibody binding sites coincide with known infectivity enhancing sites, such as the H69V70 deletion

that emerged during an example of intra-host evolution (Kemp et al., 2021) and in Alpha (Meng et al.,

2021) and Omicron variants (Meng et al., 2022). It is therefore plausible that the NTD plays an active

role in virus entry by engaging with host cofactors and triggering conformational changes of the RBD.

Despite over 20 mutations documented in the NTD, the role of those mutations in infectivity and

their impact on the immune response elicited by vaccines is less clear. We reported that the H69V70

deletion found in Alpha was positively selected to increase its infectivity with a modest decrease in

immune evasion (Meng et al., 2021). Here we hypothesised that the NTD plays a regulatory role that

impacts S1/S2 cleavage and ACE2 usage. We constructed a panel of chimeric spike proteins with NTD

from different VOCs in a variety of VOC backbones. We examined those chimeras alongside the

parental VOCs in pseudovirus based entry assays (Mlcochova et al., 2020) and investigated spike-

mediated fusogenicity. Our data are consistent with a model whereby the NTD regulates virus entry

and cell-cell fusion in a variant context-dependent manner.

Results

NTD increases Delta infectivity in lung cells and airway organoids

The most dramatic changes in spike between Kappa and Delta lie in the NTD. Both Kappa and Delta

spike are efficiently cleaved in the producer cells (Mlcochova et al., 2021). We sought to assess the

contribution of the NTD in spike cleavage in purified PV by western blot. We included a deletion

mutant in the NTD (delH69/V70) as a control due to its known efficient spike cleavage (Kemp et al.,

2021; Meng et al., 2021). Plasmids encoding HIV Gag/pol, a genome flanked by LTRs encoding

luciferase, as well as the corresponding spike were transfected into 293T producer cells. The

supernatants were harvested and pelleted through ultracentrifugation for western blot analysis. Our

data show that the H69/V70 deletion increased S1/S2 cleavage compared to Wu-1 as expected

(Figure S1A). Kappa and Delta spike were efficiently cleaved with a more pronounced cleavage

observed in Delta (Figure S1B). We additionally observed that the Kappa spike was prone to S1

shedding evidenced by a higher S2/S1 ratio compared to that of the WT (Figure S1C). In contrast,

Delta spikes were more stable. We further noticed that the increased efficiency in spike cleavage

requires a cognate NTD as the spike bearing the RBD mutations alone failed to be cleaved efficiently

(Figure S1D), suggesting NTD on its own or together with the RBD influences the cleavage activity.

We conclude that the Delta spike has evolved to be optimal in efficient spike cleavage whilst

maintaining spike stability, reminiscent of the emergence of D614G in the early pandemic (Gobeil et

al., 2021; Yurkovetskiy et al., 2020; Zhang et al., 2021b, 2020).

We previously observed enhanced cell-free infectivity for Delta over Kappa in Calu3 cells in the PV

system (Mlcochova et al., 2021). Importantly this difference was also evident for airway organoids

(Mlcochova et al., 2021; Youk et al., 2020) suggesting that at endogenous expression levels of ACE2

and TMPRSS2, Delta has enhanced ability over Kappa in virus entry. Calu-3 and airway organoids

express high levels of TMPRSS2 (Meng et al., 2022). Given that cleavage efficiency positively

correlates with the infectivity in lung cells (Meng et al., 2022), we sought to examine whether the

Delta NTD also contributes to higher infectivity over Kappa in vitro.

To test this we constructed chimeras harboring either the Delta NTD or the Kappa NTD in Delta or

Kappa backbones (Figure 1A). We then used the PV bearing different spikes to transduce cell lines to

examine the entry efficiency. Interestingly we observed elevated cell-free infectivity in a Kappa

chimera bearing the Delta NTD in Calu3 cells and primary human airway organoids but not in other

cell lines, indicating that the NTD confers the specificity for this increase (Figure 1B). Consistent with

this model, a decrease in infectivity in Calu3 cells and primary human airway organoids was

observed when the Kappa NTD was exchanged with that of Delta. We further extended this

observation to the WT backbone where a hybrid WT spike bearing either Kappa or Delta NTD was

expressed (Figure 1C and S2). Consistent with these observations, both Kappa and Delta NTD when

fused with WT increased cell-free infectivity in Calu3 lung cells.

The NTD influences virus entry efficiency

We next sought to delineate the underlying mechanism for this increased infectivity in Delta over

Kappa. Spike cleavage correlates with SARS-CoV-2 virus entry pathway preference (Meng et al., 2022;

V'kovski et al., 2021). Western blots using the cell lysates from the 293T virus producer cells showed

that the expression levels of chimeric spikes were comparable (Figure 2A&B). However, the cleavage

of spike in the chimeras phenocopied the cleavage patterns from which their NTD were derived

(Figure 2C). We extended this observation in WT chimeras where increased spike cleavage was also

evident following addition of either Kappa NTD or Delta NTD (Figure S2B). Structural data indicate

that the NTD, RBD and PBCS are cooperatively regulated (Gobeil et al., 2021). Hence we reasoned

that the observed increase in the accessibility to the furin cleavage site may be regulated by the

allostery involving the NTD.

SARS-CoV-2 primarily enters Calu-3 cells through membrane fusion due to an abundant expression

of TMPRSS2 and ACE2 at the plasma membrane. The primary cleavage at S1/S2 is a prerequisite for a

secondary cleavage at S2' site by proteases such as TMPRSS2 at the plasma membrane (V'kovski et

al., 2021), triggering the conformational rearrangement of S2 for the exposure of the fusion peptide.

Due to more efficient cleavage of the Delta spike (Figure S1), we reasoned that the Delta spike may

be more efficiently primed to use TMPRSS2 at the plasma membrane for its entry. To test this

hypothesis, we transduced the virus bearing either authentic spike of Delta or Kappa or their NTD-

chimeric counterparts into TMPRSS2 overexpressing 293T cells or parental 293T cells (Figure 2D). As

expected, all the viruses including chimeras increased entry efficiency when TMPRSS2 was present

with an even more pronounced increase observed in Delta, consistent with the observation that

Delta is more efficient in utilising TMPRSS2. Intriguingly, we also observed an increased transduction

efficiency for the Kappa chimera that was comparable to that of Delta, suggesting that the Delta NTD

enables the entry of chimeric Kappa by efficient TMPRSS2 usage. Consistent with this model, the

ratio of entry for the Delta chimera containing Kappa NTD entry in TMPRSS2 overexpressing cells

versus normal expressing cells was reduced.

To further investigate the dependence of TMPRSS2 on virus entry, we pretreated A549-

ACE2/TMPRSS2 cells or airway organoids with either camostat (a TMPRSS2 inhibitor) or E64D (a

cathepsin inhibitor). We observed that the IC50 of camostat in Delta was approximately 2 fold higher

than Kappa, whilst E46D had little effect on either virus (Figure 2E). Reassuringly, a similar

observation was made in airway organoids (Figure 2F), suggesting that Delta is more resistant to

camostat and hence is more efficient in utilising available TMPRSS2 for virus entry. Indeed the Kappa

chimera bearing Delta NTD showed a similar drug sensitivity profile to Delta implying that the NTD is

accountable for this shift in TMPRSS2 sensitivity (Figure 3S).

It is reported that the NTD modulates RBD conformation through allosteric effects (Qing et al., 2021).

We speculated the accessibility to ACE2 is altered in the chimeric spikes. We next went on to

examine this by transducing parental 293T cells or their isogenic counterparts where the expression

of TMPRSS2 is abrogated and ACE2 is overexpressed (A2delT2). Indeed, we found a 30 fold increase

in Delta when ACE2 is overexpressed whereas Kappa only had 4 fold increase (Figure 4G). More

strikingly, the Kappa chimera, bearing the Delta NTD in the Kappa spike backbone, showed a similar

magnitude of increase as Delta. Conversely, the Delta chimera bearing a Kappa NTD manifested a

decrease in ACE2 dependence compared to Delta, albeit to a lesser extent. Taken together, our data

support the notion that the Delta NTD modulates the use of TMPRSS2 more efficiently due to spike

cleavage and that the Delta NTD allosterically regulates the RBD to increase efficiency of ACE2 usage.

NTD mutations and deletions impact entry efficiency

To probe the mutations in the Delta NTD that may contribute to the enhanced infectivity in Calu3

cells, we constructed a series of spike bearing PV by reverting a cluster of amino acids (142D, 156G,

del157F and del158R) to WT Wuhan-1 residues individually. Interestingly, each individually reverted

mutant showed a decrease in infectivity in Calu3 cells with the greatest reduction seen following re-

insertion of the deleted 157R/158R (3 fold). This is in agreement with the infectivity reduction in the

Delta NTD bearing Kappa chimera (Figure 1B). The decrease in infectivity was only observed in Calu3

cells but not in Hela-ACE2 cells, consistent with the cell specificity in virus entry conferred by the

NTD. Since the NTD harbors the antigenic supersites that are mutated in Delta (Cerutti et al., 2021;

McCallum et al., 2021b; McCarthy et al., 2021; Suryadevara et al., 2021), we predicted that the

reversion of such mutations would at least partially rescue the loss of sensitivity to the vaccine sera.

To test this, PV were harvested and used to transduce HeLa-ACE2 cells in the presence of a dilution

series of vaccine sera from vaccinees who had received two doses of BNT162b2 at least 1 month

prior to sampling. D142G alone did not alter the sensitivity of neutralisation (Figure 3). However,

G156E or repairing the deletion of 157F/158R increased the sensitivity of neutralization by 2 fold,

consistent with the notion that the NTD is important for immune evasion and infectivity.

Delta NTD confers faster cell-cell fusion kinetics in a context-dependent manner

Syncytium formation, mediated by spike and the host receptor ACE2, has been found in SARS-CoV-2

infected patients and is thought to be important for disease progression (Braga et al., 2021). We and

others had previously demonstrated that variants bearing furin cleavage site mutations exhibit a

pronounced increase in fusogenicity, with the exceptions of Omicron (Figure 4A) (Buchrieser et al.,

2021; Meng et al., 2021, 2022; Mlcochova et al., 2021; Rajah et al., 2021). Given that spike cleavage

largely correlates with fusion and we observed an enhanced cleavage in the Kappa chimera, we

sought to examine whether spike mediated fusion is altered. We first confirmed that efficient

cleavage is required for the syncytia formation as the mutants bearing the WT proline at position

681 (681P) decreased fusion efficiency in both parental and chimeras whereas the mutation to

histidine (681H), the other cleavage site mutation firstly found in Alpha and then in Omicron, did not

(Figure 4B). We then went on to explore whether or not the fusogenicity is different between Delta

and Kappa and their chimeras. Concordant with the published studies we found a marked increase in

fusion for Delta and Kappa compared to WT (Figure 4C) (Mlcochova et al., 2021; Rajah et al., 2021).

Unexpectedly, we found the fusion kinetics between Delta and Kappa were noticeably different

despite manifesting a comparably maximal fusion activity at the steady-state (around 70%; Figure

3C). More specifically, Delta was more fusogenic at earlier time points whereas Kappa required two

more hours to reach a similar intensity. Intriguingly when the Kappa NTD was fused into Delta the

fusion phenotype shifted from Delta to Kappa, with a slowing of kinetics (Figure 3C). Conversely, we

observed a fast-fusing Delta phenotype when the Delta NTD was swapped into Kappa. Of note, we

also observed a faster fusion phenotype when the Delta NTD was fused into the WT backbone,

though this effect was less pronounced compared to that observed in the Kappa chimera (Figure 4C).

Taken together we have demonstrated that the Delta NTD can drive faster fusion in both WT and

Kappa spike backgrounds.

Epidemiological studies suggest that the more recent Omicron (BA.1 and BA.2) variants are distantly

related to all the other early pandemic variants (Simon-Loriere and Schwartz, 2022). It therefore

remains possible that the ability of the Delta NTD to drive faster fusion is dependent on their spike

background. We found that in comparison to BA.1 BA.2 was more efficient in entering Calu3 cells,

though still not as efficient as Delta (Figure 5A). An elevated level of entry of BA.2 over BA.1 was

observed in H1299 cells where the entry of Delta was the least efficient. These data suggest that

BA.2 is capable of infecting both TMPRSS2 low (H1299) and high (Calu3) cells efficiently.

We then introduced the Delta NTD into the both BA.1 and BA.2 to examine whether the Delta NTD

could also confer increased fusion to Omicron, a known poor cell-cell fusion spike (Figure 5B).

Although BA.2 showed faster fusion than BA.1, consistent with a recently published study (Yamasoba

et al., 2022), both Omicron and the Delta-NTD fused Omicron chimeras showed poor spike-mediated

fusion and that most importantly the difference between Omicron and its counterpart was negligible.

Intriguingly, this phenotype correlates with inefficient cleavage of spike even after the cleavage-

enhancing Delta NTD was fused into an Omicron spike background.

Omicron has evolved to avoid the early entry route via TMPRSS2 at the plasma membrane by

entering the cells through endosomal route (Meng et al., 2022). Given spike cleavage was not

affected by domain swapping, we predicted that the Omicron chimera bearing the Delta NTD would

fail to confer increased sensitivity to TMPRSS2, in contrast to our findings for the B.1.617 lineages

Delta and Kappa. Indeed there was only a moderate increase in entry (2-3 fold) when TMPRSS2 was

overexpressed, compared to 9 fold for Delta, reinforcing the notion that the virus entry of Omicron

is TMPRSS2 independent and that altering the NTD is not sufficient to confer TMPRSS2 usage to

Omicron.

We next sought to examine whether the BA.2 NTD could rescue the poor fusogenicity of BA.1 and/or

whether the BA.1 NTD impaired the fusogenicity of BA.2. In contrast to the fast fusion activity

conferred by the Delta NTD in Kappa or WT (Figure 4C), BA.2 NTD failed to increase the fusion

kinetics of the BA.1 chimeric spike bearing BA.2 NTD (Figure 5D). The BA.1 NTD modestly

suppressed the fusion of BA.2 chimeric spike bearing BA.1 NTD. Interestingly, when the sensitivity to

TMPRSS2 was probed (Figure 5E), we found little difference between BA.1 and BA.2 and their

chimeras (8-10 fold) in comparison to Delta (over 50 fold), indicating that in Omicron the NTD on its

own is not sufficient to affect TMPRSS2 utilisation. Taken together, these data strongly suggest that

the enhanced fusogenicity of the Delta NTD and the spike cleavage is context-dependent and that

non-NTD regions may be involved to suppress efficient fusion activity of Omicron spike.

Discussion

Cooperativity between the NTD and RBD is reflected by the finding that NTD deletions were

previously discovered after the emergence of the RBD mutations; for example, N439K and Y453F

neutralising antibody escape mutations were identified first before the acquisition of delH69/V70 in

the NTD (Meng et al., 2021). We previously showed that some escape mutations reduced virus entry

efficiency and the delH69/V70 NTD deletion restored entry whilst maintaining immune evasion

(Kemp et al., 2021; Meng et al., 2021). Therefore, viruses have selected various mutations in the RBD

due to the immune pressure imposed on this immunodominant site by the host. However these

RBD-mutated viruses have suboptimal fitness, thus placing selection pressure to acquire

compensatory mutations in the NTD. We propose that NTDs in VOC have evolved under such

pressure.

The NTD has been proposed to modulate the RBD to a receptor accessible mode (Qing et al., 2021).

We speculated that by alternating the NTD of Delta into Kappa the receptor engagement with ACE2

is affected. Indeed we observed an enhanced virus entry in the Kappa chimera bearing the Delta

NTD (Figure 2G). However an increased accessibility to ACE2 cannot solely account for the enhanced

infectivity observed in Calu3 as such an increase is not evident for the cell lines where ACE2 is also
abundantly expressed. A marked increase for chimeric Kappa spike to use TMPRSS2 at the plasma
membrane provides a plausible explanation for this specificity; the chimeric Kappa spike is as
efficient as Delta on TMPRSS2 usage (Figure 2D). This observation is in agreement with our inhibitor
experiment, whereby the addition of the TMPRSS2 protease inhibitor camostat has a more
pronounced inhibitory effect for Kappa than Delta (Figure 2E&F). However, the Delta NTD cannot
enhance the cleavage of Omicron spike nor its dependence on TMPRSS2 (Figure 5B&C), suggesting
that the presence of the Delta NTD is not the sole determinant of TMPRSS2 usage. Furthermore,
although allosteric conformational changes involving NTD and other regions may contribute to
phenotype, binding of the NTD to a secondary cofactor at the plasma membrane cannot be
completely excluded (McCallum et al., 2021b).

The NTD of VOCs, including poorly fusogenic Omicron (Meng et al., 2022; Suzuki et al., 2022), once
fused with WT exhibits an increased fusion in the presence of trypsin indicating the accessibility of
spike cleavage by host protease is affected through the allostery of the NTD (Qing et al., 2022). We
have proposed that the sensitivity to TMPRSS2 correlates with cleavage status at S1/S2, which has
an additional impact on the cell to cell fusion (Meng et al., 2022). Consistent with this notion, we
found the Delta NTD confers faster fusion kinetics in both WT and Kappa which coincides with an
elevated S1/S2 cleavage in WT and Kappa spike bearing the Delta NTD. However S1/S2 cleavage is
unaltered in the Delta NTD bearing Omicron which also has poor fusogenicity. These data
demonstrated that regions other than NTD may also be required for efficient cleavage. Indeed a
recent study using a domain swapping approach has suggested the RBD of Omicron contributes to
inefficient spike cleavage and inferior fusogenicity (doi.org/10.1101/2022.04.03.486864). The
specific site in S2 preceding to fusion peptide has also been implicated in facilitating an efficient
cleavage in accordance with the NTD (Qing et al., 2022).

We observed that the Kappa spike is less stable than that of Delta (Figure S1C). We suspect that
when Delta first emerged with the RBD mutations the intermediate strain was less fit leading to a
premature S1 shedding. However, the subsequently acquired mutations in the NTD allosterically
foster non-covalent interactions between S1 and S2 forming more stabilised protomers before
engagement with ACE2. This process ultimately leads to active S1 shedding and exposure of the
fusion peptide. D614G emerged as a positively selected mutation in an early stage of the pandemic
persisting in subsequent lineages by stabilising trimeric spike. The more stable form of S1 is
therefore better positioned to engage with ACE2 (Daniloski et al., 2021; Díaz-Salinas et al., 2022;
Yurkovetskiy et al., 2020; Zhang et al., 2021b, 2020). It is reasonable to speculate that the effect of
the NTD mutations is analogous to the acquisition of D614G by strengthening the intramolecular

interactions to prevent S1 from being loosened prematurely while bolstering the RBD in an ACE2-
accessible form.

Finally we showed that Delta NTD revertant mutations (G156E and 157F/158R repair) were able to
compromise cell entry and increase sensitivity to neutralisation by vaccine sera. Reassuringly, a
similar finding was recently reported in breakthrough B.1.617 viruses where the same mutations in
the NTD accounted for the attenuated neutralisation sensitivity (Mishra et al., 2022).

In summary, our data support the cooperativity between NTD, RBD and the furin cleavage site. We
propose that the NTD allosterically affects the conformation of the RBD for ACE2 binding and the
furin cleavage site for spike cleavage. The pre-cleaved spike confers specificity on virus entry via
TMPRSS2 usage and drives a faster virus spread with more efficient spike-mediated fusion. Our
model explains the dominance of Delta over Kappa by being highly immune evasive on the one hand
through the acquisition of known escape mutations, while on the other hand gaining S1/S2 stability
and increased infectivity in lung cells. Our study highlights the importance of the continuous effort in
monitoring both RBD and NTD mutations in order to understand the biology of variants, and possibly
explains the lack of dominant SARS-CoV-2 inter-variant recombinants bearing breakpoints within
spike. On the translational/therapeutic side, combination antibody therapy targeting both NTD and
RBD might therefore be less prone to resistance than monotherapy or combinations targeting the
RBD alone.

Acknowledgments

RKG is supported by a Wellcome Trust Senior Fellowship in Clinical Science (WT108082AIA). This
study was supported by the Cambridge NIHRB Biomedical Research Centre, Addenbrooke's
Charitable Trust and the Rosetrees Trust. We would like to thank Paul Lehner for Calu-3, James Voss
for HeLa ACE2, Simon Cook for H1299 and Suzanne Rihn for the A549 cells. We are grateful to Leo
James for 293T-ACE2-ΔTMPRSS2, 293T-TMPRSS2, Vero-GFP1-10 and 293T-GFP11 cells and to Guido
Papa and Anna Albecka for the help on imaging. The authors acknowledge plasmids from the G2P-
UK National Virology consortium funded by MRC/UKRI (grant ref: MR/W005611/1), and thank Tom
Peacock and Wendy Barclay. The views expressed are those of the author(s) and not necessarily
those of the NIHR.

Methods

Cells and plasmids

Calu-3 (a human lung epithelial cell line; a gift from Paul Lehner) cells were maintained in Eagle's minimum essential medium containing 10% FBS and 1% PS. Vero-ACE2/TMPRSS2 cells (a gift from Emma Thomson), Hela-ACE2 (a gift from James Voss) and A549-ACE2/TMPRSS2 (a gift from Massimo Palmarini were maintained in Dulbecco's modified Eagle's medium (DMEM) containing 10% FBS and 1% PS. 293T (CRL-3216) and its derivative cell lines including 293TACE2ΔTMPRSS2, 293T-TMPRSS2 and 293T-GFP11 have been described previously (Papa et al., 2021). All the 293T cell lines as well as Vero-GFP1-10 were maintained in DMEM with 10% FBS and 1% PS. All cells were regularly tested and are mycoplasma free. Airway epithelial organoids were obtained and maintained as previously described (Meng et al., 2022; Youk et al., 2020). Human distal lung parenchymal tissues were obtained from adult donors with no background lung pathologies from Papworth Hospital Research Tissue Bank (T02233). Airway organoids were cultured in 48-well plate and were passaged every 2 weeks as previously reported (Meng et al., 2022).

pCDNA_SARS_CoV2_D416G_S WT, Delta, Kappa, Omicron BA.1 and Omicron BA.2 plasmids with 19 amino acid deletion at the C-terminus were generated by gene synthesis. For the construction of the chimeras, the region encompassing the NTD was digested with Bsu36I and HindIII (both NEB) before being gel purified and ligated back to the respective backbone that was cut with the same pair of restriction enzymes. Amino acid substitutions in the Delta NTD (D142G, G156E and repair of 157F and 158R) were introduced into the pCDNA_SARS-CoV-2_ D614G_Delta_S plasmid using the QuikChange Lightning Site-Directed Mutagenesis kit, following the manufacturer's instructions (Agilent Technologies). Sequences were checked by Sanger sequencing.

Pseudotype virus preparation and infectivity titration

Plasmids encoding the spike protein of SARS-CoV-2_D614G with a C terminal 19 amino acid deletion were used. Viral vectors were prepared by transfection of 293T cells by using Fugene HD transfection reagent (Promega) as described previously (Kemp et al., 2021). In brief, 293T cells were transfected with a mixture of 11ul of Fugene HD, 1µg of pCDNAΔ19 spike, 1ug of p8.91 HIV-1 gag-pol expression vector and 1.5µg of pCSFLW (expressing the firefly luciferase reporter gene with the HIV-1 packaging signal). Viral supernatant was collected at 48 h after transfection, filtered through 0.45um filter and stored at -80°C. Infectivity was measured by luciferase detection (Bright Glo; Promega) in target cells. The raw readings (in relative light unit (RLU)) were then normalised with the SG-PERT and plotted using GraphPad Prism.

PV SG-PERT

The SARS-CoV2 spike-pseudotyped viruses containing supernatants were standardised using a SYBR Green-based product-enhanced PCR assay (SG-PERT) as described previously (Pizzato et al., 2009). Briefly, 10-fold dilutions of virus supernatant were lysed in a 1:1 ratio in a 2x lysis solution (made up of 40% glycerol v/v 0.25% Triton X-100 v/v 100mM KCl, RNase inhibitor 0.8 U/ml, TrisHCL 100mM, buffered to pH7.4) for 10 minutes at room temperature. Sample lysates (12 µl) were added to 13 µl of SYBR Green master mix (containing 0.5µM of MS2-RNA Fwd and Rev primers, 3.5pmol/ml of MS2-RNA, and 0.125U/µl of Ribolock RNAse inhibitor and cycled in a QuantStudio (Thermofisher). Relative amounts of reverse transcriptase activity were determined as the rate of transcription of bacteriophage MS2 RNA, with absolute RT activity calculated by comparing the relative

393 amounts of RT to an RT standard of known activity.

Western blot

For cell lysates, 293 cells were washed and lysed in lysis buffer (Cell Signalling)
and lysates were diluted with 4 × sample buffer (Biorad) and boiled for 10 m before
subjected to western blotting. For virion purification, clarified supernatants were loaded onto a thin
layer of 8.4% optiprep density gradient medium (Sigma-Aldrich) and placed in a TLA55 rotor
(Beckman Coulter) for ultracentrifugation for 2 hours at 20,000 rpm. The pellet was then
resuspended for western blotting. For protein detection, the following antibodies were used:
rabbit anti-SARS-CoV-2 S monoclonal antibody (PA1-41165; Thermofisher), mouse anti-
SARS-CoV-2 S1 (MAB105403, R&D systems), rabbit anti-GAPDH polyclonal antibody (10494-1-AP;
Proteintech), horseradish peroxidase (HRP)-conjugated anti-rabbit and anti-mouse IgG polyclonal
antibody (Cell Signalling). Chemiluminescence was detected using ChemiDoc Touch Imaging System
(Bio-Rad). The cleavage ratio of S1 or S2 to FL in virions was determined by densitometry using
ImageJ (NIH).

Drug assay

A549-ACE2-TMPRSS2 (A549-A2T2) cells or human airway organoids were either E64D (Tocris) or
camostat (Sigma-Aldrich) treated for 2 hours at each drug concentration before the addition of a
comparable amount of input viruses pseudotyped with Delta, Kappa or chimeras (approx. 100,000
RLU). The cells were then left for 48 hours before the addition of substrate for luciferase
(Promega) and read on a Glomax plate reader (Promega). The RLU was normalised against the no-
drug control which was set as 100%.

Cell-cell fusion assay

Cell-cell fusion assays were described previously (Meng et al., 2022). Briefly, 293T GFP11 and Vero-
GFP1-10 cells were seeded at 80% confluence in a 1:1 ratio in 48 multiwell plate the day before.
Cells were co-transfected with 250 ng of spike expression plasmids using Fugene
6 following the manufacturer's instructions (Promega). Cell-cell fusion was measured using
an Incucyte and determined as the proportion of green area to total phase area over time. Data
were normalised to non-transfected control. Graphs were generated using Prism 9 software.

Neutralisation assay

426 This was performed as previously described (Collier et al., 2021b, 2021a).

References

Barton, M.I., MacGowan, S.A., Kutuzov, M.A., Dushek, O., Barton, G.J., and van der Merwe, P.A.
(2021). Effects of common mutations in the SARS-CoV-2 Spike RBD and its ligand, the human ACE2
receptor on binding affinity and kinetics. Elife *10*. https://doi.org/10.7554/eLife.70658.

Braga, L., Ali, H., Secco, I., Chiavacci, E., Neves, G., Goldhill, D., Penn, R., Jimenez-Guardeño, J.M.,
Ortega-Prieto, A.M., Bussani, R., et al. (2021). Drugs that inhibit TMEM16 proteins block SARS-CoV-2
spike-induced syncytia. Nature *594*, 88–93. https://doi.org/10.1038/s41586-021-03491-6.

Buchrieser, J., Dufloo, J., Hubert, M., Monel, B., Planas, D., Rajah, M.M., Planchais, C., Porrot, F.,
Guivel-Benhassine, F., van der Werf, S., et al. (2021). Syncytia formation by SARS-CoV-2-infected cells.
The EMBO Journal *40*. https://doi.org/10.15252/embj.2020107405.

Cerutti, G., Guo, Y., Zhou, T., Gorman, J., Lee, M., Rapp, M., Reddem, E.R., Yu, J., Bahna, F., Bimela, J.,
et al. (2021). Potent SARS-CoV-2 neutralizing antibodies directed against spike N-terminal domain
target a single supersite. Cell Host & Microbe *29*, 819-833.e7.
https://doi.org/10.1016/j.chom.2021.03.005.

Chi, X., Yan, R., Zhang, J., Zhang, G., Zhang, Y., Hao, M., Zhang, Z., Fan, P., Dong, Y., Yang, Y., et al.
(2020). A neutralizing human antibody binds to the N-terminal domain of the Spike protein of SARS-
CoV-2. Science (1979) *369*, 650–655. https://doi.org/10.1126/science.abc6952.

Collier, D.A., de Marco, A., Ferreira, I.A.T.M., Meng, B., Datir, R.P., Walls, A.C., Kemp, S.A., Bassi, J.,
Pinto, D., Silacci-Fregni, C., et al. (2021a). Sensitivity of SARS-CoV-2 B.1.1.7 to mRNA vaccine-elicited
antibodies. Nature *593*, 136–141. https://doi.org/10.1038/s41586-021-03412-7.

Collier, D.A., Ferreira, I.A.T.M., Kotagiri, P., Datir, R.P., Lim, E.Y., Touizer, E., Meng, B., Abdullahi, A.,
Baker, S., Dougan, G., et al. (2021b). Age-related immune response heterogeneity to SARS-CoV-2
vaccine BNT162b2. Nature *596*, 417–422. https://doi.org/10.1038/s41586-021-03739-1.

Daniloski, Z., Jordan, T.X., Ilmain, J.K., Guo, X., Bhabha, G., Tenoever, B.R., and Sanjana, N.E. (2021).
The spike d614g mutation increases sars-cov-2 infection of multiple human cell types. Elife *10*, 1–16.
https://doi.org/10.7554/eLife.65365.

Dhar, M.S., Marwal, R., VS, R., Ponnusamy, K., Jolly, B., Bhoyar, R.C., Sardana, V., Naushin, S.,
Rophina, M., Mellan, T.A., et al. (2021). Genomic characterization and epidemiology of an emerging
SARS-CoV-2 variant in Delhi, India. Science (1979) *374*, 995–999.
https://doi.org/10.1126/science.abj9932.

Díaz-Salinas, M.A., Li, Q., Ejemel, M., Yurkovetskiy, L., Luban, J., Shen, K., Wang, Y., and Munro, J.B.
(2022). Conformational dynamics and allosteric modulation of the SARS-CoV-2 spike. Elife *11*.
https://doi.org/10.7554/eLife.75433.

Ferreira, I.A.T.M., Kemp, S.A., Datir, R., Saito, A., Meng, B., Rakshit, P., Takaori-Kondo, A., Kosugi, Y.,
Uriu, K., Kimura, I., et al. (2021). SARS-CoV-2 B.1.617 Mutations L452R and E484Q Are Not
Synergistic for Antibody Evasion. The Journal of Infectious Diseases
https://doi.org/10.1093/infdis/jiab368.

Gobeil, S.M.-C., Janowska, K., McDowell, S., Mansouri, K., Parks, R., Manne, K., Stalls, V., Kopp, M.F.,
Henderson, R., Edwards, R.J., et al. (2021). D614G Mutation Alters SARS-CoV-2 Spike Conformation
and Enhances Protease Cleavage at the S1/S2 Junction. Cell Reports *34*, 108630.
https://doi.org/10.1016/j.celrep.2020.108630.

Hoffmann, M., Kleine-Weber, H., and Pöhlmann, S. (2020a). A Multibasic Cleavage Site in the Spike
Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. Molecular Cell *78*, 779-784.e5.
https://doi.org/10.1016/j.molcel.2020.04.022.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S.,
Herrler, G., Wu, N.-H., Nitsche, A., et al. (2020b). SARS-CoV-2 Cell Entry Depends on ACE2 and
TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell *181*, 271-280.e8.
https://doi.org/10.1016/j.cell.2020.02.052.

Jackson, C.B., Farzan, M., Chen, B., and Choe, H. (2022). Mechanisms of SARS-CoV-2 entry into cells.
Nature Reviews Molecular Cell Biology *23*, 3–20. https://doi.org/10.1038/s41580-021-00418-x.

Johnson, B.A., Xie, X., Bailey, A.L., Kalveram, B., Lokugamage, K.G., Muruato, A., Zou, J., Zhang, X.,
Juelich, T., Smith, J.K., et al. (2021). Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis.
Nature *591*, 293–299. https://doi.org/10.1038/s41586-021-03237-4.

Kemp, S.A., Collier, D.A., Datir, R.P., Ferreira, I.A.T.M., Gayed, S., Jahun, A., Hosmillo, M., Rees-Spear,
C., Mlcochova, P., Lumb, I.U., et al. (2021). SARS-CoV-2 evolution during treatment of chronic
infection. Nature *592*, 277–282. https://doi.org/10.1038/s41586-021-03291-y.

Lempp, F.A., Soriaga, L.B., Montiel-Ruiz, M., Benigni, F., Noack, J., Park, Y.-J., Bianchi, S., Walls, A.C.,
Bowen, J.E., Zhou, J., et al. (2021). Lectins enhance SARS-CoV-2 infection and influence neutralizing
antibodies. Nature *598*, 342–347. https://doi.org/10.1038/s41586-021-03925-1.

Li, D., Edwards, R.J., Manne, K., Martinez, D.R., Schäfer, A., Alam, S.M., Wiehe, K., Lu, X., Parks, R.,
Sutherland, L.L., et al. (2021). In vitro and in vivo functions of SARS-CoV-2 infection-enhancing and
neutralizing antibodies. Cell *184*, 4203-4219.e32. https://doi.org/10.1016/j.cell.2021.06.021.

Liu, Y., Soh, W.T., Kishikawa, J., Hirose, M., Nakayama, E.E., Li, S., Sasai, M., Suzuki, T., Tada, A.,
Arakawa, A., et al. (2021). An infectivity-enhancing site on the SARS-CoV-2 spike protein targeted by
antibodies. Cell *184*, 3452-3466.e18. https://doi.org/10.1016/j.cell.2021.05.032.

Liu, Y., Liu, J., Plante, K.S., Plante, J.A., Xie, X., Zhang, X., Ku, Z., An, Z., Scharton, D., Schindewolf, C.,
et al. (2022). The N501Y spike substitution enhances SARS-CoV-2 infection and transmission. Nature
602, 294–299. https://doi.org/10.1038/s41586-021-04245-0.

McCallum, M., Walls, A.C., Sprouse, K.R., Bowen, J.E., Rosen, L.E., Dang, H. v., de Marco, A., Franko,
N., Tilles, S.W., Logue, J., et al. (2021a). Molecular basis of immune evasion by the Delta and Kappa
SARS-CoV-2 variants. Science (1979) *374*, 1621–1626. https://doi.org/10.1126/science.abl8506.

McCallum, M., de Marco, A., Lempp, F.A., Tortorici, M.A., Pinto, D., Walls, A.C., Beltramello, M., Chen,
A., Liu, Z., Zatta, F., et al. (2021b). N-terminal domain antigenic mapping reveals a site of
vulnerability for SARS-CoV-2. Cell *184*, 2332-2347.e16. https://doi.org/10.1016/j.cell.2021.03.028.

McCarthy, K.R., Rennick, L.J., Nambulli, S., Robinson-McCarthy, L.R., Bain, W.G., Haidar, G., and
Duprex, W.P. (2021). Recurrent deletions in the SARS-CoV-2 spike glycoprotein drive antibody
escape. Science (1979) *371*, 1139–1142. https://doi.org/10.1126/science.abf6950.

Meng, B., Kemp, S.A., Papa, G., Datir, R., Ferreira, I.A.T.M., Marelli, S., Harvey, W.T., Lytras, S.,
Mohamed, A., Gallo, G., et al. (2021). Recurrent emergence of SARS-CoV-2 spike deletion H69/V70
and its role in the Alpha variant B.1.1.7. Cell Reports *35*.
https://doi.org/10.1016/j.celrep.2021.109292.

Meng, B., Abdullahi, A., Ferreira, I.A.T.M., Goonawardane, N., Saito, A., Kimura, I., Yamasoba, D.,
Gerber, P.P., Fatihi, S., Rathore, S., et al. (2022). Altered TMPRSS2 usage by SARS-CoV-2 Omicron
impacts infectivity and fusogenicity. Nature *603*, 706–714. https://doi.org/10.1038/s41586-022-
04474-x.

Mishra, T., Dalavi, R., Joshi, G., Kumar, A., Pandey, P., Shukla, S., Mishra, R.K., and Chande, A. (2022).
SARS-CoV-2 spike E156G/Δ157-158 mutations contribute to increased infectivity and immune escape.
Life Science Alliance *5*, e202201415. https://doi.org/10.26508/lsa.202201415.

Mlcochova, P., Collier, D., Ritchie, A., Assennato, S.M., Hosmillo, M., Goel, N., Meng, B., Chatterjee,
K., Mendoza, V., Temperton, N., et al. (2020). Combined Point-of-Care Nucleic Acid and Antibody

Testing for SARS-CoV-2 following Emergence of D614G Spike Variant. Cell Reports Medicine *1*,
100099. https://doi.org/10.1016/j.xcrm.2020.100099.

Mlcochova, P., Kemp, S., Dhar, M.S., Papa, G., Meng, B., Ferreira, I.A.T.M., Datir, R., Collier, D.A.,
Albecka, A., Singh, S., et al. (2021). SARS-CoV-2 B.1.617.2 Delta variant replication and immune
evasion. Nature https://doi.org/10.1038/s41586-021-03944-y.

Ou, T., Mou, H., Zhang, L., Ojha, A., Choe, H., and Farzan, M. (2021a). Hydroxychloroquine-mediated
inhibition of SARS-CoV-2 entry is attenuated by TMPRSS2. PLOS Pathogens *17*, e1009212.
https://doi.org/10.1371/journal.ppat.1009212.

Ou, T., Mou, H., Zhang, L., Ojha, A., Choe, H., and Farzan, M. (2021b). Hydroxychloroquine-mediated
inhibition of SARS-CoV-2 entry is attenuated by TMPRSS2. PLOS Pathogens *17*, e1009212.
https://doi.org/10.1371/journal.ppat.1009212.

Papa, G., Mallery, D.L., Albecka, A., Welch, L.G., Cattin-Ortolá, J., Luptak, J., Paul, D., McMahon, H.T.,
Goodfellow, I.G., Carter, A., et al. (2021). Furin cleavage of SARS-CoV-2 Spike promotes but is not
essential for infection and cell-cell fusion. PLOS Pathogens *17*, e1009246.
https://doi.org/10.1371/journal.ppat.1009246.

Park, J.E., Li, K., Barlan, A., Fehr, A.R., Perlman, S., McCray, P.B., and Gallagher, T. (2016). Proteolytic
processing of middle east respiratory syndrome coronavirus spikes expands virus tropism. Proc Natl
Acad Sci U S A *113*, 12262–12267. https://doi.org/10.1073/pnas.1608147113.

Peacock, T.P., Goldhill, D.H., Zhou, J., Baillon, L., Frise, R., Swann, O.C., Kugathasan, R., Penn, R.,
Brown, J.C., Sanchez-David, R.Y., et al. (2021). The furin cleavage site in the SARS-CoV-2 spike protein
is required for transmission in ferrets. Nature Microbiology *6*, 899–909.
https://doi.org/10.1038/s41564-021-00908-w.

Peng, R., Wu, L.-A., Wang, Q., Qi, J., and Gao, G.F. (2021). Cell entry by SARS-CoV-2. Trends in
Biochemical Sciences *46*, 848–860. https://doi.org/10.1016/j.tibs.2021.06.001.

Pizzato, M., Erlwein, O., Bonsall, D., Kaye, S., Muir, D., and McClure, M.O. (2009). A one-step SYBR
Green I-based product-enhanced reverse transcriptase assay for the quantitation of retroviruses in
cell culture supernatants. Journal of Virological Methods *156*, 1–7.
https://doi.org/10.1016/j.jviromet.2008.10.012.

Qing, E., Kicmal, T., Kumar, B., Hawkins, G.M., Timm, E., Perlman, S., and Gallagher, T. (2021).
Dynamics of sars-cov-2 spike proteins in cell entry: Control elements in the amino-terminal domains.
MBio *12*. https://doi.org/10.1128/mBio.01590-21.

Qing, E., Li, P., Cooper, L., Schulz, S., Jäck, H.-M., Rong, L., Perlman, S., and Gallagher, T. (2022). Inter-
domain communication in SARS-CoV-2 spike proteins controls protease-triggered cell entry. Cell
Reports *39*, 110786. https://doi.org/10.1016/j.celrep.2022.110786.

Rajah, M.M., Hubert, M., Bishop, E., Saunders, N., Robinot, R., Grzelak, L., Planas, D., Dufloo, J.,
Gellenoncourt, S., Bongers, A., et al. (2021). SARS-CoV-2 Alpha, Beta, and Delta variants display
enhanced Spike-mediated syncytia formation. The EMBO Journal *40*.
https://doi.org/10.15252/embj.2021108944.

Ramanathan, M., Ferguson, I.D., Miao, W., and Khavari, P.A. (2021). SARS-CoV-2 B.1.1.7 and B.1.351
spike variants bind human ACE2 with increased affinity. Lancet Infect Dis *21*, 1070.
https://doi.org/10.1016/S1473-3099(21)00262-0.

Saville, J.W., Mannar, D., Zhu, X., Srivastava, S.S., Berezuk, A.M., Demers, J.-P., Zhou, S., Tuttle, K.S.,
Sekirov, I., Kim, A., et al. (2022). Structural and biochemical rationale for enhanced spike protein
fitness in delta and kappa SARS-CoV-2 variants. Nature Communications *13*, 742.
https://doi.org/10.1038/s41467-022-28324-6.

Simon-Loriere, E., and Schwartz, O. (2022). Towards SARS-CoV-2 serotypes? Nature Reviews
Microbiology *20*, 187–188. https://doi.org/10.1038/s41579-022-00708-x.

Supasa, P., Zhou, D., Dejnirattisai, W., Liu, C., Mentzer, A.J., Ginn, H.M., Zhao, Y., Duyvesteyn, H.M.E.,
Nutalai, R., Tuekprakhon, A., et al. (2021). Reduced neutralization of SARS-CoV-2 B.1.1.7 variant by
convalescent and vaccine sera. Cell *184*, 2201-2211.e7. https://doi.org/10.1016/j.cell.2021.02.033.

Suryadevara, N., Shrihari, S., Gilchuk, P., VanBlargan, L.A., Binshtein, E., Zost, S.J., Nargi, R.S., Sutton,
R.E., Winkler, E.S., Chen, E.C., et al. (2021). Neutralizing and protective human monoclonal
antibodies recognizing the N-terminal domain of the SARS-CoV-2 spike protein. Cell *184*, 2316-
2331.e15. https://doi.org/10.1016/j.cell.2021.03.029.

Suzuki, R., Yamasoba, D., Kimura, I., Wang, L., Kishimoto, M., Ito, J., Morioka, Y., Nao, N., Nasser, H.,
Uriu, K., et al. (2022). Attenuated fusogenicity and pathogenicity of SARS-CoV-2 Omicron variant.
Nature *603*, 700–705. https://doi.org/10.1038/s41586-022-04462-1.

Ulrich, L., Halwe, N.J., Taddeo, A., Ebert, N., Schön, J., Devisme, C., Trüeb, B.S., Hoffmann, B., Wider,
M., Fan, X., et al. (2022). Enhanced fitness of SARS-CoV-2 variant of concern Alpha but not Beta.
Nature *602*, 307–313. https://doi.org/10.1038/s41586-021-04342-0.

V'kovski, P., Kratzel, A., Steiner, S., Stalder, H., and Thiel, V. (2021). Coronavirus biology and
replication: implications for SARS-CoV-2. Nature Reviews Microbiology *19*, 155–170.
https://doi.org/10.1038/s41579-020-00468-6.

Whittaker, G.R., Daniel, S., and Millet, J.K. (2021). Coronavirus entry: how we arrived at SARS-CoV-2.
Current Opinion in Virology *47*, 113–120. https://doi.org/10.1016/j.coviro.2021.02.006.

Yamasoba, D., Kimura, I., Nasser, H., Morioka, Y., Nao, N., Ito, J., Uriu, K., Tsuda, M., Zahradnik, J.,
Shirakawa, K., et al. (2022). Virological characteristics of the SARS-CoV-2 Omicron BA.2 spike. Cell
https://doi.org/10.1016/j.cell.2022.04.035.

Youk, J., Kim, T., Evans, K. v., Jeong, Y.-I., Hur, Y., Hong, S.P., Kim, J.H., Yi, K., Kim, S.Y., Na, K.J., et al.
(2020). Three-Dimensional Human Alveolar Stem Cell Culture Models Reveal Infection Response to
SARS-CoV-2. Cell Stem Cell *27*, 905-919.e10. https://doi.org/10.1016/j.stem.2020.10.004.

Yurkovetskiy, L., Wang, X., Pascal, K.E., Tomkins-Tinch, C., Nyalile, T.P., Wang, Y., Baum, A., Diehl,
W.E., Dauphin, A., Carbone, C., et al. (2020). Structural and Functional Analysis of the D614G SARS-
CoV-2 Spike Protein Variant. Cell *183*, 739-751.e8. https://doi.org/10.1016/j.cell.2020.09.032.

Zhang, J., Xiao, T., Cai, Y., Lavine, C.L., Peng, H., Zhu, H., Anand, K., Tong, P., Gautam, A., Mayer, M.L.,
et al. (2021a). Membrane fusion and immune evasion by the spike protein of SARS-CoV-2 Delta
variant. Science (1979) *374*, 1353–1360. https://doi.org/10.1126/science.abl9463.

Zhang, J., Cai, Y., Xiao, T., Lu, J., Peng, H., Sterling, S.M., Walsh, R.M., Rits-Volloch, S., Zhu, H.,
Woosley, A.N., et al. (2021b). Structural impact on SARS-CoV-2 spike protein by D614G substitution.
Science (1979) *372*, 525–530. https://doi.org/10.1126/science.abf2303.

Zhang, L., Jackson, C.B., Mou, H., Ojha, A., Peng, H., Quinlan, B.D., Rangarajan, E.S., Pan, A.,
Vanderheiden, A., Suthar, M.S., et al. (2020). SARS-CoV-2 spike-protein D614G mutation increases

virion spike density and infectivity. Nature Communications *11*. https://doi.org/10.1038/s41467-
020-19808-4.

Figure 1

Figure 1: Delta exhibits increased infectivity over Kappa in Calu3 cells and is dependent on the NTD. (A): Schematic diagrams of WT, Kappa and Delta with their chimeras bearing swapped NTD. The consensus mutations between Kappa and Delta are annotated in blue. The monomeric spikes shown on the right-hand side are for illustration purposes. (B): PV bearing Delta or Kappa or chimeric spike was used to transduce Calu3 and organoids expressing endogenous levels of ACE2 and TMPRSS2 and ACE2/TMPRSS2 overexpressing cell lines including Hela-ACE2, Vero-ACE2/TMPRSS2 and A549-ACE2/TMPRSS2. Unpaired t test. (C): WT, WT with KappaNTD and WT with DeltaNTD were used to transduce Calu3 cells. In B&C, mean +/- SEM are shown for technical replicates (n=2-4). Two-sided unpaired Student t test, ns: not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Data are representative of at least two experiments.

Figure 2

Figure 2: The NTD modulates the usage of TMPRSS2 and ACE2 for virus entry. (A): A representative western blot of cell lysates showing spike cleavage in Delta, Kappa and its chimeras. GAPDH was probed as a loading control. (B): The band intensity from (A) was densitometrically calculated using ImageJ. Total S2 associated spike proteins (S2 and FL) were then normalized against GAPDH across Delta, Kappa and NTD bearing chimeras. (C): The ratio of S2/FL+S2 from the band intensity of S2 and FL shown in (A) was plotted. n≥3. (D): Delta, Kappa or chimeras was transduced into either parental 293 cells or 293T cells overexpressing TMPRSS2. The fold increase of the virus entry in T2 overexpressing cells over parental cells is shown above the scatter plots. mean +/- SEM are shown for technical replicates (n=4). Two-sided unpaired Student t test, ***p<0.001. Data are representative of two experiments. (E): The entry efficiency of Delta and Kappa in A549-ACE2/TMPRSS2 cells in the presence of TMPRSS2 specific inhibitor camostat or cathepsin specific inhibitor E64D. The lentivirus pseudotyped with VSV-G was used as a control. The RLU was normalized with non-drug added control giving a percentage of infection. The data showing the mean±SD from 4 experiments were plotted. (F): The entry efficiency of Delta and Kappa in the presence of camostat in airway organoids. Data are representative of two experiments. (G): Delta, Kappa or chimeras was transduced into either parental 293T cells or 293T cells overexpressing ACE2 with abrogated TMPRSS2 expression (A2delT2). The fold increase of the virus entry in A2delT2 overexpressing cells over parental cells is shown above the scatter plots. mean +/- SEM are shown for technical replicates (n=4). Two-sided unpaired Student t test, ***p<0.001, ****p<0.0001. Data are representative of two experiments.

Figure 3

Figure 3: Reversing the mutations in the Delta NTD reduces the infectivity in lung cells and increases the neutralisation sensitivity to the vaccine-elicited antibody. (A): PV bearing Delta and its reversions were used to transduce Calu3 and Hela-ACE2 cells. mean +/- SEM are shown for technical replicates (n=4). Two-sided unpaired Student t test. Data are representative of two experiments. (B): Examples of neutralisation curves from ID32, 63 and 105 vaccinees with PVs bearing the reversion at 142, 156 or 157/8. (C): The 50% serum neutralisation was plotted across ten sera showing the geometric mean with geometric SD. Paired Wilcoxon was used for analysis. Data are representative of two experiments. ns: not significant, *p<0.05, **p<0.01.

Figure 4

Figure 4: The SARS-CoV-2 Delta NTD increases the fusion kinetics of Kappa and WT spikes. (A): A schematic diagram showing the split GFP system for spike-ACE2 mediated cell fusion. (B): 681R or 681H is required for the enhanced fusogenicity in Delta and its chimera bearing Kappa NTD. (C): The fused Delta NTD in Kappa and WT increased the fusion kinetics of their counterparts, respectively. The line graphs on the right show the percentage of the positive GFP area at 12, 14, 16, 20, 22 and 23 hr post transfection. The data showing the SEM at each point were averaged from two experiments. The heatmap at each time point shows the mean of the GFP positive area over the field of view from two experiments.

Figure 5

Figure 5: The SARS-CoV-2 Delta NTD or BA.2 NTD does not alter spike fusion or sensitivity to TMPRSS2 of BA.1. (A): PV bearing Delta, BA.1, BA.2 or chimeric forms of BA.1 and BA.2 spike were used to transduce Calu3, H1299 and 293T expressing endogenous levels of ACE2 and TMPRSS2 and TMPRSS2 overexpressing 293T cells. (B): The fusion kinetics of the fused Delta NTD in BA.1 and BA.2 along with their parental spikes. The western blot showing cleavage of spike is directly underneath the heatmap. (C): BA.1, BA.2 or chimeras with Delta were transduced into either parental 293 cells or 293T cells overexpressing TMPRSS2. The fold increase of the virus entry in TMPRSS2 overexpressing cells over parental cells is shown above the scatter plots. (D): The fusion assay of BA.1, BA.2 and their chimeric BA.1 bearing BA.2 NTD and BA.2 bearing BA.1 NTD. The line graphs show the percentage of the positive GFP area at 1 hr interval post transfection. The data showing the SEM at each point were averaged from two experiments. (E): BA.1, BA.2 or chimeras bearing BA.2 NTD or BA.1 NTD together with Delta were transduced into either parental 293 cells or 293T cells overexpressing TMPRSS2. The fold increase of the virus entry in T2 overexpressing cells over parental cells is shown above the scatter plots. In A, C&E, the plots are representative from two experiments. mean +/- SEM are shown for technical replicates (n=4). Two-sided unpaired Student t test, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure S1

A

Virions

B

C

D

Virions

Supplementary Figure 1: The NTD is required for efficient cleavage of SARS-CoV-2 spike. (A): Western blots of purified PV bearing either H69V70 deletion, WT, Kappa or Delta spikes. The sizes of protein markers were labelled to the left of the blot and the corresponding bands were labelled to the right. (B): The intensity of the spike-associated bands on the western blots was densitometrically quantified (ImageJ) before the ratio was calculated for cleavage (S2/FL, paired t test.) (B) or spike stability (S1/S2; one sample t test) (C). In both (B) and (C) each dot represents one PV preparation. (D): A representative western blot of the purified PVs bearing RBD mutations only in Kappa and Kappa spike, together with WT and non-spike control. *p<0.05.

Figure S2

A

Supplementary Figure 2: (A) The infectivity of WT, WT with Kappa NTD or Delta NTD chimeras in A549-A2/T2 cells (A) and western blot showing spike cleavage (B).

Figure S3

Supplementary Figure 3: The Delta NTD fused with Kappa increases camostat resistance to a similar level as Delta.