Generalizable strategy to analyze domains in the context of parent protein architecture: A CheW case study

Luke R. Vass[1,#a], Katie M. Branscum[#b], Robert B. Bourret[1*], Clay A. Foster[1,#b]

[1]Department of Microbiology & Immunology, University of North Carolina, Chapel Hill, North Carolina, United States of America

[#a]Current Address: Department of Pathology, University of Virginia, Charlottesville, Virginia, United States of America

[#b]Current Address: Department of Pediatrics, Section Hematology/Oncology, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, United States of America

Running Title: Domain variation with protein architecture

*Corresponding author

Email: bourret@med.unc.edu

Co-author email addresses:

Luke R. Vass: lukev3@vt.edu

Katie M. Branscum: katief@tribaldiagnostics.com

Clay A. Foster: clay-foster@ouhsc.edu

Data Availability Statement:

The data that support the findings of this study were derived from the following resource

available in the public domain: Pfam (release v.33), http://pfam.xfam.org.

Conflict of Interest Statement:

The authors declare no conflict of interest.

ABSTRACT

Domains are the three-dimensional building blocks of proteins. An individual

domain can occur in a variety of domain architectures that perform unique functions and

are subject to different evolutionary selective pressures. We describe an approach to

evaluate the variability in amino acid sequences of a single domain across architectural

contexts. The ability to distinguish different evolutionary outcomes of one protein

domain can help determine whether existing knowledge about a specific domain will

apply to an uncharacterized protein, lead to insights and hypotheses about function, and

guide experimental priorities.

We developed and tested our approach on CheW-like domains (PF01584), which

mediate protein/protein interactions and are difficult to compare experimentally. CheW-

like domains occur in CheW scaffolding proteins, CheA kinases, and CheV proteins that

regulate bacterial chemotaxis. We analyzed 16 domain architectures that included 94%

of all CheW-like domains found in nature. We identified six *Classes* of CheW-like

domains with presumed functional differences. CheV and most CheW proteins

contained *Class* 1 domains, whereas some CheW proteins contained *Class* 6 (~20%) or

Class 2 (~1%) domains instead. Most CheA proteins contained *Class* 3 domains. CheA

proteins with multiple Hpt domains contained *Class* 4 domains. CheA proteins with two

CheW-like domains contained one *Class* 3 and one *Class* 5.

We also created SimpLogo, an innovative method for visualizing amino acid

composition across large sets of multiple sequence alignments of arbitrary length.

SimpLogo offers substantial advantages over standard sequence logos for comparison

and analysis of related protein sequences. The R package for SimpLogo is freely

available.

KEYWORDS

protein domain, domain architecture, CheW, CheA, CheV, SimpLogo

1 | INTRODUCTION

Many proteins consist of a chain of discrete domains. Protein domains are typically compact, independently folding units. Originally, protein domains were defined based on protease resistance; however, they are now recognized by characteristic amino acid sequence features. Individual domains possess structures and functions that are subject to natural selection and thus are discrete units of evolution. Domains were assembled in many permutations during evolution to create different proteins. Therefore, domains can be considered modular "building blocks" of proteins. The Protein Family (Pfam) database currently contains almost 20,000 different domains.[1]

Identifying domains is useful because once nature has evolved the solution to a problem (e.g., a protein domain that performs a specific function), the solution is frequently repeated. Thus, domain identity can provide useful general predictions about the structure and function of domains in experimentally uncharacterized proteins, as well as providing an effective means by which to compare proteins of known function. In recent years, more complex techniques have been proposed for comparing and quantifying homology between entire multi-domain proteins, including numerous alignment-free methodologies.[2,3] Such approaches can successfully group functional proteins (such as kinases) at higher resolutions than standard sequence analyses by incorporating information on both catalytic domains and on attached accessory domains. However, much of the practical value of such predictions and our overall understanding of protein function still relies on the accuracy of discrete, individual domain descriptions (like those found in Pfam). Such domain families are often considered to be monolithic. While useful in some respects, in actuality substantial

differences can be found between domains of the same type in different proteins. For

example, response regulator proteins are formed by linking a receiver domain

(Response_reg, PF00072) to a wide variety of output domains. At least three types of

evidence demonstrate the differences between receiver domains in different types of

response regulators: (i) phylogenetic analysis shows receiver domains co-evolve with

their output domains;[4,5] (ii) receiver domain:output domain interaction surfaces and

receiver domain dimerization surfaces vary as a function of binding partner and whether

a receiver domain exerts positive or negative regulation on its output domains;[6] and (iii)

receiver domains support phosphorylation and dephosphorylation reaction kinetics that

differ by many orders of magnitude depending on the timescale of the associated

biological function.[7-10] Thus, receiver domains, though termed the same, differ

substantially depending upon the domain architecture in which they are found.

Variation between different instances of a single specific protein domain raises

multiple questions. If the structure and/or function of a domain are experimentally

characterized in one protein, then how well does that information translate into useful

insights about the same individual domain in another protein? In other words, how do

protein domains themselves vary in the contexts of different domain architectures?

Genome sequencing has far outstripped experimental capacity and provides a rich

source of relevant information. Can sophisticated methods of amino acid sequence

analysis identify contexts (i.e., location within domain architectures) in which current

knowledge about a specific protein domain applies? Conversely, given the enormous

scope of uncharacterized proteins and domain architectures, can we identify contexts

where current knowledge about a specific domain is not applicable? Such a method

would provide a way to inspire and focus future experimental investigations of protein

domains in a manner that is most likely to be informative and may also lead to useful

hypotheses and testable predictions. We describe here a generally applicable approach

to analyze the variation in a specific protein domain as a function of the contexts in

which the domain is found, revealing residue-level insights at a substantially higher

resolution than a typical Pfam domain assignment.

As a test case to illustrate our method, we chose CheW-like domains (CheW,

PF01584), which are part of the signal transduction pathways that regulate bacterial

chemotaxis.[11] CheW-like domains are known to act as scaffolds for various

protein/protein interactions.[12,13] The absences of an active site, substrate or ligand

binding, or enzymatic activity make CheW-like domains challenging to study

experimentally, because no simple assay exists by which to compare the properties of

different CheW-like domains. One might envision measuring binding constants, but it

would be difficult to untangle the contributions of the CheW-like domains and their

partner proteins. Furthermore, CheW-like domains are used to assemble very large

multi-protein complexes that couple chemoreceptors to kinases, so the biological

relevance of the binding constant between an isolated CheW-like domain and its

receptor or kinase partner is not immediately obvious. Finally, evolutionary constraints

on CheW function are such that sequences can diverge substantially and still be

functionally interchangeable.[14] Thus, an objective method to classify CheW-like

domains within different contexts would be particularly helpful to clarify the

circumstances in which current knowledge is likely to be applicable and those which

would benefit from further investigation.

CheW-like domains are found in three broad lineages of domain architectures:

CheW scaffold proteins, which consist only of CheW-like domains; CheA kinases, which

combine CheW-like domains with domains that provide kinase activity and

phosphorylation sites; and CheV proteins, which combine CheW-like domains with

receiver domains. Notably, the function(s) of CheV proteins are poorly understood[13,15].

A recent review of the diversity of bacterial chemotaxis systems asked, "Can efficient

computational approaches be developed to distinguish … CheW from other CheW

domain-containing proteins?".[16] Our results suggest that the answer is yes, and that not

all CheW-like domains are alike.

2 | MATERIALS AND METHODS

2.1 | Protein sequence database

Protein sequences were initially obtained from the Pfam database (v.33).[1] Any sequence found within the Representative Proteome 35 (RP35) to contain the CheW-like domain (PF01584) was extracted for further use, retaining only the CheW-like domain sequence.[17] "35" refers to the co-membership threshold used to control the granularity of the sequence space represented. An RP set best represents the known sequence diversity of a group of similar proteomes/organisms. To construct the RP35 data set, available protein sequences for completed proteomes within the UniProtKB database were grouped based on their UniRef50 cluster membership. UniRef, or UniProt Reference Clusters, are sets of proteins grouped together by sequence in such a way as to produce tight clusters of orthologs/inparalogs, maximizing sequence space while minimizing redundancy; 50 refers to the sequence identity threshold used for the clustering procedure.[18] We chose RP35 as the best compromise between information diversity and sequence space reduction. CheW-like domain sequences were grouped into *Architectures* based on the domain organization of the parent protein (Figure 2). Rare *Architectures* exhibited by fewer than 40 sequences were excluded from the analysis for practical reasons and because trends identified using such sparse data would likely be difficult to interpret with accuracy.

2.2 | Sequence analysis and initial clustering

CheW-like domain sequences were collectively aligned to the accepted CheW-like domain HMM (PF01584) using HMMER3.[19] The removal of poorly aligned sequences or subsequences has been shown to be beneficial for large-scale,

downstream sequence analysis and phylogenetic profiling. To that end, we used trimAl

(version 1.2) to remove spuriously aligned regions (using a column gap-threshold of

90%, assuming that the top 10% most "gappy" columns were mostly noise).[20]

Sequences with >50% gap-characters were also removed.

To identify the presence of the sequence subpopulations within our data set, we

performed an initial cluster analysis on our full sequence set. Clustering of protein

sequences can be performed in a number of ways, most commonly by calculating

pairwise dissimilarity scores between sequences in an alignment and clustering the

resulting matrix. Estimating protein sequence (dis)similarity requires the use of a

substitution matrix to describe the rate or frequency at which a residue is substituted for

another over time, simulating the process of evolution. We tested four commonly used

evolutionary substitution models on our data set: GONNET, point accepted mutation

250 (PAM250; number of mutations per 100 amino acids), BLOcks SUbstitution Matrix

30 (BLOSUM30), and Jones-Taylor-Thornton (JTT).[21-24] Each is tailored towards higher

diversity sequence groups (the average sequence identity reported in the full Pfam

alignment for CheW-like domain PF01584 is only ~20%). We used the mat.dis() function

from the bios2mds package in R (version 1.2.3) to calculate four pairwise-dissimilarity

matrices from our full sequence alignment to use as inputs for k-medoids clustering.[25]

Clustering techniques like k-medoids requires an initial target number of clusters

(the k-value) as input. An optimal k-value can be estimated by various metrics, such as

the average silhouette width (a measure of how similar a point is to its own cluster

compared to other clusters), but choosing the "correct" number of clusters is

challenging, subjective and dependent on the overall goals of the analysis.[26] *A priori*, we

chose $k = 21$ (the total initial number of distinct architectural *Contexts*, including multiple

domains in the same protein, used to partition the CheW-like domain sequences). This

was an intentional overestimation of the true number of subpopulations. An optimal *k*

value (~3-4) was also estimated *a posteriori* using the average silhouette width method

implemented in the pamk() function from the fpc package in R (version 2.2-9; *k*=3-10).[27]

Initial *k*-medoids clustering was performed using the pam() function from the cluster

package within R (version 2.1.2; *k*-medoid clustering is sometimes called Partitioning

Around Medoids).[28]

For convenience, we only show solutions derived from the GONNET substitution

matrix in the main text. Solutions derived from the other three models are featured in the

Supporting Information. The similarity of the partitions generated using the four

substitution models was quantified using both the Adjusted Rand Index (ARI)[29,30] and

Normalized Mutual Information (NMI),[31,32] commonly used metrics when comparing

clustering schemes. Each of the four solutions was highly similar, with ARI scores

ranging from 0.6-0.7 and NMI scores of 0.7-0.8, suggesting a relatively stable and

generalizable underlying data structure.

2.3 | Multidimensional scaling analysis

The pairwise dissimilarities between the medoids for each of the 21 identified

Clusters were extracted and used as input for analysis with multidimensional scaling

(MDS). MDS is a multivariate dimensionality reduction and visualization technique that

takes pairwise similarities/dissimilarities/distances for a data set and attempts to

reconstruct a simplified map that maintains the relative positions of the data points and

is easily interpretable (i.e., in a lower *k*-dimensional space). Two main types of MDS

exist: metric MDS, which focuses on the "distances" in the matrix being analyzed, and

non-metric MDS, which assumes that onl the relative rankings of the distances are

known.

Due to its simplicity and previous success when applied to protein sequences,

we chose metric MDS[25] to visualize the results of the initial *k*-medoid clustering

procedure. For the subsequent subsampling steps (described below), we first tested an

implementation of the metric MDS algorithm featured in the wcmdscale() function from

the vegan package in R (version 2.5-7).[33] However, other forms of MDS, such as the

non-metric approach, may offer additional advantages under the appropriate

circumstances. Examination of our MDS solutions indicated that a significant proportion

of the total eigenvalues were negative (~15.4-15.7%). Though the individual negative

eigenvalues were miniscule compared to the positive entries, it was possible that our

initial dissimilarities were non-Euclidean in nature, and non-metric MDS might offer

additional insight. Consequently, we tested a non-metric implementation of the MDS

algorithm using the metaMDS() function, also from the vegan package (using

parameters: *trymax = 2000, noshare = F, maxit = 9999, sratmax = 0.99999999, k = 3,*

model = "local", autotransform = F, wascores = F, center = T, pc = T) [33]. The resulting

MDS plots (included in the Supporting Information) were visualized using the ggplot2

package in R (version 3.3.3).[34] While we encountered significant challenges achieving

convergence (likely due to the still relatively large sequence set), the non-metric MDS

solutions were largely reminiscent of the basic layouts generated using the metric

algorithm, even when sampling smaller subsets of sequences.

2.4 | Sequence subsampling strategy

A primary technical disadvantage of the MDS technique is its substantial

computational cost. The time complexity of standard metric MDS is proportional to the

square of the input size, indicating significant temporal and technical requirements when

analyzing a high-dimensional data set with thousands of data points.[35] Thus, while we

could effectively apply MDS to the cluster medoids from our limited initial analysis, the

approach becomes impractical when analyzing a larger, more complete data set like our

full dissimilarity matrix featuring ~8500 CheW-like domains.

One popular approach to dealing with large, unwieldy data is random

subsampling, called stratified sampling when working with distinct groups or strata

within the original population. However, for this strategy to be effective, adequate

representation of each stratum is essential. Guided by our initial *k*-medoids clustering

results (which effectively revealed any strata within the sequence data), each

architectural *Context* was treated as a separate homogeneous class, except for CheW.I,

which was split into CheW.IA, CheW.IB and CheW.IC *Types*. CheW-like domains from

parent *Architectures* featuring multiple domains were split into separate *Contexts* with

appropriate suffixes added relative to their N- to C-terminal position in the parent

protein. Two main approaches to stratified subsampling exist: randomly selecting a

predefined number of observations from each defined subgroup; or randomly selecting

a number of observations proportional to the original subgroup sizes. Given the wide

disparity in the number of CheW-like domain sequences of each *Type*, we chose the

former to facilitate computational efficiency. The mat.dis() function from the bios2mds R

package was again utilized for the iterative sampling procedure.[25] Each subsampling

round randomly selected 25 sequences from each of the 23 *Types* (without

replacement; smaller subsets were also explored with similar results). Sampling was

repeated 10 times (575 distinct sequences per iteration), generating a total of 10

replicate sequence sets for each of the four previously described substitution models, to

assess the stability of the various randomized MDS solutions. We also included the

CheW-like domains of 18 well-studied reference proteins as an internal constant for

each iteration (data not shown). Pairwise dissimilarity matrices were estimated for each

replicate set for further use during the MDS analysis described above.

2.5 | Conservation analysis and SimpLogo creation

All CheW-like domain sequences were first aligned collectively to the conserved

CheW-like Hidden Markov Model (HMM) profile (PF01584) obtained from the Pfam

database. Separate sequence alignments were then segmented based on

corresponding *Types* of CheW-like domains. For each alignment, the *Escherichia coli*

CheW sequence was inserted at the top for use as a reference. Columns introducing

gaps (relative to the *E. coli* CheW sequence) were removed. Position-wise count

matrices and corresponding frequencies were generated for each alignment using the

Biostrings and motifStack packages in R.[36,37] Physicochemically similar residue types

were combined to reduce visual complexity of the final representation, creating nine

amino acid groups (including gap characters): positively-charged side chains

(His/Lys/Arg), negatively-charged side chains (Asp/Glu), small volume side chains

(Ala/Gly), polar side chains (Gln/Asn/Ser/Thr), large hydrophobic side chains

(Met/Val/Leu/Ile), gaps (-), aromatic side chains (Phe/Tyr/Trp), Pro, and Cys. Each

amino acid category was assigned an "idealized" color and converted to tristimulus

values in the CIE XYZ color space. XYZ space is linear and non-negative (as opposed

to traditional RGB space), making it simple to mix colors. Final values were calculated

by weighting (multiplying) the idealized colors of each amino acid group by their relative

frequencies and summing the results for each position. Logos were visualized using

geom_tile objects (from the ggplot2 suite), with colors and heights for each position

based on the aforementioned blended tristimulus values and their overall gap character

frequency, respectively. "Idealized" (i.e., color at 100% frequency) color strips for the top

two most frequent residue types were added to each position (at the top and bottom of

each tile) to mitigate uncertainty between true qualitative colors and blended hues.

Position-wise information content (in bits) was estimated for each alignment again using

motifStack and visualized with a line trace at the top.[37] To create traditional sequence

logos (with similar dimensions, for comparison purposes), the Logolas R package was

used with default settings.[38] An R package for SimpLogo was created and is freely

available (along with an example and the sequence alignments used to generate Figure

5) in the Github repository https://www.github.com/clayfos/SimpLogo (v.0.1.1,

https://doi.org/10.5281/zenodo.6323714).

2.6 | Protein homology modeling and structural analysis

Crystal structures of the CheW-like domains from experimentally well-studied *E.*

coli CheW or CheA proteins are not available. To visualize the relevant functional

surfaces of the reference *E. coli* CheW-like domains CheA-P5 and full CheW protein,

we modeled trimeric CheW:CheA-P5:receptor complexes (capturing both interface 1

and 2) based on the crystal structure of the *Thermotoga maritima* multimer featuring

CheW, CheA-P5 fragment, and the chemoreceptor TM14 (PDB ID 4JPB).[39] We first

generated homology models of *E. coli* CheA-P5 (residues 514-640; using the crystal

structure of *T. maritima* CheA PDB ID 1B3Q[40] as a template) and CheW (residues 18-

154; using the solution structure of *E. coli* CheW PDB ID 2HO9 as a template) with the

Phyre2 web portal.[41] We then superimposed the top models against the corresponding

T. maritima molecules in the crystal structure (PDB ID 4JPB[39]) using PyMOL (v.2.3.3).[42]

3 | RESULTS

3.1 | CheW-like domains are incorporated into dozens of distinct domain *Architectures* in nature

We surveyed the known domain architectures into which CheW-like domains are incorporated, which provided initial sequence groupings for further investigation. We extracted the sequences of all proteins containing a CheW-like domain (PF01584; 8484 sequences containing 9094 unique CheW-like domains) from the Pfam database (v.33)[1] using the Representative Proteome 35 (RP35) sequence set.[17] The use of RP35 provided an efficient and representative data set while substantially reducing the sequence space and complexity required for the analysis. CheW-like domain sequences from rare architectures (<40 member sequences) were excluded because we anticipated that their interpretation would be too speculative. The remaining sample contained 8581 CheW-like domains (~94% of original) in 8061 proteins (~95% of original). Additional curation of sequences described in Methods (e.g., for excessive gap character) resulted in final analysis of 8528 sequences from CheW-like domains.

We employ the capitalized and italicized terms *Architecture*, *Context*, *Cluster*, *Group*, *Type*, and *Class* to refer to the specific hierarchy used to categorize sequences of CheW-like domains in this study. The relationships between the classification terms, as well as the flow of our sorting scheme, are outlined in Figure 1. We first categorized CheW-like domain sequences into three "lineages" based on the architecture and putative role(s) of their parent protein (Figure 2). Our nomenclature for *Architectures* containing CheW-like domains used the accepted lineage of the parent protein (e.g., CheA) as a prefix and a Roman numeral as a suffix. CheA-lineage *Architectures* were

numbered by decreasing membership size, starting with Roman numeral I. Additionally,

Architectures containing two or more CheW-like domains (e.g., CheA.V from Figure 2)

were split into their constituent CheW-like domains and analyzed as separate sequence

Contexts. An additional suffix of an Arabic numeral identified the "position" of the CheW-

like domain relative to the N-terminus and the other CheW-like domains in the parent

protein, e.g., CheA.V.1 (*Contexts* are not labeled in Figure 2). The CheA-lineage was

the most diverse, with a dozen major *Architectures*. In addition to CheW-like domains,

most CheA *Architectures* also contained HATPase_c domains to bind ATP and catalyze

phosphorylation, and histidine phosphotransferase (Hpt) domains to provide a

phosphorylation site.

Although several CheV-lineage architectures (named for the prototypical *Bacillus*

subtilis CheV protein[43]) were detected in the RP35 set, the only organization

incorporating an appreciable number of sequences (>40) was the prototypical form,

comprised of a N-terminal CheW-like domain and a C-terminal receiver domain, which

was termed the CheV.I *Architecture* for consistency. The CheV-lineage was the least

abundant of the three architectural lineages.

CheW-lineage architectural variants constituted the largest collection of

sequences, with over half of the entire data set attributable to prototypical CheW

proteins comprised of a single CheW-like domain, from which the *Architecture* gets its

name (CheW.I). Additional rare (~1 to 3% the number of CheW.I) *Architectures* featured

two or three CheW-like domains fused together (termed CheW.II and CheW.III

respectively) with no other detectable domains present. Individual CheW-like domains

of CheW.II and CheW.III *Architectures* were separated into distinct *Contexts* for further

analysis, doubling or tripling the number of sequences for the corresponding

Architecture (i.e., CheW.II.1, for the first position CheW-like domain, CheW.II.2 for the

second position CheW-like domain, etc.).

Although we focused only on the more abundantly represented architectural

variants, CheW-like domains are ubiquitous and can be found embedded in dozens of

distinct *Architectures* encompassing nearly every component and layer of the microbial

chemotaxis system.[44] We briefly investigated the sequence groups that failed to pass

our chosen thresholds. The majority of these "rare" *Architectures* fell within the CheA-

lineage (containing anywhere from zero to 10 Hpt domains). Atypical, full-length CheA-

lineage proteins were also present, fused to CheB, CheC, CheX, or MCP domains. In

addition to a CheW.IV *Architecture*, CheW-like domains were detected fused to CheC,

CheR, CheX, CheZ, Hpt, or receiver domains, as well as PAS domains that might

provide input information to chemosensory systems. Finally, CheW-like domains were

infrequently found in proteins with a variety of other domains that have no obvious

connection to chemotaxis.

3.2 | CheW-like domains segregated into distinct sequence *Groups* as a function

of parent domain *Architecture*

We expected that CheW-like domains extracted from a common architectural

Context would more closely resemble each other rather than the same domains

obtained from contrasting architectural variants. We based our assumption on existing

knowledge of microbial response regulator receiver domains, which cluster

phylogenetically predicated on the presence and/or type of an attached output

domain.[4,5] However, the full dataset contained >8500 CheW-like domain sequences,

making analysis and interpretation challenging. Ideally, each CheW-like domain *Context*

would behave as a homogeneous sequence set, and a basic stratified sampling

strategy could be used to select a more representative and manageable population for

study. While the assumption of homogeneity seemed reasonable (shared architecture

typically implies common function), we wished to first gain insight into the underlying

structure of the sequence space by performing unsupervised clustering on the full list of

sequences. This allowed us to more accurately assess and incorporate the

heterogeneity of the data into our approach. Essentially, our overall approach

incorporated a simplified multistage sampling scheme, first clustering the entire

population to reveal major strata within each architectural *Context*, then randomly

sampling from within these strata, ensuring that all major sequence groups were

represented. We chose to use a *k*-medoids approach for the initial clustering procedure

to mitigate the difficulty of visualizing >8500 data points. The *k*-medoids algorithm is a

classical clustering technique that partitions the data into *k* number of *Clusters*,

attempting to minimize the distance between data points and the center of their relevant

Cluster. Each *Cluster* is represented by a single central data point called a cluster

medoid, which allowed us to visualize the *Clusters* using just the medoid sequences,

rather than the entire data set.

A representative multi-dimensional scaling (MDS) plot describing just the 21

extracted medoids (based on the dissimilarities calculated using the GONNET model) is

shown in Figure 3. Additional plots generated using dissimilarities calculated with three

other substitution models are included in Figure S1 (all four partitions were highly

similar). Letters naming the *Clusters* for each solution were assigned based on

decreasing number of sequences in the *Cluster*. There were three substantial *Groups* of

Clusters of CheW-like domains (labeled 1, 3, or 4 in Figure 3), and another smaller,

intermediate *Group* separating *Groups* 1 and 3 (labeled 2 in Figure 3, more apparent

when considering only the first dimension).

We next examined the composition of the individual *Clusters* to determine if the

original architectural *Contexts* of CheW-like domains separated into multiple stable

strata. Although each medoid was representative of the dissimilarity data, a *Cluster* may

contain multiple domain *Contexts*. Understanding the composition of each *Cluster* would

allow us to properly organize sequences for analysis on the more complete set, rather

than on just the representative medoids, ensuring that no distinct *Contexts* would be

underrepresented. Table 1 assigns *Clusters* to the four main *Groups* indicated in Figure

3 and contains a detailed breakdown of the architectural *Contexts* of the sequences in

each *Cluster.*

Multiple patterns are apparent in Table 1. First, sequences form *Clusters* and

Clusters form *Groups* primarily based on parent protein lineage. Most or all sequences

within a *Cluster* are from either CheA proteins or CheW/CheV proteins, not both. Most

CheW-like domains from CheA proteins are in *Group* 1 and most CheW-like domains

from CheW and CheV proteins are in *Group* 4. Closer inspection reveals additional

structure in the data:

- 408 The "second" position CheW-like domains from *Architectures* CheA.V and

409   CheA.VI formed their own distinct sequence *Group* 2, made up of *Cluster* P.

- 410 CheW-like domains from *Architectures* CheA.VII and CheA.X comprised the bulk

411   of *Cluster* R, the smallest *Cluster* in *Group* 1.

- About 80% of domains from CheW.I *Architectures* and greater than 90% of domains from CheW.II and CheW.III *Architectures* were in *Group* 4.

- About 20% of domains from CheW.I *Architectures* and 5% of domains from CheW.II and CheW.III *Architectures* were in *Group* 3, possibly distinct from the majority of CheW-lineage domains.

- About 1% of domains from CheW.I *Architectures* and 2% of domains from CheW.II and CheW.III *Architectures* were in *Group* 1, suggesting similarity to the CheA-lineage.

- About 99% of CheW-like domains from CheV.I *Architectures* were in *Group* 4, suggesting that the presence of the attached receiver domain has little effect on the primary sequence of the CheW-like domain in CheV proteins.

- Similar to CheW proteins, about 1% of CheW-like domains in CheV.1 *Architectures* were in *Group* 1, suggesting similarity to the CheA-lineage.

Based on the analysis of Table 1, we assumed for further processing that each architectural *Context* could be reasonably treated as a single homogeneous entity, except for CheW.I. We split CheW.I into three distinct *Types* designated CheW.IA (sequence *Group* 3), CheW.IB (sequence *Group* 4), and CheW.IC (sequence *Group* 1). Remaining outlier sequences in each architectural *Context* (typically a small number of sequences in each *Context* assigned to substantially different *Clusters* than the majority) were excluded from the analysis as noise. We believe that our evaluation produced an accurate summary of the underlying data structure, though we acknowledge the somewhat subjective nature of cluster analysis and that alternative

perspectives may offer additional valid insights. To explore the data at higher resolution

and understand sequence variation within the *Clusters*, we next performed MDS

analysis on a larger, randomly sampled subset of our sequence data, now organized

into 23 *Types* (due to CheW.I sub-designations) instead of 21 *Contexts*.

3.3 | Random stratified subsampling facilitates efficient multi-dimensional scaling

analysis on large protein sequence sets and revealed *Classes* of CheW-like

domains

Our initial cluster analysis (Figure 3 and Table 1) revealed how to organize

sequences of CheW-like domains into 23 *Types* that should ensure all major

subpopulations receive appropriate representation in a random stratified subsampling

scheme described in Methods. Figure 4 shows the top three dimensions of one

representative solution derived from the GONNET substitution matrix. The remaining

nine iterations, as well as solutions calculated using randomly sampled subsets for the

other three substitution matrices described in Methods, are in Figures S2-S13. The

solution shown in Figure 4 revealed similar relationships to those found in Figure

3/Table 1, which we now designate as *Classes* (sub-sampling MDS) rather than *Groups*

(*k*-medoid MDS). On the right area of the plot (Left and Center Panels), most CheW-

lineage and CheV-lineage sequence *Types* (*Class* 1) clustered in a densely packed

region. A large agglomeration of sequences from CheA-lineage *Types* (*Class* 3) was

near the left side of the plot area (Left and Center panels). The CheA.V.2 and

CheA.VI.2 *Types* (*Class* 5) were markedly different from the other CheA-lineage *Types*,

consistent with previous assignment to *Cluster* P as the sole member of *Group* 2 in

Figure 3/Table 1. Additional differentiation within the CheA-derived *Types* could be

observed more effectively in the Center and Right Panels (incorporating Dimension 3),

with the CheA.VII and CheA.X *Types* (*Class* 4) separating from the main body. This

supports the clustering results presented in Table 1 and Figure 3, with the majority of

CheA.VII and CheA.X sequence *Types* found in the somewhat independent *Cluster* R.

Sequences belonging to the rare CheW.IC *Type* (*Class* 2) were spread across Figure 4,

alongside *Class* 1 CheW/CheV sequences, *Class* 3 CheA sequences, and in between

(Left and Center Panels). In Figure 3/Table 1, a large subset of CheW-lineage

sequences (*Group* 3) separated from most CheW- and CheV-lineage sequences

(*Group* 4). The *Group* 3 sequences are termed *Type* CheW.IA (*Class* 6) in the analysis

of Figure 4 and surprisingly did not separate from the main CheW/CheV *Types* (*Class*

1).

We also tried non-metric MDS as described in Methods and shown in Figures

S14-S25. We concluded that the data structures were likely consistent irrespective of

the MDS algorithm. Given the clear sequence heterogeneity within the CheW-like

domains, we next sought to determine factors within the domain itself that distinguished

the various *Types*.

3.4 | SimpLogo: A simplified method for visualizing large numbers of distinct

protein sequences aligned over many positions

Arguably the most popular method for comparing sequence groups and their

alignments is the sequence logo. In this context, a logo is a graphical representation

depicting sequence conservation by stacking relevant characters, such as single-letter

codes for amino acids or nucleotides, at each position in an alignment.[45] The relative

heights of the characters are proportional to their frequency at the given position.

Additionally, the overall height of a stack is often scaled based on the information

content stored at that position within the alignment. Typical applications for amino acid

sequence logos include visualizing functional motifs or binding sites within a protein

family/group. However, the volume of information and the graphical complexity of a logo

often makes it challenging to interpret when examining regions longer than two dozen

positions. Sequence logos are less practical when comparing a large group of separate

alignments (>12) with many positions (>50). A more ideal and complementary

visualization technique would allow us to first evaluate the macroscopic characteristics

of each sequence group. Subtle differences could then be examined in greater detail for

specific regions using a more traditional approach like a true sequence logo, or even the

original sequence alignment. Here, we describe a visualization format (colloquially

referred to as SimpLogo) designed to compare the CheW-like domain sequences from

the numerous *Types* described above. We sought to reduce the 23 individual sequence

alignments (*Types*) into simplified, pseudo-one-dimensional representations. We

accomplished this by weighting and blending qualitative colors assigned to specific

amino acid residue types (grouped by physicochemical properties into eight classes)

based on their frequency at a given position within the alignments (Figure 5). We also

modified the overall height based on its proportion of gap-character (details in Methods).

This produced a single row of colored "boxes" for each sequence *Type*, reducing a

complex problem to basic pattern recognition. To differentiate between secondary

colors and a near-equal mixing of primary colors, colored strips were included on the

top and bottom of each box, indicating first and second most frequent residue types.

The simplified scheme allowed for the simultaneous comparison of dozens of distinct

sequence families with lengths over many positions (>100) and encoding substantially

more information about intermediate frequencies than simple consensus sequences.

Additionally, we created traditional sequence logos (using the Logolas package in R) for

the same alignments using similar image dimensions for comparison (see Figure S26).

Highly conserved/informative positions are identifiable in the traditional logo, but the

overall lack of space and reduced character sizes often make interpretation of specific

positions challenging, especially for the lower frequency/less conserved positions with

smaller stack heights. The patterns of gap characters (indicated by dashes) are also

more challenging to interpret when compared to the SimpLogo representation in Figure

5. One could imagine how difficult interpretation would become with double the number

of sequence groups or sequence lengths using traditional methods. Therefore, we

envision the new SimpLogo visualization technique being used not as a replacement for

traditional sequence logos, but as a supplementary technique for large scale exploratory

analyses.

3.5 | *Classes* of CheW-like domains display characteristic gap patterns and

residue identities

The SimpLogo representation in Figure 5 revealed multiple trends. Overall, the

relationships displayed in Figure 5 mimicked the trends seen in Figure 4, with most of

the CheA-lineage *Types* clustered tightly (*Class* 3), the exceptions being the

CheA.VII/CheA.X (*Class* 4) and CheA.V.2/CheA.VI.2 (*Class* 5) *Types*. Most CheW-

lineage *Types* (*Classes* 1 and 6) and the CheV-lineage *Type* (*Class* 1) appeared

similar, except for the CheW.IC *Type* (*Class* 2), which was intermediate in character

between CheA and CheW *Types*.

Figure 5 is information dense, so interpretation is facilitated by provision of

appropriate context. *E.* coli CheW and CheA proteins are experimentally well-studied,

but crystal structures of the corresponding CheW-like domains are not available. To

provide a structural frame of reference for our analysis, we constructed simple

homology models of *E. coli* CheW and the CheW-like domain (termed P5) of CheA

based on experimentally determined crystal structures of the corresponding *T. maritima*

domains as described in Methods. For convenient cross-referencing, Table S1 contains

a detailed list of putative functions for relevant residues of CheW-like domains based on

various literature sources and our homology models. We assumed, for the sake of

speculation, that the analogous interaction surfaces and behaviors are largely

maintained throughout the various CheW-like domain topologies found in nature and

represented in Figure 5. A glance reveals that the gap characters (box heights) of the

CheA-, CheV-, and CheW-lineages were different, which likely played an important role

in differentiating the sequence *Types*, particularly at positions 45-56 and 120-131.

Similarly, there were clear differences in residue identities, especially involving *Class* 5

domains. In Text S1, we describe observations concerning gap patterns and residue

identities at length, and also suggest possible functional implications. These

descriptions illustrate the detailed information and hypotheses that can be derived from

analysis of SimpLogos in conjunction with existing experimental data. However, the

primary purpose of this report is to describe our generally applicable strategy to analyze

protein domains in different architectural contexts and, as summarized in the

Discussion, the higher order conclusions that can be drawn from sorting a single Pfam

domain into different *Classes*.

4 | DISCUSSION

4.1 | Implications of six CheW-like domain *Classes*

The existence of *Class* 1, which encompasses CheV and most CheW proteins, has multiple implications. First, evidence exists that CheV proteins can function as scaffold proteins similarly to CheW proteins[12,46-48]; however, more detailed information about the functions and mechanisms of CheV is sparse.[13,15] It is tempting to extrapolate what is known about response regulator proteins (composed of N-terminal receiver and C-terminal output domains) to formulate hypotheses about CheV (composed of C-terminal receiver and N-terminal CheW-like domains). However, our observation that CheW-like domains in CheV proteins were similar to those in CheW proteins constrains such hypotheses by suggesting that any interactions between the receiver and CheW-like domains in CheV proteins are not sufficiently specific to be subject to selective pressure for evolutionary optimization. There are at least three categories of hypotheses about CheV inspired by response regulators:

- The receiver domain might alter the conformation of or occlude access to the CheW-like domain, depending on the phosphorylation state of the receiver. Alexander *et al.* suggested that such phosphorylation-mediated changes could explain the experimentally established role of CheV phosphorylation[49,50] in adaptation by changing the conformation of the CheW-like domain to alter the degree of coupling between receptor and kinase function.[15] Our findings make this otherwise reasonable mechanism now seem unlikely. In *Bacillus subtilis* CheV, the CheW-like domain inhibits phosphotransfer from CheA-P to CheV and stabilizes CheV-P from autodephosphorylation.[49] Our findings suggest that these

effects are more likely the result of steric effects than conformational coupling

between the two domains of CheV. The determination of CheV structures in the

presence and absence of the stable phosphoryl group analog BeF_3^- would be

highly informative.

• Alexander *et al.*[15] and Ortega & Zhulin[46] proposed that the phosphorylation site in

the receiver domain of CheV might act as a phosphate sink, while the CheW-like

domain localizes CheV appropriately. Our results are consistent with such a

hypothesis because no interaction between the two domains of CheV is required

that might lead to divergence of CheW-like domains in CheW and CheV proteins.

In a phosphate sink mechanism, when environmental circumstances change and

phosphoryl groups stop flowing into the system from ATP, phosphoryl groups

flow backwards from the CheY response regulator to CheA to the sink and are

lost to hydrolysis. Localizing the sink near CheA would enhance efficiency. We

note that another possible role of CheV is as a reservoir that retains phosphoryl

groups in a buffering capacity.

• In a more dynamic role, phosphorylation-mediated dimerization of the receiver

domain of CheV might control the ability of the CheW-like domain to bind with

receptor and/or kinase partners. To the best of our knowledge, the multimeric

state of CheV and any effect of phosphorylation on multimeric state have not

been investigated.

Second, five (CheA.III, CheA.IV, CheA.V, CheA.VII, CheA.X) of the 12 CheA

Architectures in Figure 2 contained a receiver domain immediately C-terminal to a

CheW-like domain, which is reminiscent of CheV proteins. However, the fact that

CheW-like domains in CheV and CheA proteins were in different *Classes* (1 vs. 3, 4, or

5) strongly suggests that adjacent CheW-like and receiver domains in CheA proteins do

not act as CheV analogs. Consistent with this prediction, the adjacent CheW-like

domain did not affect autodephosphorylation of the receiver domain from *Azorhizobium*

caulinodans CheA,[51] in contrast to *B. subtilis* CheV.[49]

Third, as far as we know, obvious questions about CheW-lineage proteins with

multiple domains have not been explored experimentally. It is not known if all CheW-like

domains of multi-domain CheW proteins are required for function, or if there are

differences in functional roles between the domains. The fine structure of the data in

Table 1 (specifically the number of sequences in a *Cluster* varied substantially as a

function of *Context* within CheW.II and CheW.III *Architectures*) strongly suggests that

the domains in a single multidomain CheW protein often belong to different sequence

Clusters or even *Groups*, which could suggest domain specialization. On the other

hand, because experimentally characterized single domain CheW proteins belonged to

Class 1, which also includes ~93% of domains in multi-domain CheW proteins, our

results suggest that there is likely little to be learned about general CheW function by

exploring multi-domain CheW proteins. Presumably, multi-domain CheW proteins

support formation of complexes with specific components. This conjecture is supported

by the observation that CheW.III proteins form cytoplasmic double layered arrays with

CheA and a special type of double domain chemoreceptor in *Vibrio cholerae*.[52]

Approximately 1% of CheW-like domains in CheW proteins were CheA-like

(CheW.IC *Type*) and formed a *Class* of their own. Some members of *Class* 2 appeared

to span the gap between CheA proteins and CheW/CheV proteins (Figures 4 and S2-

S25). The CheW-like domain is the most recently evolved fold in proteins that support

bacterial chemotaxis.[44] The birth of CheW-like domains allowed the separation of

receptor and kinase functions into different proteins, in turn facilitating the integration of

multiple stimuli sensed by arrays of different receptors into a single kinase. We are

unaware of any evidence establishing whether CheW-like domains evolved first in

CheW, CheA, or CheV proteins. Because CheV proteins are less common and less

widely distributed than CheA or CheW proteins,[44] it seems reasonable that CheV

proteins came after CheA and CheW. Because members of *Class* 2 segregated with

Class 1, *Class* 3 and in between in our analyses (Figure 4), *Class* 2 may represent a

"missing link" in the evolution of CheW-like domains from CheW to CheA or from CheA

to CheW. Closer examination of sequences in *Class* 2 may provide new insights into the

evolution of chemotaxis systems.

CheW-like domains in most CheA proteins with one CheW-like domain were

similar (*Class* 3), even though there were 10 different major *Architectures* for such

proteins, including one (CheA.IX) that lacked a phosphorylation site (Hpt domain) and

two (CheA.XI, CheA.XII) that lacked ATP binding sites (HATPase_c domain). Such

"split" CheA *Architectures* are not degenerate, but can work together to reconstitute a

functional unit.[53] N-terminal CheW-like domains in CheA *Architectures* with two CheW-

like domains (CheA.V.1, CheA.VI.1) were similar to CheW-like domains in CheA

Architectures with a single CheW-like domain. The existence of *Class* 3 suggests that

what is known about CheW-like domains in experimentally well-characterized CheA

proteins is likely to be applicable to the majority of CheW-like domains in CheA proteins.

The exceptions would be *Class* 4 and 5 CheW-like domains.

Class 4 CheW-like domains were found in CheA.VII and CheA.X *Architectures*,

which have three Hpt domains and a C-terminal receiver domain. There were many

minor CheA-lineage *Architectures* that contained up to 10 N-terminal Hpt domains and a

C-terminal receiver domain. To the best of our knowledge, the only such protein to have

been experimentally characterized is *Pseudomonas aeruginosa* ChpA, which controls

Type IV pili rather than flagella. ChpA has six Hpt domains and supports an

intramolecular phosphorelay.[54] Phosphotransfer within ChpA proceeds from ATP to the

Hpt domains close to the HATPase_c domain to the receiver domain to Hpt domains

further from the HATPase_c domain. Thus, a plausible hypothesis is that a distinct

Class of CheW-like domains is required to support intramolecular phosphorelays in

CheA proteins with multiple Hpt domains and a receiver domain. Perhaps the receiver

domain (immediately adjacent to the CheW-like domain) must be held in a particular

orientation to be accessible to multiple Hpt domains.

Class 5 featured the C-terminal CheW-like domain in CheA proteins with dual

CheW-like domains. *A. caulinodans* CheA,[51] *Azospirillum brasilense* CheA1,[55] and

Rhodospirillum centenum CheA1[56] have been explored experimentally and all belong to

the CheA.V *Architecture* with two CheW-like domains. In each case, the C-terminal

receiver domain is functionally important. However, the only experimental investigation

of a *Class* 5 CheW-like domain of which we are aware was in *A. caulinodans* CheA,

where removal of the C-terminal CheW-like domain diminishes chemotaxis but does not

reveal the functional role of the *Class* 5 domain.[51] *Class* 5 CheW-like domains were

substantially different from experimentally characterized CheW-like domains. Therefore,

because investigation of *Class* 5 is especially likely to reveal new information, *Class* 5

arguably represents the highest priority for future research on CheW-like domains.

Class 6 was a *Type* (CheW.IA) encompassing ~20% of CheW proteins that

appeared different from most CheW proteins (*Class* 1) in some analyses and similar in

others. *Class* 6 CheW-like domains appeared to be different from the rest by *k*-medoid

analysis (see Figure 3/Table 1). *Class* 6 appeared indistinguishable from *Class* 1 in the

metric MDS analyses of random stratified subsampling in Figure 4 and Figures S2-S13,

and the differences were subtle in the SimpLogo of Figure 5. *Class* 6 showed some

separation from *Class* 1 in the non-metric MDS analyses of random stratified

subsampling in Figures S14-S25. Method-dependent classification suggests that *Class*

6 is unlikely to represent a high priority for investigation. However, a companion

manuscript reports additional analyses that may shed light on *Class* 2 and *Class* 6

CheW-like domains.[57]

4.2 | Generally applicable methods to analyze protein domains as a function of

architectural context

We have described a workflow (Figure 1) to analyze variation between

sequences of a single protein domain as a function of architectural context within

proteins. To ensure a robust workflow, we tested various approaches to optimize our

method, including using multiple different amino acid substitution models and

multidimensional scaling techniques to estimate and visualize relevant domain

sequence subpopulations. Furthermore, the similarity of our partition results using these

different models was compared using ARI and NMI analyses. We first applied a basic *k*-

medoid clustering step to quantify the relative homogeneity of a given domain sequence

set. This allowed us to identify constituent strata for representation in a more complete

sequence analysis utilizing a random stratified sub-sampling scheme. We demonstrated

our approach using CheW-like domains as a general case study. The results were

largely independent of methods chosen, suggesting robust conclusions. Grouping

sequence *Types* of CheW-like domains into six major *Classes* led to a variety of useful

observations, insights, hypotheses, and suggested areas of future experimental

investigation.

It might appear challenging to apply our approach to domains found in many

distinct domain architectures. However, domain architecture distributions tend to have a

long tail, with only a few organizations representing the majority of cases[58]. Here, 16 out

of 111 architectures in Pfam that contained CheW-like domains incorporated ~94% of

CheW-like domains. Of course, individual "rare" architectures of interest could always

be manually included with minimal difficulty. Another concern could involve the number

of sequences to analyze. Our case study involved analysis of ~8500 sequences.

Significantly more or less sequences may be available for other domains. However, our

approach should still be practical. The number of starting sequences to be analyzed can

be managed by choice of an appropriate input data set. We chose (and recommend)

the use of Representative Proteome sets as a starting point, where the sequence space

size and granularity can be modified by changing the co-membership threshold of the

relevant RP set cutoff (we used 35% as our threshold, but RP sets generated using

15%, 55%, and 75% are also available).[17] The subsampling used in our analysis (see

Figure 4) is a standard strategy to deal with impractically large data sets.

Finally, we developed and have made available (see Methods) SimpLogo, a new

means of visualizing amino acid sequence conservation patterns across many subtypes

of the same domain and for sequences of arbitrary length. Comparison of the

SimpLogos in Figure 5 and traditional sequence logos in Figure S26 quickly reveal the

utility of SimpLogos. Differences in gap patterns are immediately obvious. Identifying

differences in amino acid composition at individual positions is more time consuming,

but feasible. Traditional sequence logos are useful to communicate information about

reasonably conserved positions but of little utility for most other positions, whereas

SimpLogo provides useable information about all aligned positions within a given

sequence.

ACKNOWLEDGEMENTS

We thank Emily N. Kennedy and Sarah A. Barr for their insightful input and helpful discussions, including Sarah's idea that receiver domains of CheV proteins might serve as reservoirs rather than sinks for phosphoryl groups.

This work was funded by National Institutes of Health grant GM050860 to Robert B. Bourret. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of General Medical Sciences or the National Institutes of Health.

REFERENCES

1. Mistry J, Chuguransky S, Williams L, et al. Pfam: The protein families database in 2021. *Nucleic Acids Res.* 2020;49(D1):D412-D419.

2. Martin J, Anamika K, Srinivasan N. Classification of protein kinases on the basis of both kinase and non-kinase regions. *PLoS One.* 2010;5(9):e12460.

3. Syamaladevi DP, Joshi A, Sowdhamini R. An alignment-free domain architecture similarity search (ADASS) algorithm for inferring homology between multi-domain proteins. *Bioinformation.* 2013;9(10):491-499.

4. Koretke KK, Lupas AN, Warren PV, Rosenberg M, Brown JR. Evolution of two-component signal transduction. *Mol Biol Evol.* 2000;17(12):1956-1970.

5. Pao GM, Saier MH, Jr. Response regulators of bacterial signal transduction systems: selective domain shuffling during evolution. *J Mol Evol.* 1995;40(2):136-154.

6. Gao R, Bouillet S, Stock AM. Structural basis of response regulator function. *Annu Rev Microbiol.* 2019;73:175-197.

7. Immormino RM, Silversmith RE, Bourret RB. A variable active site residue Iinfluences the kinetics of response regulator phosphorylation and dephosphorylation. *Biochemistry.* 2016;55(39):5595-5609.

8. Page SC, Immormino RM, Miller TH, Bourret RB. Experimental analysis of functional bariation within protein families: Receiver domain autodephosphorylation kinetics. *J Bacteriol.* 2016;198(18):2483-2493.

9. Straughn PB, Vass LR, Yuan C, Kennedy EN, Foster CA, Bourret RB. Modulation of

response regulator CheY reaction kinetics by two variable residues that affect

conformation. *J Bacteriol.* 2020;202(15).

10. Thomas SA, Immormino RM, Bourret RB, Silversmith RE. Nonconserved active site

residues modulate CheY autophosphorylation kinetics and phosphodonor

preference. *Biochemistry.* 2013;52(13):2262-2273.

11. Bi S, Sourjik V. Stimulus sensing and signal processing in bacterial chemotaxis. *Curr*

Opin Microbiol. 2018;45:22-29.

12. Abedrabbo S, Castellon J, Collins KD, Johnson KS, Ottemann KM. Cooperation of

two distinct coupling proteins creates chemosensory network connections. *Proc Natl*

Acad Sci U S A. 2017;114(11):2970-2975.

13. Huang Z, Pan X, Xu N, Guo M. Bacterial chemotaxis coupling protein: Structure,

function and diversity. *Microbiol Res.* 2019;219:40-48.

14. Alexandre G, Zhulin IB. Different evolutionary constraints on chemotaxis proteins

CheW and CheY revealed by heterologous expression studies and protein sequence

analysis. *J Bacteriol.* 2003;185(2):544-552.

15. Alexander RP, Lowenthal AC, Harshey RM, Ottemann KM. CheV: CheW-like

coupling proteins at the core of the chemotaxis signaling network. *Trends Microbiol.*

2010;18(11):494-503.

16. Gumerov VM, Andrianova EP, Zhulin IB. Diversity of bacterial chemosensory

systems. *Curr Opin Microbiol.* 2021;61:42-50.

17. Chen C, Natale DA, Finn RD, et al. Representative proteomes: A stable, scalable

and unbiased proteome set for sequence analysis and functional annotation. *PLoS*

One. 2011;6(4):e18910-e18910.

18. Consortium TU. UniProt: The universal protein knowledgebase in 2021. *Nucleic*

Acids Res. 2020;49(D1):D480-D489.

19. Eddy SR. Accelerated Profile HMM Searches. *PLOS Comp Biol.*

2011;7(10):e1002195.

20. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics.*

2009;25(15):1972-1973.

21. Dayhoff M, Schwartz R, Orcutt B. A model of evolutionary change in proteins. *Atlas*

of Protein Sequence and Structure. 1978;5:345-352.

22. Gonnet GH, Cohen MA, Benner SA. Exhaustive matching of the entire protein

sequence database. *Science.* 1992;256(5062):1443-1445.

23. Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. *Proc*

Natl Acad Sci U S A. 1992;89(22):10915-10919.

24. Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data matrices

from protein sequences. *Comput Appl Biosci.* 1992;8(3):275-282.

25. Pelé J, Bécu J-M, Abdi H, Chabbert M. Bios2mds: An R package for comparing

orthologous protein families by metric multidimensional scaling. *BMC Bioinform.*

2012;13(1):133.

26. Rousseeuw PJ. Silhouettes: A graphical aid to the interpretation and validation of

cluster analysis. *J Comput Appl Math.* 1987;20:53-65.

27. *fpc: Flexible procedures for clustering* [computer program]. Version 2.2-92015.

28. *cluster: Cluster analysis basics and extensions* [computer program]. Version

2.1.22012.

29. Hubert L, Arabie P. Comparing partitions. *J Classif.* 1985;2(1):193-218.

30. Steinley D. Properties of the Hubert-Arable Adjusted Rand Index. *Psychol Methods.*

2004;9(3):386-396.

31. Cover TM, Thomas JA. *Elements of information theory.* Wiley-Interscience; 2006.

32. Strehl A, Ghosh J. Cluster ensembles - A knowledge reuse framework for combining

multiple partitions. *J Mach Learn Res.* 2003;3(null):583–617.

33. *vegan: Community Ecology Package. R package version 2.5-6. 2019* [computer

program]. Version 2.5-72020.

34. Wickham H. *ggplot2: Elegant graphics for data analysis.* 3rd ed: Springer-Verlag

New York; 2016.

35. Tzeng J, Lu HH-S, Li W-H. Multidimensional scaling for large genomic data sets.

BMC Bioinform. 2008;9(1):179.

36. *Biostrings: Efficient manipulation of biological strings* [computer program]. Version

2.60.12015.

37. Ou J, Wolfe SA, Brodsky MH, Zhu LJ. motifStack for the analysis of transcription

factor binding site evolution. *Nat Methods.* 2018;15(1):8-9.

38. Dey KK, Xie D, Stephens M. A new sequence logo plot to highlight enrichment and

depletion. *BMC Bioinform.* 2018;19(1):473.

39. Li X, Fleetwood AD, Bayas C, et al. The 3.2 Å resolution structure of a

receptor:CheA:CheW signaling complex defines overlapping binding sites and key

residue interactions within bacterial chemosensory arrays. *Biochemistry.*

2013;52(22):3852-3865.

40. Bilwes AM, Alex LA, Crane BR, Simon MI. Structure of CheA, a signal-transducing

histidine kinase. *Cell.* 1999;96(1):131-141.

41. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. The Phyre2 web portal

for protein modeling, prediction and analysis. *Nat Protoc.* 2015;10(6):845-858.

42. *PyMOL* [computer program]. Version 2.3.3. http://www.pymol.org/pymol2020.

43. Fredrick KL, Helmann JD. Dual chemotaxis signaling pathways in *Bacillus subtilis*: a

sigma D-dependent gene encodes a novel protein with both CheW and CheY

homologous domains. *J Bacteriol.* 1994;176(9):2727-2735.

44. Wuichet K, Zhulin IB. Origins and diversification of a complex signal transduction

system in prokaryotes. *Sci Signal.* 2010;3(128):ra50-ra50.

45. Schneider TD, Stephens RM. Sequence logos: A new way to display consensus

sequences. *Nucleic Acids Res.* 1990;18(20):6097-6100.

46. Ortega DR, Zhulin IB. Evolutionary genomics suggests that CheV Is an additional

adaptor for accommodating specific chemoreceptors within the chemotaxis signaling

complex. *PLOS Comp Biol.* 2016;12(2):e1004723.

47. Walukiewicz HE, Tohidifar P, Ordal GW, Rao CV. Interactions among the three

adaptation systems of *Bacillus subtilis* chemotaxis as revealed by an *in vitro*

receptor-kinase assay. *Mol Microbiol.* 2014;93(6):1104-1118.

48. Yang W, Alvarado A, Glatter T, Ringgaard S, Briegel A. Baseplate variability of

Vibrio cholerae chemoreceptor arrays. *Proc Natl Acad Sci U S A.*

2018;115(52):13365-13370.

49. Karatan E, Saulmon MM, Bunn MW, Ordal GW. Phosphorylation of the response

regulator CheV is required for adaptation to attractants during *Bacillus subtilis*

chemotaxis. *J Biol Chem.* 2001;276(47):43618-43626.

50. Rosario MM, Fredrick KL, Ordal GW, Helmann JD. Chemotaxis in *Bacillus subtilis*

requires either of two functionally redundant CheW homologs. *J Bacteriol.*

1994;176(9):2736-2739.

51. Kennedy EN, Barr SA, Liu X, et al. *Azorhizobium caulinodans* chemotaxis is

controlled by an unusual phosphorelay network. *J Bacteriol.* 2021:JB0052721.

52. Briegel A, Ortega DR, Mann P, Kjaer A, Ringgaard S, Jensen GJ. Chemotaxis

cluster 1 proteins form cytoplasmic arrays in *Vibrio cholerae* and are stabilized by a

double signaling domain receptor DosM. *Proc Natl Acad Sci U S A.*

2016;113(37):10412-10417.

53. Porter SL, Wadhams GH, Armitage JP. *Rhodobacter sphaeroides*: complexity in

chemotactic signalling. *Trends Microbiol.* 2008;16(6):251-260.

54. Silversmith RE, Wang B, Fulcher NB, Wolfgang MC, Bourret RB. Phosphoryl group

flow within the *Pseudomonas aeruginosa* Pil-Chp chemosensory system: Differential

function of the eight phosphotransferase and three receiver domains. *J Biol Chem.*

2016;291(34):17677-17691.

55. Bible AN, Stephens BB, Ortega DR, Xie Z, Alexandre G. Function of a chemotaxis-

like signal transduction pathway in modulating motility, cell clumping, and cell length

in the alphaproteobacterium *Azospirillum brasilense*. *J Bacteriol.* 2008;190(19):6365-

6375.

56. Jiang ZY, Gest H, Bauer CE. Chemosensory and photosensory perception in purple

photosynthetic bacteria utilize common signal transduction components. *J Bacteriol.*

1997;179(18):5720-5727.

57. Vass LR, Bourret RB, Foster CA. Analysis of CheW-like domains provides insights

into organization of prokaryotic chemotaxis systems. *Proteins.* 2022;Submitted.

58. Naveenkumar N, Kumar G, Sowdhamini R, Srinivasan N, Vishwanath S. Fold

combinations in multi-domain proteins. *Bioinformation.* 2019;15(5):342-350.

Table 1. Distribution of CheW-like domain *Context*s across *k*-medoid *Clusters*[a]

Architectural Context		Group 1							Group 2	Group 3						Group 4							# Seqs
		C	E	H	I	L	M	R	P	N	O	Q	S	T	U	A	B	D	F	G	J	K	
CheA-lineage	CheA.I	444	127	64	257	170																	1062
	CheA.II	140	312	3	2	8	167																632
	CheA.III	75	104	18	18	26	29	8								1							279
	CheA.IV	74	79	48	6	22		4															233
	CheA.V.1		4	88																			92
	CheA.V.2	2	3						89														94
	CheA.VI.1	10	3	73	2																		88
	CheA.VI.2	2	3		1				78								1	2	1				88
	CheA.VII	3	8		1			68															80
	CheA.VIII	65																					65
	CheA.IX	14	8	26	5	3		1															57
	CheA.X	5			1			50															56
	CheA.XI	29	7	2	10	3	1																52
	CheA.XII	23	22	1	1			1															48
CheV-lineage	CheV.I	4			1			1								50	11	1	4	455	290		817
CheW-lineage	CheW.I	20	4	5	11	3		2		189	157	135	117	115	92	1093	979	605	485	5	7	234	4258
	CheW.II.1	3		2	1						3	1				8	11	20	5				54
	CheW.II.2	2					1			2	1				1	12		32	2			1	54
	CheW.III.1	1								2	3				1	62	21	35	13			2	140
	CheW.III.2	1				1					3			1	1	107	4	13	8				139
	CheW.III.3	1									6					91	17	12	13				140
# Seqs		918	684	330	318	235	197	135	168	193	173	136	117	116	95	1425	1045	719	530	460	297	237	8528

[a]*Clusters* and *Groups* taken from analysis shown in Figure 3. Table entries show the number of sequences of each architectural *Context* found within each *Cluster*. *Clusters* are named in order of decreasing size.

FIGURE LEGENDS

FIGURE 1. Hierarchy of classification terms and sorting workflow. The initial steps were derived purely from domain *Architectures*, which may contain CheW-like domains in more than one *Context*. The number of *Contexts* defined (dashed arrow) the number of *Clusters* into which sequences of CheW-like domains were initially sorted. However, sequences in a given *Context* sorted into different *Clusters* and vice versa. The *Clusters* sorted into larger *Groups* of sequences. Comparison of architectural *Contexts* with sequence *Groups* revealed that all but one *Context* included *Clusters* within a single *Group*. One *Context* contained *Clusters* that spanned three *Groups* and so was split into three *Types*, whereas all other *Contexts* became a single *Type*. Minor *Clusters* of sequences associated with each *Context* were discarded from further analysis during the transition to *Types*. The *Types* sorted into larger *Classes* of sequences associated with one or more architectural *Contexts*. Figure 5 illustrates the final *Types* and *Classes* but was not part of the classification process. See text and indicated Figures or Table for details.

FIGURE 2. Schematic of the most abundant CheW-containing domain *Architectures* seen in nature. Sixteen distinct architectural variants containing the CheW-like domain were identified within the Pfam database (≥40 sequences each, ordered by decreasing abundance within each shaded group). Sequences for their corresponding CheW-like domains were extracted and initially classified according to the schematic. For parent proteins containing multiple CheW-like domains, CheW-like domains were initially classified according to the position (*Context*) of the domain, resulting in one sequence entry per CheW-like domain. Representative diagrams are

shown for each *Architecture* (from left to right, N- to C-terminal). Pfam domain IDs

included are as follows: CheW-like (PF01584); H-kinase dimerization (PF02895); Hpt

(PF01627); Response regulatory (PF00072); CheY-binding (PF09078); HATPase_c

(PF02518); P2 (PF07194). Individual schematics are not necessarily drawn to scale.

FIGURE 3. **Cluster analysis and multidimensional scaling of representative**

sequences of CheW-like domains reveals *Groups*. Multidimensional scaling plots

showing representative medoids from each of the 21 *Clusters*, generated using

dissimilarity matrix calculated with the GONNET substitution model. Each clustering

attempt reconstituted roughly the same data structure, consisting of three main *Groups*

(numbered 1, 3 and 4) of *Clusters* along with a weaker, more intermediate *Group*

(numbered 2) intersecting *Groups* 1 and 3. While the colors and specific *Cluster*

designations may shift between analyses (overall *Cluster* sizes were slightly altered

depending on the substitution matrix used, affecting *Cluster* names), the actual

partitioning of CheW sequences was highly consistent, regardless of the substitution

model used (see Figure S1).

FIGURE 4. Metric multidimensional scaling plots showing 25 randomly

subsampled sequences from each of the 23 CheW-like domain *Types*. The top

three dimensions (Dim1, Dim2, Dim3) were included for visualization purposes. Analysis

was performed on dissimilarities calculated using the GONNET substitution model. The

location of *Types* belonging to each of six *Classes* are indicated. Additional sampling

iterations are shown in Figures S2-S4. Sampling iterations with other substitution

models are shown in Figures S5-S13.

FIGURE 5. **SimpLogo representation of major CheW-like domain *Types*.** Rows

correspond to domain *Types*. Sequences were aligned to the CheW-like domain HMM

(Pfam ID PF01584) and separated by *Type*. Residues with physicochemical similarities

were merged and assigned colors. Idealized colors (legend) were weighted by residue

frequency and mixed to produce a colored box at each position. Heights were scaled by

gap percentage. To differentiate between secondary colors and a near-equal mixing of

primary colors, colored strips were included on the top and bottom of each box,

indicating first and second most frequent residue types respectively. *Types* were

organized by *Class* in the figure. Position-wise information content (in bits) was

calculated for each alignment (top). Numbering is for *E. coli* CheW, framing our

observations with reference to the well-studied prototypical protein.

FIGURE 1. Hierarchy of classification terms and sorting workflow. The initial steps were derived purely from domain *Architectures*, which may contain CheW-like domains in more than one *Context*. The number of *Contexts* defined (dashed arrow) the number of *Clusters* into which sequences of CheW-like domains were initially sorted. However, sequences in a given *Context* sorted into different *Clusters* and vice versa. The *Clusters* sorted into larger *Groups* of sequences. Comparison of architectural *Contexts* with sequence *Groups* revealed that all but one *Context* included *Clusters* within a single *Group*. One *Context* contained *Clusters* that spanned three *Groups* and so was split into three *Types*, whereas all other *Contexts* became a single *Type*. Minor *Clusters* of sequences associated with each *Context* were discarded from further analysis during the transition to *Types*. The *Types* sorted into larger *Classes* of sequences associated with one or more architectural *Contexts*. Figure 5 illustrates the final *Types* and *Classes* but was not part of the classification process. See text and indicated Figures or Table for details.

**FIGURE 2. Schematic of the most abundant CheW-containing domain
Architectures seen in nature.** Sixteen distinct architectural variants containing the
CheW-like domain were identified within the Pfam database (≥40 sequences each,
ordered by decreasing abundance within each shaded group). Sequences for their
corresponding CheW-like domains were extracted and initially classified according to
the schematic. For parent proteins containing multiple CheW-like domains, CheW-like
domains were initially classified according to the position (*Context*) of the domain,
resulting in one sequence entry per CheW-like domain. Representative diagrams are
shown for each *Architecture* (from left to right, N- to C-terminal). Pfam domain IDs
included are as follows: CheW-like (PF01584); H-kinase dimerization (PF02895); Hpt
(PF01627); Response regulatory (PF00072); CheY-binding (PF09078); HATPase_c
(PF02518); P2 (PF07194). Individual schematics are not necessarily drawn to scale.

Cluster Medoids - GONNET (mMDS)

Cluster Assignment

Cluster A ●   Cluster E ○   Cluster I ▲   Cluster M ▲   Cluster Q ■   Cluster U ■
Cluster B ●   Cluster F ○   Cluster J ▲   Cluster N ▲   Cluster R ■
Cluster C ●   Cluster G ●   Cluster K ▲   Cluster O ■   Cluster S ■
Cluster D ●   Cluster H ▲   Cluster L ▲   Cluster P ■   Cluster T ■

FIGURE 3. **Cluster analysis and multidimensional scaling of representative sequences of CheW-like domains reveals *Groups*.** Multidimensional scaling plots showing representative medoids from each of the 21 *Clusters*, generated using dissimilarity matrix calculated with the GONNET substitution model. Each clustering attempt reconstituted roughly the same data structure, consisting of three main *Groups* (numbered 1, 3 and 4) of *Clusters* along with a weaker, more intermediate *Group* (numbered 2) intersecting *Groups* 1 and 3. While the colors and specific *Cluster* designations may shift between analyses (overall *Cluster* sizes were slightly altered depending on the substitution matrix used, affecting *Cluster* names), the actual partitioning of CheW sequences was highly consistent, regardless of the substitution model used (see Figure S1).

FIGURE 4. Metric multidimensional scaling plots showing 25 randomly
subsampled sequences from each of the 23 CheW-like domain *Types*. The top
three dimensions (Dim1, Dim2, Dim3) were included for visualization purposes. Analysis
was performed on dissimilarities calculated using the GONNET substitution model. The
location of *Types* belonging to each of six *Classes* are indicated. Additional sampling
iterations are shown in Figures S2-S4. Sampling iterations with other substitution
models are shown in Figures S5-S13.

FIGURE 5. **SimpLogo representation of major CheW-like domain *Types*.** Rows correspond to domain *Types*.
Sequences were aligned to the CheW-like domain HMM (Pfam ID PF01584) and separated by *Type*. Residues with
physicochemical similarities were merged and assigned colors. Idealized colors (legend) were weighted by residue
frequency and mixed to produce a colored box at each position. Heights were scaled by gap percentage. To differentiate
between secondary colors and a near-equal mixing of primary colors, colored strips were included on the top and bottom
of each box, indicating first and second most frequent residue types respectively. *Types* were organized by *Class* in the
figure. Position-wise information content (in bits) was calculated for each alignment (top). Numbering is for *E. coli* CheW,
framing our observations with reference to the well-studied prototypical protein.