Generalizable strategy to analyze domains in the context of parent protein

architecture: A CheW case study

Luke R. Vass[1,#a], Katie M. Branscum[#b], Robert B. Bourret[1*], Clay A. Foster[1,#b]

[1]Department of Microbiology & Immunology, University of North Carolina, Chapel Hill,

North Carolina, United States of America

[#a]Current Address: Department of Pathology, University of Virginia, Charlottesville,

Virginia, United States of America

[#b]Current Address: Department of Pediatrics, Section Hematology/Oncology, University

of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, United States of

America

Running Title: Domain variation with protein architecture

*Corresponding author

Email: bourret@med.unc.edu

Co-author email addresses:

Luke R. Vass: lukev3@vt.edu

Katie M. Branscum: katief@tribaldiagnostics.com

Clay A. Foster: clay-foster@ouhsc.edu

Data Availability Statement:

22   The data that support the findings of this study were derived from the following resource

23   available in the public domain: Pfam, http://pfam.xfam.org.

Conflict of Interest Statement:

25   The authors declare no conflict of interest.

ABSTRACT

Domains are the three-dimensional building blocks of proteins. An individual domain can be found in a variety of protein architectures that perform unique functions and are subject to different evolutionary selective pressures. We describe an approach to evaluate the variability in amino acid sequences of a single domain across architectural contexts. The ability to distinguish the different evolutionary paths of one protein domain can help determine whether existing knowledge about a specific domain will apply to an uncharacterized protein. Such discrimination can lead to insights and hypotheses about function, as well as guide experimental priorities.

We developed and tested our approach on CheW-like domains (PF01584), which mediate protein/protein interactions and are difficult to compare experimentally. CheW-like domains occur in CheW scaffolding proteins, CheA kinases, and CheV proteins that regulate bacterial chemotaxis. We chose 16 protein *Architectures* that included 94% of all CheW-like domains found in nature. Because some *Architectures* had more than one CheW-like domain, CheW-like domains were divided into 21 distinct *Contexts*. The CheW-like domain sequences were closely related within most *Contexts*; however, one *Context* was subdivided into three *Types*. The resulting 23 sequence *Types* coalesced into five or six *Classes* of CheW-like domains, which we described in detail.

In addition, we created SimpLogo, an innovative method for visualizing amino acid composition across large sets of multiple sequence alignments of arbitrary length. SimpLogo offers substantial advantages over standard sequence logos for comparison and analysis of related protein sequences. The R package for SimpLogo is freely available.

50 **KEYWORDS**

51   protein domain, protein architecture, CheW, CheA, CheV, SimpLogo

1 | INTRODUCTION

Many proteins consist of a chain of discrete domains. Protein domains are typically compact, independently folding units. Originally, protein domains were defined based on protease resistance; however, they are now recognized by characteristic amino acid sequence features. Individual domains possess structures and functions that are subject to natural selection and thus are discrete units of evolution. Domains were assembled in many permutations during evolution to create different proteins. Therefore, domains can be considered modular "building blocks" of proteins. The Protein Family (Pfam) database currently contains almost 20,000 different domains.[1]

Identifying domains is useful because once nature has evolved the solution to a problem (e.g., a protein domain that performs a specific function), the solution is used repeatedly. Thus, domain identity often provides useful general predictions about the structure and function of domains in experimentally uncharacterized proteins, as well as providing an effective means by which to compare proteins of known function. The success of such predictions relies on the accuracy of domain descriptions, which are often considered to be monolithic. While useful in some respects, in actuality there can be substantial differences between domains of the same type that are found in different proteins. For example, response regulator proteins are formed by linking a receiver domain (Response_reg, PF00072) to a wide variety of output domains. At least three types of evidence demonstrate the differences between receiver domains in different types of response regulators: (i) phylogenetic analysis shows receiver domains co-evolve with their output domains,[2,3] (ii) receiver domain:output domain interaction surfaces and receiver domain dimerization surfaces vary as a function of binding

partner and whether a receiver domain exerts positive or negative regulation on its

output domains,[4] and (iii) receiver domains support phosphorylation and

dephosphorylation reaction kinetics that differ by many orders of magnitude depending

on the timescale of the associated biological function.[5-8] Thus, receiver domains, though

termed the same, differ depending upon the protein architecture in which they are

found.

Variation between different instances of a particular domain raises multiple

questions. If the structure, function, and/or other characteristics of a domain are

experimentally characterized in one protein, then how well does that information

translate into useful insights about the same domain in another protein? In other words,

how does a specific protein domain vary when found in the contexts of different protein

architectures? Genome sequencing has far outstripped experimental capacity but

provides a rich source of relevant information. Can sophisticated methods of amino acid

sequence analysis identify contexts (i.e., location within protein architectures) in which

current knowledge about a specific protein domain applies? Conversely, given the

enormous scope of uncharacterized proteins and protein architectures, can we identify

contexts where current knowledge about a specific domain may not apply? Such a

method would provide a way to inspire and focus future experimental investigations of

protein domains in a manner that is most likely to be informative and may also lead to

useful hypotheses and testable predictions. We describe here our development of a

generally applicable approach to analyze the variation in a specific protein domain as a

function of the contexts in which the domain is found, revealing insights at a

substantially higher resolution than a typical imprecise Pfam domain assignment.

As a test case to develop, validate, and illustrate our method, we chose CheW-

like domains (CheW, PF01584), which are part of the signal transduction pathways that

control bacterial chemotaxis.[9] The known function of CheW-like domains is to act as

scaffolds for protein/protein interactions.[10,11] The absences of an active site, substrate

or ligand binding, or enzymatic activity make CheW-like domains challenging to study

experimentally because no simple assay exists by which to compare the properties of

different CheW-like domains. One might envision measuring binding constants, but it

would be difficult to untangle the contributions of the CheW-like domains and their

partner proteins. Furthermore, CheW-like domains are used to assemble very large

multi-protein complexes that couple transmembrane chemoreceptors to cytoplasmic

kinases, so the biological relevance of the binding constant between an isolated CheW-

like domain and its receptor or kinase partner is not immediately obvious. Finally,

evolutionary constraints on CheW function are such that sequences can diverge

substantially and still be functionally interchangeable.[12] Thus, an objective method to

classify CheW-like domains from different contexts would be particularly helpful to

clarify the circumstances in which current knowledge is likely to be applicable and those

which would likely benefit from further investigation.

CheW-like domains are found in three broad lineages of protein architectures:

CheW scaffold proteins, which consist only of CheW-like domains; CheA kinases, which

combine CheW-like domains with domains that provide kinase activity and

phosphorylation sites; and CheV proteins, which combine CheW-like domains with

receiver domains. Notably, the function of CheV proteins is poorly understood. A recent

review of the diversity of bacterial chemotaxis systems asked, "Can efficient

computational approaches be developed to distinguish … CheW from other CheW

domain-containing proteins?".[13] Our results suggest that the answer is yes, and that not

all CheW proteins are alike.

2 | MATERIALS AND METHODS

2.1 | Protein sequence database

The Pfam database (version 33, obtained May 2020) was used for the purposes

of this study.[1] Any sequence found within the Representative Proteome 35 (RP35) to

contain the CheW-like domain (PF01584) was extracted for further use, retaining only

the CheW-like domain sequence.[14] "35" refers to the co-membership threshold used to

control the granularity of the sequence space represented. An RP set best represents

the known sequence diversity of a group of similar proteomes/organisms. To construct

the RP35 data set, available protein sequences for completed proteomes within the

UniProtKB database were grouped based on their UniRef50 cluster membership.

UniRef, or UniProt Reference Clusters, are sets of proteins grouped together by

sequence in such a way as to produce tight clusters of orthologs/inparalogs, maximizing

sequence space while minimizing redundancy; 50 refers to the sequence identity

threshold used for the clustering procedure.[15] We chose RP35 as the best compromise

between information diversity and sequence space reduction. CheW-like domain

sequences were grouped into *Architectures* based on the domain organization of the

parent protein (Figure 2). *Architectures* capturing fewer than 40 sequences were

excluded from the analysis as uninformative.

2.2 | Sequence analysis and initial clustering

CheW-like domain sequences were collectively aligned to the current CheW-like

domain HMM (PF01584) using HMMER3 (to facilitate cross-architecture

comparisons).[16] The removal of poorly aligned sequences or subsequences has been

shown to be beneficial for large-scale, downstream sequence analysis and phylogenetic

profiling. To that end, we used trimAl (version 1.2) to remove spuriously aligned regions

within our CheW-like domain sequences (using a column gap-threshold of 90%,

assuming that the top 10% most "gappy" columns were mostly noise).[17] Entries

with >50% gap-characters were also removed.

We then used the mat.dis() function from the bios2mds package in R (version

1.2.3) to generate symmetric, pairwise-dissimilarity matrices from our sequence

alignment for further clustering (with zero values along the diagonal, representing a

sequence's dissimilarity to itself)[18] using four different evolutionary substitution models:

GONNET, point accepted mutation 250 (PAM250; number of mutations per 100 amino

acids), BLOcks SUbstitution Matrix 30 (BLOSUM30), and Jones-Taylor-Thornton

(JTT).[19-22] Each is tailored towards higher diversity sequence groups (the average

sequence identity reported in the full Pfam alignment for CheW-like domain PF01584 is

only ~20%). The *k*-medoids clustering protocol was performed using the pam() function

from the cluster package within R (version 2.1.2; *k*-medoid clustering is sometimes

called Partitioning Around Medoids).[23] Our initial input parameter *k* was set to 21 (total

number expected of distinct contexts for CheW-like domain; intentionally chosen as an

overestimate). An optimal *k* value (~3-4) was also estimated using the average

silhouette width method implemented in the pamk() function from the fpc package in R

(version 2.2-9; *k*=3-10).[24] The similarity of the partitions generated using the four

substitution models was quantified using both the Adjusted Rand Index (ARI) and

Normalized Mutual Information (NMI) metrics.

ARI is a popular nonparametric statistic used in cluster validation to provide a

quantitative measurement of the similarity between two partition schemes (regardless of

cluster nomenclature). It involves producing a general Rand Index (RI)[25] by calculating

the ratio of the number of common element groupings between solutions using the

following equation:

175
$$RI = \frac{a + b}{\binom{n}{2}} = \frac{correct\ same\ pairs + correct\ different\ pairs}{total\ number\ of\ pairs}$$
(1)

Here, the binomial coefficient $\binom{n}{2}$ represents the total number of unordered pairs in the

set of *n* elements (equivalent to *n(n-1)/2*). In the equation, *a* represents the number of

subset assignments common to both solutions, and *b* represents the number of

assignments that are different. RI scores range from 0 (no overlap between clustering

solutions) to 1 (perfect overlap). The more accurate Adjusted RI[26] essentially rescales

the RI value to accommodate for the fact that random chance will produce some level of

overlap between clusters using the following simplified scheme:

183
$$ARI = \frac{(RI - expected\ RI)}{(maximum\ RI - expected\ RI)}$$
(2)

This relationship ensures that for a completely random assignment of clusters and

samples, the ARI score will equal zero.

Mutual information (MI) is an information theoretic metric quantifying the mutual

dependence between two variables, revealing the amount of information obtainable for

one variable through the observation of another.[27] It is essentially a measure of

correlation and uncertainty; high MI suggests a large decrease in uncertainty about one

random variable based on data from another random variable, whereas zero MI

suggests that the two variables are independent. The measure of this uncertainty is

called entropy. The MI between random variables X and Y is given by the simplified

equation:

194
$$MI(X;Y) = H(X) - H(X|Y) \qquad (3)$$

195   Here, $H(X)$ is the entropy of the single random variable X and $H(X|Y)$ is the conditional

196   entropy of variable X conditional upon the knowledge of variable Y. A normalized variant

197   of this (NMI) is often used to compare clustering performance.[28] The normalized form of

198   MI is defined by the following equation:

199
$$NMI(X;Y) = \frac{H(X) + H(Y) - H(X|Y)}{\left(H(X) + H(Y)\right)\big/2} \qquad (4)$$

NMI has the advantage of scaling the MI measurement within an upper bound of 1

(perfect correlation) and a lower bound of 0 (no correlation), making interpretation more

intuitive. Two partitioning schemes with NMI scores close or equal to 1 are nearly

identical.

2.3 | Multidimensional scaling analysis

Multidimensional scaling (MDS) analysis was performed using three different

algorithms. Metric MDS methods focus on the "distances" in the matrix being analyzed,

whereas non-metric MDS methods focus on the ranking in the matrix. Due to its

simplicity and previous success when applied to protein sequences, we chose to use a

metric MDS algorithm[18] to analyze sequence data from *k*-medoid clustering. For

sequence subsampling, we used two MDS methods. The first was a implementation of

the metric algorithm featured in the wcmdscale() function from the vegan package in R

(version 2.5-7).[29] Given the large number of negligible negative eigenvalues generated

using the metric implementation, we speculated that our initial pairwise distances may

have been non-Euclidean in nature. Consequently, we tested a non-metric

implementation of the MDS algorithm using the metaMDS() function, also from the

vegan package (using parameters: *trymax = 2000, noshare = F, maxit = 9999, sratmax*

= 0.99999999, k = 3, model = "local", autotransform = F, wascores = F, center = T, pc =

T) [29]. Resulting plots were visualized using the ggplot2 package in R (version 3.3.3).[30]

2.4 | Sequence subsampling strategy

Random sequence subsampling was performed on the various CheW-containing

Types to facilitate dimensionality reduction analysis. Each architectural *Context* was

treated as a separate homogeneous class, except for CheW.I, which was split into

CheW.IA, CheW.IB and CheW.IC *Types*. CheW-like domains from parent *Architectures*

featuring multiple domains were split into separate *Contexts* with appropriate suffixes

added relative to their N- to C-terminal position in the parent protein. Two main

approaches to stratified subsampling exist: randomly selecting a predefined number of

observations from each defined subgroup; or randomly selecting a number of

observations proportional to the original subgroup sizes. Given the wide disparity in the

number of CheW-like domain sequences of each *Type*, we chose the former to facilitate

computational efficiency. The mat.dis() function from the bios2mds R package was

incorporated into an iterative sampling procedure.[18] Each subsampling round randomly

selected 25 sequences from each *Type* (without replacement; smaller subsets were

also tested with similar results). Sampling was repeated 10 times, generating a total of

10 replicate sequence sets for each of the four previously described substitution

models. The function also calculated pairwise dissimilarity matrices for each replicate

set for further use during the MDS analysis.

2.5 | Conservation analysis and SimpLogo creation

All CheW-like domain sequences were first aligned collectively to the conserved

CheW-like Hidden Markov Model (HMM) profile (PF01584) obtained from the Pfam

database. Separate sequence alignments were then segmented based on

corresponding types of CheW-like domains. For each alignment, the aligned,

standalone *Escherichia coli* CheW sequence was inserted at the top for use as a

reference point. Columns introducing gaps (relative to the *E. coli* CheW sequence) were

removed for simplicity. Position-wise count matrices and corresponding frequencies

were generated for each alignment using the Biostrings and motifStack packages in

R.[31,32] Physicochemically similar residue types were combined to reduce visual

complexity of the final representation, creating nine amino acid groups (including gap

characters): positively-charged side chains (His/Lys/Arg), negatively-charged side

chains (Asp/Glu), small volume side chains (Ala/Gly), polar side chains

(Gln/Asn/Ser/Thr), large hydrophobic side chains (Met/Val/Leu/Ile), gaps (-), aromatic

side chains (Phe/Tyr/Trp), Pro, and Cys. Each amino acid category was assigned an

"idealized" color and converted to tristimulus values in the CIE XYZ color space. XYZ

space is linear and non-negative (as opposed to traditional RGB space), making it

simple to mix colors. Final values were calculated by weighting (multiplying) the

idealized colors of each amino acid group by their relative frequencies and summing the

results for each position. Logos were visualized using simple geom_tile objects (from

the ggplot2 suite), with colors and heights for each position based on the

aforementioned blended tristimulus values and their overall gap character frequency,

respectively. "Idealized" (i.e., color at 100% frequency) color strips for the top two most

frequent residue types were added to each position (at the top and bottom of each tile)

to mitigate uncertainty between true qualitative colors and blended hues. Position-wise

information content (in bits) was also estimated for each alignment again using

motifStack and visualized with a line trace at the top.[32] To create traditional sequence

logos (for comparison purposes), the Logolas R package was used with default settings

to generate figures with similar dimensions.[33] An R package for SimpLogo was created

and is available (along with an example and the sequence alignments used to generate

Figure 5) in the Github repository https://www.github.com/clayfos/SimpLogo. We used

Zenodo to assign a DOI to the repository: https://doi.org/10.5281/zenodo.6323714.

2.6 | Protein homology modeling and structural analysis

To visualize the relevant functional surfaces of the reference *E. coli* CheW-like

domains CheA-P5 and standalone CheW, we modeled trimeric CheW:CheA-

P5:receptor complexes (capturing both interface 1 and 2) based on the crystal structure

of the *T. maritima* multimer featuring CheW, CheA-P5 fragment, and the chemoreceptor

TM14 (PDB ID 4JPB).[34] We first generated homology models of *E. coli* CheA-P5

(residues 514-640; using the crystal structure of *T. maritima* CheA PDB ID 1B3Q[35] as a

template) and CheW (residues 18-154; using the solution structure of *E. coli* CheW PDB

ID 2HO9 as a template) with the Phyre2 web portal.[36] We then superimposed the top

models against the corresponding *T. maritima* molecules in the crystal structure (PDB

ID 4JPB[34]) using PyMOL (v.2.3.3).[37]

3 | RESULTS

3.1 | CheW-like domains are incorporated into dozens of distinct protein *Architectures* in nature

We first surveyed the known protein architectures into which CheW-like domains are incorporated, which provided initial sequence groupings for further investigation. We extracted the sequences of all proteins containing a CheW-like domain (PF01584; 8484 sequences containing 9094 unique CheW-like domains) from the Pfam database (version 33, accessed in May 2020)[1] using the Representative Proteome 35 (RP35) sequence set.[14] The use of RP35 provided an efficient and representative data set while substantially reducing the sequence space and complexity required for the analysis. CheW-like domain sequences from rare architectures (<40 member sequences) were excluded because we anticipated that interpretation would be too speculative. The remaining sample contained 8581 CheW-like domains (94% of original) in 8061 proteins (95% of original). Additional curation of sequences described in Materials & Methods (e.g., for excessive gap character) resulted in final analysis of 8528 sequences from CheW-like domains (94% of original).

We use the capitalized and italicized terms *Architecture*, *Context*, *Cluster*, *Group*, *Type*, and *Class* to indicate specific ways of categorizing sequences of CheW-like domains in this study. The relationships between the classification terms, as well as the flow of our sorting scheme, are outlined in Figure 1. We first categorized CheW-like domain sequences into three "lineages" based on the architecture and putative role(s) of their parent protein (Figure 2). Our nomenclature for *Architectures* containing CheW-like domains used the accepted lineage of the parent protein (e.g., CheA) as a prefix

and a Roman numeral as a suffix. CheA-lineage *Architectures* were numbered by

decreasing membership size, starting with Roman numeral I. Additionally, *Architectures*

containing two or more CheW-like domains (e.g., CheA.V from Figure 2) were split into

their constituent CheW-like domains and analyzed as separate sequence *Contexts*. An

additional suffix of an Arabic numeral identified the "position" of the CheW-like domain

relative to the N-terminus and the other CheW-like domains in the parent protein, e.g.,

CheA.V.1 (*Contexts* are not labeled in Figure 2). The CheA-lineage was the most

diverse, with a dozen major *Architectures*. In addition to CheW-like domains, most

CheA *Architectures* also contained HATPase_c domains to bind ATP and catalyze

phosphorylation, and histidine phosphotransferase (Hpt) domains to provide a

phosphorylation site.

Although several CheV-lineage architectures (named for the canonical *Bacillus*

subtilis CheV protein) were detected in the RP35 set, the only organization

incorporating an appreciable number of sequences was the canonical CheV protein

(termed the CheV.I *Architecture* for consistency). The CheV-lineage was the least

abundant of the three architectural lineages.

CheW-lineage architectural variants constituted the largest collection of

sequences, with over half of the entire data set attributable to canonical, standalone

CheW proteins from which the *Architecture* gets its name (CheW.I). Additional rare (~1

to 3% the number of CheW.I) *Architectures* featured two or three CheW-like domains

fused together (termed CheW.II and CheW.III respectively) with no other detectable

domains present. Individual CheW-like domains of CheW.II and CheW.III *Architectures*

were separated into distinct *Contexts* for further analysis, doubling or tripling the number

of sequences for the corresponding *Architecture* (i.e., CheW.II.1, for the first position

CheW-like domain, CheW.II.2 for the second position CheW-like domain, etc.).

3.2 | CheW-like domains segregate into distinct sequence *Groups* as a function of

parent protein *Architecture*

We expected that CheW-like domains extracted from a common architectural

Context would more closely resemble each other rather than the same domains

obtained from contrasting architectural variants. We based our assumption on existing

knowledge of microbial response regulator receiver domains, which cluster

phylogenetically predicated on the presence and/or type of an attached output

domain.[2,3] However, the full dataset contained over 8500 CheW-like domain sequences,

making analysis and interpretation challenging. Ideally, each CheW-like domain *Context*

would behave as a homogeneous sequence set, and a basic stratified sampling

strategy could be used to select a representative and more manageable population for

study. While the assumption of homogeneity seemed reasonable (shared architecture

typically implies common function), we wished to first gain insight into the underlying

structure of the sequence space by performing unsupervised clustering on the full list of

sequences. Unsupervised clustering would allow us to intelligently select the best

representative sample from our data set. In essence, our approach would be a

simplified multistage sampling scheme, first clustering the entire population to reveal

major strata within each architectural *Context*, then randomly sampling from within

these strata. This would ensure that all major sequence groups were represented. We

chose to use a *k*-medoids approach for the initial clustering procedure to mitigate the

difficulty of visualizing over 8500 data points. The *k*-medoids algorithm is a classical

clustering technique that partitions the data into *k* number of *Clusters*, attempting to

minimize the distance between data points and the center of their relevant *Cluster*. Each

Cluster is represented by a single central data point called a cluster medoid, which

allowed us to visualize the *Clusters* using just the medoid sequences, rather than the

entire data set.

Cluster analysis of protein sequences can be performed in a number of ways,

though it most commonly involves using a multiple sequence alignment to calculate

dissimilarity scores between all possible pairs of sequences and clustering the resulting

matrix. We generated symmetric, pairwise-dissimilarity matrices from our sequence

alignment for further clustering as described in Methods. Estimating protein sequence

(dis)similarity requires the use of a substitution matrix to describe the rate or frequency

at which a residue is substituted for another over time, simulating the process of

evolution. Many types of substitution matrices exist. Because we were uncertain which

would be most appropriate for our CheW-like domain data set, we tested four distinct

substitution models as described in Methods. We calculated four distinct dissimilarity

matrices for our sequence alignment using the four substitution models as inputs for the

k-medoids cluster analysis. For convenience, we only show solutions derived from the

GONNET substitution matrix in the main text, due to the documented ability of GONNET

to handle sequences of greater evolutionary distance.

The *k*-medoid technique requires an initial target number of clusters (the *k*-value)

as input. An optimal *k*-value can be estimated with various metrics, such as the average

silhouette width (ASW; a measure of how similar a point is to its own cluster compared

to other clusters), but choosing the "correct" number of clusters is challenging,

subjective and dependent on the overall goals of the analysis.[38] *A priori*, we chose $k =$

21 (the total initial number of distinct architectural *Contexts*, including multiple domains

in the same protein, used to partition the CheW-like domain sequences) as our *k*-value.

This was an intentional overestimation of the true number of subpopulations (optimal *k*-

value estimation using the ASW suggested 3-4 clusters was ideal, data not shown). We

then performed clustering as described in Methods. Pairwise dissimilarities between the

medoids for each of the 21 identified *Clusters* were extracted and used as input for

further analysis with a metric implementation of multidimensional scaling (MDS) as

described in Methods. MDS is a multivariate dimensionality reduction and visualization

technique that takes pairwise similarities/dissimilarities/distances for a data set and

attempts to reconstruct a simplified map that maintains the distances between data

points and is easily interpretable (i.e., in a lower *k*-dimensional space). In metric MDS

(sometimes more generally referred to as Principal Coordinates Analysis), we attempt to

visualize the (dis)similarity data along the dimensions that maximize the amount of

variation (or information) within the data, similar to Principal Component Analysis.

A representative MDS plot describing just the 21 extracted medoids (based on

the dissimilarities calculated using the GONNET model) is shown in Figure 3. Additional

plots generated using dissimilarities calculated with the remaining substitution matrices

are in Figure S1. Letters naming the *Clusters* for each solution were assigned based on

decreasing number of sequences in the *Cluster*. There were three substantial *Groups* of

Clusters of CheW-like domains (labeled 1, 3, or 4 in Figure 3), and another smaller,

intermediate *Group* separating *Groups* 1 and 3 (labeled 2 in Figure 3, more apparent

when considering only the first dimension). The partitions were highly similar between

the four clustering solutions (Figures 3 and S1), with pairwise Adjusted Rand Index

(ARI) scores ranging from 0.6-0.7 and Normalized Mutual Information (NMI) scores of

0.7-0.8 calculated as described in Methods. Consistently high ARI and NMI values

across solutions derived using four different substitution matrices suggested a relatively

stable underlying data structure.

We next examined the composition of the individual *Clusters* to determine if the

original architectural *Contexts* of CheW-like domains separated into multiple stable

strata. Although each medoid was representative of the dissimilarity data, a *Cluster* may

contain multiple domain *Contexts*. Understanding the composition of each *Cluster* would

allow us to properly organize sequences for analysis on the more complete sequence

set, rather than on just the representative medoids, ensuring that no distinct *Contexts*

would be underrepresented. Table 1 assigns *Clusters* to the four main *Groups* indicated

in Figure 3 and contains a detailed breakdown of the architectural *Contexts* of the

sequences in each *Cluster*.

Multiple patterns are apparent in Table 1. First, sequences form *Clusters* and

Clusters form *Groups* primarily based on parent protein lineage. Most or all sequences

within a *Cluster* are from either CheA proteins or CheW/CheV proteins, not both. Most

CheW-like domains from CheA proteins are in *Group* 1 and most CheW-like domains

from CheW and CheV proteins are in *Group* 4. Closer inspection reveals additional

structure in the data:

• Sequences of CheW-like domains from most CheA-derived *Contexts* were found

in *Group* 1.

- The "second" position CheW-like domains from *Architectures* CheA.V and CheA.VI formed their own distinct sequence *Group* 2, made up of *Cluster* P.

- CheW-like domains from *Architectures* CheA.VII and CheA.X comprised the bulk of *Cluster* R, the smallest *Cluster* in *Group* 1.

- About 80% of domains from CheW.I *Architectures* and greater than 90% of domains from CheW.II and CheW.III *Architectures* were in *Group* 4.

- About 20% of domains from CheW.I *Architectures* and 5% of domains from CheW.II and CheW.III *Architectures* were in *Group* 3, possibly distinct from the majority of CheW-lineage domains.

- About 1% of domains from CheW.I *Architectures* and 2% of domains from CheW.II and CheW.III *Architectures* were in *Group* 1, suggesting similarity to the CheA-lineage.

- About 99% of CheW-like domains from CheV.I *Architectures* were in *Group* 4, suggesting that the presence of the attached receiver domain has little effect on the primary sequence of the CheW-like domain in CheV proteins.

- Similar to CheW proteins, about 1% of CheW-like domains in CheV.1 *Architectures* were in *Group* 1, suggesting similarity to the CheA-lineage.

Based on the analysis of Table 1, we assumed for further processing that each architectural *Context* could be reasonably treated as a single homogeneous entity, except for CheW.I. We split CheW.I into three distinct *Types* designated CheW.IA (sequence *Group* 3), CheW.IB (sequence *Group* 4), and CheW.IC (sequence *Group* 1). Remaining outlier sequences in each architectural *Context* (typically a small number of sequences in each *Context* assigned to substantially different *Clusters* than the

majority) were excluded from the analysis as noise. We believe that our evaluation

produced an accurate summary of the underlying data structure, though we

acknowledge the somewhat subjective nature of cluster analysis and that alternative

perspectives may offer equally valid insights. We found the visualization in Figure 3 to

be limited in interpretability when searching for biologically relevant trends. To address

this shortcoming, we next performed MDS analysis on a larger, randomly sampled

subset of our sequence data, now organized into 23 *Types* (due to CheW.I sub-

designations) instead of 21 *Contexts*.

3.3 | Random stratified subsampling facilitates efficient multi-dimensional scaling

analysis on large protein sequence sets and reveals *Classes* of CheW-like

domains

One technical disadvantage of the MDS technique is its substantial

computational cost. The time complexity of standard metric MDS is proportional to the

square of the input size, which indicates significant temporal and technical requirements

when analyzing a high-dimensional data set with thousands of data points.[39] Thus, the

MDS technique is impractical when analyzing a data set such as our original ~8500 x

8500 dissimilarity matrix. One popular approach to dealing with large, unwieldy data is

random subsampling, called stratified sampling when working with distinct groups or

strata within the original population. For this strategy to be effective, adequate

representation of each stratum is essential. Our initial cluster analysis (Figure 3 and

Table 1) revealed how to organize sequences of CheW-like domains into 23 *Types* that

should ensure all major subpopulations receive appropriate representation in such a

sampling scheme.

We randomly chose 25 sequences from each of the 23 previously described

Types to analyze with metric MDS as described in Methods. We repeated this stratified

random sampling procedure for 10 iterations (575 distinct sequences per iteration) to

assess the stability of the various randomized MDS solutions. We also included the

CheW-like domains of 18 well-studied reference proteins as an internal constant for

each iteration (data not shown).

Figure 4 shows the top three dimensions of one representative solution derived

from the GONNET substitution matrix. The remaining nine iterations, as well as

solutions calculated using randomly sampled subsets for the other three previously

described substitution matrices, are in Figures S2-S13. The solution shown in Figure 4

revealed similar relationships to those found in Figure 3/Table 1, which we now

designate as *Classes* (sub-sampling MDS) rather than *Groups* (*k*-medoid MDS). On the

right area of the plot (Left and Center Panels), most CheW-lineage and CheV-lineage

sequence *Types* (*Class* 1) clustered in a densely packed region. A large agglomeration

of sequences from CheA-lineage *Types* (*Class* 3) was near the left side of the plot area

(Left and Center panels). The CheA.V.2 and CheA.VI.2 *Types* (*Class* 5) were markedly

different from the other CheA-lineage *Types*, consistent with previous assignment to

Cluster P as the sole member of *Group* 2 in Figure 3/Table 1. Additional differentiation

within the CheA-derived *Types* could be observed more effectively in the Center and

Right Panels (incorporating Dimension 3), with the CheA.VII and CheA.X *Types* (*Class*

4) separating from the main body. This supports the clustering results presented in

Table 1 and Figure 3, with the majority of CheA.VII and CheA.X sequence *Types* found

in the somewhat independent *Cluster* R. Sequences belonging to the rare CheW.IC

Type (*Class* 2) were spread across Figure 4, alongside *Class* 1 CheW/CheV

sequences, *Class* 3 CheA sequences, and in between (Left and Center Panels). In

Figure 3/Table 1, a large subset of CheW-lineage sequences (*Group* 3) separated from

most CheW- and CheV-lineage sequences (*Group* 4). The *Group* 3 sequences are

termed *Type* CheW.IA (*Class* 6) in the analysis of Figure 4 and surprisingly did not

separate from the main CheW/CheV *Types* (*Class* 1).

Other forms of MDS may offer additional advantages under the appropriate

circumstances. Examination of our randomly sampled MDS solutions indicated that a

significant proportion of the total eigenvalues were negative (~15.4-15.7%). Though the

individual negative eigenvalues were miniscule compared to the positive entries, it was

possible that our initial dissimilarities were non-Euclidean in nature, and another method

such as non-metric MDS might offer additional insight. As a test, we attempted to derive

further solutions using a non-metric implementation of MDS as described in Methods.

These can be found in Figures S14-S25 in Supporting Information. We encountered

significant challenges related to solution convergence, likely due to the still relatively

large and complex data set (despite our subsampling strategy). However, the non-

metric MDS solutions were largely reminiscent of the basic layouts generated using the

metric algorithm, even when sampling smaller subsets of sequences. Thus, we

concluded that the data structures were likely consistent irrespective of the MDS

algorithm. Given the clear sequence heterogeneity within the CheW-like domains, we

next sought to determine factors within the domain itself that distinguished the various

Types.

3.4 | SimpLogo: A simplified method for visualizing large numbers of distinct protein sequences aligned over many positions

Arguably the most popular method for comparing sequence groups and their

alignments is the sequence logo. In this context, a logo is a graphical representation

depicting sequence conservation by stacking relevant characters, such as single-letter

codes for amino acids or nucleotides, at each position in an alignment.[40] The relative

heights of the characters are proportional to their frequency at the given position.

Additionally, the overall height of a stack is often scaled based on the information

content stored at that position within the alignment. Typical applications for amino acid

sequence logos include visualizing functional motifs or binding sites within a protein

family/group. However, the volume of information and the graphical complexity of a logo

often makes it challenging to interpret when examining regions longer than two dozen

positions. Sequence logos are less practical when comparing a large group of separate

alignments (>12) with many positions (>50). A more ideal and complementary

visualization technique would allow us to first evaluate the macroscopic characteristics

of each sequence group. Subtle differences could then be examined in greater detail for

specific regions using a more traditional approach like a true sequence logo, or even the

original sequence alignment. Here, we describe a visualization format (colloquially

referred to as SimpLogo) designed to compare the CheW-like domain sequences from

the numerous *Types* described above. We sought to reduce the 23 individual sequence

alignments (*Types*) into simplified, pseudo-one-dimensional representations. We

accomplished this by weighting and blending qualitative colors assigned to specific

amino acid residue types (grouped by physicochemical properties into eight classes)

based on their frequency at a given position within the alignments. We also modified the

overall height based on its proportion of gap-character (details in Methods). This

produced a single row of colored "boxes" for each sequence *Type*, reducing a complex

problem to basic pattern recognition. To differentiate between secondary colors and a

near-equal mixing of primary colors, colored strips were included on the top and bottom

of each box, indicating first and second most frequent residue types. The simplified

scheme allowed for the simultaneous comparison of dozens of distinct sequence

families with lengths over many positions (>100) and encoding substantially more

information about intermediate frequencies than simple consensus sequences.

Additionally, we created traditional sequence logos (using the Logolas package in R) for

the same alignments using similar image dimensions for comparison (see Figure S26).

Highly conserved/informative positions are identifiable in the traditional logo, but the

overall lack of space and reduced character sizes often make interpretation of specific

positions challenging, especially for the lower frequency/less conserved positions with

smaller stack heights. The patterns of gap characters (indicated by dashes) are also

more challenging to interpret when compared to the SimpLogo representation in Figure

5. One could imagine how difficult interpretation would become with double the number

of sequence groups or sequence lengths using traditional methods. Therefore, we

envision the new SimpLogo visualization technique being used not as a replacement for

traditional sequence logos, but as a supplementary technique for large scale exploratory

analyses.

3.5 | *Classes* **of CheW-like domains display characteristic gap patterns**

The SimpLogo representation in Figure 5 revealed multiple trends. Overall, the

relationships displayed in Figure 5 mimicked the trends seen in Figure 4, with most of

the CheA-lineage *Types* clustered tightly (*Class* 3), the exceptions being the

CheA.VII/CheA.X (*Class* 4) and CheA.V.2/CheA.VI.2 (*Class* 5) *Types*. Most CheW-

lineage *Types* (*Classes* 1 and 6) and the CheV-lineage *Type* (*Class* 1) appeared

similar, except for the CheW.IC *Type* (*Class* 2), which was intermediate in character

between CheA and CheW *Types*. The gap character of the three canonical parent

protein lineages likely played an important role in differentiating the sequence *Types*,

particularly at positions 45-56 and 120-131. Here, we provide observations related to

the corresponding gap patterns and offer possible functional implications. We assumed,

for the sake of speculation, that the analogous interaction surfaces and behaviors are

largely maintained throughout the various CheW-like domain topologies found in nature.

All positions described are relative to *E. coli* CheW unless otherwise specified. Crystal

structures of the CheW-like domains from well-studied *E.* coli CheW or CheA proteins

are not available. To provide a structural reference for our observations, we constructed

simple homology models of *E. coli* standalone CheW (residues 18-154) and the CheW-

like domain (termed P5) of CheA (residues 514-640) using the Phyre2 web service.[36]

Various structural alignments were performed against an existing crystal structure

featuring CheW, CheA P5 domain, and the chemoreceptor TM14 from *Thermatoga*

maritima (PDB ID 4JPB)[34] to provide insight into the functional consequences of the

characteristics described in the following paragraphs. Table S1 contains a detailed list

of relevant putative residue functions for convenient cross-referencing, based on

various literature sources and our homology models.

At positions 45, 55, and 56, the CheA-lineage *Types* generally exhibited

sequence gaps, consistent with the observation that sequences of CheW-like domains

from CheA proteins typically appeared shorter overall than from single domain CheW

proteins when using the standard Pfam model (corroborated in manual CheW-like

domain sequence alignments, data not shown). Experimental binding data suggest that

positions 45-56 are involved in the interaction between the standalone CheW and

CheA, at least in *E. coli.*[41,42] The CheW-like domain topology resembles that of a SRC

Homology 3 (SH3) domain and is made up of two separate, pseudo-symmetric

subdomains (1 and 2).[43] Subdomain 1 of the CheW-like P5 domain in *E. coli* CheA

(approximately residues 514-531 and 596-640) interacts with subdomain 2 of

standalone *E. coli* CheW (approximately residues 36-113; termed interface 1), primarily

through the β3-β4-β5 regions of each individual subdomain (approximately residues

608-633 on P5 and 44-67 on CheW, based on homology modeling).[34] However, the

complex signaling array/lattice formed by chemoreceptor:CheW:CheA oligomers is

facilitated by the formation of a secondary interface between subdomain 1 of standalone

CheW and subdomain 2 of P5 (approximately residues 114-134 on CheW and 539-556

on P5, based on homology modeling; termed interface 2), an interaction that contributes

substantially to the cooperativity of the signaling array.[44] Positions 45-56 did not overlap

with known chemoreceptor binding surfaces on CheA or CheW, implying that

differences in gap character seen in Figure 5 likely affect the primary and/or secondary

associations between the CheW-like domains of CheA- and CheW-lineage

Architectures and not direct interactions with the chemoreceptor. Because subdomain 2

serves distinct purposes in CheA and CheW, differences in the corresponding functional

residues seemed reasonable.

There were two exceptions to the described dichotomy between CheA-lineage

and CheW-/CheV-lineage *Types* regarding gaps at positions 45, 55, and 56. First,

Types CheA.V.1 and CheA.VI.1 exhibited only a partial gap pattern at position 45.

Recall that *Types* CheA.V.I and CheA.VI.1 fell into *Cluster* H (Figure 3/Table 1).

Although ultimately considered part of *Group* 1, *Cluster* H showed significant deviation

from the rest of the CheA-lineage *Types*, with increased resemblance to *Group* 2

(*Cluster* P, made up of *Types* CheA.V.2 and CheA.VI.2 from the same *Architectures*) in

Figure 3. However, *Cluster P* exhibited gaps at position 45 (Figure 4). Second, the gap

patterns at positions 45, 55, and 56 exhibited by *Type* CheW.IC were more reminiscent

of the CheA-lineage sequences than the rest of the CheW-lineage *Types*. This is

consistent with observations that (i) the CheW.IC *Type* at least partially resembled

CheW-like domains from CheA-lineages (both belonged to *Group* 1 in Table 1) and (ii)

as *Class* 2 spanned the sequence "gap" between CheA (*Class* 3) and CheW (*Class* 1)

regions of Figure 4.

CheA-lineage *Types* (*Classes* 3-5) exhibited stretches of gaps at positions 120-

131, which corresponds to subdomain 1 in CheW-like domains. Therefore, sequence

differences here could also affect the association between CheA- and CheW-/CheV-

lineage proteins. CheV- and CheW-lineage *Classes* 1 and 6 shared the gap pattern

near position 120 with CheA-lineage *Classes* 3-5 but lacked the stretches of gaps in

subdomain 1 near positions 124 and 130. In contrast, CheW.IC (*Class* 6) again more

closely resembled the gap profile seen in the CheA-lineage *Classes* 3-5 at positions

120-131.

Type CheA.V.2 and CheA.VI.2 domains (*Class* 5) displayed a slightly modified

gap profile at positions 122-130 and exhibited a more diffuse level of gap character at

positions just upstream of the region (indicated by subtly "shrunken" boxes stretching

back to residue position 110). Although the role(s) played by positions 109-110 in array

formation is unclear, structural and biochemical data suggest that residue 108 is directly

involved in chemoreceptor binding.[45] In the dual CheW-like domain CheA.V and

CheA.VI *Architectures*, the "first" position CheW-like domains (*Types* CheA.V.1 and

CheA.VI.1) did not exhibit the same deviation in gap character at residues 110-119 as

Class 5, suggesting that the "second" position CheW-like domains may bind

chemoreceptors that are distinct from those of their N-terminal counterparts, or perhaps

bind entirely different partners through this region.

3.6 | *Classes* of CheW-like domains display characteristic residue identities

Careful inspection of Figure 5 also revealed several characteristic deviations in

residue frequencies at key positions, many of which involved *Class* 5 domains:

The entire region between positions 108-134 was filled with conserved charged

and polar residues that are likely involved in intermolecular/interdomain interactions in

CheA-lineage *Types* but that were absent in the other *Types*. At position 110, several of

the CheA-lineage *Types* encoded a negatively charged (red) residue (particularly in

CheA.II/III/IV/V.1/VI.1/VII/X), whereas the position was less conserved in the CheV- and

CheW-lineage *Classes* 1 and 6. The gap character was increased at position 110 in

Class 5. Based on our homology models and alignment against another crystal

structure of dimeric *T. maritima* CheA (residues 290-671 relative to the *T. maritima*

protein; PDB ID 1B3Q), residue 110 likely interacts with the helical dimerization domain

of the CheA kinase, possibly explaining the change in conservation in CheA-related

Types compared to the other lineages. Similarly, position 113 was strongly hydrophobic

(green), position 114 was strongly positively charged (blue), and position 115 was

polar/Pro (yellow/white) in almost all the CheA-lineage *Types* except *Class* 5, but not in

the CheV- and CheW-lineage *Classes* 1 and 6. These positions are likely related to

formation of the interface between the CheW-like domains of CheA and CheW.

Class 5 domains exhibited distinct patterns of charged amino acids. Position 38

was positively charged (blue) in *Class* 5 but negatively charged (red) in all other

Classes except 4. *Class* 5 also exhibited a pair of conserved, negatively charged (red)

residues at positions 40-41, a unique trait not exhibited by any other *Class*. In *E. coli,*

the region equivalent to position 40 is involved in chemoreceptor binding,[41,42,45] again

suggesting different sensor/partner specificity for *Class* 5 domains.

At position 49, all four of the CheW-like domains in the CheA.V and CheA.VI

Architectures shared a conserved negatively charged (red) residue. The remaining

CheA-lineage *Types* exhibited substantially lower conservation, leading to brownish

coloration, whereas the CheV- and CheW-lineage *Types* primarily exhibited a

conserved Pro residue (white). Equivalent residues in *E. coli* are involved in the

association of CheW-like domains in CheA and CheW, based on the homology models.

Class 4 *Types* (CheA.VII and CheA.X) did not appear obviously different than

Class 3 *Types* (most CheA *Types*) in Figure 5 except for Pro at position 87. The only

Type that consistently stood out in *Classes* 3 and 4 was *Type* CheA.VIII. At position 27,

most *Types* exhibited a low level of conservation, leading to a reddish-brown color.

However, the CheA.VIII *Type* featured almost exclusively negatively charged (red)

residue types. Closer examination of the traditional sequence logo and alignment for the

CheA.VIII *Type* revealed a nearly invariant Glu. In our *E. coli* CheW homology model,

the negatively charged Glu side chain of position 27 formed a hydrogen bond with the

hydroxyl group of Tyr29 (brown) near the site of interface 2, which is essential for

higher-order oligomer formation.

At position 41, there was an increased conservation of positively charged (blue)

residues in many of the *Types* from CheA- and CheW-lineages, including CheA.V.1,

CheA.XI, CheA.XII, CheA.VII, CheA.X, and to a lesser extent CheW.IB, CheW.III.1 and

CheW.III.3. However, *Class* 5 domains exhibited an increase in negative charge (red) at

the same position. *Type* CheA.VIII featured a highly conserved Gln (yellow) at position

41 followed by a similarly conserved Pro (white). The Gln Pro pair was not found in

other *Types*. Experimental data and our homology models suggested that position 41 is

involved in both CheA/CheW association and chemoreceptor binding.[42]

We did not observe any positions where *Class* 6 (CheW.IA), representing ~20%

of CheW-lineage domains, differed from *Class* 1 (most CheW domains and CheV

domains). However, we did observe some variation within *Class* 1:

• Position 56 exhibited gaps in most CheA-lineage *Types* and a strong

hydrophobic (green) character in most CheW-lineage *Types* but exhibited

increased negative charge (red) in CheA.VIII, CheV.I and CheW.III.3, and a

positively charged ([blue) residue in CheW.IB. *In E. coli* CheW, experimental data

indicate that position 56 is involved in the formation of interface 1, forming a

hydrogen bond with Tyr81. Our homology models suggested that the region is

likely substantially altered in CheA-lineage structures, as indicated by the

consistent gaps at positions 55-56,[41] with the exception of CheW-like domains

from *Type* CheA.VIII (which includes the P5 subdomain of *E. coli* CheA). The

deviations seen at position 56 were likely a result (or cause) of differences in

primary CheW/CheA association interactions between various combinations of

CheA and CheW *Types*.

- 696 At position 32, most *Types* exhibited Pro (white), except *Types* CheW.IB,

CheW.II.1, CheW.II.2, and CheV.I, which featured increased negatively charged

(red) residues in the first three, and almost exclusively Asn (yellow) in the latter.

Position 32 is proximal to the chemoreceptor binding interface (according to our

homology model for the CheW-like domain of *E. coli* CheA) and is likely involved

in the positioning and/or formation of corresponding intermolecular interactions.

The high conservation of a Pro residue in most *Types* suggested an important

structural feature.

- 704 Position 74 featured a seemingly buried hydrophobic (green) residue in most

CheW-like domains, with the exceptions of *Types* CheW.IB, CheW.II.1 and

CheW.III.3, which instead displayed positively charged (blue) residues (primarily

Lys). The entire region between positions 72-74 was significantly more positively

charged in the CheW-lineage *Types*, with the possible exception of CheW.IA.

The implications of this unique characteristic are unknown, because the positions

are not known to be involved in receptor binding or interface formation.

• At positions 86-87 (a critical residue pair involved in chemoreceptor binding in *E.*

coli CheW[41,45]), most of the CheW-lineage *Types* exhibited polar (yellow) and

positively charged (blue) residues, except for CheW.II.1 and CheW.IC, which

featured negatively charged/polar (red/yellow) and small-volume/hydrophobic

(dark grey/green) residue pairs, respectively.

Finally, CheA.VI.1, CheW.II.2 and CheW.III.2 were the only *Types* with a strongly

conserved Cys (cyan) residue, found at positions 24, 50-51, or 135 respectively. To the

best of our knowledge, no naturally occurring disulfide bond formation has been

reported involving a CheW-like domain, though conservation implied that the Cys

residues were functionally or structurally significant in some way. The role of position 24

is currently unclear. Positions 50 and 135 are thought to be involved in the association

of CheA- and CheW-lineage domains.

 4 | DISCUSSION

 4.1 | CheW-like domains sort into six *Classes*

 We started with three lineages and 16 protein *Architectures* (Figure 2) that

 contained 94% of CheW-like domains. Due to multiple domains in four *Architectures*,

 there were a total of 21 *Contexts* for CheW-like domains. Cluster analysis (Figure 3)

 revealed that the largest *Context* (single domain CheW proteins) split into three *Types*

 (Table 1). Random stratified subsampling (Figure 4) showed that the 23 *Types* formed

 five or six *Classes* of CheW-like domains, a conclusion that was supported by visual

 inspection of SimpLogos (Figure 5):

 Class 1 contained seven *Types* of CheW-like domains (CheV.I, CheW.IB,

 CheWII.1, CheWII.2, CheWIII.1, CheWIII.2, CheWIII.3) that comprised CheV proteins

 and most CheW proteins. CheW-like domains in CheW proteins were generally similar,

 regardless of whether the protein had one, two, or three CheW-like domains. CheW-like

 domains in CheV proteins were similar to CheW-like domains in CheW proteins.

 Class 2 contained the ~1% of CheW-like domains in CheW proteins (*Type*

 CheW.IC) that shared similarity with CheW-like domains in CheA proteins (Table 1,

 Figure 5). In MDS analyses, members of *Class* 2 landed in the CheW region, the main

 CheA region, or in between (see Figures 4 and Figures S2-S25).

 Class 3 contained 10 *Types* of CheW-like domains (CheA.I, CheA.II, CheA.III,

 CheA.IV, CheA.V.1, CheA.VI.1, CheA.VIII, CheA.IX, CheA.XI, CheA.XII) that

 encompassed almost all CheA proteins. CheW-like domains in most CheA proteins with

 one CheW-like domain were similar, even though there were 10 different major

 Architectures for such proteins, including one (CheA.IX) that lacked a phosphorylation

site (Hpt domain) and two (CheA.XI, CheA.XII) that lacked ATP binding sites

(HATPase_c domain). Such "split" CheA *Architectures* are not degenerate, but can work

together to reconstitute a functional unit.[46] N-terminal CheW-like domains in CheA

Architectures with two CheW-like domains (CheA.V.1, CheA.VI.1) were similar to

CheW-like domains in CheA *Architectures* with a single CheW-like domain.

Class 4 was made up of two *Types* of CheW-like domains (CheA.VII & CheA.X)

from CheA proteins with three Hpt domains and a receiver domain. *Class* 4 CheW-like

domains consistently diverged from other CheW-like domains in CheA proteins with a

single CheW-like domain (Figure 4).

Class 5 consisted of C-terminal CheW-like domains in CheA proteins with two

CheW-like domains (CheA.V.2 & CheA.VI.2). *Class* 5 CheW-like domains were different

from all other CheW-like domains in CheA proteins (*Classes* 3 and 4), CheW proteins

(*Classes* 1 and 6), and CheV proteins (*Class* 1) (Figure 4).

Visualization of *Class* 6 depended on the method used. Approximately 20% of

CheW-like domains in CheW proteins (*Type* CheW.IA) appeared to be different from the

rest by *k*-medoid analysis (see Figure 3/Table 1). *Class* 6 appeared indistinguishable

from *Class* 1 in the metric MDS analyses of random stratified subsampling in Figure 4

and Figures S2-S13, and the differences were subtle in the SimpLogo of Figure 5. *Class*

6 showed some separation from *Class* 1 in the non-metric MDS analyses of random

stratified subsampling in Figures S14-S25.

4.2 | Implications of CheW-like domain *Classes*

The existence of *Class* 1, which encompasses CheV and most CheW proteins,

has multiple implications. First, evidence exists that CheV proteins can function as

scaffold proteins similarly to CheW proteins; however, more detailed information about

the functions and mechanisms of CheV is sparse.[11,47,48] It is tempting to extrapolate

what is known about response regulator proteins (composed of N-terminal receiver and

C-terminal output domains) to formulate hypotheses about CheV (composed of C-

terminal receiver and N-terminal CheW-like domains). However, our observation that

CheW-like domains in CheV proteins were similar to those in CheW proteins constrains

such hypotheses by suggesting that any interactions between the receiver and CheW-

like domains in CheV proteins are not sufficiently specific to be subject to selective

pressure for evolutionary optimization. There are at least three categories of hypotheses

about CheV inspired by response regulators:

• The receiver domain might alter the conformation of or occlude access to the

CheW-like domain, depending on the phosphorylation state of the receiver.

Alexander *et al.* suggested that such phosphorylation-mediated changes could

explain the role of CheV in adaptation by changing the conformation of the

CheW-like domain to alter the degree of coupling between receptor and kinase

function.[47] Our findings make this otherwise reasonable hypothesis now seem

unlikely. In *Bacillus subtilis* CheV, the CheW-like domain inhibits phosphotransfer

from CheA-P to CheV and stabilizes CheV-P from autodephosphorylation.[49] Our

findings suggest that these effects are more likely the result of steric effects than

conformational coupling between the two domains of CheV. The determination of

CheV structures in the presence and absence of the stable phosphoryl group

analog BeF_3^- would be highly informative.

• The phosphorylation site in the receiver domain of CheV might act as a

phosphate sink, while the CheW-like domain localizes CheV appropriately.[47,48]

Our results are consistent with such a hypothesis because no interaction

between the two domains of CheV is required that might lead to divergence of

CheW-like domains in CheW and CheV proteins. In a phosphate sink

mechanism, when environmental circumstances change and phosphoryl groups

stop flowing into the system from ATP, phosphoryl groups flow backwards from

the CheY response regulator to CheA to the sink and are lost to hydrolysis.

Localizing the sink near CheA would enhance efficiency. We note that another

possible role of CheV is as a reservoir that retains phosphoryl groups in a

buffering capacity.

• In a more dynamic role, phosphorylation-mediated dimerization of the receiver

domain of CheV might control the ability of the CheW-like domain to bind with

receptor and/or kinase partners. To the best of our knowledge, the multimeric

state of CheV and any effect of phosphorylation on multimeric state have not

been investigated.

Second, five (CheA.III, CheA.IV, CheA.V, CheA.VII, CheA.X) of the 12 CheA

Architectures in Figure 2 contained a receiver domain immediately C-terminal to a

CheW-like domain, which is reminiscent of CheV proteins. However, the fact that

CheW-like domains in CheV and CheA proteins were in different *Classes* strongly

suggests that adjacent CheW-like and receiver domains in CheA proteins do not act as

CheV analogs. Consistent with this prediction, the adjacent CheW-like domain did not

affect autodephosphorylation of the receiver domain from *Azorhizobium caulinodans*

CheA,[50] in contrast to *B. subtilis* CheV.[49]

Third, as far as we know, obvious questions about multi-domain CheW proteins

have not been explored experimentally. It is not known if all domains of multi-domain

CheW proteins are required for function, or if there are differences in functional roles

between the domains. The fine structure of the data in Table 1 (specifically the number

of sequences in a *Cluster* varied substantially as a function of *Context* within CheW.II

and CheW.III *Architectures*) strongly suggests that the domains in a single multidomain

CheW protein often belong to different sequence *Clusters* or even *Groups*, which could

suggest domain specialization. On the other hand, because experimentally

characterized single domain CheW proteins belonged to *Class* 1, which also includes

~93% of domains in multi-domain CheW proteins, our results suggest that there is likely

little to be learned about general CheW function by exploring multi-domain CheW

proteins. Presumably, multi-domain CheW proteins support formation of complexes with

specific components. This conjecture is supported by the observation that CheW.III

proteins form cytoplasmic double layered arrays with CheA and a special type of double

domain chemoreceptor in *Vibrio cholerae*.[51]

Approximately 1% of CheW-like domains in CheW proteins were CheA-like

(CheW.IC *Type*) and formed a *Class* of their own. Some members of *Class* 2 appeared

to span the gap between CheA proteins and CheW/CheV proteins (Figure 4). The

CheW-like domain is the most recently evolved fold in proteins that support bacterial

chemotaxis.[52] The birth of CheW-like domains allowed the separation of receptor and

kinase functions into different proteins, in turn facilitating the integration of multiple

stimuli sensed by arrays of different receptors into a single kinase. We are unaware of

any evidence establishing whether CheW-like domains evolved first in CheW, CheA, or

CheV proteins. Because CheV proteins are less common and less widely distributed

than CheA or CheW proteins,[52] it seems reasonable that CheV proteins came after

CheA and CheW. Because members of *Class* 2 segregated with *Class* 1, *Class* 3 and in

between in our analyses (Figure 4), *Class* 2 may represent a "missing link" in the

evolution of CheW-like domains from CheW to CheA or from CheA to CheW. Closer

examination of sequences in *Class* 2 may provide new insights into the evolution of

chemotaxis systems.

The existence of *Class* 3, which covered CheW-like domains in most CheA

proteins, suggests that what is known about CheW-like domains in experimentally well-

characterized CheA proteins is likely to be applicable to the majority of CheW-like

domains in CheA proteins. The exceptions would be *Class* 4 and 5 CheW-like domains.

Class 4 CheW-like domains were found in CheA.VII and CheA.X *Architectures*,

which have three Hpt domains and a C-terminal receiver domain and to the best of our

knowledge have not been experimentally characterized. However, *Pseudomonas*

aeruginosa ChpA, which has six Hpt domains but controls Type IV pili rather than

flagella, supports an intramolecular phosphorelay.[53] Phosphotransfer within ChpA

proceeds from ATP to the Hpt domains close to the HATPase_c domain to the receiver

domain to Hpt domains further from the HATPase_c domain. Thus, a plausible

hypothesis is that a distinct *Class* of CheW-like domains is required to support

intramolecular phosphorelays in CheA proteins with multiple Hpt domains and a

receiver domain. Perhaps the receiver domain must be held in a particular orientation to

be accessible to multiple Hpt domains.

Class 5 was the C-terminal CheW-like domain in CheA proteins with dual CheW-

like domains. *A. caulinodans* CheA,[50] *Azospirillum brasilense* CheA1,[54] and

Rhodospirillum centenum CheA1[55] have been explored experimentally and all belong to

the CheA.V *Architecture* with two CheW-like domains. In each case, the C-terminal

receiver domain is functionally important. However, the only experimental investigation

of a *Class* 5 CheW-like domain of which we are aware was in *A. caulinodans* CheA,

where removal of the C-terminal CheW-like domain diminishes chemotaxis but does not

reveal the functional role of the *Class* 5 domain.[50] *Class* 5 CheW-like domains were

substantially different from experimentally characterized CheW-like domains. Therefore,

because investigation of *Class* 5 is especially likely to reveal new information, *Class* 5

arguably represents the highest priority for future research on CheW-like domains.

Class 6 was a *Type* (CheW.IA) encompassing ~20% of CheW proteins that

appeared different from most CheW proteins (*Class* 1) in some analyses and similar in

others. Method-dependent classification suggests that *Class* 6 is unlikely to represent a

high priority for investigation.

4. 3 | Generally applicable methods to analyze protein domains as a function of

architectural context

We have described a workflow (Figure 1) to analyze variation between

sequences of a single protein domain as a function of architectural context within

proteins. To ensure a robust workflow, we tested various approaches to optimize our

method, including using multiple different amino acid substitution models and

multidimensional scaling techniques to estimate and visualize relevant domain

sequence subpopulations. Furthermore, the similarity of our partition results using these

different models was compared using ARI and NMI analyses. We first applied a basic *k*-

medoid clustering step to quantify the relative homogeneity of a given domain sequence

set. This allowed us to identify constituent strata for representation in a more complete

sequence analysis utilizing a random stratified sub-sampling scheme. We demonstrated

our approach using CheW-like domains as a general case study. The results were

largely independent of methods chosen, suggesting robust conclusions. Grouping

sequence *Types* of CheW-like domains into five or six major *Classes* led to a variety of

useful observations, insights, hypotheses, and suggested areas of future experimental

investigation.

It might appear challenging to apply our approach to domains found in many

distinct protein architectures. However, protein architecture distributions tend to have a

long tail, with only a few organizations representing the majority of the cases. Here, 16

out of 111 architectures in Pfam that contained CheW-like domains incorporated ~94%

of CheW-like domains. Of course, individual "rare" architectures of interest could always

be manually included with minimal difficulty. Another concern could involve the number

of sequences to analyze. Our case study involved analysis of ~8500 sequences.

Significantly more or less sequences may be available for other domains. However, our

approach should still be practical. The number of starting sequences to be analyzed can

be managed by choice of an appropriate input data set. We chose (and recommend)

the use of Representative Proteome sets as a starting point, where the sequence space

size and granularity can be modified by changing the co-membership threshold of the

relevant RP setcutoff (we used 35% as our threshold, but RP sets generated using

15%, 55%, and 75% are also available).[14] The subsampling used in our analysis (see

Figure 4) is a standard strategy to deal with impractically large datasets.

Finally, we developed and have made available (see Methods) SimpLogo, a new

means of visualizing amino acid sequence conservation patterns across many subtypes

of the same domain and for sequences of arbitrary length. Comparison of the

SimpLogos in Figure 5 and traditional sequence logos in Figure S26 quickly reveal the

utility of SimpLogos. Differences in gap patterns are immediately obvious. Identifying

differences in amino acid composition at individual positions is more time consuming,

but feasible. Traditional sequence logos are useful to communicate information about

reasonably conserved positions but of little utility for most other positions, whereas

SimpLogo provides useable information about all aligned positions within a given

sequence.

ACKNOWLEDGEMENTS

We thank Emily N. Kennedy and Sarah A. Barr for their insightful input and

helpful discussions, including Sarah's idea that receiver domains of CheV proteins

might serve as reservoirs rather than sinks for phosphoryl groups.

This work was funded by National Institutes of Health grant GM050860 to Robert

B. Bourret. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institute of General Medical

Sciences or the National Institutes of Health.

REFERENCES

1. Mistry J, Chuguransky S, Williams L, et al. Pfam: The protein families database in 2021. *Nucleic Acids Res.* 2020;49(D1):D412-D419.

2. Koretke KK, Lupas AN, Warren PV, Rosenberg M, Brown JR. Evolution of two-component signal transduction. *Mol Biol Evol.* 2000;17(12):1956-1970.

3. Pao GM, Saier MH, Jr. Response regulators of bacterial signal transduction systems: selective domain shuffling during evolution. *J Mol Evol.* 1995;40(2):136-154.

4. Gao R, Bouillet S, Stock AM. Structural basis of response regulator function. *Annu Rev Microbiol.* 2019;73:175-197.

5. Immormino RM, Silversmith RE, Bourret RB. A variable active site residue Iinfluences the kinetics of response regulator phosphorylation and dephosphorylation. *Biochemistry.* 2016;55(39):5595-5609.

6. Page SC, Immormino RM, Miller TH, Bourret RB. Experimental analysis of functional bariation within protein families: Receiver domain autodephosphorylation kinetics. *J Bacteriol.* 2016;198(18):2483-2493.

7. Straughn PB, Vass LR, Yuan C, Kennedy EN, Foster CA, Bourret RB. Modulation of response regulator CheY reaction kinetics by two variable residues that affect conformation. *J Bacteriol.* 2020;202(15).

8. Thomas SA, Immormino RM, Bourret RB, Silversmith RE. Nonconserved active site residues modulate CheY autophosphorylation kinetics and phosphodonor preference. *Biochemistry.* 2013;52(13):2262-2273.

9. Bi S, Sourjik V. Stimulus sensing and signal processing in bacterial chemotaxis. *Curr*
Opin Microbiol. 2018;45:22-29.

10. Abedrabbo S, Castellon J, Collins KD, Johnson KS, Ottemann KM. Cooperation of
two distinct coupling proteins creates chemosensory network connections. *Proc Natl*
Acad Sci U S A. 2017;114(11):2970-2975.

11. Huang Z, Pan X, Xu N, Guo M. Bacterial chemotaxis coupling protein: Structure,
function and diversity. *Microbiol Res.* 2019;219:40-48.

12. Alexandre G, Zhulin IB. Different evolutionary constraints on chemotaxis proteins
CheW and CheY revealed by heterologous expression studies and protein sequence
analysis. *J Bacteriol.* 2003;185(2):544-552.

13. Gumerov VM, Andrianova EP, Zhulin IB. Diversity of bacterial chemosensory
systems. *Curr Opin Microbiol.* 2021;61:42-50.

14. Chen C, Natale DA, Finn RD, et al. Representative proteomes: A stable, scalable
and unbiased proteome set for sequence analysis and functional annotation. *PLoS*
One. 2011;6(4):e18910-e18910.

15. Consortium TU. UniProt: The universal protein knowledgebase in 2021. *Nucleic*
Acids Res. 2020;49(D1):D480-D489.

16. Eddy SR. Accelerated Profile HMM Searches. *PLOS Comp Biol.*
2011;7(10):e1002195.

17. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics.*
2009;25(15):1972-1973.

18. Pelé J, Bécu J-M, Abdi H, Chabbert M. Bios2mds: An R package for comparing

orthologous protein families by metric multidimensional scaling. *BMC Bioinform.*

2012;13(1):133.

19. Dayhoff M, Schwartz R, Orcutt B. A model of evolutionary change in proteins. *Atlas*

of Protein Sequence and Structure. 1978;5:345-352.

20. Gonnet GH, Cohen MA, Benner SA. Exhaustive matching of the entire protein

sequence database. *Science.* 1992;256(5062):1443-1445.

21. Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. *Proc*

Natl Acad Sci U S A. 1992;89(22):10915-10919.

22. Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data matrices

from protein sequences. *Comput Appl Biosci.* 1992;8(3):275-282.

23. *cluster: Cluster analysis basics and extensions* [computer program]. Version

2.1.22012.

24. *fpc: Flexible procedures for clustering* [computer program]. Version 2.2-92015.

25. Hubert L, Arabie P. Comparing partitions. *J Classif.* 1985;2(1):193-218.

26. Steinley D. Properties of the Hubert-Arable Adjusted Rand Index. *Psychol Methods.*

2004;9(3):386-396.

27. Cover TM, Thomas JA. *Elements of information theory.* Wiley-Interscience; 2006.

28. Strehl A, Ghosh J. Cluster ensembles - A knowledge reuse framework for combining

multiple partitions. *J Mach Learn Res.* 2003;3(null):583–617.

29. *vegan: Community Ecology Package. R package version 2.5-6. 2019* [computer

program]. Version 2.5-72020.

30. Wickham H. *ggplot2: Elegant graphics for data analysis.* 3rd ed: Springer-Verlag
New York; 2016.

31. *Biostrings: Efficient manipulation of biological strings* [computer program]. Version
2.60.12015.

32. Ou J, Wolfe SA, Brodsky MH, Zhu LJ. motifStack for the analysis of transcription
factor binding site evolution. *Nat Methods.* 2018;15(1):8-9.

33. Dey KK, Xie D, Stephens M. A new sequence logo plot to highlight enrichment and
depletion. *BMC Bioinform.* 2018;19(1):473.

34. Li X, Fleetwood AD, Bayas C, et al. The 3.2 Å resolution structure of a
receptor:CheA:CheW signaling complex defines overlapping binding sites and key
residue interactions within bacterial chemosensory arrays. *Biochemistry.*
2013;52(22):3852-3865.

35. Bilwes AM, Alex LA, Crane BR, Simon MI. Structure of CheA, a signal-transducing
histidine kinase. *Cell.* 1999;96(1):131-141.

36. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. The Phyre2 web portal
for protein modeling, prediction and analysis. *Nat Protoc.* 2015;10(6):845-858.

37. *PyMOL* [computer program]. Version 2.3.3. http://www.pymol.org/pymol2020.

38. Rousseeuw PJ. Silhouettes: A graphical aid to the interpretation and validation of
cluster analysis. *J Comput Appl Math.* 1987;20:53-65.

39. Tzeng J, Lu HH-S, Li W-H. Multidimensional scaling for large genomic data sets.
BMC Bioinform. 2008;9(1):179.

40. Schneider TD, Stephens RM. Sequence logos: A new way to display consensus
sequences. *Nucleic Acids Res.* 1990;18(20):6097-6100.

41. Boukhvalova M, VanBruggen R, Stewart RC. CheA kinase and chemoreceptor

interaction surfaces on CheW. *J Biol Chem.* 2002;277(26):23596-23603.

42. Boukhvalova MS, Dahlquist FW, Stewart RC. CheW binding interactions with CheA

and Tar: Importance for chemotaxis signaling in *Escherichia coli. J Biol Chem.*

2002;277(25):22251-22259.

43. Griswold IJ, Zhou H, Matison M, et al. The solution structure and interactions of

CheW from *Thermotoga maritima. Nat Struct.* 2002;9(2):121-125.

44. Piñas GE, Frank V, Vaknin A, Parkinson JS. The source of high signal cooperativity

in bacterial chemosensory arrays. *Proc Natl Acad Sci U S A.* 2016;113(12):3335-

3340.

45. Liu JD, Parkinson JS. Genetic evidence for interaction between the CheW and Tsr

proteins during chemoreceptor signaling by *Escherichia coli. J Bacteriol.*

1991;173(16):4941-4951.

46. Porter SL, Wadhams GH, Armitage JP. *Rhodobacter sphaeroides*: complexity in

chemotactic signalling. *Trends Microbiol.* 2008;16(6):251-260.

47. Alexander RP, Lowenthal AC, Harshey RM, Ottemann KM. CheV: CheW-like

coupling proteins at the core of the chemotaxis signaling network. *Trends Microbiol.*

2010;18(11):494-503.

48. Ortega DR, Zhulin IB. Evolutionary genomics suggests that CheV Is an additional

adaptor for accommodating specific chemoreceptors within the chemotaxis signaling

complex. *PLOS Comp Biol.* 2016;12(2):e1004723.

49. Karatan E, Saulmon MM, Bunn MW, Ordal GW. Phosphorylation of the response

regulator CheV is required for adaptation to attractants during *Bacillus subtilis*

chemotaxis. *J Biol Chem.* 2001;276(47):43618-43626.

50. Kennedy EN, Barr SA, Liu X, et al. *Azorhizobium caulinodans* chemotaxis is

controlled by an unusual phosphorelay network. *J Bacteriol.* 2021:JB0052721.

51. Briegel A, Ortega DR, Mann P, Kjaer A, Ringgaard S, Jensen GJ. Chemotaxis

cluster 1 proteins form cytoplasmic arrays in *Vibrio cholerae* and are stabilized by a

double signaling domain receptor DosM. *Proc Natl Acad Sci U S A.*

2016;113(37):10412-10417.

52. Wuichet K, Zhulin IB. Origins and diversification of a complex signal transduction

system in prokaryotes. *Sci Signal.* 2010;3(128):ra50-ra50.

53. Silversmith RE, Wang B, Fulcher NB, Wolfgang MC, Bourret RB. Phosphoryl group

flow within the *Pseudomonas aeruginosa* Pil-Chp chemosensory system: Differential

function of the eight phosphotransferase and three receiver domains. *J Biol Chem.*

2016;291(34):17677-17691.

54. Bible AN, Stephens BB, Ortega DR, Xie Z, Alexandre G. Function of a chemotaxis-

like signal transduction pathway in modulating motility, cell clumping, and cell length

in the alphaproteobacterium *Azospirillum brasilense*. *J Bacteriol.* 2008;190(19):6365-

6375.

55. Jiang ZY, Gest H, Bauer CE. Chemosensory and photosensory perception in purple

photosynthetic bacteria utilize common signal transduction components. *J Bacteriol.*

1997;179(18):5720-5727.

1060

Table 1. Distribution of CheW-like domain *Context*s across *k*-medoid *Clusters*[a]

Architectural Context		C	E	H	I	L	M	R	P	N	O	Q	S	T	U	A	B	D	F	G	J	K	# Seqs
					1				2			3							4				
CheA-lineage	CheA.I	444	127	64	257	170																	1062
	CheA.II	140	312	3	2	8	167																632
	CheA.III	75	104	18	18	26	29	8								1							279
	CheA.IV	74	79	48	6	22		4															233
	CheA.V.1		4	88																			92
	CheA.V.2	2	3							89													94
	CheA.VI.1	10	3	73	2																		88
	CheA.VI.2	2	3		1					78						1	2	1					88
	CheA.VII	3	8		1			68															80
	CheA.VIII	65																					65
	CheA.IX	14	8	26	5	3	1																57
	CheA.X	5			1	50																	56
	CheA.XI	29	7	2	10	3	1																52
	CheA.XII	23	22	1	1			1															48
CheV-lineage	CheV.I	4			1			1								50	11	1	4	455	290		817
CheW-lineage	CheW.I	20	4	5	11	3		2		189	157	135	117	115	92	1093	979	605	485	5	7	234	4258
	CheW.II.1	3		2	1						3	1				8	11	20	5				54
	CheW.II.2	2					1			2	1				1	12		32	2			1	54
	CheW.III.1	1								2	3				1	62	21	35	13			2	140
	CheW.III.2	1				1					3			1	1	107	4	13	8				139
	CheW.III.3	1									6					91	17	12	13				140
# Seqs		918	684	330	318	235	197	135	168	193	173	136	117	116	95	1425	1045	719	530	460	297	237	8528

1062

1063 [a]*Clusters* and *Groups* taken from analysis shown in Figure 3. Table entries show the
1064 number of sequences of each architectural *Context* found within each *Cluster*.
1065 *Clusters* are named in order of decreasing size.
1066

 FIGURE LEGENDS

FIGURE 1. Hierarchy of classification terms and sorting workflow. The initial steps

were derived purely from protein *Architectures*, which may contain CheW-like domains

in more than one *Context*. The number of *Contexts* defined (dashed arrow) the number

of *Clusters* into which sequences of CheW-like domains were initially sorted. However,

sequences in a given *Context* sorted into different *Clusters* and vice versa. The *Clusters*

sorted into larger *Groups* of sequences. Comparison of architectural *Contexts* with

sequence *Groups* revealed that all but one *Context* included *Clusters* within a single

Group. One *Context* contained *Clusters* that spanned three *Groups* and so was split into

three *Types*, whereas all other *Contexts* became a single *Type*. Minor *Clusters* of

sequences associated with each *Context* were discarded from further analysis during

the transition to *Types*. The *Types* sorted into larger *Classes* of sequences associated

with one or more architectural *Contexts*. Figure 5 illustrates the final *Types* and *Classes*

but was not part of the classification process. See text and indicated Figures or Table

for details.

FIGURE 2. Schematic of the most abundant CheW-containing protein

***Architectures* seen in nature.** Sixteen distinct architectural variants containing the

CheW-like domain were identified within the Pfam database (≥40 sequences each,

ordered by decreasing abundance within each shaded group). Sequences for their

corresponding CheW-like domains were extracted and initially classified according to

the schematic. For parent proteins containing multiple CheW-like domains, CheW-like

domains were initially classified according to the position (*Context*) of the domain,

resulting in one sequence entry per CheW-like domain. Representative diagrams are

shown for each *Architecture* (from left to right, N- to C-terminal). Pfam domain IDs

included are as follows: CheW-like (PF01584); H-kinase dimerization (PF02895); HPt

(PF01627); Response regulatory (PF00072); CheY-binding (PF09078); HATPase_c

(PF02518); P2 (PF07194). Individual schematics are not necessarily drawn to scale.

FIGURE 3. **Cluster analysis and multidimensional scaling of representative**

sequences of CheW-like domains reveals *Groups*. Multidimensional scaling plots

showing representative medoids from each of the 21 *Clusters*, generated using

dissimilarity matrix calculated with the GONNET substitution model. Each clustering

attempt reconstituted roughly the same data structure, consisting of three main *Groups*

(numbered 1, 3 and 4) of *Clusters* along with a weaker, more intermediate *Group*

(numbered 2) intersecting *Groups* 1 and 3. While the colors and specific *Cluster*

designations may shift between analyses (overall *Cluster* sizes were slightly altered

depending on the substitution matrix used, affecting *Cluster* names), the actual

partitioning of CheW sequences was highly consistent, regardless of the substitution

model used (see Figure S1).

FIGURE 4. Metric multidimensional scaling plots showing 25 randomly

subsampled sequences from each of the 23 CheW-like domain *Types*. The top

three dimensions (Dim1, Dim2, Dim3) were included for visualization purposes. Analysis

was performed on dissimilarities calculated using the GONNET substitution model. The

location of *Types* belonging to each of six *Classes* are indicated. Additional sampling

iterations are shown in Figures S2-S4. Sampling iterations with other substitution

models are shown in Figures S5-S13.

FIGURE 5. **SimpLogo representation of major CheW-like domain *Types*.** Rows

correspond to domain *Types*. Sequences were aligned to the CheW-like domain HMM

(Pfam ID PF01584) and separated by *Type*. Residues with physicochemical similarities

were merged and assigned colors. Idealized colors (legend) were weighted by residue

frequency and mixed to produce a colored box at each position. Heights were scaled by

gap percentage. To differentiate between secondary colors and a near-equal mixing of

primary colors, colored strips were included on the top and bottom of each box,

indicating first and second most frequent residue types respectively. *Types* were

organized by *Class* in the figure. Position-wise information content (in bits) was

calculated for each alignment (top). Numbering is for *E. coli* CheW, framing our

observations with reference to the well-studied standalone protein.

1123

1124

FIGURE 1. Hierarchy of classification terms and sorting workflow. The initial steps
were derived purely from protein *Architectures*, which may contain CheW-like domains
in more than one *Context*. The number of *Contexts* defined (dashed arrow) the number
of *Clusters* into which sequences of CheW-like domains were initially sorted. However,
sequences in a given *Context* sorted into different *Clusters* and vice versa. The *Clusters*
sorted into larger *Groups* of sequences. Comparison of architectural *Contexts* with
sequence *Groups* revealed that all but one *Context* included *Clusters* within a single
Group. One *Context* contained *Clusters* that spanned three *Groups* and so was split into
three *Types*, whereas all other *Contexts* became a single *Type*. Minor *Clusters* of
sequences associated with each *Context* were discarded from further analysis during
the transition to *Types*. The *Types* sorted into larger *Classes* of sequences associated
with one or more architectural *Contexts*. Figure 5 illustrates the final *Types* and *Classes*
but was not part of the classification process. See text and indicated Figures or Table
for details.
1139

1140

FIGURE 2. Schematic of the most abundant CheW-containing protein
Architectures **seen in nature.** Sixteen distinct architectural variants containing the
CheW-like domain were identified within the Pfam database (≥40 sequences each,
ordered by decreasing abundance within each shaded group). Sequences for their
corresponding CheW-like domains were extracted and initially classified according to
the schematic. For parent proteins containing multiple CheW-like domains, CheW-like
domains were initially classified according to the position (*Context*) of the domain,
resulting in one sequence entry per CheW-like domain. Representative diagrams are
shown for each *Architecture* (from left to right, N- to C-terminal). Pfam domain IDs
included are as follows: CheW-like (PF01584); H-kinase dimerization (PF02895); HPt
(PF01627); Response regulatory (PF00072); CheY-binding (PF09078); HATPase_c
(PF02518); P2 (PF07194). Individual schematics are not necessarily drawn to scale.

1153

1154

FIGURE 3. **Cluster analysis and multidimensional scaling of representative**
sequences of CheW-like domains reveals _Groups_. Multidimensional scaling plots
showing representative medoids from each of the 21 _Clusters_, generated using
dissimilarity matrix calculated with the GONNET substitution model. Each clustering
attempt reconstituted roughly the same data structure, consisting of three main _Groups_
(numbered 1, 3 and 4) of _Clusters_ along with a weaker, more intermediate _Group_
(numbered 2) intersecting _Groups_ 1 and 3. While the colors and specific _Cluster_
designations may shift between analyses (overall _Cluster_ sizes were slightly altered
depending on the substitution matrix used, affecting _Cluster_ names), the actual
partitioning of CheW sequences was highly consistent, regardless of the substitution
model used (see Figure S1).
1166

FIGURE 4. **Metric multidimensional scaling plots showing 25 randomly subsampled sequences from each of the 23 CheW-like domain *Types*.** The top three dimensions (Dim1, Dim2, Dim3) were included for visualization purposes. Analysis was performed on dissimilarities calculated using the GONNET substitution model. The location of *Types* belonging to each of six *Classes* are indicated. Additional sampling iterations are shown in Figures S2-S4. Sampling iterations with other substitution models are shown in Figures S5-S13.

FIGURE 5. **SimpLogo representation of major CheW-like domain *Types*.** Rows correspond to domain *Types*.
Sequences were aligned to the CheW-like domain HMM (Pfam ID PF01584) and separated by *Type*. Residues with
physicochemical similarities were merged and assigned colors. Idealized colors (legend) were weighted by residue
frequency and mixed to produce a colored box at each position. Heights were scaled by gap percentage. To differentiate
between secondary colors and a near-equal mixing of primary colors, colored strips were included on the top and bottom
of each box, indicating first and second most frequent residue types respectively. *Types* were organized by *Class* in the
figure. Position-wise information content (in bits) was calculated for each alignment (top). Numbering is for *E. coli* CheW,
framing our observations with reference to the well-studied standalone protein.