Pupil size as a robust marker of attentional capture by nicotine-related stimuli in smokers

Elvio Blini[1] and Marco Zorzi[1,2]

1.   Department of General Psychology, University of Padova, Italy

2.   IRCCS San Camillo Hospital, Venice-Lido, Italy

Correspondence:

Elvio Blini

Department of General Psychology, University of Padova

via Venezia 12, 35131 Padova (Italy)

Tel: +39 (0)49 827 6149

Email: elvio.blini (at) unipd.it

Keywords: Attentional capture; Pupil dilation; Autonomic response; Smokers; Reward; Motivation; Spatial attention;

Abstract

Spatial attention can be magnetically attracted by behaviorally salient stimuli. This phenomenon occasionally conflicts with behavioral goals, leading to maladaptive consequences, as in the case of behavioral addictions, in which attentional capture has been often described and linked with clinically meaningful variables, such as craving level or dependence intensity. Here, we sought to probe the markers of attentional priority in smokers through eye-tracking measures, by leveraging the established link between eye movements and spatial attention. We were interested, in particular, in potential markers related to pupil dilation, because pupil size reflects a range of autonomic, affective, and cognitive/attentional reactions to behaviorally significant stimuli and is a robust marker of appetitive and aversive learning. We found that, indeed, the pattern of pupil dilation and constriction to nicotine-related visual stimuli could reliably predict, in crossvalidated logistic regression, the smoking status of female and male smokers better than more traditional proxy measures based on response times or eye position. Pupil dilation was more sensitive at lower nicotine dependence levels, supporting previous claims that the incentive strength of conditioned stimuli gradually decreases and leaves space for a more automated, habit-driven processing mode. Additionally, it was more sensitive with increased abstinence time, suggesting that it may reflect craving levels. We conclude that pupillometry can provide a robust marker for attentional priority computation and provide useful indications regarding motivational states and the individual attitudes toward conditioned stimuli.

1. Introduction

Intrinsically rewarding stimuli strongly drive spatial attention (Anderson et al., 2011; Awh et al., 2012; Chelazzi et al., 2013; Hickey et al., 2010). Although motivation has been scarcely considered in cognitive models of attention, animal models aside, it has recently been integrated in the wider concept of priority maps (Bisley & Mirpour, 2019; Fecteau & Munoz, 2006; Sprague & Serences, 2013). Priority maps do represent the physical, low-level properties of stimuli but, the higher the hierarchy level, the more maps start to encode behavioral relevance beyond those features (Bisley & Mirpour, 2019; Fecteau & Munoz, 2006). Traces of behavioral salience can be found in the brain as early as the superior colliculus (White, Berg, et al., 2017; White, Kan, et al., 2017) and the primary visual cortex (Li, 2002), although still confined to the physical

features that they encode (Bisley & Mirpour, 2019). Recent evidence highlights how, similarly, grid cells in the entorhinal cortex do not rigidly and objectively encode spatial locations, but rather incorporate learned reward positions (Boccara et al., 2019; Butler et al., 2019). Overall, the concept of priority maps helped moving beyond classic dichotomies such as bottom-up vs. top-down attentional processing (Awh et al., 2012). Value-driven attentional capture (Anderson et al., 2011, 2014; Bourgeois et al., 2016; Camara et al., 2013), on the one hand, and statistical learning or selection history (Duncan & Theeuwes, 2020; Failing & Theeuwes, 2018; Jiang et al., 2015), on the other hand, appear to escape these labels, all while representing powerful determinants of attentional and exploration biases. This is arguably efficient and evolutionarily convenient, in that both attention and reward serve to filter the most relevant sensory information in order to optimize cognitive and behavioral adaptation (Driver, 2001; Manohar et al., 2015; Maunsell, 2004). As such, they are strictly intertwined processes, to the point that they can easily be confounded (Anderson et al., 2011; Maunsell, 2004; Raymond & O'Brien, 2009). Yet, studying them, and their interaction, is not only relevant for understanding human cognition in general, but also provides a framework to study instances in which value conflicts with behavioral goals, leading to maladaptive consequences. An example is the case of behavioral addictions.

Addiction disorders are multifaceted conditions. Among their many features, dysfunctions to the processing of rewards have been described, and consist of an increased sensitivity to selected reinforcers at the expense of alternative ones (Goldstein & Volkow, 2002; Noël et al., 2013). These stimuli gain high attentional priority in everyday life situations, creating a vicious loop believed to help maintaining addictive behaviors (Volkow et al., 2010), and to the point of hampering inhibitory and monitoring processes (Blini et al., 2020; Cox et al., 2006; Field et al., 2007; Field & Cox, 2008; Goldstein & Volkow, 2011; Noël et al., 2013). Not surprisingly, interventions aimed at modifying attentional biases have been proposed. Their main goal is to divert attention away from disorder-relevant stimuli, especially when these are competing with other, neutral ones, and this is chiefly achieved by leveraging on statistical learning procedures. However, the effectiveness of these interventions remains to date, inconclusive (Cox et al., 2014; Mogoaşe et al., 2014). The results are seemingly inconsistent, as are studies attempting to relate the degree of attentional bias to the chances of relapse (Christiansen et al., 2015). Therefore, the causal role of attentional bias in addiction is still debated, while attentional bias itself remains a valid marker of the underlying motivational state (Christiansen et al., 2015),

which naturally fluctuates over time. Even if considered a byproduct, thus, it can still represent a useful proxy for clinically meaningful information such as current craving level (Field et al., 2014), current concerns about behavioral addiction, or the broad affective reaction to these meaningful stimuli. As such, it is important to deepen our knowledge and characterization of attentional biomarkers, in that these measures could be useful and objective indices for longitudinal and interventional studies.

An established paradigm to study attentional capture is the Dot-Probe Task (DPT; Chanon et al., 2010; Ehrman et al., 2002; Waters et al., 2003). In the case of smokers, two pictures are shown on either side of a computer screen, one depicting a Nicotine-Related Stimulus (NRS), one a physically matched neutral object. After a variable delay, and along the tradition of Posner-like cueing tasks (Posner, 1980), the pictures disappear, and one to-be-discriminated target is presented on either side. Differently from non-smokers or former smokers, active smokers present a response times advantage whenever the target is presented on the same side of the NRS, showing that spatial attention is magnetically attracted by the cue (Ehrman et al., 2002). This effect appears to go beyond the increased familiarity of smokers with NRS (Chanon et al., 2010), and it is highly dependent on Stimulus-Onset Asynchronies (Della Libera et al., 2019). Crucially, it is stronger for direct (i.e., overt) measures of attention (Field et al., 2009). Electroencephalographic studies, for example, have shown that masked NRS elicit a larger N2pc component (that is, an index of lateralized spatial attention) than masked control stimuli (Harris et al., 2018), suggesting that the former, at least in conditions of reduced perceptibility, capture visual attention more strongly very early (i.e., around 250 ms). Another important proxy of overt attentional capture is eye movements, because eye movements and spatial attention are profoundly linked one another (Blini, Pitteri, & Zorzi, 2019; Casarotti et al., 2012; Rizzolatti et al., 1987). When stimuli are given in the visual modality, the oculomotor system codes and is biased by their value (Camara et al., 2013; Manohar & Husain, 2015; Muhammed et al., 2020; Takikawa et al., 2002; Theeuwes & Belopolsky, 2012). This is also seen, in smokers administered with the DPT, as longer time spent fixating NRS and/or first fixations more often directed toward these stimuli (Field et al., 2004; Mogg et al., 2003, 2005).

Pupil dilation, in particular, is a more special measure among those that can be inferred by recording the eyes of the participants. Pupil diameter changes first and foremost following changes in the amount of light entering the eyes (see, e.g., Binda & Murray, 2015). However, pupil dilation has been associated with phasic activation of the locus coeruleus (Aston-Jones & Cohen, 2005), a major noradrenergic hub involved in the integration of

the attentional systems in the brain, as well as in balancing bottom-up and top-down aspects of perception (Reynaud et al., 2021). Pupil dilation reflects, beyond that, the activation of the autonomic system, and as such it ideally probes: affective and emotional responses to visual stimuli (Bogdanova et al., 2022; Dureux et al., 2021); the amount of cognitive effort deployed for a task (e.g., working memory load, Lisi et al., 2015; Mathôt, 2018); and, perhaps in light of the strict dialogue between the locus coeruleus and limbic areas subserving motivation, the subjective value of a given stimulus (Muhammed et al., 2016; Pietrock et al., 2019). Accordingly, a recent meta-analysis concluded that pupil dilation is a robust marker of Pavlovian-like conditioning (Finke et al., 2021). Despite these premises, however, studies attempting to characterize attentional capture by means of pupillometry are still scarce (but see Chae et al., 2008). The current study aims to fill this gap, by assessing: i) whether pupil can, in a fast event-related design, reliably assess attentional capture for NRS; ii) how this measure compares with more common ones (e.g., as obtained from a DPT task); iii) whether it can be used for a reliable classification of smokers; iv) whether it correlates with either smoking intensity or craving urges.

2. Methods

All materials, raw data, and analysis scripts for this study are available at the following link (referred to as Supplementary materials): https://osf.io/6r5ch/

2.1 Participants

In this work, we planned to acquire multiple measures of attentional and autonomic engagement with Nicotine-Related Stimuli (NRS) in smokers vs. controls. Instead of calibrating the recruitment plan on each individual measure, we reasoned that a multivariate approach – attempting to classify the group on the basis of these measures – would be a more appropriate and powerful approach. Thus, we conducted an a priori power analysis for a one-sided binomial test (i.e., testing that classification performance is superior to chance level). We assumed the minimum effect size of interest (and practical relevance) to be a classification accuracy of 70%. With a type 1 error rate set to alpha= 0.05, 80% statistical power for this design is reached at N= 40 (**Fig. 1**).

Power analysis was conducted through simulations, and the relative scripts are available in the Supplementary Materials. Note that, at N= 40, a classification accuracy of 65% would be significantly superior to chance (p= 0.04) according to this test; however, this is to be considered the underlying, "true" effect size, which does not invariantly lead to the same classification accuracy due to random fluctuations, hence the concept of statistical power of a test. Therefore, we planned to enroll a minimum of 20 smokers and 20 matched controls. The assumption of our elective multivariate classifier was that all participants must have all the pre-identified behavioral predictors, without missing data. Based on our set of data quality (explained below), this led to the recruitment of 51 participants in total.

The sample was composed of 28 smokers (age: 24.04 ± 2.69 years; 18 were women, 4 were left handed) and 23 non-smokers (age: 23.13 ± 2.75 years; 16 were women, 1 was left handed). All participants were recruited from students of the University of Padova (Italy). Inclusion criteria were normal (or corrected-to-normal) vision and no history of neurological or psychiatric diseases. Occasional (i.e., non-daily) smokers were excluded and all smokers in the final sample smoked a self-reported mean of 8.1 cigarettes a day and for an average duration of 5 years. Most of them were light smokers, however, as seen by low average scores (1.96 ± 2.1) at the Fagerström test for Nicotine Dependence (FTND; Heatherton et al., 1991). Because nicotine can considerably reduce pupil size (Wardhani et al., 2020), and because we sought to evaluate the tasks in conditions of relative craving, they were all asked to refrain from smoking for at least 90 minutes prior to the experiment (mean: 7.5 hours; indeed, half of them last smoked the evening before the experiment). In addition to the FTND, smokers were asked to fill in the Severity of Dependence Scale (SDS; Gossop et al., 1992) and the Questionnaire of Smoking Urges (QSU-brief; Cox et al., 2001), to assess self-reported craving. The study was approved by the Ethics Committee of the University of Padova (protocol n° 3568). It was carried out during a period of restrictions due to the covid-19 epidemic and as such all the necessary sanitary precautions were taken.

2.2 Materials and Methods

Participants were tested in a dimly lit, quiet room, their head comfortably restrained by a chinrest. They faced a remote infrared-based eye-tracker (TOBII™ Spectrum), with an embedded 24 inches monitor, at a distance

of approximately 57 cm. The session started with a 9-points calibration of the eye-tracker, which was then set to monitor participants' gaze continuously at a 600 Hz sampling rate. The open-source software OpenSesame (Mathôt et al., 2012) was used to display experimental stimuli on the screen and record the subjects' responses. Participants provided responses by means of keyboard presses (on a standard QWERTY keyboard) using the index and middle fingers of their dominant hand. All participants performed two tasks, in a fixed order: Dot-Probe Task (DPT) and Passive Viewing (PV) task (**Fig. 1**).

Fig. 1. Left panel: structure and time course of the two experimental tasks, Dot-Probe task (DPT, upper row) and Passive Viewing (PV, bottom row). Note that stimuli depicted in the image are only examples, both tasks involved pictures of people smoking (see main text). Right panel: nominal power for our design depicted as a function of sample size and varying degrees of a priori effect sizes (i.e., classification accuracy to a binomial test).

DPT. The first task was meant to measure spontaneous (spatial) attentional capture by nicotine-related stimuli (NRS). Each trial started with a fixation cross appearing at the center of the screen for 500 ms. Then, two images (6° x 4.5°) were always presented concurrently on the left and right sides of the screen, about 2° away from fixation horizontally. One image depicted a Nicotine-Related Stimulus (NRS; e.g., a person smoking), and one a perceptually-matched control stimulus (e.g., a person with otherwise similar features, see the paragraph below). The side of NRS appearance (i.e., left, right) was equiprobable. We choose a relatively long

on-screen duration for these images (i.e., SOA, 1.5s). Previous studies have highlighted how attentional capture effects can unfold at different temporal windows, e.g. differently at 200 ms than 800 ms (Della Libera et al., 2019). Using only one, extra-long SOA is guaranteed to miss this fine-grained dissociation. However, our choice was motivated based on pilot testing and ultimately our stress on overt attentional measures (i.e., eye movements) over response times, which have been found to be more sensitive to craving (Field et al., 2009) although requiring an extended time window (Field et al., 2004; Mogg et al., 2003). After this long SOA, the target appeared either above the NRS or the control image (balanced). The target was a circle (1.5° diameter) with a gap either in its upper or lower part, and participants had to indicate the position of the gap through the corresponding arrow keys. The target remained on screen until the participants' response, up to a maximum of 1.5s. Participants were first administered 12 practice trials, later discarded from analyses, followed by 240 experimental trials. A break was foreseen halfway through the task in order to allow participants to rest.

PV. The second task was a passive viewing task of NRS or control images presented centrally on the screen (4.8° x 3.6°), and the main objective was to acquire indices of autonomic activation through patterns of pupil dilation and constriction. Each trial started with 1s foveal presentation of a scrambled version of the target image; then, each image was presented on screen for 3s. Pupil diameter was measured continuously in both these phases. In a minority of trials (30%), a visual cue (a gray dot, 1° diameter) followed the presentation of the target image. The aim was to encourage sustained attention to the images, in absence of an otherwise active behavioral task. The cue remained on screen until the participants' detection (response given with the spacebar) up to a maximum of 1.5s. During the inter-trial interval, we also presented scrambled images on screen, as to keep all sources of luminance constant. Participants were first administered 8 practice trials, later discarded from analyses, followed by 200 experimental trials. A break was foreseen halfway through the task in order to allow participants to rest.

Images selection. Images were drawn from the SmoCuDa database (Manoliu et al., 2020), a validated and well-described collection of smoking-related images. Following pilot testing, we ended up selecting 10 images all depicting people smoking (i.e., social stimuli). Each stimulus was paired, to the best of our possibilities, to a control image, matched for all the relevant features but the smoke-related content. All stimuli were transformed into grayscale images, and were processed in order to equate their mean luminance. Images were all originally 800 x 600 pixels large, and were rescaled to fit the needs of each task.

2.3 Data processing

DPT. We discarded practice trials and anticipations (responses faster than 100 ms, <0.005%). All participants performed well in this task (minimum accuracy: 80.4%), thus, there were no exclusions related to this measure of behavioral performance. Furthermore, because accuracy was on average very high (M= 98.4%, SD= 2.95%), with little variability, this measure was not further considered in the analyses. We analyzed, instead, the mean Response Times (RTs) for correct responses. We computed RTs for each Condition (NRS, control) and then subtracted the former from the latter so that positive values index enhanced attentional engagement with NRS (i.e., faster responses when probes appear over NRS).

We then moved to assess eye movements. We only retained X and Y positions of the gaze when data were available for both eyes, and took their average value. As a cautionary measure, we discarded the trials in which more than 40% of the data were missing (this includes blinks, artifacts, lost connection with the eye-tracker, etc.); furthermore, we established, prior to testing, to discard from the sample participants remaining with fewer than 50% of valid trials. This led to the exclusion of 6 participants from the main analysis (note: other participants, for a total of 11, were discarded for the same reason applied to the PV task). Missing trials for the remaining participants were very few (for smokers: 2.5% ± 5.2%; for controls: 5.2% ± 8.2%). The gaps in the remaining recordings were then linearly interpolated for both the X and Y axes. We focused the analyses on the 1500 ms window in which the cue was presented, prior to the target probe. We reconstructed, in this window, the pattern of eye fixations through an automated, velocity-based algorithm (von der Malsburg, 2015), and only considered fixation events lasting at least 40 ms. As a first measure, we computed, for each participant, the proportion of first fixations falling within the NRS region vs the control region. This variable was meant to index a very early, automatic capture of attention and eye movements by NRS. Then, we computed dwell time, for each trial, as the cumulative duration of fixations in each image Condition. This variable is also in line with previous studies, and was meant to index a more or less thorough visual scanning of the different image classes. Both these variables were transformed into bias scores by subtraction so that negative values reflect a bias towards control images, and positive towards NRS.

PV. We first discarded practice trials. Missed responses to the infrequent probe were very rare (0.002%), showing some engagement with the presented images. As a first measure, we computed the RTs to the probe

for NRS vs. control images, and then subtracted the former from the latter so that positive values index enhanced attentional engagement with NRS.

We then moved to assess the time course of pupil dilation, mostly capitalizing on preprocessing steps adopted in previous studies (Dureux et al., 2021). First, we took the average pupil diameter of the left and right eyes, but only when both signals were valid as flanked by the eyetracker. We focused our analyses on the baseline period (1s, scrambled images) and the 3s window in which images were presented. Implausible values for pupil diameter (<2 mm and > 7mm), although infrequent (<0.0002%), were omitted. For missing samples within each trial, we followed an interpolation strategy akin to that used for the DPT: we discarded trials in which more than 40% of the data were missing (this includes blinks, artifacts, lost connection with the eye-tracker, etc.), and linearly interpolated gaps in the remaining ones. Participants with fewer than 50% of valid trials (N= 11) were discarded from the main analyses. Missing trials for the remaining participants were few (for smokers: 8.4% ± 13.6%; for controls: 10% ± 13%). We applied a low-pass filter to the raw traces and ultimately down-sampled the data to 25 ms epochs by taking the median pupil diameter for each time bin. The average pupil diameter was similar in the two groups (4.7mm for smokers vs 4.56mm for controls, $t_{(37.66)}$= 0.462, p= 0.647). Note that the acute effects of nicotine intake are known to globally constrict the pupils (Wardhani et al., 2020), but we asked all participants to refrain from smoking prior to the experiment. At any rate, in order to better cope with individual (and group) differences we z-tranformed pupil diameter values separately for each participant (Dureux et al., 2021). This way, a value of 0 represents the subject-specific mean pupil diameter and, regardless of baseline values, scores represent dilation (positive values) or constriction (negative values) expressed as a fraction of the overall participant's variability. Finally, all series were realigned to the beginning of the target image phase (either NRS or control) via subtraction of the first sample in this epoch. We assessed the traces in full when analyses focused on the PV task alone. For the main analyses, however, exploiting a multivariate classifier, we decided to avoid the extremely high dimensionality of these data, which yields overfitting, and rather focused on a time window identified by a cluster-based permutation test. Cluster-based permutation test can handle very elegantly the problem of multiple comparisons in autocorrelated data, although this may come at the price of a more imprecise estimation of the temporal features of the reported effects (e.g., latency, see Sassenhagen & Draschkow, 2019); however, since pupil dilation is a much slower physiological signal with respect to, for example, the electric or magnetic

signals captured by EEG or MEG, we did not anticipate this parameter to be the most relevant. Traces from NRS vs control images were subtracted so that positive values indexed enhanced pupil dilation to NRS, and negative values a relative constriction (see **Fig. 2**); the values used as predictors were therefore the mean, cluster-wide differences in these curves for each participant.

2.4 Analyses

Analyses were performed with R 4.1.2 (The R Core Team, 2018). Each measure of interest was first analyzed individually by means of two-sample t-tests with Welch's correction for unequal variances (two-tailed). The main interest of this work, however, was to probe the generalizability of these measures and their ability to predict smoking status of out-of-the-box, new individuals. This was to probe the global capability of eye-tracking and autonomic measures to generalize and possibly represent a viable biomarker of smoking behavior. We started with computing general (logistic) linear models with a Best Subset Regression approach. All possible combinations of the predictors were tested in glms, and the overall best model was selected only based on the lowest Bayesian and Akaike Information Criteria, which mitigates the problem of multiple comparisons. This approach was used for the initial features selection step, and the best features were then probed in follow-up crossvalidated logistic regressions. We adopted a Leave-One-(Subject)-Out (LOO) crossvalidation setup, where just one participant was circularly included in the test set. Then, we computed sensitivity, specificity, and area under the curve of the classifier, together with an overall classification accuracy measure, which was submitted to a one-tailed binomial test.

Finally, we performed correlations between the variables of interest, obtained from the two tasks, the individual measures related to the intensity of smoking (e.g., number of cigarettes per day) or subjective craving (e.g., QSU-brief, abstinence duration), and the predictions of the classifier for each participant (i.e., the estimated probability to belong to one group or another). Correlation analyses should be taken with caution due to the small sample size.

3. Results

3.1 Individual results

DPT: Response Times. The difference in RTs when the probe appeared on the NRS, with respect to control images, did not differ between groups (smokers: 8.79 ± 12.18 ms; non-smokers: 2.3 ± 14.85 ms; $t_{(36.59)}$= 1.51, p= 0.139, two-tailed).

DPT: Dwell time. Dwell time to NRS vs. control images did not differ between groups ($t_{(28.89)}$= 0.485, p= 0.631, two-tailed).

DPT: proportion of first fixations. This index of early attentional capture did not differ between groups as well ($t_{(37.88)}$= 0.169, p= 0.867, two-tailed).

PV: RTs to the probe. RTs did not differ between the two groups when the probe appeared over NRS vs. control images (smokers: 0.66 ± 22.06 ms; non-smokers: -3.35 ± 23.65 ms; $t_{(37.82)}$= 0.55, p= 0.58, two-tailed).

PV: Pupil dilation. Results are depicted in **Fig. 2**. The changes in pupil diameter were dominated by a sustained, steep pattern of pupil dilation during the course of the trial. This can be ascribed to the presence of the infrequent (30% of the trials) probe, which could follow the presentation of images. This feature of the task prompted temporal predictions and expectations, on one hand, and motor preparation for the manual response to produce, which are both factors known to affect pupil's dilation (Akdoğan et al., 2016; Wang et al., 2016). The effect on pupil dilation was so large that an increased inter-trial interval would probably have been beneficial, in that pupil diameter return to baseline levels appears to occur still during the time window prior to target presentation, in which scrambled images are presented. In this regard, our choice to realign pupil dilation to the first sample of target presentation is meant to be conservative in that the observed differences would have been magnified by accounting for the whole baseline window. That said, this carry-over effect, and indeed the overall pupil dilation during the course of the trials, appears larger for non-smokers. We computed the temporal derivative of pupil diameter (i.e., Gain, see Dureux et al., 2021) to assess whether steepness differs between the two groups. Despite some trend suggesting steeper pupil dilation for non-smokers, there was no conclusive evidence in this regard ($F_{(1, 38)}$= 3.63, p= 0.064); there were no effects of Condition or interaction between Group and Condition either when assessing the Gain (all Fs< 0.07, all ps>

0.356). However, the interaction between Group and Condition unfolded with different time-windows during the trial. Cluster-based permutation test revealed a large temporal cluster (800-2850 ms) in which pupil dilation to NRS vs control images significantly differed between groups ($p<0.001$, 5000 permutations). We thus computed the mean, cluster-wide difference between pupil dilation to NRS vs control images for each participant between 800 and 2850 ms. In this window, smokers presented a pattern of overall constriction of the pupil (M= -0.025, SD= 0.075) which, when compared to the overall pattern of pupil dilation observed in non-smokers (M= 0.036, SD= 0.067) resulted significantly different ($t_{(37.58)}$= 2.7, p= 0.01, two-tailed). This variable was then used for classification (see below).

Fig. 2. Time course of pupil dilation during the baseline and target presentation windows (left panel). Pupil dilated steeply over time, likely due to the expectation of the appearance of a visual probe, and the relative motor preparation. Differences between NRS and perceptually matched images differed between the two groups (right panel: shadowed areas represent 95% confidence interval for the data). Cluster-based permutations highlighted, in a data-driven fashion, a rather extended window (800-2850 ms) in which smokers presented, on average, a pattern of pupil constriction with respect to non-smokers.

3.2 Multivariate classification

We started by assessing the most relevant features with a Best Subset Regression approach. BSR modeling selected the logistic model with only one predictor, pupil dilation, as the best compromise between complexity and fit to the data. The BIC of this model was 55.57 (AIC= 52.2), far better than the null model (BIC= 59.14, AIC= 57.45), but only marginally better than the model also including RTs to the DPT task (BIC= 57.5, AIC= 52.43). We therefore kept these two features for follow-up, crossvalidated logistic regressions.

The best model included both pupil dilation to NRS images and RTs: here, the crossvalidation accuracy reached 75%, with excellent sensitivity and specificity (both 75%), and AUC of 0.745. The performance was significantly superior to chance according to a one-tailed binomial test (p= 0.001, CI$_{95\%}$[58.8% – 87.3%]). The second best model only included pupil dilation as predictor of smoking status. The performance of the classifier in this case was good, with an accuracy of 65% (both sensitivity and specificity were 65%, AUC was 0.723). This performance was also significantly superior to chance according to a one-tailed binomial test (p= 0.04, CI$_{95\%}$[48.3% – 79.4%]). When compared against a chance level of 65%, the model including both predictors was not significantly better (p= 0.12). For comparison, the model only including RTs performed very poorly, with an accuracy in crossvalidation of 52.5%. Sensitivity was 55%, specificity was 50%, and the area under the curve (AUC) was 0.535.

Overall, these results suggest very good classification capability for measures of attentional capture and in particular the autonomic response conveyed by the pupil, which appears necessary and sufficient for classification of smoking status.

3.3 Explorative analyses and correlations

This part only reports a selection of the results, for brevity, although the results can be assessed in full in the supplementary materials. These analyses should be taken with caution in light of the small sample size considered.

Two measures from the DPT task were correlated with the amount of pupil dilation during the subsequent PV task. This is interesting because it suggests that the observed pattern of pupil constriction in smokers may be

partially related to behavioral measures that have been more classically associated with attentional capture as assessed by the DPT task. First, there was a negative correlation with dwell time (r= -0.42, $CI_{95\%}$[-0.65, -0.12], $t_{(38)}$= 2.83, p= 0.0073), which was of similar magnitude in the two groups. A more thorough visual exploration of NRS images was correlated with a more vigorous *constriction* of the pupil in the PV task. There was, on the other hand, a positive correlation between pupil dilation to NRS and the proportion of first fixations towards the NRS stimuli in the DPT task (r= 0.29, $CI_{95\%}$[-0.02, 0.55], $t_{(38)}$= 1.89, p= 0.066) which was, however, driven by smokers (r= 0.54, $CI_{95\%}$[0.12, 0.79], $t_{(18)}$= 2.69, p= 0.015; for non smokers: r= 0.07, p= 0.76, n.s.). Smokers presenting a more vigorous pupil constriction in the PV were those with a more pronounced tendency to direct the initial fixation *away* from NRS. Overall, dwell time and the direction of the initial fixation were positively correlated in non-smokers (r= 0.46, $CI_{95\%}$[0.02, 0.75], $t_{(18)}$= 2.186, p= 0.042), but not in smokers (r= -0.02, p= 0.95, n.s.). Note that the direction of the first fixation is likely to reflect an initial, automatic, fast attentional response to NRS, but because each trial lasted (in the DPT) 1500 ms this initial bias had plenty of room to be potentially overturned. Previous studies have shown that behavioral performance at short vs. long SOAs can be anticorrelated in heavy smokers (della Libera et al., 2019). The disruption of this link may perhaps suggest a two-stages mechanism, in smokers, consisting in a first, early phase in which attention is diverted away from NRS, followed by compensatory mechanisms, unfolding during the course of the trial, favoring a more thorough processing of NRS, although likely modulated by individual traits which are worth considering (della Libera et al., 2019). We thus moved our focus to the group of smokers to assess potential correlations with demographic factors. We first observed that all the three questionnaires were correlated among them, though not with the behavioral measures drawn from the task. We assessed the classifier's predictions for each smoker – predictions are scored in terms of the probability to belong to the smokers' group (values >0.5) vs the non-smokers group (values <0.5, i.e. classification errors). These values were – unsurprisingly, given the results of the modeling detailed above – vastly driven by pupil dilation. More surprising, instead, was noting that the classifier performed best (and the pupil constricted more vigorously) for the participants who smoked less (r= -0.48, $CI_{95\%}$[-0.76, -0.05], $t_{(18)}$= 2.33, p= 0.03). This is interesting, albeit counterintuitive, and will be coped with in the discussion. As last remark, we also note that there was a correlation between the classifier's performance and abstinence time, i.e. hours since the last cigarette. The longer the abstinence, the better the classification (r= 0.55, $CI_{95\%}$[0.15, 0.8], $t_{(18)}$= 2.82, p= 0.01). In other words, false negatives were more

common in individuals who smoked more recently. Also accounting for the fact that those who smoked less were more likely to refrain from smoking for longer times, this latter finding may indicate that some role for craving urges in the emergence of attentional capture effects (**Fig. 3**).

Fig. 3. The relation between the classifier's performance (data points are either circles, i.e. the classification was correct, or triangles, i.e. the classification was not correct), and measures of smoking intensity (self-reported cigarettes per day) and abstinence time (hours). Data points only depict smokers, thus errors are all false negatives. The classifier's predictions were mostly driven by pupil dilation, although there was a minor role for RTs in the DPT task as well. The classifier worked best, and the pupil constricted more vigorously, with light smokers and with participants who refrained from smoking for longer periods prior to the experiment (which are partly the same subgroup of participants).

4. Discussion

In this study, we sought to probe the markers of attentional priority in smokers through eye-tracking measures. In contrast with previous reports, we did not find that behavioral (e.g., RTs) and eye fixation measures were sensitive enough to index attentional capture, let alone accurately predict smoking status. On the other hand, we found differences in the time course of pupil dilation to NRS between smokers and non-smokers, which proved to be a reliable predictor of smoking status. We used a passive viewing task in which participants were simply asked to fixate smoke-related or neutral images for 3 seconds. However, to ensure minimal engagement with all images, rare probes were presented in a minority of trials, after the presentation of images, and participants had to quickly detect them. As a result, the main pattern observed at the level of pupil diameter was a very large, steep, and sustained dilation along the entire temporal window. This can be readily interpreted in terms of mounting temporal expectations about the appearance of the probe, i.e. temporal predictions, and the inherent preparation for the motor response, both of which affect the size of the pupil (Akdoğan et al., 2016; Wang et al., 2016). Within this global pattern, however, pupil dilation to NRS stimuli compared to neutral ones differed as a function of smoking status. We found an interaction, starting from about 800 ms and sustained up to 2850 ms, in which, on average, smokers pupils tended to *constrict* more than non-smokers ones. This is interesting, albeit not entirely expected on the basis of evidence pointing to a robust role of pupil dilation in both appetitive and aversive conditioning (Finke et al., 2021). Moreover, the only previous study that assessed pupil size in response to NRS (Chae et al., 2008) reported pupil dilation in a sample of 7 smokers (against 12 controls), similarly to when viewing highly arousing images (regardless of their valence). However, their task was a pure passive viewing task in which images were presented on screen for a very long duration (30 s). One possibility is that – being in our case the main task awaiting and reacting quickly to the infrequent probe, and being pupil chiefly reflecting that with a strong dilation – relative constriction may suggest attentional resources are slightly diverted from this main goal. The pupil constriction observed in this setting may thus be task-dependent. Although this remains speculative, it is also supported by several observations. First, the latency of this effect (i.e., about 800 ms) – considered that pupil dilation is a relatively slow physiological signal, especially when compared, for example, to EEG – is quite fast and compatible with an early attentional capture by NRS. Second, there was a correlation with the results of the DPT task, in terms of dwell times, which are more classically attributed to spatial attentional biases. Even though there was no

significant difference in dwell times for NRS in smokers, i.e. longer visual exploration of NRS images, at least in our setting, the preference for NRS correlated with a more vigorous *constriction* of the pupil in a subsequent, independent task (PV). The opposite was true, however, when assessing the directional bias toward NRS (i.e., the tendency to fixate NRS stimuli first), for which the correlation with pupil dilation was positive, and for smokers only. In other words, pupil constriction was stronger for participants who favored, in the DPT task, gazing away from NRS stimuli at first. Previous studies have shown that behavioral performance at short vs. long SOAs can be anticorrelated (della Libera et al., 2019), showing that different mechanisms of attentional engagement can unfold at a different pace. One possibility is that the autonomic signal conveyed by the pupil may reflect the combination of all these underlying processes, i.e. both short-term NRS avoidance and long-term NRS preference. With this respect, the term "attentional capture" may be to some extent misleading, in that it implies a discrete, triggering event in time, which acts in a dichotomous fashion; attentional processes are, instead, likely to build up gradually in time, in ways that are not as straightforward, hence viewing them as the ongoing computation of behavioral priority appears more appropriate (see Anderson, 2021, for discussion). That said, we cannot rule out the alternative possibility that results may be due to different affective reactions in the two groups, for example with non-smokers implicitly judging NRS stimuli as more aversive.

Whilst possibly task-dependent, pupil constriction to NRS in this setting retained some predictive validity. In fact, the autonomic signal conveyed by pupil size was, in our study, a far better predictor of smoking status than more traditional proxy measures of attentional capture. The information conveyed by pupil size appears necessary and sufficient for a good classification of smoking status (65% accuracy), although additional proxy measures (i.e., RTs gain to NRS stimuli) may additionally improve the classifier's performance up to 75%. There is, of course, a lot of room for improvement, both in terms of the choice of the modeling approach and the range of behavioral measures to feed to the classifier. For example, several different Stimulus-Onset-Asynchronies may be needed to exploit the full potential of this paradigm (e.g. Della Libera et al., 2019). However, our results put forward pupil dilation as a particularly robust and sensitive candidate to this aim. Furthermore, we maintain that the focus on prediction performance instead of classic inference and significance testing represents a leap forward in the attempt to characterize robust biomarkers of addiction as well as cognitive effects (Yarkoni & Westfall, 2017). In addition, it is equally interesting to focus on the

instances in which the classifier does not correctly predict the smoking status of individuals (i.e., false negatives). Interestingly, albeit counterintuitively, this happened more often with individuals who smoked more cigarettes per day, i.e. a proxy measure for the intensity of smoking. In other words, the classifier worked best, and pupil constricted more vigorously, for participants who smoked less. This result is, however, not the first of its kind (e.g., Mogg et al., 2003, 2005). It has been proposed that, as nicotine dependence increases, the strength of NRS as incentives actually diminishes and leaves space for more habit-driven, automated modes of consumption (Mogg et al., 2005). As a result, previous studies found increased dwell times for NRS at lower levels of nicotine dependence as well (Mogg et al., 2003, 2005); our results fit well with these seminal works and theoretical accounts (Di Chiara, 2000), though ours extend to the autonomic response conveyed by pupil size. The major drawback is that, if this is true, the quest for potential biomarkers or objective indices of dependence should move past indices of attentional capture, as these may be less relevant in heavy(er) smokers. Similarly, this may partly detract from the idea that treatments aimed at modifying attentional bias can be effective. Besides the nontrivial choice of a paradigm that is capable to bias attention for a sustained period of time, in ecological conditions and everyday life situations (i.e., outside the lab), and all while maintaining a short administration time, attentional bias may not be the most critical feature of smokers seeking treatment. On the other hand, however, attentional capture may remain useful in characterizing the stage of dependence, potentially triggering different management strategies. Furthermore, we did find some nonconclusive evidence for attentional bias being stronger for individuals who refrained from smoking for longer before the experiment, which may suggest some relationship with craving. Thus, a more ecumenical conclusion would be that attentional bias may be a useful, objective measure that can be acquired in longitudinal and interventional studies, although a complete overview would only be achieved by collecting all the remaining pieces of the puzzle. In this regard, eye-tracking is a relatively easy, portable, and inexpensive technology, has been extensively studied and associated with cognitive processes in health and disease, and therefore appears a promising approach for future studies. Pupil dilation, more specifically, appears to be in a more than ideal position to this aim, in that it is capable to jointly reflect affective, perceptual/attentional, and cognitive processes, thus returning an appropriately multifaceted picture.

5. Acknowledgments

We are grateful to Dilara Aladağ for help in data collection. This study was carried out within the scope of the research program Dipartimenti di Eccellenza (art.1, commi 314-337 legge 232/2016), which was supported by a grant from MIUR to the Department of General Psychology, University of Padua.

6. Declarations

Funding. This study was carried out within the scope of the research program Dipartimenti di Eccellenza (art.1, commi 314-337 legge 232/2016), which was supported by a grant from MIUR to the Department of General Psychology, University of Padua.

Conflict of interest. The authors declare no conflict of interest.

Ethics approval. The study was approved by the Ethics Committee of the University of Padua (protocol n° 3568). Informed consent was obtained from all individual participants included in the study, in accordance with the declaration of Helsinki.

Availability of data, material, and code. The data, code, and materials for this manuscript are available in the Open Science Framework repository: https://osf.io/6r5ch/

7. References

Akdoğan, B., Balcı, F., & Rijn, H. van. (2016). Temporal Expectation Indexed by Pupillary Response. *Timing & Time Perception*, *4*(4), 354–370. https://doi.org/10.1163/22134468-00002075

Anderson, B. A., Laurent, P. A., & Yantis, S. (2011). Value-driven attentional capture. *Proceedings of the National Academy of Sciences*, *108*(25), 10367–10371. https://doi.org/10.1073/pnas.1104047108

Anderson, B. A., Laurent, P. A., & Yantis, S. (2014). Value-driven attentional priority signals in human basal ganglia and visual cortex. *Brain Research*, *1587*, 88–96. https://doi.org/10.1016/j.brainres.2014.08.062

Anderson, B. A. (2021). Time to stop calling it attentional "capture" and embrace a mechanistic understanding of attentional priority. *Visual Cognition*, 29(9), 537-540.

Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. *Annu. Rev. Neurosci.*, 28, 403-450.

Awh, E., Belopolsky, A. V., & Theeuwes, J. (2012). Top-down versus bottom-up attentional control: A failed theoretical dichotomy. *Trends in Cognitive Sciences*, *16*(8), 437–443. https://doi.org/10.1016/j.tics.2012.06.010

Binda, P., & Murray, S. O. (2015). Spatial attention increases the pupillary response to light changes. *Journal of Vision*, *15*(2), 1. https://doi.org/10.1167/15.2.1

Bisley, J. W., & Mirpour, K. (2019). The neural instantiation of a priority map. *Current Opinion in Psychology*, *29*, 108–112. https://doi.org/10.1016/j.copsyc.2019.01.002

Blini, E., Pitteri, M., & Zorzi, M. (2019). Spatial grounding of symbolic arithmetic: an investigation with optokinetic stimulation. *Psychological research*, 83(1), 64-83.

Blini, E., Tilikete, C., Chelazzi, L., Farnè, A., & Hadj-Bouziane, F. (2020). The role of the vestibular system in value attribution to positive and negative reinforcers. *Cortex*, *133*, 215–235. https://doi.org/10.1016/j.cortex.2020.09.004

Boccara, C. N., Nardin, M., Stella, F., O'Neill, J., & Csicsvari, J. (2019). The entorhinal cognitive map is attracted to goals. *Science*, *363*(6434), 1443–1447. https://doi.org/10.1126/science.aav4837

Bogdanova, O. V., Bogdanov, V. B., Miller, L. E., & Hadj-Bouziane, F. (2022). Simulated proximity enhances perceptual and physiological responses to emotional facial expressions. *Scientific Reports*, *12*(1), 109. https://doi.org/10.1038/s41598-021-03587-z

Bourgeois, A., Chelazzi, L., & Vuilleumier, P. (2016). Chapter 14—How motivation and reward learning modulate selective attention. In B. Studer & S. Knecht (A c. Di), *Progress in Brain Research* (Vol. 229, pagg. 325–342). Elsevier. https://doi.org/10.1016/bs.pbr.2016.06.004

Butler, W. N., Hardcastle, K., & Giocomo, L. M. (2019). Remembered reward locations restructure entorhinal spatial maps. *Science*, *363*(6434), 1447–1452. https://doi.org/10.1126/science.aav5297

Camara, E., Manohar, S., & Husain, M. (2013). Past rewards capture spatial attention and action choices. *Experimental Brain Research*, *230*(3), 291–300. https://doi.org/10.1007/s00221-013-3654-6

Casarotti, M., Lisi, M., Umiltà, C., & Zorzi, M. (2012). Paying Attention through Eye Movements: A Computational Investigation of the Premotor Theory of Spatial Attention. *Journal of Cognitive Neuroscience*, *24*(7), 1519–1531. https://doi.org/10.1162/jocn_a_00231

Chae, Y., Lee, J.-C., Park, K.-M., Kang, O.-S., Park, H.-J., & Lee, H. (2008). Subjective and Autonomic Responses to Smoking-Related Visual Cues. *The Journal of Physiological Sciences*, *advpub*, 0803240039–0803240039. https://doi.org/10.2170/physiolsci.RP014207

Chanon, V. W., Sours, C. R., & Boettiger, C. A. (2010). Attentional bias toward cigarette cues in active smokers. *Psychopharmacology*, *212*(3), 309–320. https://doi.org/10.1007/s00213-010-1953-1

Chelazzi, L., Perlato, A., Santandrea, E., & Della Libera, C. (2013). Rewards teach visual selective attention. *Vision Research*, *85*, 58–72. https://doi.org/10.1016/j.visres.2012.12.005

Christiansen, P., Schoenmakers, T. M., & Field, M. (2015). Less than meets the eye: Reappraising the clinical relevance of attentional bias in addiction. *Addictive Behaviors*, *44*, 43–50. https://doi.org/10.1016/j.addbeh.2014.10.005

Cox, L. S., Tiffany, S. T., & Christen, A. G. (2001). Evaluation of the brief questionnaire of smoking urges (QSU-brief) in laboratory and clinical settings. *Nicotine & Tobacco Research*, *3*(1), 7–16. https://doi.org/10.1080/14622200124218

Cox, W. M., Fadardi, J. S., Intriligator, J. M., & Klinger, E. (2014). Attentional bias modification for addictive behaviors: Clinical implications. *CNS Spectrums*, *19*(3), 215–224. https://doi.org/10.1017/S1092852914000091

Cox, W. M., Fadardi, J. S., & Pothos, E. M. (2006). The Addiction-Stroop Test: Theoretical Considerations and Procedural Recommendations. *Psychological Bulletin*, *132*(3), 443–476.

Della Libera, C., Zandonai, T., Zamboni, L., Santandrea, E., Sandri, M., Lugoboni, F., Chiamulera, C., & Chelazzi, L. (2019). Revealing Dissociable Attention Biases in Chronic Smokers Through an Individual-Differences Approach. Scientific Reports, 9(1), 4930. https://doi.org/10.1038/s41598-019-40957-0

Di Chiara, G. (2000). Role of dopamine in the behavioural actions of nicotine related to addiction. *European Journal of Pharmacology*, *393*(1), 295–314. https://doi.org/10.1016/S0014-2999(00)00122-9

Driver, J. (2001). A selective review of selective attention research from the past century. *British Journal of Psychology*, *92*(1), 53–78. https://doi.org/10.1348/000712601162103

Duncan, D., & Theeuwes, J. (2020). Statistical learning in the absence of explicit top-down attention. *Cortex*, *131*, 54–65. https://doi.org/10.1016/j.cortex.2020.07.006

Dureux, A., Blini, E., Grandi, L. C., Bogdanova, O., Desoche, C., Farnè, A., & Hadj-Bouziane, F. (2021). Close facial emotions enhance physiological responses and facilitate perceptual discrimination. *Cortex*. https://doi.org/10.1016/j.cortex.2021.01.014

Ehrman, R. N., Robbins, S. J., Bromwell, M. A., Lankford, M. E., Monterosso, J. R., & O'Brien, C. P. (2002). Comparing attentional bias to smoking cues in current smokers, former smokers, and non-smokers using a dot-probe task. *Drug and Alcohol Dependence*, *67*(2), 185–191. https://doi.org/10.1016/S0376-8716(02)00065-0

Failing, M., & Theeuwes, J. (2018). Selection history: How reward modulates selectivity of visual attention. *Psychonomic Bulletin & Review*, *25*(2), 514–538. https://doi.org/10.3758/s13423-017-1380-y

Fecteau, J. H., & Munoz, D. P. (2006). Salience, relevance, and firing: A priority map for target selection. *Trends in Cognitive Sciences*, *10*(8), 382–390. https://doi.org/10.1016/j.tics.2006.06.011

Field, M., Christiansen, P., Cole, J., & Goudie, A. (2007). Delay discounting and the alcohol Stroop in heavy drinking adolescents. *Addiction*, *102*(4), 579–586. https://doi.org/10.1111/j.1360-0443.2007.01743.x

Field, M., & Cox, W. (2008). Attentional bias in addictive behaviors: A review of its development, causes, and consequences. *Drug and Alcohol Dependence*, *97*(1–2), 1–20. https://doi.org/10.1016/j.drugalcdep.2008.03.030

Field, M., Marhe, R., & Franken, I. H. A. (2014). The clinical relevance of attentional bias in substance use disorders. *CNS Spectrums*, *19*(3), 225–230. https://doi.org/10.1017/S1092852913000321

Field, M., Mogg, K., & Bradley, B. P. (2004). Eye movements to smoking-related cues: Effects of nicotine deprivation. *Psychopharmacology*, *173*(1), 116–123. https://doi.org/10.1007/s00213-003-1689-2

Field, M., Munafò, M. R., & Franken, I. H. A. (2009). A meta-analytic investigation of the relationship between attentional bias and subjective craving in substance abuse. *Psychological Bulletin*, *135*(4), 589–607. https://doi.org/10.1037/a0015843

Finke, J. B., Roesmann, K., Stalder, T., & Klucken, T. (2021). Pupil dilation as an index of Pavlovian conditioning. A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, *130*, 351–368. https://doi.org/10.1016/j.neubiorev.2021.09.005

Goldstein, R. Z., & Volkow, N. D. (2002). Drug Addiction and Its Underlying Neurobiological Basis: Neuroimaging Evidence for the Involvement of the Frontal Cortex. *American Journal of Psychiatry*. https://doi.org/10.1176/appi.ajp.159.10.1642

Goldstein, R. Z., & Volkow, N. D. (2011). Dysfunction of the prefrontal cortex in addiction: Neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*, *12*(11), 652–669. https://doi.org/10.1038/nrn3119

Gossop, M., Griffiths, P., Powis, B., & Strang, J. (1992). Severity of dependence and route of administration of heroin, cocaine and amphetamines. *British Journal of Addiction*, *87*(11), 1527–1536. https://doi.org/10.1111/j.1360-0443.1992.tb02660.x

Harris, J. A., Donohue, S. E., Ilse, A., Ariel Schoenfeld, M., Heinze, H.-J., & Woldorff, M. G. (2018). EEG measures of brain activity reveal that smoking-related images capture the attention of smokers outside of awareness. *Neuropsychologia*, *111*, 324–333. https://doi.org/10.1016/j.neuropsychologia.2018.02.005

Heatherton, T. F., Kozlowski, L. T., Frecker, R. C., & Fagerstrom, K.-O. (1991). The Fagerström Test for Nicotine Dependence: A revision of the Fagerstrom Tolerance Questionnaire. *British Journal of Addiction*, *86*(9), 1119–1127. https://doi.org/10.1111/j.1360-0443.1991.tb01879.x

Hickey, C., Chelazzi, L., & Theeuwes, J. (2010). Reward Changes Salience in Human Vision via the Anterior Cingulate. *The Journal of Neuroscience*, *30*(33), 11096–11103. https://doi.org/10.1523/JNEUROSCI.1026-10.2010

Jiang, Y. V., Sha, L. Z., & Remington, R. W. (2015). Modulation of spatial attention by goals, statistical learning, and monetary reward. *Attention, Perception, & Psychophysics*, *77*(7), 2189–2206. https://doi.org/10.3758/s13414-015-0952-z

Li, Z. (2002). A saliency map in primary visual cortex. *Trends in Cognitive Sciences*, *6*(1), 9–16. https://doi.org/10.1016/S1364-6613(00)01817-9

Lisi, M., Bonato, M., & Zorzi, M. (2015). Pupil dilation reveals top–down attentional load during spatial monitoring. *Biological Psychology*, *112*, 39–45. https://doi.org/10.1016/j.biopsycho.2015.10.002

Manohar, S. G., Chong, T. T.-J., Apps, M. A. J., Batla, A., Stamelou, M., Jarman, P. R., Bhatia, K. P., & Husain, M. (2015). Reward Pays the Cost of Noise Reduction in Motor and Cognitive Control. *Current Biology*, *25*(13), 1707–1716. https://doi.org/10.1016/j.cub.2015.05.038

Manohar, S. G., & Husain, M. (2015). Reduced pupillary reward sensitivity in Parkinson's disease. *Npj Parkinson's Disease*, *1*, 15026. https://doi.org/10.1038/npjparkd.2015.26

Manoliu, A., Haugg, A., Sladky, R., Hulka, L., Kirschner, M., Brühl, A. B., Seifritz, E., Quednow, B., Herdener, M., & Scharnowski, F. (2020). SmoCuDa: A Validated Smoking Cue Database to Reliably Induce Craving in Tobacco Use Disorder. *European Addiction Research*, 1–8. https://doi.org/10.1159/000509758

Mathôt, S. (2018). Pupillometry: Psychology, Physiology, and Function. *Journal of Cognition*, *1*(1), 16. https://doi.org/10.5334/joc.18

Mathôt, S., Schreij, D., & Theeuwes, J. (2012). OpenSesame: An open-source, graphical experiment builder for the social sciences. *Behavior Research Methods*, *44*(2), 314–324. https://doi.org/10.3758/s13428-011-0168-7

Maunsell, J. H. R. (2004). Neuronal representations of cognitive state: Reward or attention? *Trends in Cognitive Sciences*, *8*(6), 261–265. https://doi.org/10.1016/j.tics.2004.04.003

Mogg, K., Bradley, B. P., Field, M., & Houwer, J. D. (2003). Eye movements to smoking-related pictures in smokers: Relationship between attentional biases and implicit and explicit measures of stimulus valence. *Addiction*, *98*(6), 825–836. https://doi.org/10.1046/j.1360-0443.2003.00392.x

Mogg, K., Field, M., & Bradley, B. P. (2005). Attentional and approach biases for smoking cues in smokers: An investigation of competing theoretical views of addiction. *Psychopharmacology*, *180*(2), 333–341. https://doi.org/10.1007/s00213-005-2158-x

Mogoaşe, C., David, D., & Koster, E. H. W. (2014). Clinical Efficacy of Attentional Bias Modification Procedures: An Updated Meta-Analysis. *Journal of Clinical Psychology*, *70*(12), 1133–1157. https://doi.org/10.1002/jclp.22081

Muhammed, K., Dalmaijer, E., Manohar, S., & Husain, M. (2020). Voluntary modulation of saccadic peak velocity associated with individual differences in motivation. *Cortex*, *122*, 198–212. https://doi.org/10.1016/j.cortex.2018.12.001

Muhammed, K., Manohar, S., Ben Yehuda, M., Chong, T. T.-J., Tofaris, G., Lennox, G., Bogdanovic, M., Hu, M., & Husain, M. (2016). Reward sensitivity deficits modulated by dopamine are associated with apathy in Parkinson's disease. *Brain*, *139*(10), 2706–2721. https://doi.org/10.1093/brain/aww188

Noël, X., Brevers, D., & Bechara, A. (2013). A neurocognitive approach to understanding the neurobiology of addiction. *Current Opinion in Neurobiology*, *23*(4), 632–638. https://doi.org/10.1016/j.conb.2013.01.018

Pietrock, C., Ebrahimi, C., Katthagen, T. M., Koch, S. P., Heinz, A., Rothkirch, M., & Schlagenhauf, F. (2019). Pupil dilation as an implicit measure of appetitive Pavlovian learning. *Psychophysiology*, *56*(12), e13463. https://doi.org/10.1111/psyp.13463

Posner, M. I. (1980). Orienting of attention. *The Quarterly Journal of Experimental Psychology*, *32*(1), 3–25.

Raymond, J. E., & O'Brien, J. L. (2009). Selective Visual Attention and Motivation: The Consequences of Value Learning in an Attentional Blink Task. *Psychological Science*, *20*(8), 981–988. https://doi.org/10.1111/j.1467-9280.2009.02391.x

Reynaud, A. J., Blini, E., Koun, E., Macaluso, E., Meunier, M., & Hadj-Bouziane, F. (2021). Atomoxetine modulates the contribution of low-level signals during free viewing of natural images in rhesus monkeys. *Neuropharmacology*, *182*, 108377. https://doi.org/10.1016/j.neuropharm.2020.108377

Rizzolatti, G., Riggio, L., Dascola, I., & Umiltá, C. (1987). Reorienting attention across the horizontal and vertical meridians: Evidence in favor of a premotor theory of attention. *Neuropsychologia*, *25*(1, Part 1), 31–40. https://doi.org/10.1016/0028-3932(87)90041-8

Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. Psychophysiology, 56(6), e13335.

Sprague, T. C., & Serences, J. T. (2013). Attention modulates spatial priority maps in the human occipital, parietal and frontal cortices. *Nature Neuroscience*, *16*(12), 1879–1887. https://doi.org/10.1038/nn.3574

Takikawa, Y., Kawagoe, R., Itoh, H., Nakahara, H., & Hikosaka, O. (2002). Modulation of saccadic eye movements by predicted reward outcome. *Experimental Brain Research*, *142*(2), 284–291. https://doi.org/10.1007/s00221-001-0928-1

The R Core Team. (2018). *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.* https://www.r-project.org/

Theeuwes, J., & Belopolsky, A. V. (2012). Reward grabs the eye: Oculomotor capture by rewarding stimuli. *Vision Research*, *74*, 80–85. https://doi.org/10.1016/j.visres.2012.07.024

Volkow, N. D., Wang, G.-J., Fowler, J. S., Tomasi, D., Telang, F., & Baler, R. (2010). Addiction: Decreased reward sensitivity and increased expectation sensitivity conspire to overwhelm the brain's control circuit. *BioEssays*, *32*(9), 748–755. https://doi.org/10.1002/bies.201000042

von der Malsburg, T. (2015). Saccades: Detection of fixations in eye-tracking data. R package version 0.1-1. URL http://CRAN. R-project. org/package= saccades.

Wang, C.-A., McInnis, H., Brien, D. C., Pari, G., & Munoz, D. P. (2016). Disruption of pupil size modulation correlates with voluntary motor preparation deficits in Parkinson's disease. *Neuropsychologia*, *80*, 176–184. https://doi.org/10.1016/j.neuropsychologia.2015.11.019

Wardhani, I. K., Mathôt, S., Boehler, C. N., & Laeng, B. (2020). Effects of nicotine on pupil size and performance during multiple-object tracking in non-nicotine users. *International Journal of Psychophysiology*, *158*, 45–55. https://doi.org/10.1016/j.ijpsycho.2020.09.005

Waters, A. J., Shiffman, S., Bradley, B. P., & Mogg, K. (2003). Attentional shifts to smoking cues in smokers. *Addiction*, *98*(10), 1409–1417. https://doi.org/10.1046/j.1360-0443.2003.00465.x

White, B. J., Berg, D. J., Kan, J. Y., Marino, R. A., Itti, L., & Munoz, D. P. (2017). Superior colliculus neurons encode a visual saliency map during free viewing of natural dynamic video. *Nature Communications*, *8*(1), 14263. https://doi.org/10.1038/ncomms14263

White, B. J., Kan, J. Y., Levy, R., Itti, L., & Munoz, D. P. (2017). Superior colliculus encodes visual saliency before the primary visual cortex. *Proceedings of the National Academy of Sciences*, *114*(35), 9451–9456. https://doi.org/10.1073/pnas.1701003114

Yarkoni, T., & Westfall, J. (2017). Choosing prediction over explanation in psychology: Lessons from machine learning. *Perspectives on Psychological Science*, 12(6), 1100-1122.