Chromosome-scale assembly of the lablab genome - A model for inclusive orphan crop genomics

Isaac Njaci[1,2†], Bernice Waweru[1†], Nadia Kamal[3†], Meki Shehabu Muktar[4], David Fisher[5], Heidrun Gundlach[3], Collins Muli[1], Lucy Muthui[1], Mary Maranga[6], Davies Kiambi[7], Brigitte L Maass[8], Peter MF Emmrich[2,9], Jean-Baka Domelevo Entfellner[1], Manuel Spannagl[3], Mark A Chapman[5*], Oluwaseyi Shorinola[1,2*] and Chris S Jones[1*]

[1]International Livestock Research Institute, Nairobi 00100, Kenya;
[2]John Innes Centre, Norwich Research Park Norwich, NR4 7UH, UK
[3]Helmholtz Zentrum München, German Research Center for Environmental Health, Plant Genome and Systems Biology, Ingolstädter Landstr. 1, 85764 Neuherberg, Germany
[4]International Livestock Research Institute, Addis Ababa, Ethiopia.
[5]University of Southampton, School of Biological Sciences, Southampton, SO17 1BJ, UK
[6]Malopolska Centre of Biotechnology, Jagiellonian University, Gronostajowa 7A, Kraków, Poland
[7]Department of Microbiology and Biochemistry, Hochschule Geisenheim University, Von-Lade-Straße 1, 65366 Geisenheim, Germany
[8]Department of Crop Sciences, Georg-August-University Göttingen, Grisebachstr. 6, 37077 Göttingen, Germany
[9]Department for International Development, University of East Anglia. Norwich, NR4 7TJ, United Kingdom

[†]These authors contributed equally.
*Corresponding authors.

Abstract

Orphan crops (also described as underutilised and neglected crops) hold the key to diversified and climate-resilient food systems. After decades of neglect, the genome sequencing of orphan crops is gathering pace, providing the foundations for their accelerated domestication and improvement. Recent attention has however turned to the gross under-representation of researchers in Africa in the genome sequencing efforts of their indigenous orphan crops. Here we report a radically inclusive approach to orphan crop genomics using the case of *Lablab purpureus* (L.) Sweet (syn. *Dolichos lablab*, or hyacinth bean) – a legume native to Africa and cultivated throughout the tropics for food and forage. Our Africa-led South-North plant genome collaboration produced a high-quality chromosome-scale assembly of the lablab genome – the first chromosome-scale plant genome assembly locally sequenced in Africa. We also re-sequenced cultivated and wild accessions of lablab from Africa confirming two domestication events and examined the genetic diversity in lablab germplasm conserved in Africa. Our approach provides a valuable resource for lablab improvement and also presents a model that could be explored by other researchers sequencing indigenous crops particularly from Low and middle income countries (LMIC).

Introduction

Three major crops currently provide more than 40% of global calorie intake[1]. This over-dependence
on a few staple crops increases the vulnerability of global food systems to environmental and social
instabilities. One promising strategy to diversify food systems is to improve the productivity and
adoption of climate-resilient but underutilised orphan crops through genome-assisted breeding[2].

Genome-assisted breeding offers hope of a new green revolution by helping to uncover and unlock
novel genetic variation for crop improvement. Over the last 20 years, the genomes of 135
domesticated crops have been sequenced and assembled[3], including those of orphan crops[2].
However, it has recently been acknowledged that researchers from Africa are grossly under-
represented in the genome sequencing efforts of their indigenous orphan crops[3,4]. None of the
assemblies of native African plant species released till date were sequenced in Africa. The acute lack
of sequencing facilities and high-performance computing infrastructures as well as bioinformatics
capacity to handle big genome data, has meant that researchers in Africa have historically taken the
back seat in most genome sequencing efforts[5].

Here we present a model to overcome this under-representation through an inclusive orphan crop
genomics approach. We applied an Africa-led, internationally collaborative approach to the genome
sequencing of lablab (*Lablab purpureus* L. Sweet) - a tropical legume native to Africa (Figure 1A).
Lablab is remarkably drought-resilient and thrives in a diverse range of environments, as such it is
widely cultivated throughout the tropical and subtropical regions of Africa and Asia[6]. Lablab is a
versatile multipurpose crop that contributes towards food, feed, nutritional and economic security, and
is also rich in bioactive compounds with pharmacological potential, including against SARS-Cov2[7–10].
Climate change is driving researchers to investigate crops like lablab for its outstanding drought
tolerance[11].

Our Africa-led genome collaboration produced a chromosome-scale assembly of lablab – the first
chromosome-scale plant genome assembly sequenced in Africa. We also discuss the main features
and benefits from our inclusive approach, and suggest this can serve as a roadmap for future
genomic investigations of indigenous African crops.

Results

Genome sequencing

High acquisition and maintenance cost of sequencing platforms is a major limiting factor to genomics
research in Africa. To circumvent this limitation, we used the portable and low-cost Oxford Nanopore
Technology (ONT) MinION platform for in-country sequencing of the genome of lablab (cv.
Highworth). We generated 4.7 M reads with a mean read length of 6.1 Kbp (Table S1). This amounted
to 28.4 Gbp of sequences and 67x coverage of the lablab genome based on a previously estimated
genome size of 423 Mbp[12]. The reads were initially assembled into 2,260 contigs with an N50 of 11.0

Mbp. The assembly was polished for error correction using ~380x of publicly available Illumina short
reads (NCBI Bioproject PRJNA474418).
82
Using high-throughput Chromosome Conformation Capture (Hi-C), we clustered and oriented the
contigs into 11 pseudomolecules covering 417.8 Mbp (98.6% of the estimated genome size) with an
N50 of 38.1 Mbp (Figure 1B, Table S2, Supplemental methods). Our chromosome-scale assembly of
the lablab genome has 61-fold improvement in contiguity compared to the previously published short
read assembly[12]. For consistency with published legume genome sequences, we assigned
chromosome names based on syntenic relationship with *Phaseolus vulgaris* (common bean[13]) and
Vigna unguiculata (cowpea[14]) (Figure S1, Supplemental methods).

Genome annotation and gene family analyses

To annotate the genome, we established an international collaboration comprising locally trained
African researchers (see discussion) and international partners with established genome annotation
pipelines. We used an automated pipeline based on protein homology, transcript evidence and *ab*
initio predictions to identify protein coding genes in the lablab genome. This resulted in a total of
30,922 gene models (79,512 transcripts). A subset of 24,972 of these gene models show no
homology to transposable elements (TEs) and can be confidently considered as high quality protein-
coding non-TE gene models (Figure 1B, Table S3). BUSCO scores of the non-TE gene models were
97.3%, 96.4%, and 94.9% against the universal single copy genes from the embryophyta, eudicots,
and fabales lineages, respectively, suggesting a high level of completeness of the gene space (Figure
1C). The number of non-TE protein-coding genes identified in our study is 19.2% greater than in the
previous short-read assembly[12]. A functional description could be assigned to 28,927 (93.3%) of the
genes.
103
A total of 168,174 TE sequences, occupying 28.1% of the genome, were identified in the lablab
genome (Figure 1B). Of these, 89.6% were classified into 13 superfamilies and 2,353 known families
(Table S4, Figure S2). Long Terminal Repeat - RetroTransposons (LTR-RTs) were the most abundant
TEs, with 85,149 sequences occupying 83 Mb (19.9%) of the genome (Figure 1B). Copia were the
most abundant LTR-RT superfamily, occupying 13.2% of the genome compared to gypsy elements
that occupied only 4.7%. We also report an average LTR Assembly Index (LAI) of 19.8 (Figure 1D).
DNA transposons were smaller in number and size relative to LTR-RTs, and were distributed more
evenly across the chromosomes (Table S4, Figure S2).
112
A further 100,741 repetitive sequences were identified but could not be classified as TEs. Combining
the annotated TEs and unclassified repeats reveals an overall repeat content of 43.4% of the
genome. We also identified 142,302 tandem repeats (TRs) covering 43 Mb (11.2%) of the genome
(Figure 1B, Table S5). Most of these were minisatellites (10-99 bp), while satellite repeats (>100 bp)
make up the largest total proportion of TRs in the genome (7.4% of the genome; Table S5). Both the
tandem and unclassified repeats were found to concentrate within a distinct, overlapping cluster at the
point of peak repeat density on each chromosome, indicating that they are likely centromeric repeats
(Figure S2A).
121
Gene family analysis and comparison to other legumes (*P. vulgaris, V. angularis*, *Cajanus cajan*,
Medicago truncatula)*,* and using *Arabidopsis thaliana* as an outgroup, placed 24,397 (97.7%) of the
24,972 non-TE lablab genes into orthogroups. Comparison of the five legumes (Figure 2A) revealed
14,047 orthogroups in common, and identified 417 (1.7%) lablab genes in 119 species-specific
orthogroups that were absent from the other four legumes. These lablab-unique gene families were

enriched for fatty acid biosynthesis, arabinose metabolism gene ontology (GO) classifiers while
several were involved in pollen-pistil interactions and general plant development (Table S6). Using the
phylogenetic relationships between the species, 448 gene families were significantly expanded in
lablab compared to other legumes and *Arabidopsis*, while 899 were contracted (Figure 2B). Expanded
gene families were enriched for lignin and pectin metabolism, photosynthesis among others (Table
S7; Figure 2C).

Evidence for two domestications of lablab

Understanding the transition from wild species to domesticated crop can provide insight into the
location of domestication, the strength of genetic bottleneck (and identification of wild alleles not
present in the domesticated gene pool) and can lead to identifying candidate genes underlying
domestication traits. Previous work has suggested that lablab domestication occurred at least twice,
separately in the two-seeded and four-seeded gene pools[15,16]. Using our chromosome-scale assembly
as a reference, we examined whether this is indeed the case by resequencing a panel of two-seeded
and four-seeded wild (ssp. *uncinatus*) and domesticated (ssp. *purpureus*) lablab accessions (Table
S8). We also gathered publicly available short read data for cv. Highworth and nine species from
three related genera (*Vigna*, *Phaseolus* and *Macrotyloma*, Table S8) as outgroups to determine the
phylogenetic position of lablab. All lablab samples had a >95% mapping against the lablab reference
genome at a depth of 7.0 - 11.2x while the related genera had considerably lower mapping of 30 -
54% at a depth of 3.5 - 10.9x; Table S8). A total of 39,907,704 SNPs were identified across all 22
samples and 15,428,858 across the 13 lablab samples.
147
A filtered SNP data set of 67,259 SNPs (see Methods) was used for phylogenetic and diversity
analyses. Neighbor Joining phylogenetic analysis rooted with two *Macrotyloma* samples revealed that
all lablab samples formed one group separate from the *Vigna* and *Phaseolus* samples which are each
reciprocally monophyletic. A clear division between the two- and four-seeded lablab samples could be
observed (100% bootstrap support) with wild and domesticated samples found in both groups (Figure
3). Our study thus confirms the previous hypotheses of two origins of domesticated lablab. Genetic
diversity (π per 100 Kb window) within each gene pool was relatively low and significantly greater
(unpaired T-test, t = 8.2415, df = 2651, P < 0.0001) in the two-seeded group (5.79 x 10^{-6} (+/- 2.51 x
10^{-6} [SD]) than the four-seeded group (5.04 x 10^{-6} (+/- 2.15 x 10^{-6} [SD]). Divergence between the two-
and four-seeded gene pools was high (mean Fst per 100kb window = 0.43 +/- 0.32 [SD]) which could
suggest that these gene pools should be taxonomically re-evaluated as separate species.

Genetic diversity in a global lablab collection

To assess within and between accession diversity in the global lablab gene pool, we genotyped 1,860
individuals from 166 lablab accessions using DArTseq genotyping-by-sequencing (GBS) (Table S9).
We identified 41,718 genome-wide SNP and 73,211 SilicoDArT markers, of which 91% and 57%
mapped onto the lablab genome, respectively (Figure S3). The two-seeded and wild samples mapped
with a significant amount of missing data (due to the high genetic divergence described above),
therefore we excluded these and report only results for the widespread four-seeded cultivated group.
In addition, only individuals that were considered true-to-type or progeny (see Supplementary
Information) were included. This resulted in 1,462 individuals from 138 accessions being retained for
the final analysis.
169
170 Using a subset of 2,460 quality-filtered genome-wide SNPs (see Methods) for STRUCTURE[17]
171 analysis, we identified four populations (cluster I - IV) in the lablab germplasm collection (Figure 4A).

Similar clustering and population stratification were detected by hierarchical clustering and PCA (Figure 4B and C). The clustering shows some correspondence with the geographical origin of the genotypes. Accessions in cluster I were mainly from outside Africa and included all the accessions of ssp. *bengalensis*, which has long, relatively narrow pods with up to seven seeds and a particular seed arrangment in the pod. More than 85% of the accessions in clusters II, III and IV are from Africa or were originally collected by the Grassland Research Station in Kitale (Kenya, but most have uncertain origin, Table S9).

The pairwise Fst values among the four clusters varied from 0.31 between clusters I and III to 0.91 between clusters II and IV (Table S10). Analysis of molecular variance (AMOVA) further showed presence of higher genetic variation between the four clusters (62.44%) than within the clusters (37.56%) (Table S11). Within group genetic distance between accessions, Nei's D[18], was lowest within cluster IV (mean D = 0.003) and highest for cluster I (mean D = 0.164; Table S12). Mean Nei's D between progenies of the 41 accessions with ≥2 progenies per accession ranged from 0.0015 to 0.1516 indicating that within accession genetic diversity is generally low, as expected for a predominantly self-pollinating species such as lablab[19].

We found that the population clusters often differed in their mean phenotypes based on historical data describing phenology and morpho-agronomic traits[20]. Twelve of 13 quantitative traits (Figure S4; Table S13) and five of eight qualitative characters (Figure S5; Table S14) differed among the four clusters despite a certain level of phenotypic variation within every cluster. Cluster I accessions are phenotypically variable, containing early-flowering, short plants and includes the only three erect accessions and all ssp. *bengalensis* in a sub-cluster. Plants had four to six relatively large seeds per pod. Cluster II contains the earliest, only colored-flowering accessions, with high flowering node density, and most producing up to four black, mottled seeds per pod. Plants were rather short and had the smallest leaves. Cluster III also includes diverse phenotypes; overall plants were relatively tall, broad, leafy and intermediate to late-flowering with the largest leaves and shortest pods with up to four rather small seeds. Cluster IV comprises the most homogeneous phenotypes; it had the latest, only white-flowering accessions and plants were rather tall, broad and leafy with long flower peduncles, a high number of flowering nodes and four relatively small tan-colored seeds per pod.

Discussion

Africa has a rich plant biodiversity that includes 45,000 species[21], most of which are under-studied and under-utilised. To fully explore these genetic resources, it is important to develop inclusive research models that enable and empower local researchers to study these species under a resource-limited research setting. Our work describes an inclusive African-led effort to produce high-quality genome resources for a climate-resilient and multipurpose native African orphan crop - lablab. Our chromosome-scale reference assembly of lablab improves on the previous assembly in several ways and also highlights some interesting features about lablab's genome, domestication and diversity.

With the use of long-reads and Hi-C scaffolding, we achieved 61-fold improvement in contiguity, and identified a further 34 Mbp of repetitive sequences and 19.2% more gene content compared to the short-read based assembly[12]. In addition, the high average LTR Assembly Index (LAI)[22] (19.8; Figure 1C), comparable to the LAI of a PacBio-based assembly of common bean[23], indicates a high-level of completeness of the repeat space in our assembly. As has been found in other legumes, LTR-RTs were the predominant TE class in our lablab assembly[13,14,24]. In contrast to findings from lablab's close relatives, however, we found copia LTR-RTs to be more abundant than gypsy LTR-RTs. It is

219 uncommon to see a greater abundance of copia LTR-RTs when compared to gypsy LTR-RTs in plant
220 genomes[25,26], and although the biological significance of elevated copia abundances remains to be
221 seen, further genome sequencing will determine whether this finding is indeed a distinguishing feature
222 of lablab.
223
Lablab has a smaller genome size than other sequenced legumes and also has a smaller number of
species-specific orthogroups. Nevertheless, the orthogroup analysis identified several GO categories
enriched in the lablab-specific orthogroups; of particular interest are those involved in fatty acid
metabolism, which could underlie seed oil content and composition. In addition arabinose metabolism
genes were enriched in the lablab-unique genes and several other cell wall-related GO terms
(specifically related to pectin and lignin) in the orthogroups expanded in lablab. Cell wall modification
could be related to protection from pathogens[27] or drought tolerance[28].
231
A dual origin of domesticated lablab was confirmed, with the localised (to Ethiopia) two-seeded and
the widespread four-seeded types being genetically distinct and domestication events occurring in
both of these groups. This therefore adds lablab to the relatively 'exclusive' list of crops with more
than one origin, which includes common bean[13], lychee[29], Tartary buckwheat[30] and, potentially, rice[31]
and barley[32]. Data on reproductive isolation between the gene pools is unclear, and crosses are only
known between four-seeded samples[33–35], thus any taxonomic reassessment (first suggested by
Maass et al. 2005[15]) should begin with assessing reproductive compatibility between the gene pools.
239
Importantly, our project provides a model for increasing the representation of local researchers in the
sequencing of their indigenous crops. Recent studies and commentaries have highlighted the
disconnect between the species origin and the location of the institutions leading their sequencing[3–5].
This is particularly true for Africa, where none of its sequenced indigenous crops were sequenced on
the continent[4]. We surveyed 31 publications describing the genome sequencing of 24 indigenous
African crops. More than 85% of these publications do not have first or corresponding authors with
affiliations in Africa and 42% do not have any authors with an African affiliation (Table S15). Our
project breaks this trend because sequencing and coordination efforts were done or led from within
Africa, while still recruiting international partners where complementary expertise was beneficial to the
project. Thus we encourage contribution of the international community in African orphan crop
genomics while supporting more active involvement from local researchers.
251
Three main features characterised our inclusive genome collaboration model - access to low-cost
portable sequencing, in-depth capacity building and equitable international collaboration. The high
acquisition and maintenance costs of genome sequencing technologies has historically limited the
participation of researchers working in LMIC in genome collaborations. Low-cost and portable
sequencing platforms such as the ONT MinION, are now making long-read sequencing accessible to
researchers in LMIC, thus "democratising" genome sequencing. We spent less than $4,000 to procure
the MinION sequencer and kits (ONT starter pack and extra flow cells) used to sequence our lablab
genome. This low cost is partly due to the small genome size of lablab, but it nonetheless
demonstrates how accessible modern portable sequencing platforms can be for researchers in
resource-limited research settings. Despite these low costs, there are, however, still logistical
challenges to overcome in getting needed reagents to local labs.
263
Secondly, our project benefited from efforts to build in-depth bioinformatics skills in Africa[5]. Four of the
African authors in our study, including two of the first authors, benefited from a residential 8-month
bioinformatics training in Africa. We posit that such in-country and long-term training, as opposed to
short training, are more effective in developing the high-competence bioinformatics skills that the

continent needs. Once trained, these researchers will feel empowered to participate or lead genomic
projects, and importantly use such projects as opportunities to train many more researchers, thus
creating a continuous stream of human resources equipped to explore the rich genetic resources on
the continent.
272
Lastly, establishing an international collaboration helped us to take advantage of existing expertise
and already developed pipelines for genome analyses. With over 20 years of plant genome
sequencing, the global plant science community have developed tools, pipelines and protocols for
plant genome analyses. This means African researchers do not have to 'reinvent the wheel' for
orphan crop genomics, but instead can form strategic collaborations to access needed expertise and
networks. To fully benefit from big-data and a suite of readily-available genomic tools, it is also vital
that African institutions are supported to build or access physical or cloud computing infrastructure for
high-throughput data analytics. This will also ensure that genomic data produced on the continent are
locally managed and made readily accessible to local researchers and the global community.
282
283 Our lablab genome assembly and collaboration provides a roadmap for improving agronomic, yield
284 and nutritional traits in other African orphan crops. Given the Africa-centred and inclusive nature of
285 our work, this could be used as a model by individual labs and multinational genome consortia
286 including the Africa Biogenome Initiative[3] to generate high-quality genomic resources for many
287 indigenous species across the continent.
288

Methods

Reference genome DNA extraction and sequencing

L. purpureus (L.) Sweet cv. Highworth[36] seeds were germinated in a petri dish on filter papers
moistened with tap water. The sprouted seedlings were transferred to soil and allowed to grow for
one month in the greenhouse facility at the International Livestock Research Institute (ILRI, Kenya).
Two grams of young trifoliate leaves were harvested, flash frozen in liquid nitrogen and stored at -
80^0C. The leaves were ground in liquid nitrogen using a pestle and mortar and High Molecular Weight
(HMW) DNA extracted with Carlson lysis buffer (100 mM Tris-HCl, pH 9.5, 2% CTAB, 1.4 M NaCl, 1%
PEG 8000, 20 mM EDTA) followed by purification using the Qiagen Genomic-tip 100/G based on the
Oxford Nanopore Technologies (ONT) HMW plant DNA extraction protocol. The library was prepared
following the ONT SQK-LSK109 ligation sequencing kit protocol. A total of 1 µg of genomic DNA was
repaired and 3'-adenylated with the NEBNext FFPE DNA Repair Mix and the NEBNext® Ultra™ II
End Repair/dA-Tailing Module and sequencing adapters ligated using the NEBNext Quick Ligation
Module (NEB). After library purification with AMPure XP beads, sequencing was conducted at ILRI
(Kenya) using the R9.4.1 flow cells on a MinION sequencer platform.

Genome Assembly

Guppy basecaller (v4.1.1)[37] was used for base calling the reads using the high accuracy basecalling
model and the resulting fastq files were used for genome assembly. Flye *de novo* long reads
assembler[38] (ver 2.7.1) was used for the assembly with the default parameters. The draft assembly
was polished with lablab Illumina shorts reads[12] using HyPo hybrid polisher[39]. The draft genome
assembly quality was assessed using QUAST[40] and its completeness evaluated using BUSCO (ver.

4.0.6)[41]. The Hi-C library for genome scaffolding was prepared, sequenced and assembled by phase
genomics, USA (Supplemental Information).

Gene Annotation

Protein sequences from five closely related species (*P. vulgaris, V. angularis*, *C. cajan*, and *M.*
truncatula) as well as *Arabidopsis thaliana* were used as protein homology evidence. RNAseq data
from *Lablab purpureus* cv. Highworth leaves, stem, sepals, and petals[12] was used in *de novo*
transcript assembly with Trinity[42] (ver 2.8.5) and provided as transcript evidence. The Funannotate
pipeline[43] (ver 1.8.7) was used for gene prediction using RNA-Seq reads, *de novo* assembled
transcripts and soft-masked genome as input to generate an initial set of gene models using PASA[44]
(ver 2.4.1). Next, the gene models and protein homology evidence were used to train Augustus[45] (ver
3.3.3), SNAP[46] (ver 2006-07-28) and Glimmerhmm[47] (ver 3.0.4) *ab initio* gene predictors and predicted
genes passed to Evidence modeller[48] (ver 1.1.1) with various weights for integration. tRNAscan-SE[49]
(ver 2.0.9) was used to predict non-overlapping tRNAs. Transcript evidence was then used to correct,
improve and update the predicted gene models and refine the 5'- and 3'-untranslated regions (UTRs).
324
The plant.annot pipeline (github.com/PGSB-HMGU/plant.annot) was also used for the prediction of
protein coding genes and incorporated homology information and transcript evidence as well. In the
evidence-based step, RNA-Seq data from cv. Highworth leaf, stem, sepal and petal[12] was used for the
genome-guided prediction of gene structures. HISAT2[50] (version 2.1.0, parameter –dta) was used to
map RNA-Seq data to the reference genome and the transcripts assembled with Stringtie[51] (version
1.2.3, parameters -m 150 -t -f 0.3). For the homology-based step, homologous proteins from the
closely related species were mapped to the reference genome using the splice-aware mapper
GenomeThreader[52] (version 1.7.1, parameters: -startcodon -finalstopcodon -species medicago -
gcmincoverage 70 -prseedlength 7 -prhdist 4). Transdecoder[53] (version 3.0.0) was used to predict
protein sequences and to identify potential open reading frames. The predicted protein sequences
were compared to a protein reference database (UniProt Magnoliophyta, reviewed/Swiss-Prot) using
BLASTP[54] (-max_target_seqs 1 -evalue 1e^{-05}). Conserved protein family domains for all proteins were
identified with hmmscan[55] version 3.1b2. Transdecoder-predict was run on the BLAST and hmmscan
results and the best translation per transcript was selected. Results from the homology and transcript-
based gene prediction approaches were combined and redundant protein sequences were removed.
340
The results from both the funannotate and plant.annot pipelines were combined and redundant
protein sequences as well as non-coding genes removed. The functional annotation of transcripts as
well as the assignment of Pfam[56]- and InterPro[57]-domains, and GO[58,59] terms, were performed using
AHRD   (Automatic   assignment   of   Human   Readable   Descriptions,
https://github.com/groupschoof/AHRD; version 3.3.3). AHRD assesses homology information to other
known proteins using BLASTP searches against Swiss-Prot, The Arabidopsis Information Resource
(TAIR), and TrEMBL. The functional annotations are defined using the homology information and the
domain search results from InterProScan and Gene Ontology terms. In order to distinguish
transposon related genes from other genes, the functional annotation was used to tag TE-related
genes in the genome annotation file. BUSCO[41] v5.2.2 was used to assess the completeness of the
genome annotation, with sets of universal single copy gene orthologs from embryophyta, fabales, and
eudicots odb10 lineages[41].
353

Repeat Annotation

Repeat annotations for transposable elements (TE) and tandem repeats were conducted independently. For TE annotation, a novel Lablab TE library was constructed using the Extensive de novo TE Annotator (EDTA v1.9.7) pipeline[60]. EDTA incorporates both structure and homology-based detection programs to annotate the predominant TE classes found in plant genomes. EDTA utilises LTRharvest[61], LTR_FINDER[62], LTR_retriever[63], TIR-Learner[64], HelitronScanner[65], RepeatModeler2[66] and RepeatMasker[67] for identification of novel TE sequences. The outputs of each module are then combined and filtered to compile a comprehensive, non-redundant TE library. EDTA's inbuilt whole genome annotation function was then used to produce a non-overlapping TE annotation for lablab using the TE library as input. Further calculation of metrics and data visualisation were carried out in R[68] using the tidyverse suite[69] of packages.

Tandem repeats were identified with TandemRepeatFinder[70] under default parameters and subjected to an overlap removal by prioritising higher scores. Higher scoring matches were assigned first. Lower scoring hits at overlapping positions were either shortened or removed. Removal was triggered if the lower scoring hits were contained to ≥ 90% in the overlap or if less than 50 bp of rest length remained.

Gene family and expansion analysis

Gene families were identified using a genome-wide phylogenetic comparison of the lablab protein sequences and four other legumes. This comprised *P. vulgaris* (PhaVulg1_0)*, V. angularis* (Vigan1.1), *C. cajan* (V1.0)*,* and *M. truncatula (*MtrunA17r5)*.* Orthofinder[71] (Version 2.4) was used to identify orthologs and co-orthologs between these species and to group them into gene families. *Arabidopsis thaliana* (Araport 11) was used as an outgroup. The longest transcript was selected for genes with multiple splice variants.

In order to analyse gene family expansion and contraction in lablab, the gene family file produced by Orthofinder was further analysed with CAFE5[72]. An ultrametric tree was built with Orthofinder (r=160) and CAFE5 [72] was run with -k 3. Enrichment analysis using a fisher's exact test (padj ≤ 0.05) of significantly (p-value of gene family sizes[73] ≤ 0.05) expanded gene families was performed with TopGO[74].

Resequencing and Phylogenetic Analyses

Lablab seeds (obtained from ILRI) for the resequencing were germinated in a 1:1 mixture of vermiculite and Levingtons's M2+S compost in a greenhouse (22°C and 16 hour day) at the University of Southampton. Young leaf tissue was harvested from one-month old seedlings and snap frozen in liquid nitrogen. DNA was extracted from leaf tissue using a CTAB-based protocol[75] with minor modifications. In total, 12 samples from two and four-seeded wild and domesticated lablab accessions were sequenced using 2 x 150 bp PE sequencing on an Illumina platform at Novogene (Cambridge, UK) (Table S6). Short read data from lablab cv. Highworth[12], three *Phaseolus*, four *Vigna*, and two *Macrotyloma* species were downloaded from the NCBI Sequence Read Archive (Table S8). A maximum of 100 M read pairs were downloaded.

The reads were trimmed using Trimmomatic[76] (ver 0.32) with the parameters; ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10, LEADING:5, TRAILING:5, SLIDINGWINDOW:4:15, MINLEN:72. Between 21.9 and 97.1 M reads remained after trimming. The trimmed reads were mapped to the chromosome-scale lablab assembly (excluding unmapped contigs) using Bowtie2[77]

(ver 2.2.3) and *--very-sensitive-local* settings. SAMtools[78] (ver 1.1) was used to convert .sam to .bam
files which were then sorted, and duplicated reads were removed using the Picard toolkit[79] (ver 2.8.3,
VALIDATION_STRINGENCY=LENIENT). Depth was estimated using SAMtools[78] (Table S6). Using
mpileup from bcftools[80] (ver 1.6.0), the individual sorted bam files were combined into a multi-sample
VCF using the settings *-Q 13* and *-q 10* and variant detection was performed with "bcftools call".
Variants were subsequently filtered using "bcftools filter", *-i'QUAL>20 & DP>6'.* The proportion of
missing data per individual was calculated using vcftools[81] (ver 0.1.14; Table S6). Finally, vcftools was
used to trim the filtered VCF, removing SNPs that were missing in more than two samples and those
with a minor allele frequency of <5%. Finally, only SNPs that were at least 2 Kbp apart were included.
The final file contained 67,259 SNPs. VCF2Dis (github.com/BGI-shenzhen/VCF2Dis/; ver 1.36) was
used to create a distance matrix which was submitted to the FAST-ME server
(atgc-montpellier.fr/fastme) to generate a NJ tree. A total of 1000 replicate matrices were generated in
VCF2Dis and the phylip commands "*neighbor"* and "*consense"* were used to calculate bootstrap
values. Genetic diversity for the two subpopulations and Fst between the subpopulations were
calculated from the final VCF file using vcftools in 100kb windows.

Population structure and diversity

A total of 1,860 seedlings from 166 *Lablab purpureus* accessions, that have been maintained at the
ILRI forage genebank were grown from seed under screen house conditions at ILRI, Ethiopia.
Genomic DNA was extracted from leaves using a DNeasy® Plant Mini Kit (Qiagen Inc., Valencia, CA).
The DNA samples were genotyped by the DArTseq genotyping platform at Diversity Arrays
Technology, Canberra, Australia[82]. A subset of 2,460 robust SNP markers was filtered based on the
marker's minor allele frequency (MAF ≥ 2 %), missing values (NA ≤ 10 %), independence from each
other (Linkage disequilibrium-LD ≤ 0.7), and their distribution across the genome.

A pairwise IBD (Identity-By-Descent) analysis was conducted using PLINK[83] and contaminants
excluded from the following analyses (see Supplemental Information) Genetic diversity was estimated
using pairwise Nei's genetic distance[18]. Population stratification was assessed using the Bayesian
algorithm implemented in STRUCTURE[17], in which the burn-in time and number of iterations were
both set to 100,000 with 10 repetitions, testing the likelihood of 1-10 subpopulations in an admixture
model with correlated allele frequencies. Using Structure Harvester[84] the most likely number of
subpopulations was determined by the Evanno ΔK method[85]. Accessions with less than 60%
membership probability were considered admixed. Hierarchical clustering, principal component
analysis (PCA), fixation index (Fst), and analysis of molecular variance (AMOVA) were conducted
using the R-packages *Poppr*[86], adegenet[87], and APE[88].

Data availability

431 The lablab genome is available from NCBI BioProject (PRJNA824307) and at
432 https://hpc.ilri.cgiar.org/~bngina/lablab_longread_sequencing_March_2022/. Raw sequencing
433 reads for the resequencing are available from the NCBI SRA under project number
434 PRJNA834808.

References

1. FAOSTAT. *Food balance sheets,* http://faostat.fao.org (2022).
2. Chapman, M. A., He, Y. & Zhou, M. Beyond a reference genome: pangenomes and population genomics of underutilized and orphan crops for future food and nutrition security. *New Phytol.* (2022) doi:10.1111/nph.18021.
3. Marks, R. A., Hotaling, S., Frandsen, P. B. & VanBuren, R. Representation and participation across 20 years of plant genome sequencing. *Nat Plants* **7**, 1571–1578 (2021).
4. Ebenezer, T. E. *et al.* Africa: sequence 100,000 species to safeguard biodiversity. *Nature* **603**, 388–392 (2022).
5. Ghazal, H. *et al.* Plant genomics in Africa: present and prospects. *Plant J.* **107**, 21–36 (2021).
6. Maass, B. L. *et al.* Lablab purpureus-A Crop Lost for Africa? *Trop. Plant Biol.* **3**, 123–135 (2010).
7. Habib, H. M., Theuri, S. W., Kheadr, E. E. & Mohamed, F. E. Functional, bioactive, biochemical, and physicochemical properties of the Dolichos lablab bean. *Food Funct.* **8**, 872–880 (2017).
8. Weldeyesus, G. Forage productivity system evaluation through station screening and intercropping of lablab forage legume with maize under Irrigated lands of smallholder farmers. *Afr. J. Agric. Res.* **12**, 1841–1847 (2017).
9. Minde, J. J., Venkataramana, P. B. & Matemu, A. O. Dolichos Lablab-an underutilized crop with future potentials for food and nutrition security: a review. *Crit. Rev. Food Sci. Nutr.* **61**, 2249–2261 (2021).
10. Liu, Y.-M. *et al.* A Carbohydrate-Binding Protein from the Edible Lablab Beans Effectively Blocks the Infections of Influenza Viruses and SARS-CoV-2. *Cell Rep.* **32**, 108016 (2020).
11. Missanga, J. S., Venkataramana, P. B. & Ndakidemi, P. A. Recent developments in Lablab purpureus genomics: A focus on drought stress tolerance and use of genomic resources to develop stress-resilient varieties. *Legume Science* **3**, (2021).
12. Chang, Y. *et al.* The draft genomes of five agriculturally important African orphan crops. *Gigascience* **8**, (2019).
13. Schmutz, J. *et al.* A reference genome for common bean and genome-wide analysis of dual domestications. *Nat. Genet.* **46**, 707–713 (2014).
14. Lonardi, S. *et al.* The genome of cowpea (Vigna unguiculata [L.] Walp.). *Plant J.* **98**, 767–782 (2019).
15. Maass, B. L., Jamnadass, R. H., Hanson, J. & Pengelly, B. C. Determining Sources of Diversity in Cultivated and Wild Lablab purpureus Related to Provenance of Germplasm by using Amplified Fragment Length Polymorphism. *Genet. Resour. Crop Evol.* **52**, 683–695 (2005).
16. Maass, B. L., Robotham, O. & Chapman, M. A. Evidence for two domestication events of hyacinth bean (Lablab purpureus (L.) Sweet): a comparative analysis of population genetic data. *Genet. Resour. Crop Evol.* **64**, 1221–1230 (2017).
17. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of Population Structure Using Multilocus Genotype Data. *Genetics* vol. 155 945–959 (2000).
18. Nei, M. Genetic distance between populations. *Am. Nat.* (1972).
19. Daszkowska-Golec, A. The landscape of plant genomics after 20 years. *Trends Genet.* **38**, 310–311 (2022).
20. Pengelly, B. C. & Maass, B. L. Lablab purpureus (L.) Sweet–diversity, potential use and determination of a core collection of this multi-purpose tropical legume. *Genet. Resour. Crop Evol.* (2001).
21. Linder, H. P. The evolution of African plant diversity. *Front. Ecol. Evol.* **2**, (2014).
22. Ou, S., Chen, J. & Jiang, N. Assessing genome assembly quality using the LTR Assembly Index (LAI). *Nucleic Acids Res.* **46**, e126 (2018).

23.   Phytozome v13. http://phytozome.jgi.doe.gov.

24.   Schmutz, J. *et al.* Genome sequence of the palaeopolyploid soybean. *Nature* **463**, 178–183
(2010).

25.   Jaillon, O. *et al.* The grapevine genome sequence suggests ancestral hexaploidization in major
angiosperm phyla. *Nature* **449**, 463–467 (2007).

26.   Argout, X. *et al.* The genome of Theobroma cacao. *Nat. Genet.* **43**, 101–108 (2011).

27.   Houston, K., Tucker, M. R., Chowdhury, J., Shirley, N. & Little, A. The Plant Cell Wall: A Complex
and Dynamic Structure As Revealed by the Responses of Genes under Stress Conditions. *Front.*
Plant Sci. **7**, 984 (2016).

28.   Wang, L. *et al.* A role for a cell wall localized glycine-rich protein in dehydration and rehydration
of the resurrection plant Boea hygrometrica. *Plant Biol.* **11**, 837–848 (2009).

29.   Hu, G. *et al.* Two divergent haplotypes from a highly heterozygous lychee genome suggest
independent domestication events for early and late-maturing cultivars. *Nat. Genet.* **54**, 73–83
(2022).

30.   Zhang, K. *et al.* Resequencing of global Tartary buckwheat accessions reveals multiple
domestication events and key loci associated with agronomic traits. *Genome Biol.* **22**, 23 (2021).

31.   Civáň, P., Craig, H., Cox, C. J. & Brown, T. A. Three geographically separate domestications of
Asian rice. *Nat Plants* **1**, 15164 (2015).

32.   Morrell, P. L. & Clegg, M. T. Genetic evidence for a second domestication of barley (Hordeum
vulgare) east of the Fertile Crescent. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3289–3294 (2007).

33.   Sultana, N., Ozaki, Y., Okubo, H. & Others. The use of RAPD markers in lablab bean (Lablab
purpureus (L.) Sweet) phylogeny. *Bulletin of the Institute of Tropical Agriculture, Kyushu*
University **23**, 45–51 (2000).

34.   Liu, C. J. Genetic diversity and relationships among Lablab purpureus genotypes evaluated
using RAPD as markers. *Euphytica* **90**, 115–119 (1996).

35.   Bohra, A., Jha, U. C., Kishor, P. B. K., Pandey, S. & Singh, N. P. Genomics and molecular
breeding in lesser explored pulse crops: current trends and future opportunities. *Biotechnol. Adv.*
32, 1410–1428 (2014).

36.   Habte, E., Gari, A., Lire, H. & Jones, C. Field trial of Lablab (Lablab purpureus) genotypes under
rain   fed   conditions   in   Ethiopia.
https://cgspace.cgiar.org/bitstream/handle/10568/116652/lablab_report.pdf?sequence=2 (2021).

37.   Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat. Biotechnol.*
35, 316–319 (2017).

38.   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads using
repeat graphs. *Nat. Biotechnol.* **37**, 540–546 (2019).

39.   Kundu, R., Casey, J. & Sung, W.-K. HyPo: Super Fast & Accurate Polisher for Long Read
Genome Assemblies. *bioRxiv* 2019.12.19.882506 (2019) doi:10.1101/2019.12.19.882506.

40.   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for genome
assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

41.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics **31**, 3210–3212 (2015).

42.   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

43.   Palmer, J. & Stajich, J. *nextgenusfs/funannotate: funannotate v1.5.3.* (2019).
doi:10.5281/zenodo.2604804.

44.   Haas, B. J. *et al.* Improving the Arabidopsis genome annotation using maximal transcript
alignment assemblies. *Nucleic Acids Res.* **31**, 5654–5666 (2003).

45.   Stanke, M., Steinkamp, R., Waack, S. & Morgenstern, B. AUGUSTUS: a web server for gene

531  finding in eukaryotes. *Nucleic Acids Res.* **32**, W309–12 (2004).

532 46. Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59 (2004).

533 47. Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: two open source ab
534  initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878–2879 (2004).

535 48. Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using EVidenceModeler and
536  the Program to Assemble Spliced Alignments. *Genome Biol.* **9**, R7 (2008).

537 49. Chan, P. P. & Lowe, T. M. tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences.
538  *Methods Mol. Biol.* **1962**, 1–14 (2019).

539 50. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment
540  and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

541 51. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq
542  reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

543 52. Gremme, G., Brendel, V., Sparks, M. E. & Kurtz, S. Engineering a software tool for gene
544  structure prediction in higher organisms. *Information and Software Technology* **47**, 965–978
545  (2005).

546 53. Haas, B., Papanicolaou, A. & Others. TransDecoder (find coding regions within transcripts).
547  *Google Scholar* (2016).

548 54. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

549 55. Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity
550  searching. *Nucleic Acids Res.* **39**, W29–37 (2011).

551 56. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–D419
552  (2021).

553 57. Blum, M. *et al.* The InterPro protein families and domains database: 20 years on. *Nucleic Acids*
554  *Res.* **49**, D344–D354 (2021).

555 58. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res.* **49**, D325–D334
556  (2021).

557 59. Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nature Genetics* vol. 25
558  25–29 (2000).

559 60. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a
560  streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

561 61. Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible software for de novo
562  detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

563 62. Ou, S. & Jiang, N. LTR_FINDER_parallel: parallelization of LTR_FINDER enabling rapid
564  identification of long terminal repeat retrotransposons. *Mob. DNA* **10**, 48 (2019).

565 63. Ou, S. & Jiang, N. LTR_retriever: A Highly Accurate and Sensitive Program for Identification of
566  Long Terminal Repeat Retrotransposons. *Plant Physiol.* **176**, 1410–1422 (2018).

567 64. Su, W., Gu, X. & Peterson, T. TIR-Learner, a New Ensemble Method for TIR Transposable
568  Element Annotation, Provides Evidence for Abundant New Transposable Elements in the Maize
569  Genome. *Mol. Plant* **12**, 447–460 (2019).

570 65. Xiong, W., He, L., Lai, J., Dooner, H. K. & Du, C. HelitronScanner uncovers a large overlooked
571  cache of Helitron transposons in many plant genomes. *Proc. Natl. Acad. Sci. U. S. A.* **111**,
572  10263–10268 (2014).

573 66. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable element
574  families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457 (2020).

575 67. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0. 2013--2015. (2015).

576 68. R Core Team. R: A Language and Environment for Statistical Computing. (2019).

577 69. Wickham, H. *et al.* Welcome to the tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

578 70. Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res.*
579  **27**, 573–580 (1999).

580   71. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
581   genomics. *Genome Biol.* **20**, 238 (2019).
582   72. Mendes, F. K., Vanderpool, D., Fulton, B. & Hahn, M. W. CAFE 5 models variation in
583   evolutionary   rates   among   gene   families.   *Bioinformatics*   (2020)
584   doi:10.1093/bioinformatics/btaa1022.
585   73. De Bie, T., Cristianini, N., Demuth, J. P. & Hahn, M. W. CAFE: a computational tool for the study
586   of gene family evolution. *Bioinformatics* **22**, 1269–1271 (2006).
587   74. Alexa, A. & Rahnenfuhrer, J. topGO: enrichment analysis for gene ontology. *R package version*
588   (2010).
589   75. Doyle, J. J. Isolation of plant DNA from fresh tissue. *Focus* **12**, 13–15 (1990).
590   76. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
591   data. *Bioinformatics* **30**, 2114–2120 (2014).
592   77. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
593   357–359 (2012).
594   78. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
595   (2009).
596   79. Picard toolkit. *Broad Institute, GitHub repository* (2019).
597   80. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).
598   81. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).
599   82. Kilian, A. *et al.* Diversity arrays technology: a generic genome profiling technology on open
600   platforms. *Methods Mol. Biol.* **888**, 67–89 (2012).
601   83. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage
602   analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
603   84. Earl, D. A. & vonHoldt, B. M. STRUCTURE HARVESTER: a website and program for visualizing
604   STRUCTURE output and implementing the Evanno method. *Conserv. Genet. Resour.* **4**, 359–
605   361 (2012).
606   85. Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of individuals using the
607   software STRUCTURE: a simulation study. *Mol. Ecol.* **14**, 2611–2620 (2005).
608   86. Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. Poppr: an R package for genetic analysis of
609   populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281 (2014).
610   87. Jombart, T. adegenet: a R package for the multivariate analysis of genetic markers.
611   *Bioinformatics* **24**, 1403–1405 (2008).
612   88. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R
613   language. *Bioinformatics* **20**, 289–290 (2004).
614   89. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
615   display and annotation. *Nucleic Acids Res.* **49**, W293–W296 (2021).
616

Acknowledgements

This research was conducted as part of the CGIAR Research Program on Livestock, supported by
CGIAR Fund Donors. O.S. was supported by the Royal Society FLAIR award (FLR_R1_191850),
N.K., H.G. and M.S. were supported by the German Federal Ministry of Education and Research
(De.NBI, FKZ 031A536). D.F. was supported by the SoCoBio DTP (grant number BB/T008768/1;
BBSRC, UK) to carry out a PhD rotation project in the lab of MAC. DF and MAC acknowledge the use
of the IRIDIS High Performance Computing Facility, and associated support services at the University
of Southampton.

Authors' contributions

O.S., P.M.F.E., J.D.E., M.A.C., M.S. and C.S.J. conceived and planned the experiments. C.M., L.M. and O.S. performed DNA extraction and Nanopore Sequencing. I.S., B.W., M.M. and D.K. performed the genome assembly. N.K., B.W., M.S. and I.S. performed genome annotation. D.F. and H.G. annotated the transposable elements and tandem repeats. N.K., B.W. and O.S performed gene family analyses, M.A.C. analysed the re-sequencing data. M.S.M., C.S.J. and B.L.M. performed diversity analyses on global collection. I.N., B.W., N.K., M.S.M., D.F., M.A.C., O.S. and C.S.J. wrote the manuscript. All authors reviewed and approved the final manuscript.

Competing interests

All authors declare that there are no competing interests.

Materials & Correspondence.

Mark Chapman (m.chapman@soton.ac.uk), Oluwaseyi Shorinola (shorinolao@gmail.com), Chris Jones (c.jones@cgiar.org).

Figures

Figure 1: Genome Assembly of Lablab. a, Lablab purpureus plant showing flowers, leaves and pods**; b**, Gene and repeat landscape of the lablab genome**.** The tracks from the outer to the inner track show 1) Gene density, 2) Repeat density, 3) LTR-RT density, 4) Tandem repeat density. **c**, LAI index of the 11 lablab chromosomes; **d**, BUSCO scores of the Lablab purpureus genome annotation using the embryophyta, eudicots and and fabales reference lineages.

Figure 2: Gene family evolution and expansion in *Lablab purpureus*. a, Venn diagram of the number of gene families common among and unique to *Lablab purpureus, Phaseolus vulgaris, Vigna angularis, Medicago truncatula,* and *Cajanus cajan.* **b,** Cladogram of the analysed species showing the number of expanded and contracted gene families in each. Figure constructed with iTol[89]. **c,** Gene ontology terms enriched in the set of expanded gene families in *Lablab purpureus*.

Figure 3: Phylogenetics of lablab and related legumes. Neighbor Joining phylogenetic relationships among lablab samples (2-seeded and 4-seeded purpureus (domesticated) and uncinatus (wild) subspecies) and other related legumes (see Table S8 for details). Tree is rooted on Macrotyloma. All nodes received full (100%) bootstrap support. Asterisks indicate the two domestication events.

Figure 4: Clusters and subclusters of the lablab accessions used in the diversity study. a, Bar plots based on the admixture model in STRUCTURE for K = 4 (Membership of individual accessions to each subgroup is given in Table S16). **b**, Clusters detected by hierarchical clustering. **c**, Clusters detected by PCA. The colours in **b** and **c** are according to the STRUCTURE analysis in **a**.

Supplementary Information

Supplementary Methods

Hi-C Scaffolding

Chromatin conformation capture data was generated by Phase Genomics (Seattle, USA) using the Proximo Hi-C 2.0 Kit, which is a commercially available version of the Hi-C protocol. Following the manufacturer's instructions for the kit, intact cells from two samples were crosslinked using a formaldehyde solution, digested using the DPNII restriction enzyme, end repaired with biotinylated nucleotides, and proximity ligated to create chimeric molecules composed of fragments from different regions of the genome that were physically proximal in vivo, but not necessarily genomically proximal. Continuing with the manufacturer's protocol, molecules were pulled down with streptavidin beads and processed into an Illumina-compatible sequencing library. Sequencing was performed on an Illumina HiSeq, generating a total of 232,382,372 PE150 read pairs.

Reads were aligned to the draft assembly using BWA-MEM[90] with the -5SP and -t 8 options specified, and all other options default. SAMBLASTER[91] was used to flag PCR duplicates, which were later excluded from analysis. Alignments were then filtered with SAMtools[77] using the -F 2304 filtering flag to remove non-primary and secondary alignments. Putative misjoined contigs were broken using Juicebox[92] based on the Hi-C alignments. A total of 6 breaks in 6 contigs were introduced. The same alignment procedure was repeated from the beginning on the resulting corrected assembly.

Phase Genomics Proximo Hi-C genome scaffolding platform was used to create chromosome-scale scaffolds from the corrected assembly as described in Bickhart et al.[93]. As in the LACHESIS method[94], this process computes a contact frequency matrix from the aligned Hi-C read pairs, normalized by the number of DPNII restriction sites (GATC) on each contig, and constructs scaffolds in such a way as to optimize expected contact frequency and other statistical patterns in Hi-C data. Approximately 20,000 separate Proximo runs were performed to optimize the number of scaffolds and scaffold construction in order to make the scaffolds as concordant with the observed Hi-C data as possible. This process resulted in a set of 11 chromosome-scale scaffolds containing 417 Mbp of sequence (98% of the corrected assembly) with a scaffold N50 of 38.1 Mbp.

Synteny-guided Chromosome naming

We adopted a naming scheme based on synteny with closely related legumes - *P. vulgaris* (common

695 bean[13]) and *V. unguiculata* (cowpea[14]). For this, we downloaded protein sequence and GFF files of
696 PacBio-based assembly of *P. vulgaris* (v2.1) and *V. unguiculata* (v1.2) from Phytozome[23] and
697 compared this separately to lablab proteins using BLASTP[54] (settings: -max_target_seqs 1, -evalue
698 1e-10, -qcov_hsp_perc 70). MCScanX[95] was used to process the individual BLAST output and to
699 detect inter-species collinear blocks.

Filtering for true-to-type genotypes in global genebank collection.

The lablab accessions used for evaluating global diversity in this study were acquired from different
sources and conserved *ex situ* as seeds in the ILRI forage genebank, the earliest since 1982, with
periodic monitoring for viability and regeneration for renewal of the seeds. These periodic genebank
management practices involve risks to the genetic integrity of the accessions through pollen
contamination, seed contamination, segregation, mislabeling, and other factors (e.g. as described in
Chebotar et al., 2003 [96]). Hence, it was necessary to ensure the genetic integrity of plants within
accessions and avoid potential contaminants before the genetic diversity analysis. Using pairwise IBD
(Identity-By-Descent) analysis, plants within accessions were classified into "true-to-type", "progeny",
or "contaminant" based on a PI_HAT[83] value of above 0.80, between 0.125 and 0.80, or less than
0.125, respectively. Six accessions with a single plant each were excluded from the analysis.

For nine accessions, all plants were unrelated to each other, and therefore considered
"contaminants". Out of the remaining 151 accessions, 85 were 100% true-to-type, indicating that there
was no cross-pollination or seed mixing. Twenty-four accessions had a mixture of true-to-type and
their progeny, indicating that some level of cross-pollination or segregation had taken place in this
group. Another 24 accessions had a mixture of true-to-type and contaminants, and other 18
accessions had a mixture of the true-to-type, their progenies, and contaminants (Figure S6). After
removing contaminants, a total of 1680 plants were retained from these 151 accessions for genetic
diversity analysis. Of these, 1541 plants were true-to-type with 2 to 26 plants per accession, and 139
were progenies from 41 accessions (1 to 12 plants per accession).

Analysing historical lablab phenotype datasets

Phenotypic variation among the identified major molecular groups was assessed based on historical
data summarised by Pengelly and Maass (2001)[20] (127 accessions) and Wiedow (2001)[97] (95
accessions), in which morpho-agronomic traits on lablab accessions were evaluated in field trials at
Ziway site in Ethiopia, in 1998 and 2000, respectively. Seventeen accessions were analysed in both
trials, hence we could determine whether traits varied across the seasons. Where variation was low
(correlation between seasons was 80% or greater; 6 traits), data from the two trials were combined.
For the remaining 15 traits, only the 1998 phenotype data on 75 accessions was used for the analysis
of trait variation among the four genetic groups identified above. Analysis of variance (ANOVA) and
Tukey's multiple comparison test were employed to compare phenotypic variation of agro-
morphological quantitative traits with significant p values ($P < 0.01$) among clusters identified by
population structure analysis. A chi-square test was used for similar comparisons among clusters in
qualitative traits.

Supplemental References

90. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, 589–595 (2010).
91. Faust, G. G. & Hall, I. M. SAMBLASTER: fast duplicate marking and structural variant read extraction. *Bioinformatics* **30**, 2503–2505 (2014).
92. Durand, N. C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Syst* **3**, 99–101 (2016).
93. Bickhart, D. M. *et al.* Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome. *Nat. Genet.* **49**, 643–650 (2017).
94. Burton, J. N. *et al.* Chromosome-scale scaffolding of de novo genome assemblies based on chromatin interactions. *Nat. Biotechnol.* **31**, 1119–1125 (2013).
95. Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res.* **40**, e49 (2012).
96. Chebotar, S. *et al.* Molecular studies on genetic integrity of open-pollinating species rye (Secale cereale L.) after long-term genebank maintenance. *Theor. Appl. Genet.* **107**, 1469–1476 (2003).
97. Wiedow, C. Morphological and agronomical characterization of Lablab purpureus in Ethiopia. (Diploma thesis (Dipl.-Ing. agr.), University of Rostock, 2001).

Supplementary Figures

Figure S1: Chromosome-level synteny of *Lablab purpureus* with related species. *L. purpureus* chromosomes have been named according to synteny with *P. vulgaris* **(a)** and *V. unguiculata* **(b)** chromosomes.

Figure S2: Chromosomal repeat content in *Lablab purpureus*. (a) Relative densities of repeat elements along each chromosome. 1) Long Terminal Repeat RetroTransposons (LTR-RT), 2) Tandem Inverted Repeats (TIR), 3) Miniature Inverted Transposable Elements (MITE), 4) Helitron 5) Unclassified repeats, 6) Tandem repeats **(b)** Proportional abundance of identified transposable element orders on each chromosome.

Figure S3: GBS polymorphism in global lablab collection: Genome-wide distribution of SNPs (**a**) and SilicoDArT (**b**) markers across the eleven chromosomes of the lablab reference genome. The total number of SNPs or SilicoDArT markers are presented beside each chromosome. Plots produced with SRplot.

Figure S4: Quantitative phenotypic variation in global lablab collection. Boxplots showing phenotypic variation of different morpho-agronomic quantitative traits among the four genetic clusters identified in lablab**.** The colours are according to the STRUCTURE analysis with k = 4, and trait abbreviations are explained in Table S13.

Figure S5: Qualitative phenotypic variation in global lablab collection. Plots showing phenotypic variation of seven qualitative traits among the four genetic clusters identified in lablab. The colours are according to the STRUCTURE analysis with k = 4, and trait abbreviations are explained in Table S14. Points are scattered if identical values are present.

780 **Figure S6: Identity-By-Descent classification of global lablab collection.** Number of accessions
781 classified as true-to-type (TTT), true-to-type and progenies (TTT+NP), true-to-type and contaminants
782 (TTT+Cont), true-to-type and progenies and contaminants (TTT+NP+Cont), and accessions with
783 100% contaminants (Cont)**,** based on a pairwise Identity-By-Descent (IBD) analysis.

785 Supplementary Tables

Table S1: Summary of Nanopore Reads Statistics
Table S2: Table S2: Comparison of assembly statistics for the lablab genome based on short reads
and long reads.
Table S3: Summary statistics of genes in the lablab genome.
Table S4: The number of TEs, TE families and the proportion of occupied assembly length by different
classes of repeats identified and annotated in the lablab genome.
Table S5: Types, amount and proportion of tandem repeats in the lablab genome
Table S6: GO annotation of lablab-specific gene clusters
Table S7: GO annotation of gene families expanded in lablab
Table S8: Details and sequencing statistics of resequencing samples
Table S9: Population group membership
Table S10. Pairwise Fixation index (Fst) among the four major clusters (C) detected by the
STRUCTURE analysis
Table S11: AMOVA showing the genetic variance among and within clusters
Table S12: Minimum, maximum and average genetic divergence (Nei's D) between accessions within
the four clusters identified by STRUCTURE.
Table S13: Results of the analysis of variance for 13 quantitative traits among the four genetic
clusters.
Table S14: Results of the χ2 analysis for seven quantitative traits among the four genetic clusters.
Table S15: Data on inclusive crop genomics
Table S16: Membership probability of accessions from the STRUCTURE analysis
807

Figure 1: Genome Assembly of Lablab. a, Lablab purpureus plant showing flowers, leaves and pods. **b**, Gene and repeat landscape of the lablab genome. The tracks from the outer to the inner track show 1) Gene density, 2) Repeat density, 3) LTR-RT density, 4) Tandem repeat density. **c**, LAI index of the 11 lablab chromosomes. **d**, BUSCO scores of the Lablab purpureus genome annotation using the embryophyta, eudicots and and fabales reference lineages.

Figure 2: Gene family evolution and expansion in *Lablab purpureus*. a, Venn diagram of the number of gene families common among and unique to *Lablab purpureus, Phaseolus vulgaris, Vigna angularis, Medicago truncatula,* and *Cajanus cajan.* **b,** Cladogram of the analysed species showing the number of expanded and contracted gene families in each. Figure constructed with iTol[89]. **c,** Gene ontology terms enriched in the set of expanded gene families in *Lablab purpureus*.

Figure 3: Phylogenetics of lablab and related legumes. Neighbor Joining phylogenetic relationships among lablab samples (2-seeded and 4-seeded purpureus (domesticated) and uncinatus (wild) subspecies) and other related legumes (see Table S8 for details). Tree is rooted on *Macrotyloma*. All nodes received full (100%) bootstrap support. Asterisks indicate the two domestication events.

Figure 4: Clusters and subclusters of the lablab accessions used in the diversity study. a, Bar plots based on the admixture model in STRUCTURE for K = 4 (Membership of individual accessions to each subgroup is given in Table S16). **b**, Clusters detected by hierarchical clustering. **c**, Clusters detected by PCA. The colours in **b** and **c** are according to the STRUCTURE analysis in **a**.

Figure S1: Chromosome-level synteny of *Lablab purpureus* with related species. *L. purpureus* chromosomes have been named according to synteny with *P. vulgaris* **(a)** and *V. unguiculata* **(b)** chromosomes.

Figure S2: Chromosomal repeat content in *Lablab purpureus*. (a) Relative densities of repeat elements along each chromosome. 1) Long Terminal Repeat RetroTransposons (LTR-RT), 2) Tandem Inverted Repeats (TIR), 3) Miniature Inverted Transposable Elements (MITE), 4) Helitron 5) Unclassified repeats, 6) Tandem repeats **(b)** Proportional abundance of identified transposable element orders on each chromosome.

Figure S3: GBS polymorphism in global lablab collection: Genome-wide distribution of SNPs (**a**) and SilicoDArT (**b**) markers across the eleven chromosomes of the lablab reference genome. The total number of SNPs or SilicoDArT markers are presented beside each chromosome. Plots produced with SRplot.

Figure S4: Quantitative phenotypic variation in global lablab collection. Boxplots showing phenotypic variation of different morpho-agronomic quantitative traits among the four genetic clusters identified in lablab. The colours are according to the STRUCTURE analysis with k = 4, and trait abbreviations are explained in Table S13.

Figure S5: Qualitative phenotypic variation in global lablab collection. Plots showing phenotypic variation of seven qualitative traits among the four genetic clusters identified in lablab. The colours are according to the STRUCTURE analysis with k = 4, and trait abbreviations are explained in Table S14. Points are scattered if identical values are present.

Figure S6: Identity-By-Descent classification of global lablab collection. Number of accessions classified as true-to-type (TTT), true-to-type and progenies (TTT+NP), true-to-type and contaminants (TTT+Cont), true-to-type and progenies and contaminants (TTT+NP+Cont), and accessions with 100% contaminants (Cont), based on a pairwise Identity-By-Descent (IBD) analysis.