Primate sympatry shapes the evolution of their brain architecture

Benjamin Robira[1,2,✉], and Benoît Perez-Lamarque[3,4]

[1] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France.
[2] Eco-anthropologie (EA), Muséum National d'Histoire Naturelle, CNRS, Université de Paris, Musée de l'Homme, Paris, France.
[3] Institut de Biologie de l'École Normale Supérieure (IBENS), École Normale Supérieure, CNRS, INSERM, Université PSL, Paris, France.
[4] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France.

[✉] Correspondence: **Benjamin Robira <benjamin.robira@normalesup.org>**

Abstract | The main hypotheses on the evolution of animal cognition emphasise the role of conspecifics in affecting the socio-ecological environment shaping cognition. Yet, space is often simultaneously occupied by multiple species from the same ecological guild. These sympatric species can compete for food, which may thereby stimulate or hamper cognition. Considering brain size as a proxy for cognition, we tested whether species sympatry impacted the evolution of cognition in frugivorous primates. We first retraced the evolutionary history of sympatry between frugivorous primate lineages. We then fitted phylogenetic models of the evolution of the size of several brain regions in frugivorous primates, considering or not species sympatry. We found that the whole brain or brain regions used in immediate information processing were best fitted by models not considering sympatry. By contrast, models considering species sympatry best predicted the evolution of brain regions related to a long-term memory of interactions with the socio-ecological environment, with a decrease of their size the higher the sympatry. We speculate that species sympatry, by generating intense food depletion, might lead to an over-complexification of resource spatiotemporality that counteract the benefits of high cognitive abilities and/or might drive niche partitioning and specialisation, thereby inducing lower brain region sizes. In addition, we reported that primate species in sympatry diversify more slowly. This comparative study suggests that species sympatry significantly contributes to shaping primate evolution.

29   **Short title:** Sympatry shapes primates' brain architecture

30   **Keywords:** Brain size - Cognition - Diversification - Frugivory - Primates - Sympatry

INTRODUCTION

Cognition evolution is shaped by the balance between socio-ecological drivers promoting cognitive abilities (González-Forero and Gardner 2018) and physiological and energetic constraints limiting them (Navarrete, van Schaik, and Isler 2011). Primates are pivotal species for cognitive studies (Byrne 2000) because their cognition (i.e. the set of mechanisms that enable them to perceive, learn and memorise information, and make decisions, Shettleworth 2010) is thought to be promoted by interactions of individuals with conspecifics within the social unit (Byrne 2018; Dunbar and Shultz 2017), among generations (Wilson 1991; Whiten and van Schaik 2007; Reader and Laland 2002; Herrmann et al. 2007; Tomasello 2019; van Schaik and Burkart 2011), between social units (Ashton, Kennedy, and Radford 2020), or with the rest of their environment (Clutton☐Brock and Harvey 1980; Milton 1981; Rosati 2017). However, space is often occupied by many primate species, which may have dietary overlaps and compete for the same resources (Kamilar and Ledogar 2011). Because of direct and indirect interactions linked to resource availability, the presence of heterospecifics is also likely to shape the evolution of species cognition.

Retracing the evolutionary history of cognitive abilities proves to be challenging because there is still no consensual measurement for cognition applicable across all species. Up to now, a raw approximation consists in considering the (relative) brain size as a proxy for cognitive abilities, with larger sizes considered equivalent to more advanced cognitive abilities (e.g. in mammalian carnivores: Benson-Amram et al. 2016; or primates: MacLean et al. 2014). Although the relevance of this assumption is heavily limited within species, in part because of plasticity (Gonda, Herczeg, and Merilä 2013), this holds when comparing different species within taxonomically similar groups (e.g. in primates, Reader and Laland 2002). In addition, instead of considering the brain as a whole, the multifaceted aspect of animal cognition is more precisely depicted by appreciating the mosaic nature of the brain (Barton and Harvey 2000). For instance, variations in the size of some specific brain regions have been robustly associated

55   with variations in cognition related to the functions of these regions (Healy and Rowe 2007). The brain is

56   a patchwork of areas cognitively specialised that may thus follow different evolutionary trajectories.

57   Because the coexistence of primate species in the same biogeographic area, henceforth referred to

as sympatry, can affect multiple aspects of their social-ecological environment, brain regions may be

differently affected by sympatry. Considering a simplistic functional picture of the primate brain, several

hypotheses can be proposed on how species sympatry might, through direct or indirect competition or

cooperation, influence the relative benefits of cognition (i.e. the balance between its benefits and its costs).

As such, selective pressures upon cognitive abilities would affect the energy allocation toward the whole

brain and/or the within-brain allocation toward the different brain regions, resulting in changes of their

sizes relatively to the whole body and/or the whole brain, respectively.

65   First, long-term memory stands as a valuable tool to infer food availability and location when

food is rare and ephemeral but predictable (Milton 1981; Rosati 2017). Having several sympatric species

competing for the same food resource often leads to an increase in food depletion of the shared resource

compared with an environment with only one foraging species (Minot 1981). Such depletion may

therefore complexify the patterns of resource distribution and availability in space and time. This

complexification may in turn affect the selective pressures upon brain regions involved in the long-term

storing of spatiotemporal information, such as the hippocampus (Burgess, Maguire, and O'Keefe 2002,

Hypothesis 1: long-term memory is affected by sympatry). On the one hand, under reasonable food

depletion, having a better memory should be advantageous to better predict food availability and location.

In addition, resource competition between heterospecifics should promote anticipatory behaviour, hence

high cognition, as expected for within-species competition (Ashton, Kennedy, and Radford 2020). Thus,

the relative size of the hippocampus, reflecting long-term memory abilities, should be larger the higher the

sympatry (Prediction 1.1). On the other hand, intense food depletion may also increase environmental

unpredictability to a point where long-term memory is not advantageous anymore. In addition, even when

the dietary overlap between heterospecifics is inexistent, foragers may rely on phenological cross-

correlations between plant species (e.g. if plant species A produces fruits earlier in the season than plant

species B, thus the availability of B fruits can be predicted based on the availability of A fruits, even

though A is not consumed by the forager, Janmaat et al. 2012; Robira et al. 2021): Intense food depletion

may thus weaken these cross-correlations and make food availability predictions less reliable, therefore

limiting the benefits of memory (Robira et al. 2021). Due to the energy constraints of maintaining a large

brain, the hippocampus relative size could thus be smaller in species experiencing high levels of sympatry

(Prediction 1.2). Furthermore, the competition between sympatric species from a given dietary guild may

foster their specialisation toward different food resources, i.e. niche partitioning, which may impact

cognition (Aristide et al. 2016). While a species might specialise toward food that is difficult to access and

requires high cognition (e.g. through the use of tools), specialisation is generally associated with reduced

flexibility and thus lower cognitive abilities (Henke-von der Malsburg, Kappeler, and Fichtel 2020).

Therefore, because of cognitive specialisation requiring less long-term memory abilities, the hippocampus

relative size could also be smaller in sympatric species (Prediction 1.2). Alternatively, intense levels of

sympatry could also lead to disruptive selection toward cognitive abilities, with the most successful

competitors that may evolve enhanced cognition, while bad competitors may not. In this context, the

hippocampus relative sizes may either increase or decrease for the different sympatric species (Prediction

1.3; character displacement).

Second, sympatric species may enrich the landscape of visual, olfactory, or acoustic cues usable to

locate available food (e.g. Avarguès-Weber, Dawson, and Chittka 2013; Kashetsky, Avgar, and Dukas

2021). Consequently, such cues left out by heterospecifics may impact the selective pressures upon brain

regions involved in processing more immediate sensory information, such as the Main Olfactory Bulb

(MOB), the cerebellum (Koziol et al. 2014; Sokolov, Miall, and Ivry 2017), and the neocortex (Wiltgen et

al. 2004) (Hypothesis 2: cue processing is affected by sympatry). Hence, sympatry could be associated

with larger relative sizes of the MOB, the cerebellum, or the neocortex (Prediction 2). As brain regions

may be interconnected, this enlargement could be detrimental to other brain regions. For instance,

increased corticalisation could be associated with smaller hippocampus sizes (Kitamura et al., 2017).

Third, besides indirect interactions through foraging, cognition can also be triggered by direct

"social" interactions with other individuals (Byrne 2018; Dunbar and Shultz 2017), such as with the

formation of mixed-group species (Goodale et al. 2010). The striatum, a brain region stimulated during

social interactions (Báez-Mendoza and Schultz 2013), or to a lesser extent, the hippocampus (Todorov et

al. 2019), may therefore be affected by the increase of proximity, hence direct social interactions, between

heterospecifics (Hypothesis 3: sociality is affected by sympatry), leading to a larger relative size of the

striatum or hippocampus in sympatry (Prediction 3).

In any case, the evolution of brain size in primates likely impacted their dynamic of species

diversification. Larger brain sizes are indeed found to be associated with higher diversification rates in

birds (Sayol et al. 2019) and similar patterns have been suggested in primates (Melchionna et al. 2020).

However, it remains unclear how brain size and diversification are interlinked in the context of sympatry.

Here, we investigated whether species sympatry affected the evolution of cognition using

frugivorous primates as a study example. Frugivorous primates are an interesting group for such a

question because fruit is the archetype of a hard-to-find resource yet predictable (Janmaat et al. 2016), for

which cognition considerably shapes the foraging strategy (Trapanese et al. 2019). To infer the effect of

species sympatry on cognition in frugivorous primates, we evaluated the support for phylogenetic models

of brain size evolution accounting or not for species sympatry and investigated the directionality of the

selection induced by sympatry on brain size evolution. Finally, we tested for correlative patterns between

brain size or current sympatry and species diversification in all primates, to better understand the impact

of cognition and interactions between primates on their evolutionary success.

METHODS

Data processing, analyses, and plots were computed with R software (v.4.1.2, R Core Team 2020). The list and versions of packages are available in Supplementary Material, R packages used.

Data Collection

Trait data

We pooled data from previous literature surveys (Pearce et al. 2013; Willems, Hellriegel, and van Schaik 2013; Grueter 2015; DeCasien, Williams, and Higham 2017; Powell, Isler, and Barton 2017; Navarrete et al. 2018; DeCasien and Higham, 2019; Powell, Barton, and Street 2019; Todorov et al. 2019, see Supplementary Material, Trait data collection). From the global endocranial brain volume, we obtained the Encephalization Quotient (EQ, $N_{EQ,max} = 182$) such as $EQ = 1.036 \times \text{Brain size}/(0.085 \times \text{Body mass}^{0.775})$ with the brain size in cm³, 1.036 g/cm³ being the assumed homogeneous brain density, and the body mass in g. EQ indicates whether the brain size ranges above (> 1) or below (< 1) what is expected given the body mass and correcting for allometry (DeCasien, Williams, and Higham 2017). We also focused on specific regions of the brain which were chosen because they were involved in immediate sensory information processing (MOB, $N_{MOB,max} = 39$), in movement and/or general information processing and retention (neocortex, $N_{Neocortex,max} = 69$, Wiltgen et al. 2004; cerebellum, $N_{Cerebellum,max} = 70$, Koziol et al. 2014; Sokolov, Miall, and Ivry 2017), or short-term working memory and long-term spatiotemporal memory (hippocampus, $N_{Hippocampus,max} = 63$, Burgess, Maguire, and O'Keefe 2002). The striatum ($N_{Striatum,max} = 63$) supports information processing during social interaction, reward assessment, planning, or goal-oriented behaviours (Báez-Mendoza and Schultz 2013; Johnson, Meer, and Redish 2007). To investigate their evolutionary history, we first used the ratio between their volume and body mass. As such, the use of brain region sizes relative to the body mass and not raw sizes depicts the evolution of cognitive abilities in terms of differential allocation rather than

abilities per se (but see discussion in Deaner, Nunn, and van Schaik 2000). Second, we repeated the

analyses considering the ratio between the brain region size and the whole-brain size, as this might reflect

within brain energy reallocation.

For each primate species, broad dietary guilds (frugivores, folivores, nectarivores, granivores,

insectivores, omnivores) are generally considered based on the main type of food they consume. We

classified extant species as either "frugivorous" or "folivorous" based on the availability of percentage of

frugivory and folivory, prioritising frugivory over folivory. We considered a species as frugivorous if the

percentage of frugivory was at least above 20%. If not, or if the percentage of frugivory was unavailable,

we considered the species as folivorous if the percentage of folivory was above 40%. Whenever the

percentages were not available, we referred to the classification provided by DeCasien, Williams, and

Higham (2017), partly based on anatomical criteria. We discarded any other cases (e.g. species feeding on

invertebrates or tree exudates). We also replicated these diet assignments by considering a threshold of

40% for frugivory and 60% for folivory.

Ranging Data

The current biogeographic range of each primate species was assessed using ranging maps provided by the

IUCN red list (IUCN 2021). Ranging data were available for 249 species. The availability of distribution

range data is depicted in Supplementary Figure S1.

Phylogeny

We used chronogram trees of the primate taxon of the 10kTrees project (downloaded on May 2021,

version 3), as well as a consensus tree of 1000 trees for the subsequent phylogenetic analyses. The trees

contain 301 primate species. Note that in all these analyses, we discarded *Homo sapiens* and *Macaca*

sylvanus. The latter was discarded because of its complete geographic isolation and repeated intervention

of humans in population maintenance (Modolo, Salzburger, and Martin 2005). A summary of available

data per species is presented in Supplementary Figure S1.

Retracing past sympatry between primate species

Based on the biogeographic distribution of each extant primate species, we first reconstructed the history of past sympatry between primate lineages. To do so, we followed Drury et al. (2018) and first reconstructed the biogeographic history of each primate lineage to then retrace which pairs of primate lineages were likely to be simultaneously present at the same place. Leaning on Kamilar (2009), we considered that the biogeography of primates can be described by 12 discrete biogeographic areas with highly similar community structures shaped by both the environment geography and climatic correlates. These geographic areas, mapped using Google Earth Professional (v7.3.3), are represented in Figure 1. One to multiple biogeographic areas were assigned to each species as soon as 10% of their current distribution range overlapped on the surface with a given biogeographic area (see Supplementary Material, Spatial overlap assessment, Figure 2).

Given these 12 biogeographic areas, we retraced the biogeographic history of primates with the *BioGeoBEARS* package (Matzke 2013), using the biogeographic stochastic mapping algorithm (Matzke 2016). We fitted non-time-stratified dispersal-extinction-cladogenesis (DEC) models. We fixed to three biogeographic areas the maximum number of areas that a lineage can simultaneously occupy since it offers the possibility to occupy a complete mainland continent while keeping computational time reasonable. We sampled 10 histories of primate biogeographic ranges to account for the uncertainty. We assumed that two primate lineages were in sympatry at a given time whenever these two lineages simultaneously occupied the same biogeographic area. We also replicated these biogeographic assignments using instead a larger threshold of 30% of overlap (instead of 10%; see Supplementary Figure S2 for a quantification of variations in biogeographic assignments due to the use of different thresholds).

Inferring past diets of primate lineages

Next, we retraced the evolutionary history of frugivorous lineages in primates. Considering diet as a binary variable, we retraced the evolutionary history of frugivory versus folivory using extended Mk

models (Bollback 2006) implemented in the "simmap" function of the *phytools* package (Revell 2012).

We performed ancestral diet reconstructions using both combinations of dietary thresholds (20/40% and

40/60% for frugivory and folivory, respectively) and sampled 10 stochastic mapping of ancestral diet

reconstructions per run. These ancestral reconstructions were used in combination with the histories of

primate biogeographic ranges to assess whether a pair of frugivorous species was in sympatry or not

(Drury et al. 2018): we thus obtained reconstructions of the evolutionary history of sympatry between

frugivorous primate lineages.

Phylogenetic models of trait evolution: does species sympatry shape brain size evolution?

We assessed the effect of sympatry on evolution of frugivorous primate brain size using two approaches.

First, we used phylogenetic models of trait evolution to assess the role of sympatry on the evolution of

brain sizes. Second, we investigated how sympatry has influenced brain size evolution (i.e. selection

towards smaller or larger brain sizes) by evaluating correlations between current levels of sympatry and

brain sizes using linear models. In both approaches, we considered either the whole brain (using the EQ)

or the size of the aforementioned specific brain regions relative (i) to the body mass or (ii) to the whole-

brain size. Models testing the effect of species sympatry on one brain region relative to the body mass

(models (i)) assess whether the total allocation towards a specific brain region is affected by species

sympatry. In contrast, models testing the effect of species sympatry on one brain region relative to the

whole-brain size (models (ii)) assess whether the within-brain energy reallocation is affected by species

sympatry. Before fitting, relative brain sizes were log-transformed to account for allometry. Although it

does not consider the potential inter-dependence between brain regions, we modelled independently each

brain region to preserve the maximum power (i.e. the largest sample size) in each analysis.

(a) Fitting models of trait evolution

For each brain region, we independently fitted different phylogenetic models of trait evolution. Following

Drury et al. (2016), we considered three stochastic models accounting for sympatry, which expand

Brownian motion (BM) models of trait evolution. We used the "fit_t_comp" function from the *RPANDA*

package (Morlon et al. 2016) to fit: a matching competition (MC) model (Nuismer and Harmon 2015) and

linear and exponential density-dependent models (DD_{lin} and DD_{exp}, Drury et al. 2016). The MC model

considers the repulsion of brain sizes of sympatric frugivorous lineages due to competition (character

displacement), that is $z_i(t + dt) = z_i(t) + S(\mu(t) - z_i(t))dt + \sigma dB_i$ where z is the brain size of a

species i at time t, μ is the mean brain size of all sympatric species, S is a negative value that reflects the

strength of the effect of species sympatry, and σdB_i is the drift (BM) with a constant evolutionary rate σ

(Drury et al. 2016). Linear (DD_{lin}) or exponential (DD_{exp}) density-dependences (Drury et al. 2016; Weir

and Mursleen 2013) assume that the evolutionary rate, σ_l, of trait evolution, varies either positively or

negatively as a function f of the number of frugivorous sympatric lineages, such as $\sigma_l = f_{lin}(l) = \sigma_0 +$

rl and $\sigma_l = f_{exp}(l) = \sigma_0 \exp(rl)$, where σ_0 corresponds to the ancestral rate value, l indicates the number

of lineages at a given time, and r controls the speed and direction of the dependency of the rate to the

number of lineages ($r > 0$ leads to an increased rate of trait evolution in sympatry, while $r < 0$ leads to a

slowdown of trait evolution in sympatry, as observed in adaptive radiations).

We compared the support of models considering species sympatry to the support of simpler

models assuming no effect of species sympatry on the evolution of brain sizes: the Brownian motion

(BM), the Ornstein-Uhlenbeck process (OU, a model with an optimum value, see Blomberg, Rathnayake,

and Moreau 2020, for a review), or the Early-Burst model (EB), for assessing a time-dependence of the

evolutionary rate, irrespectively of the levels of species sympatry (Blomberg, Garland, and Ives 2003). We

fitted these models using the "fitContinuous" function from the *geiger* package (Slater et al. 2012; Pennell

et al. 2014). We compared the six models accounting for sympatry or not within an information-theoretic

framework (Burnham and Anderson 2002), based on the weights of Akaike Information Criteria corrected

for small samples (AICc). The model weight depicts how well the model fits the observed data compared

with the other tested models (i.e. a model set).

To account for uncertainty (in the data, Supplementary Figure S3, and fit), we fitted and compared

these models (within each model set) 10 times for 10 different ancestral reconstructions of primate

biogeography and diet considering all thresholds. Thus, for each brain region, we fitted all the models 10

(uncertainty on diet/biogeography ancestral reconstructions x 10 (uncertainty in brain/diet data) x 2

(biogeographic overlap threshold) x 2 (frugivory threshold) x 2 (folivory threshold) = 800 times. We

nonetheless stopped computations when the calculation of the likelihood was excessively long (> 1 week)

and the final sample size was of 730 model sets.

(b) Determining the effect of sympatry on brain sizes

Diversity-dependent models of trait evolution inform whether species sympatry has impacted brain size

evolution by increasing or decreasing the tempo of trait evolution. Nonetheless, they do not tell the

directionality of the effect (i.e. do brain sizes increase or decrease with sympatry?). To determine whether

species sympatry positively or negatively affected the sizes of brain regions, we independently fitted

Gaussian Pagel's lambda phylogenetic regressions for each brain region of extant frugivorous species.

This model is a derivative of the BM model, where the phylogenetic variance-covariance matrix has all

coefficients, but its diagonal ones, multiplied by lambda (therefore relaxing the hypothesis of BM). To fit

these models, we used a frequentist-based approach with the "phylolm" function from the *phylolm*

package (Ho and Ane 2014). We considered the least stringent frugivory assessment, with the frugivory

threshold fixed at 20% and the folivory threshold fixed at 40%.

The response variable was the log-transformed relative size of each brain region relative to the

body mass or the whole-brain size. Doing so controlled for allometry while being in line with the

phylogenetic models of trait evolution, which cannot account for additional variables and are thus limited

to the use of relative brain sizes. Due to data variability, we took the mean of the possible values given the

different datasets and assessed the sensitivity using non-averaged values (see Supplementary Material,

Model stability). We used as covariates (i.e. continuous predictors) two explicit measures of the level of

species sympatry for each extant frugivorous species: (1) the number of frugivorous sympatric species

(square-rooted to reach symmetrical distribution to limit leverage effects) and (2) the average percentage

of the overlapping current range (assessed based on IUCN data) with other sympatric frugivorous species.

For a given species A, sympatry with another species B was considered when at least 10% of the range of

species A overlaps with the range of species B. This was done to reduce the noise induced by coarse

identification of species range.

Body mass and sympatry

277   Body mass may itself be affected by species sympatry. Yet, as body mass was used to compute some

278   relative brain size, to control for the potential confounding effect due to a relationship between sympatry

279   and body mass, we repeated all model fitting (models of trait evolution and PGLS) with body mass as the

280   output variable.

Models of species diversification

Next, we also investigated the correlations between primate diversification rates and brain sizes or levels

of sympatry to better understand the impact of cognition and interactions between primates on their

evolutionary success. To do so, we inferred how primates diversified over time and across lineages.

Lineage-specific net diversification rates (defined as speciation minus extinction rates) were estimated

using an updated version of the *ClaDS* algorithm (Maliet, Hartig, and Morlon 2019) based on data

augmentation techniques (Maliet and Morlon 2021). Particularly, we used *ClaDS2*, a model with constant

turnover (i.e. constant ratio between extinction and speciation rates; see Supplementary Material, Primate

diversification rate over time, for further explanations). We extracted the mean diversification rates

through time and the lineage-specific diversification rate of each extant species.

We fitted Gaussian Pagel's lambda phylogenetic regressions of the different relative brain sizes

against the net diversification rates. Because assumptions for a frequentist-based approach were unmet, we

performed Bayesian-based inferences instead. We used the "MCMCglmm" function of the *MCMCglmm*

package (Hadfield 2010). Each chain had a burn-in period of 5000 iterations, a total length of 50,000

iterations, and was sampled every 50 iterations. We used the least informative priors. Fixed priors were set

to default values (Gaussian distribution of mean 0 and variance 10^8). Again, we used the mean of the

brain trait values for the main model and assessed the sensitivity by re-running the model several times

using non-averaged values.

To determine whether species sympatry was associated with lower or larger diversification rates,

we fitted frequentist-based Gaussian Pagel's lambda phylogenetic regressions with the lineage-specific

diversification rate as the output variable and used the two metrics for describing sympatry (the number of

frugivorous sympatric species and the average percentage of overlapping range with other sympatric

frugivorous species) as the tested variables.

Model implementation and stability

305   Details on the implementation, stability, and uncertainty of phylogenetic regressions are provided in

306   Supplementary Material, Model stability). Necessary assumptions on the normal distribution of residuals

307   and homoscedasticity were visually assessed and pointed to no violation (see Supplementary Material,

308   Model assumptions). In addition, we did not observe correlation issues among predictors (Variance

309   Inflation Factor, $\text{VIF}_{max} < 2$, Mundry 2014).

RESULTS

The dataset we gathered contained between 34 to 182 frugivorous primate species (depending on the brain

region considered). When considering the sizes relative to the body mass of the different brain regions, we

observed ample variations (Figure 3). For instance, the lemuriformes, which are known to prioritize smell

compared with other primate species, have the largest relative MOB size (lemuriformes: mean $\pm$ SE =

0.23 $\pm$ 0.07, other: 0.12 $\pm$ 0.04, 3). Similarly, the highest relative sizes of the striatum were found in

platyrrhini (platyrrhini: mean $\pm$ SE = 0.91 $\pm$ 0.07, other: 0.59 $\pm$ 0.07, 3), which are known to form poly-

specific associations (callitrichines in particular, Heymann and Buchanan-Smith 2000). In terms of

sympatry, we observed that on average ($\pm$ SE) the considered primate species had 52% ($\pm$ 2) of their

range overlapping with other species. That ranged from 0% of overlap (*Macaca nigra*), to 100% of

overlap (*Cercopithecus pogonias, Alouatta pigra, Loris tardigradus, Hylobates moloch, Cercocebus*

galeritus, Presbytis melalophos, Semnopithecus entellus). In terms of the distribution range, the

considered primate species co-occurred on average with 6.38 ($\pm$ 0.39) other primate species, ranging from

0 other species to 21.

To retrace past species sympatry between frugivorous lineages, we reconstructed primate

biogeographic history (Figure 1) and their diet evolution. We found that the ancestors of primates were

likely to be frugivorous and that folivory evolved more than five times, in particular in cercopithecidae

and memuridae. Estimated diet history was consistent with fossil evidence (see Supplementary Material,

Diet reconstruction, Supplementary Figure S4). We then modelled the evolution of the size of the whole

brain (EQ) or separate brain regions (neocortex, cerebellum, MOB, hippocampus, and striatum) when

considering species sympatry or not. Weighting each brain region by the whole-body mass and weighting

by whole-brain size yielded similar results (see Supplementary Material, Weighting the size of brain

regions by whole-brain size, Tables S1 and S4, Figures S5 and S6), therefore we only present the case

when weighting by body mass. We found that models not considering species sympatry best described the

evolutionary history of the EQ, the neocortex, and the cerebellum (Figures 3 and 4), two brain regions

specifically involved in immediate sensory information processing (Wiltgen et al. 2004; Koziol et al.

2014; Sokolov, Miall, and Ivry 2017), and also in memory consolidation for the neocortex (Wiltgen et al.

2004). The fact that these biggest brain regions are best described by an Ornstein-Uhlenbeck process

suggests a stabilisation towards an optimal relative size, which may illustrate the trade-off between the

costs and benefits of brain development (Isler and van Schaik 2009). Yet, given the broad range of the

study species, and the likely difference of their environments, this could hide a potentially multi-peak

stabilisation that occurred towards different optima for different clades (Mahler et al. 2013). By contrast,

density-dependent models considering species sympatry (DD_{lin} and DD_{exp}) were best supported in the

foraging-related and social-related brain regions respectively: the hippocampus, specialised in

spatiotemporal memory (Burgess, Maguire, and O'Keefe 2002), and the striatum, involved in social

interactions (Báez-Mendoza and Schultz 2013). The fact that we inferred positive rates r of density-

dependence (Figure 4) suggested an acceleration of the evolutionary tempo of trait evolution together with

increased number of frugivorous sympatric lineages for the hippocampus and the striatum. The MOB, the

brain region involved in sensory abilities, also tended to be best fitted by models considering sympatry as

a whole. Yet, Brownian motion (BM) was as likely as density-dependent or matching competition models,

preventing firm conclusions on whether sympatry affected or not MOB size evolution (Figures 3 and 4),

especially since this coincided with the most reduced sample size we had ($N_{species}$ = 34 to 39).

Next, we assessed whether higher levels of species sympatry led to "bigger" or "smaller" brain

regions. To do so, we fitted phylogenetic regressions in extant frugivorous species between the relative

sizes of the different brain regions and two measures of sympatry (1) the average percentage of

overlapping range with other frugivorous sympatric species, and (2) the number of such sympatric

frugivorous species across their current entire distribution range. When weighting each brain region by

body mass, the number of sympatric species never significantly influenced the relative brain sizes (Table

1; Supplementary Figures S7 and S10). Conversely, we found that the average percentage of overlapping

range significantly correlated, or tended to correlate, with the relative size of brain regions that were better

fitted with models considering sympatry: the hippocampus and the striatum (hippocampus: t = -1.94, p =

0.058; striatum: t = -2.26, p = 0.028). The correlations were all negative (hippocampus: est. = -0.39,

CI95% = [-0.76,-0.01]; striatum: est. = -0.4, CI95% = [-0.77,-0.04]), which means that higher range

overlap between sympatric species associates with lower relative size, insensitively to data and

phylogenetic uncertainties (Supplementary Table S3, Supplementary Figures S7 and S10). The

acceleration of the tempo of trait evolution with species sympatry ($r > 0$ in the density-dependent models)

suggests that, compared with isolated species, sympatric species are subject to a positive selection towards

smaller brains, and not to a less intense selection for advanced cognitive abilities.

Furthermore, we also investigated the influence of sympatry on body mass, which could be a

confounding explanation for the observed patterns (Smaers et al. 2021). We found that the evolutionary

history of body mass is best explained by MC models (Supplementary Figure S6). In other words, species

in sympatry tend to diverge towards being large or small. There is thus no overall linear relationship

between body mass and sympatry, explaining why we found no effect of sympatry variables on body mass

in PGLS; Supplementary Tables S2 and S7).

Finally, we investigated the evolutionary consequences of cognition and species sympatry by

evaluating whether brain sizes and levels of sympatry correlated with the lineage-specific net

diversification rates of primates (Figure 5). Overall, species diversification rates particularly boomed in

the early and late Miocene, around 25 and 11 Myr ago (Figure 5). When accounting for phylogenetic

dependence, no significant relationship between the net diversification rate and the relative size of brain

regions was found in extant primates (Table 2, Supplementary Figure S8; see robustness in Supplementary

Table S5). Although diversification was uncorrelated with brain size in frugivorous primates, it was

influenced by the sympatry context. In particular, phylogenetic regressions highlighted a negative effect of

the number of sympatric species on the diversification rate (est. = -5.04e-03, CI95% = [-0.01,1.34e-04], t

= 2.56e-03, p < 0.001, Table 3, Supplementary Figure S9, see robustness in Supplementary Table S6). In

other words, the higher the number of sympatric species, the lower the diversification rate.

DISCUSSION

Sympatry of frugivorous primate species impacts the evolution of their hippocampus and striatum sizes

Although developmental constraints can induce correlation in the evolution of different brain regions (Gómez-Robles, Hopkins, and Sherwood 2014), brain regions underpin different cognitive functions and can thus be under different, independent, selective pressures (Barton and Harvey 2000). The functional regionalisation is for instance evidenced here by the differences in relative sizes across lineages in the MOB, with larger sizes in the lemuriformes that mostly rely on smell to forage. Our analyses also highlighted such differences in evolutionary trajectories for the different brain regions by the variations in the best-fit models of size evolution. We indeed show that sympatry is one factor that affects the evolutionary regime under which only some brain regions evolve: although the brain as a whole (i.e., Encephalization Quotient, EQ), the cerebellum and the neocortex were not significantly affected by species sympatry, the latter nonetheless induced a change in the relative sizes of the hippocampus and the striatum.

Our results were similar whether the relative sizes were obtained by weighting by body mass or whole-brain size. Theoretically, the two weighting methods are expected to give insights into differences in resource allocation between body tissues or differences in resource allocation within the brain tissue, respectively. Nonetheless, we observed no differences in the response to sympatry by weighting by body mass or whole-brain size, despite weak to moderate correlations between sizes relative to the body mass and sizes relative to the whole-brain size (see Supplementary Figure S5). Therefore, our results suggest that species sympatry mostly affected within-brain energy partitioning (as weighting by body mass can depict reallocation changes towards the whole brain or only within the brain), and in particular reallocations to two relatively small brain regions: the hippocampus and the striatum.

The hippocampus and the striatum are brain regions involved in individual-based and social-based

information processing, pinpointing that these two components might be under strong selection in

primates (DeCasien, Williams, and Higham 2017; Powell, Isler, and Barton 2017; González-Forero and

Gardner 2018). The hippocampus in particular, may have played a considerable role in the evolution of

primate-like behaviours (Schilder, Petry, and Hof 2020), driven by the changes that primates faced in their

ecological environment (e.g. the spatiotemporal distribution of the food, DeCasien and Higham 2019), or

by the social environment that they faced (e.g. the number of conspecifics they interact with, Todorov et

al. 2019; DeCasien and Higham 2019).

The fact that the hippocampus, particularly relevant to processing and memorising spatiotemporal

information, is negatively sensitive to sympatry, is consistent with the idea of an effect of sympatric

species on resource spatiotemporality (Hypothesis 1). Competition is generally the first-thought factor to

describe community structures (de Almeida Rocha et al. 2015) because it might affect the environment in

which species evolve. We show that a higher level of sympatry is associated with smaller sizes of the

hippocampus (following Prediction 1.2), while the neocortex and cerebellum are not, at odd with the

hypothesis that this is the mere consequence of corticalisation (Wiltgen et al., 2004). This suggests that

indirect competition for food might contribute to convoluting the environment, and such an over-

complexification of the resource spatiotemporality may render cognitive foraging not advantageous

anymore. As a result, it might even generate a selection for smaller brain regions involved in foraging. In

parallel with the complexification of their environment, species might have also narrowed frugivorous

primates' niches, which might have synergistically affected their cognition. Indeed, the support for

matching competition when modelling the evolution of the body mass of frugivorous primates indicates

that sympatric primate species tended to diverge in terms of body mass, evolving towards lower or higher

body mass in sympatry. This is consistent with the idea of niche partitioning in sympatry, e.g. where light

primate species would occupy the canopy layer, while heavier primate species would occupy the ground.

Such niche partitioning may be accompanied by dietary specialisation (Schreier et al. 2009) and impact

cognitive abilities (Aristide et al. 2016), as dietary specialisation often requires lower cognitive abilities

and thus smaller brain region sizes.

In contrast to direct/indirect competition, potential indirect facilitation between species due to

"social" cues (Hypothesis 2), is ruled out by the absence of an effect of sympatry on brain regions

involved in immediate sensory information processing (e.g. cerebellum or neocortex). This absence of

effect can stem from two possibilities. Either foragers do not exploit cues left out by sympatric

heterospecifics, or they discarded environment cues in favour of social cues, as it has been shown that

foragers tend to use social information over environmental (i.e. personal) information, in particular in non-

perfectly predictable environments (Rafacz and Templeton 2003; Dunlap et al. 2016). In both cases,

stimulation intensity of the MOB, the cerebellum, or the neocortex would somehow remain equivalent

when in sympatry or not. Further work should explicitly test for these possibilities.

As expected (Hypothesis 3), the striatum size was relatively larger in callitrichines, particularly

known to form mixed-species groups (Heymann and Hsia 2015). Yet, overall, the striatum size was

negatively affected by sympatry. This puzzle might take root in secondary, but key, functions supported

by the striatum, namely reward expectation, goal-directed behaviour, and planning abilities (Johnson,

Meer, and Redish 2007). These three functions might as well be advantageous when foraging. As for the

hippocampus, then, the increase in environmental unpredictability could diminish the benefits of these

future-oriented skills.

Sympatry of frugivorous primate species is associated with slower diversification

Given the context-dependence of the direction of selection (towards bigger sizes when sympatry is low,

smaller sizes otherwise), there is no surprise that we do not observe a correlation between the net

diversification rate and the three brain regions affected by species sympatry. Surprisingly however, we

found no positive association between the net diversification rate and the EQ, the cerebellum, or the

neocortex, which were insensitive to species sympatry. By contrast, a positive association between brain

size and diversification was also found in birds (Sayol et al. 2019) given that bigger brains act as a buffer

to environmental challenges (Sol et al. 2007). A visual inspection of the regressions evidenced a positive

trend if not considering phylogeny (EQ and neocortex, Supplementary Figure S8). Sudden encephalisation

in primates is clearly associated with a limited number of closely related species (DeCasien, Williams, and

Higham 2017; Melchionna et al. 2020). Thus, this limits the statistical power of our analyses accounting

for phylogenetically relationships, as we cannot decipher whether larger brain size and faster species

diversification result from a true biological link or appeared simultaneously but independently. This

means that a positive association between brain size and species diversification remains a likely possibility

(as previously suggested in primates, Melchionna et al. 2020). Species sympatry, however, induced a

significant slowdown in primate diversification, a density-dependence trend frequently observed in many

tetrapod clades (Condamine, Rolland, and Morlon 2019). This frames coherently with a competitive

scenario, where the tempo of species diversification decreases when ecological niches are progressively

filled up (Rabosky and Lovette 2008). Species competing for resources are thought to contribute to

limiting competitors' range (Price and Kirkpatrick 2009), hence constraining population size and

diversification rate (Pigot and Tobias 2013).

Limits

The use of brain size as a proxy for cognition is a central debate with no optimal solution (see grounded

criticism from Deaner, Nunn, and van Schaik 2000; Healy and Rowe 2007; Logan et al. 2018; van Schaik

et al. 2021). The current flourishment of consortia, allowing for much more detailed and standardised

anatomical measurements (e.g. in primates: Milham et al. 2018), or standardised behaviourally-explicit

comparisons (e.g. on captive, Many Primates et al. 2019; or wild primates, Janmaat et al. 2021) might

alleviate biases stemming from brain size analysis, but this will take time to generate large-enough

datasets. In the meanwhile, brain size is a proxy much appreciated in practice, because of its easy

accessibility for a "large" number of species, while the multifaceted aspect of cognition can simply be

taken into account by considering the brain as a mosaic of singular and independent regionalised areas that

are cognitively specialised. Yet, this approach implies collating datasets from different sources, which can

induce a high intra- and inter-species variability, as the methodology has changed over time (discussed in

Navarrete et al. 2018). We thus sampled measures among the available datasets and repeated the analysis

several times to account for the variability of the data sets. The results remained robust to this approach.

This supports the idea that intra-species variation is negligible compared with inter-species variation, a

necessary condition for brain size to be an honest signal of cognitive differences across species, despite

differences in measurement protocols.

Not only can the nature of the data be biased, but the methods themselves can suffer from

significant limitations. In our case, these limitations are particularly true for the reconstructions of diet

history, ancestral biogeography, or species diversification that we performed, which infer the most likely

evolutionary histories based on observations at present alone. To verify the realism of these inferences, we

confronted the diet history reconstruction with fossil evidence. Dental microwear textures in tooth fossils

can indeed be used to reconstruct past diets and we found that our estimates of ancestral primate diets

were consistent with the available fossils (Ramdarshan, Merceron, and Marivaux 2012; Merceron et al.

2009, see Supplementary Figure S4). Similarly, we found that our estimates of species diversification rates

from molecular phylogenies were consistent with estimates of past primate diversity reconstructed from

all available fossil data (Springer et al. 2012). In addition, the identified breakpoints matched estimates

from previous studies (Arbour and Santana 2017; Springer et al. 2012). We did not use fossils for the

biogeographic reconstruction because, although this may help the reconstruction, the overall benefices are

minor (Wisniewski, Lloyd, and Slater 2022) and highly influenced by the fossil data quality (e.g. spatial

coverage), which is still heavily limited and biased. We nevertheless accounted for the uncertainty in all

reconstructions in our analyses by reporting results based on multiple sets of ancestral reconstructions.

Furthermore, the effectiveness of a method also depends on the proxy and associated definitions,

which may impact the insights of such large-scale analyses. For instance, this issue is clearly illustrated

with group size being used as a proxy of social complexity in social cognition (Dunbar and Shultz 2007),

despite social complexity being multifactorial. This rejoins our major assumption of brain size as an

honest signal for cognition, a multifaceted trait which is likely to be affected by diverse genes. In this

study, further simplifications were considered: brain regions were associated with one main function, an

overlap of diets was considered within the whole frugivorous primate guild, and sympatry was considered

to be based as soon as spatial overlap occurred. Instead, each brain region can be considered as a Russian

doll, with possibly redundant sub-functionalisation between spatially distant regions, frugivorous primates

can be differentially selective in the fruits they eat (Campera et al. 2019), and species may not necessarily

encounter each other at fine spatiotemporal scales (Deane et al. 2013). Although we cannot exclude that

more accurate definitions might change our results, it is important to note that such a simplification is

imposed by many constraints: from the sake of theorising to computational time. In addition, these large

definitions also have some advantages. First, they can capture a broad variety of situations. For instance,

simply defining sympatry as co-occurrence in a given biogeographic area enables considering both, direct

and indirect interactions. Second, more stringent definitions might induce strong initial constraints that

would eventually bias inferences of past history. For instance, the considered range areas to define

sympatry are large and are unlikely to capture smaller-scale spatial segregation between species (e.g. due

to habitat-specific segregation or vertical segregation). Yet, a finer mapping of species range may poorly

reflect the "realised niche," as primate populations suffer intense local extinction in recent years due to

human activity (Estrada et al. 2017; Pavoine et al. 2019). Furthermore, we also took care of varying the

stringency of our definitions (e.g. thresholds used to consider a species frugivorous based on the

percentage of frugivory, thresholds used to consider two species as sympatric based on range overlap) and

results were robust to such arbitrary choices (see Supplementary Material, Model stability).

CONCLUSION

We showed that species sympatry is an important factor shaping the brain evolutionary history of frugivorous primates. It now seems crucial to scrutinise more carefully how sympatry fits with other socio-ecological variables that also influence brain size evolution in this clade (e.g. diet, group size, home range, etc., see DeCasien, Williams, and Higham 2017; DeCasien and Higham 2019; Powell, Isler, and Barton 2017). In addition, dietary overlap and food competition might not only happen between frugivorous primates, but between any frugivores in the same area. In fact, it is very likely that any hypotheses on cognition evolution, generally discussed within species, could be broadened to a between-species context: foraging facilitation between species does exist (Olupot, Waser, and Chapman 1998; Havmøller et al. 2021), and so do polyspecific social associations (Porter 2001), as well as inter-species territory and resource defence (Drury, Cowen, and Grether 2020; Losin et al. 2016) or imitation and copying (Persson, Sauciuc, and Madsen 2018; Pepperberg 2002). Similarly, prey-predator races could shape selection on cognitive abilities (Shultz and Dunbar 2006). As Alice said, "It's a great huge game of chess that's being played—all over the world" (Carroll 1871, chap. II) and all individuals are just pieces to play with or against.

ACKNOWLEDGEMENTS

We considerably value the help provided by J. Drury in making some scripts available functions in the *RPANDA* package in *R* and helping us run them, and that of M-C. Quidoz for assistance in using the CEFE cluster. We thank S. Benhamou and M. Clairbaux for discussions on spatial projections, M. Quéroué, V. Lauret, A. Caizergues, and C. Teplistky for feedback on Bayesian computations, and H. Morlon for discussions on models of trait evolution. We also acknowledge the Editor F. Condamine, as well as P. Gonzalez, O. S. Todorov, and three other anonymous reviewers for their constructive comments on our manuscript that helped us improve it. Finally, this work could not have been possible without prior

data collection from the IUCN Red List (primate ranging), the 10kTrees project (phylogenetic trees), and

A. R. DeCasien and collaborators, L. E. Powell and collaborators, O. S. Todorov and collaborators, E. P.

Willems and collaborators, F. Pearce and collaborators, A. F. Navarrete and collaborators, and C. C.

Grueter who provided primate trait data we used as (supplementary) material with their articles. Their

indirect input is therefore tremendous.

AUTHORS' CONTRIBUTION

BR conceived the study, gathered, cleaned, and analysed the data, drew the figures, wrote the first version

of the manuscript, and subsequently revised it. BP-L implemented the ClaDS algorithm for our data,

helped with running other analyses and drawing figures, and revised the manuscript multiple times.

FUNDING

Both authors were supported by a doctoral grant from the *École Normale Supérieure*, Paris.

CONFLICT OF INTEREST

The authors declare having no conflict of interest. All authors gave final approval for publication and

agree to be held accountable for the work performed therein.

DATA, SCRIPT AND CODE AVAILABILITY

The data and codes that support the findings of this study are openly available at

https://github.com/benjaminrobira/Meta_analysis_cognition_primates.

Table 1: Species sympatry correlates negatively with the relative size of some brain regions in extant frugivorous primate species | Model estimates and significance of phylogenetic regressions to assess the relationship between relative brain sizes and species sympatry. Est.=Estimate, CI2.5%=Lower border of the CI95%, CI97.5%=Upper border of the CI95%, Se=Standard error, t=Statistics t-value. The brain regions (as well as the associated sample sizes) are indicated before each list of estimates. The transformations applied to variables are indicated between parentheses (logarithm, log, or square-root, sqrt), as well as the weighting by body mass (/body mass).

	Est.	CI2.5%	CI97.5%	Se	t	p-value
EQ (log) (N=127)						
Intercept	-0.17	-0.53	0.22	0.20	-	-
% of overlapped range	0.02	-0.08	0.13	0.05	0.41	0.68
Number of sympatric frugivores (sqrt)	0.02	-0.02	0.05	0.02	1.03	0.31
Lambda	0.98	0.94	1.00			
Hippocampus (/body mass, log) (N=50)						
Intercept	-0.92	-1.95	0.05	0.53	-	-
% of overlapped range	-0.39	-0.76	-0.01	0.20	-1.94	0.06
Number of sympatric frugivores (sqrt)	0.08	-0.06	0.20	0.07	1.21	0.23
Lambda	0.99	0.92	1.00			
Neocortex (/body mass, log) (N=56)						
Intercept	2.07	1.31	2.86	0.41	-	-
% of overlapped range	-0.23	-0.54	0.11	0.16	-1.46	0.15
Number of sympatric frugivores (sqrt)	0.02	-0.08	0.13	0.05	0.48	0.63
Lambda	0.99	0.91	1.00			

Cerebellum (/body mass, log) (N=57)

Intercept	0.60	-0.15	1.35	0.39	-	-
% of overlapped range	-0.08	-0.32	0.17	0.12	-0.7	0.49
Number of sympatric frugivores (sqrt)	-0.01	-0.1	0.07	0.04	-0.34	0.74
Lambda	1.00	0.96	1.00			

Striatum (/body mass, log) (N=50)

Intercept	-0.36	-1.18	0.44	0.44	-	-
% of overlapped range	**-0.40**	**-0.77**	**-0.04**	**0.18**	**-2.26**	**0.03**
Number of sympatric frugivores (sqrt)	0.03	-0.08	0.15	0.06	0.61	0.54
Lambda	0.98	0.85	1.00			

MOB (/body mass, log) (N=31)

Intercept	-2.76	-4.61	-0.93	1.00	-	-
% of overlapped range	-1.20	-2.65	0.35	0.80	-1.49	0.15
Number of sympatric frugivores (sqrt)	0.21	-0.18	0.56	0.19	1.12	0.27
Lambda	1.00	1e-07	1.00			

Table 2: Relative brain sizes do not correlate with primate species diversification in frugivorous primates | Model
estimates and significance of Bayesian phylogenetic regressions to assess the correlation between the net diversification
rates and the relative brain sizes. Est.=Estimate, HDP2.5%=Lower border of the 95% Highest Posterior Density,
HDP97.5%=Upper border of the 95% Highest Posterior Density, Eff. Samp.=Effective sample (adjusted for
autocorrelation). The brain regions (as well as the associated sample sizes) are indicated before each list of estimates. The
(log) indicates log-transformed variables, while the (/body mass) indicates variables weighted by body mass.

	Est.	HDP2.5%	HDP97.5%	Eff. Samp.	pMCMC
Diversification EQ (N=148)					
Intercept	0.12	0.08	0.16	900.00	-
EQ (log)	0.02	-7.91e-03	0.05	789.25	0.15
Lambda	0.83	0.76	0.9		
Diversification Hippocampus (N=61)					
Intercept	0.13	0.09	0.18	900.00	-
Hippocampus (/body mass, log)	9.10e-03	-9.48e-03	0.03	900.00	0.34
Lambda	0.73	0.6	0.85		
Diversification Neocortex (N=67)					
Intercept	0.1	0.04	0.17	991.53	-
Neocortex (/body mass, log)	7.26e-03	-0.02	0.03	900.00	0.56
Lambda	0.74	0.6	0.86		
Diversification Cerebellum (N=68)					
Intercept	0.12	0.07	0.16	900.00	-
Cerebellum (/body mass, log)	3.94e-03	-0.02	0.03	989.21	0.76
Lambda	0.74	0.6	0.86		

Diversification Striatum (N=61)					
Intercept	0.12	0.08	0.17	900.00	-
Striatum (/body mass, log)	9.11e-03	-0.01	0.03	900.00	0.44
Lambda	0.73	0.59	0.85		
Diversification MOB (N=37)					
Intercept	0.11	0.05	0.17	900.00	-
MOB (/body mass, log)	-4.79e-03	-0.02	0.01	900.00	0.59
Lambda	0.65	0.46	0.83		

	Est.	**CI2.5%**	**CI97.5%**	**Se**	**t**	**p-value**
Diversification Sympatry (N=128)						
Intercept	0.15	0.10	0.2	0.03	-	-
% of overlapped range	-5.40e-03	-0.02	9.35e-03	8.14e-03	-0.66	0.51
Number of sympatric frugivores (sqrt)	**-5.04e-03**	**-0.01**	**1.34e-04**	**2.56e-03**	**-1.97**	**0.05**
Lambda	0.96	0.89	0.99			

Figure 1: Biogeographic areas used for reconstructing the history of sympatry in frugivorous primates represented on the
Mercator projection of the world | Areas were defined as a combination of geographic and environmental criteria relative
to the primate taxonomy following results from Kamilar (2009): (1) East Madagascar (2) West Madagascar (3) West
Africa (4) Central Africa (5) East/South Africa (6) Central America (7) North South-America (8) South South-America (9)
West Asia (10) Central/East Asia (11) South Asia (12) Asian peninsula and islands. Note that northern Africa and
southern Europe were discarded because *Macaca sylvanus* was not considered.

Figure 2: The level of species sympatry varies across the primate phylogeny | Primate phylogeny from the consensus tree
of the 10kTrees project is depicted in the center, together with abbreviated species names. The corresponding non-
abbreviated names can be found using Supplementary Figure S1. Sympatric frugivorous (based on a frugivory threshold
of 20% and folivory threshold of 40%) species are linked by light grey lines. The biogeographic areas occupied by a
species are depicted by coloured rectangles. Presence was assessed given an overlap between the species range and the
biogeographic area of 10%.

Figure 3: Variations in relative brain region sizes among extant frugivorous primates | (Left) Circular plot of the sizes of the different brain regions relative to the body mass. Colours indicate the rows for the different brain regions. The darker background emphasises when values are above average, while the lighter background emphasises when values are below average. The mean value (after scaling and based on one random sampling among possible values but see Supplementary Figure S3 for visualisation of measure variability) for the Encephalization Quotient (EQ) or relative size of brain regions, when available, is depicted by a plain circle for frugivorous species. The frugivorous threshold was fixed at 20% and the folivory threshold at 40%. **(Right)** The different studied brain regions (human brain as an illustration). In short, the MOB is involved in immediate olfactory information processing, the neocortex and the cerebellum support working memory and memory consolidation of immediate sensory information processing (Wiltgen et al. 2004; Koziol et al. 2014; Sokolov, Miall, and Ivry 2017), and the hippocampus supports a working memory and a long-term spatiotemporal memory (Burgess, Maguire, and O'Keefe 2002). The striatum is involved in social information processing (Báez-Mendoza and Schultz 2013).

Figure 4: The evolution of the hippocampus and striatum in frugivorous primates are best fitted by models of trait evolution considering species sympatry | Plotted is the AICc weight, a measure of relative support for a given model, for models not considering species sympatry (BM, OU, EB) or considering species sympatry (MC, DD_{lin}, DD_{exp}). The points represent the average AICc weight obtained (when considering the six models from the same run), while the vertical bars indicate the standard deviation given all tested conditions (see Phylogenetic models of trait evolution: does species sympatry shape brain size evolution?).

Figure 5: Species sympatry in primates is negatively associated with species diversification | a) Net diversification rate over time in the primate taxon. The average diversification rate estimated based on an assumed sampling fraction of primate species ranging from 60 to 95% (at a step of 10%; then 5% from 90%) is depicted by the plain line. The grey background depicts the standard deviation. The two breakpoints, depicted by the plain dots and the vertical dotted bars, were estimated based on a three-linear regression segmentation using the *strucchange* package [Zeileis et al. (2002); Zeileis et al. (2003); Zeileis (2006); see the vignette package for statistical details]. The three fitted regressions are displayed by the dashed lines. The choice of two breakpoints was first assessed by choosing the number of breakpoints minimising the Bayesian Information Criterion. b) The branches of the primate phylogenetic trees are colored according to their speciation rates estimated using ClaDS2: red colors indicate lineages that speciate frequently, whereas blue colors indicate lineages that rarely speciate. At the tips of the tree, we reported for each extant species whether we had data on its biogeography (column 'Data availability') and if so, we indicated the number of sympatric species it has. Using PGLS, we found a significant relationship between lineage-specific diversification rates and the numbers of sympatric species at present.

REFERENCES

Arbour, J. H., and S. E. Santana. 2017. A major shift in diversification rate helps explain macroevolutionary patterns in primate species diversity. Evolution 71:1600–1613. Wiley Online Library.

Aristide, L., S. F. Dos Reis, A. C. Machado, I. Lima, R. T. Lopes, and S. I. Perez. 2016. Brain shape convergence in the adaptive radiation of new world monkeys. Proceedings of the National Academy of Sciences 113:2158–2163. National Acad Sciences.

Ashton, B. J., P. Kennedy, and A. N. Radford. 2020. Interactions with conspecific outsiders as drivers of cognitive evolution. Nature Communications 11:1–9. Nature Publishing Group.

Avarguès-Weber, A., E. H. Dawson, and L. Chittka. 2013. Mechanisms of social learning across species boundaries. 290:1–11.

Báez-Mendoza, R., and W. Schultz. 2013. The role of the striatum in social behavior. Frontiers in Neuroscience 7:233. Frontiers.

Barton, R. A., and P. H. Harvey. 2000. Mosaic evolution of brain structure in mammals. Nature 405:1055–1058. Nature Publishing Group.

Benson-Amram, S., B. Dantzer, G. Stricker, E. M. Swanson, and K. E. Holekamp. 2016. Brain size predicts problem-solving ability in mammalian carnivores. Proceedings of the National Academy of Sciences of the United States of America 113:2532–2537. National Academy of Sciences.

Blomberg, S. P., T. Garland, and A. R. Ives. 2003. Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. Evolution 57:717–745. Wiley Online Library.

Blomberg, S. P., S. I. Rathnayake, and C. M. Moreau. 2020. Beyond brownian motion and the ornstein-uhlenbeck process: Stochastic diffusion models for the evolution of quantitative characters. American Naturalist 195:145–165. The University of Chicago Press Chicago, IL.

Bollback, J. P. 2006. SIMMAP: Stochastic character mapping of discrete traits on phylogenies. BMC Bioinformatics 7:1–7. BioMed Central.

Burgess, N., E. A. Maguire, and J. O'Keefe. 2002. The human hippocampus and spatial and episodic memory. Neuron 35:625–641. Elsevier.

Burnham, K. P., and D. R. Anderson. 2002. Model selection and multimodel inference: A practical Information-theoretic approach. Second. Springer, New York.

Byrne, R. W. 2000. Evolution of primate cognition. Cognitive Science 24:543–570. Wiley Online Library.

Byrne, R. W. 2018. Machiavellian intelligence retrospective. Oxford University Press Oxford.

Campera, M., M. Balestri, M. Chimienti, V. Nijman, K. Nekaris, and G. Donati. 2019. Temporal niche separation between the two ecologically similar nocturnal primates avahi meridionalis and lepilemur fleuretae. Behavioral Ecology and Sociobiology 73:1–12. Springer.

Carroll, L. 1871. Through the looking glass. Macmillan London U.K.

Clutton‐Brock, T. H., and P. H. Harvey. 1980. Primates, brains and ecology. Journal of Zoology 190:309–323. Wiley Online Library.

Condamine, F. L., J. Rolland, and H. Morlon. 2019. Assessing the causes of diversification slowdowns: temperature-dependent and diversity-dependent models receive equivalent support. Ecology Letters 22:1900–1912. John Wiley & Sons, Ltd.

de Almeida Rocha, J. M., M. P. Pinto, J. P. Boubli, and C. E. Viveiros Grelle. 2015. The role of competition in structuring primate communities under different productivity regimes in the amazon. PLoS ONE 10:e0145699. Public Library of Science San Francisco, CA USA.

Deane, A. S., M. C. Nargolwalla, L. Kordos, and D. R. Begun. 2013. New evidence for diet and niche partitioning in rudapithecus and anapithecus from rudabánya, hungary. Journal of human evolution 65:704–714. Elsevier.

Deaner, R. O., C. L. Nunn, and C. P. van Schaik. 2000. Comparative tests of primate cognition: Different scaling methods produce different results. Brain, Behavior and Evolution 55:44–52. Karger Publishers.

DeCasien, A. R., and J. P. Higham. 2019. Primate mosaic brain evolution reflects selection on sensory and cognitive specialization. Nature Ecology and Evolution 3:1483–1493. Nature Publishing Group.

DeCasien, A. R., S. A. Williams, and J. P. Higham. 2017. Primate brain size is predicted by diet but not sociality. Nature Ecology and Evolution 1:1–7. Nature Publishing Group.

Drury, J. P., J. Clavel, M. Manceau, and H. Morlon. 2016. Estimating the effect of competition on trait evolution using maximum likelihood inference. Systematic Biology 65:700–710. Oxford University Press.

Drury, J. P., M. C. Cowen, and G. F. Grether. 2020. Competition and hybridization drive interspecific territoriality in birds. Proceedings of the National Academy of Sciences of the United States of America 117:12923–12930. National Acad Sciences.

Drury, J. P., J. A. Tobias, K. J. Burns, N. A. Mason, A. J. Shultz, and H. Morlon. 2018. Contrasting impacts of competition on ecological and social trait evolution in songbirds. PLoS Biology 16:e2003563. Public Library of Science San Francisco, CA USA.

Dunbar, R. I. M., and S. Shultz. 2007. Evolution in the social brain. Science 317:1344–1347. American Association for the Advancement of Science.

Dunbar, R. I. M., and S. Shultz. 2017. Why are there so many explanations for primate brain evolution? Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160244. The Royal Society.

Dunlap, A. S., M. E. Nielsen, A. Dornhaus, and D. R. Papaj. 2016. Foraging bumble bees weigh the reliability of personal and social information. Current Biology 26:1195–1199. Elsevier.

Estrada, A., P. A. Garber, A. B. Rylands, C. Roos, E. Fernandez-Duque, A. D. Fiore, K. Anne-Isola Nekaris, V. Nijman, E. W. Heymann, J. E. Lambert, F. Rovero, C. Barelli, J. M. Setchell, T. R. Gillespie, R. A. Mittermeier, L. V. Arregoitia, M. de Guinea, S. Gouveia, R. Dobrovolski, S. Shanee, N. Shanee, S. A. Boyle, A. Fuentes, K. C. MacKinnon, K. R. Amato, A. L. S. Meyer, S. Wich, R. W. Sussman, R. Pan, I. Kone, and B. Li. 2017. Impending extinction crisis of the world's primates: Why primates matter. Science Advances 3:e1600946. American Association for the Advancement of Science.

Gómez-Robles, A., W. D. Hopkins, and C. C. Sherwood. 2014. Modular structure facilitates mosaic evolution of the brain in chimpanzees and humans. Nature Communications 5:1–9. Nature Publishing Group.

Gonda, A., G. Herczeg, and J. Merilä. 2013. Evolutionary ecology of intraspecific brain size variation: A review. Ecology and Evolution 3:2751–2764. John Wiley; Sons Ltd.

González-Forero, M., and A. Gardner. 2018. Inference of ecological and social drivers of human brain-size evolution. Nature 557:554–557. Nature Publishing Group.

Goodale, E., G. Beauchamp, R. D. Magrath, J. C. Nieh, and G. D. Ruxton. 2010. Interspecific information transfer influences animal community structure. Trends in Ecology and Evolution 25:354–361. Elsevier Current Trends.

Grueter, C. C. 2015. Home range overlap as a driver of intelligence in primates. American Journal of Primatology 77:418–424. Wiley Online Library.

Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear mixed models: The MCMCglmm R package. Journal of Statistical Software 33:1–22.

Havmøller, L. W., J. C. Loftus, R. W. Havmøller, S. E. Alavi, D. Caillaud, M. N. Grote, B. T. Hirsch, L. L. Tórrez-Herrera, R. Kays, and M. C. Crofoot. 2021. Arboreal monkeys facilitate foraging of terrestrial frugivores. Biotropica 53:1685–1697. John Wiley; Sons Inc.

Healy, S. D., and C. Rowe. 2007. A critique of comparative studies of brain size. Proceedings of the Royal Society B: Biological Sciences 274:453–464. The Royal Society London.

Henke-von der Malsburg, J., P. M. Kappeler, and C. Fichtel. 2020. Linking ecology and cognition: Does ecological specialisation predict cognitive test performance? Behavioral Ecology and Sociobiology 74:1–26. Springer.

Herrmann, E., J. Call, M. V. Hernández-Lloreda, B. Hare, and M. Tomasello. 2007. Humans have evolved specialized skills of social cognition: The cultural intelligence hypothesis. Science 317:1360–1366. American Association for the Advancement of Science.

Heymann, E. W., and H. M. Buchanan-Smith. 2000. The behavioural ecology of mixed-species troops of callitrichine primates. Biological Reviews 75:169–190. Wiley Online Library.

Heymann, E. W., and S. S. Hsia. 2015. Unlike fellows–a review of primate–non-primate associations. Biological Reviews 90:142–156. Wiley Online Library.

Ho, L. S. T., and C. Ane. 2014. A linear-time algorithm for gaussian and non-gaussian trait evolution models. Systematic Biology 63:397–408.

Isler, K., and C. P. van Schaik. 2009. The Expensive Brain: A framework for explaining evolutionary changes in brain size. Journal of Human Evolution 57:392–400. Elsevier.

IUCN. 2021. The IUCN Red List of Threatened Species. Downloaded on the 19/01/2021.

Janmaat, K. R. L., C. Boesch, R. Byrne, C. A. Chapman, Z. B. Goné Bi, J. S. Head, M. M. Robbins, R. W. Wrangham, and L. Polansky. 2016. Spatiotemporal complexity of chimpanzee food: How cognitive adaptations can counteract the ephemeral nature of ripe fruit. American Journal of Primatology 78:626–645. Wiley Online Library.

Janmaat, K. R. L., C. A. Chapman, R. Meijer, and K. Zuberbühler. 2012. The use of fruiting synchrony by foraging mangabey monkeys: A 'simple tool' to find fruit. Animal Cognition 15:83–96. Springer.

Janmaat, K. R. L., M. de Guinea, J. Collet, R. W. Byrne, B. Robira, E. van Loon, H. Jang, D. Biro, G. Ramos-Fernández, C. Ross, A. Presotto, M. Allritz, S. Alavi, and S. Van Belle. 2021. Using natural travel paths to infer and compare primate cognition in the wild. iScience 24:102343. Elsevier.

Johnson, A., M. A. van der Meer, and A. D. Redish. 2007. Integrating hippocampus and striatum in decision-making. Current Opinion in Neurobiology 17:692–697. Elsevier.

Kamilar, J. M. 2009. Environmental and geographic correlates of the taxonomic structure of primate communities. American Journal of Physical Anthropology 139:382–393. Wiley Online Library.

Kamilar, J. M., and J. A. Ledogar. 2011. Species co-occurrence patterns and dietary resource competition in primates. American Journal of Physical Anthropology 144:131–139. Wiley Online Library.

Kashetsky, T., T. Avgar, and R. Dukas. 2021. The cognitive ecology of animal movement: Evidence from birds and mammals. Frontiers in Ecology and Evolution 0:633. Frontiers.

Kitamura, T., Ogawa, S. K., Roy, D. S., Okuyama, T., Morrissey, M. D., Smith, L. M., Redondo, R. L., and Tonegawa, S. 2017 Engrams and circuits crucial for systems consolidation of a memory. Science 356.6333: 73-78.

Koziol, L. F., D. Budding, N. Andreasen, S. D'Arrigo, S. Bulgheroni, H. Imamizu, M. Ito, M. Manto, C. Marvel, K. Parker, G. Pezzulo, N. Ramnani, D. Riva, J. Schmahmann, L. Vandervert, and T. Yamazaki. 2014. Consensus paper: The cerebellum's role in movement and cognition. Cerebellum 13:151–177. Springer.

Logan, C. J., S. Avin, N. Boogert, A. Buskell, F. R. Cross, A. Currie, S. Jelbert, D. Lukas, R. Mares, A. F. Navarrete, S. Shigeno, and S. H. Montgomery. 2018. Beyond brain size: Uncovering the neural correlates of behavioral and cognitive specialization. Comparative Cognition and Behavior Reviews 13:55–90. The Comparative Cognition Society.

Losin, N., J. P. Drury, K. S. Peiman, C. Storch, and G. F. Grether. 2016. The ecological and evolutionary stability of interspecific territoriality. Ecology Letters 19:260–267. Wiley Online Library.

MacLean, E. L., B. Hare, C. L. Nunn, E. Addessi, F. Amici, R. C. Anderson, F. Aureli, J. M. Baker, A. E. Bania, A. M. Barnard, and others. 2014. The evolution of self-control. Proceedings of the National Academy of Sciences 111:E2140–E2148. National Acad Sciences.

Mahler, D. L., Ingram, T., Revell, L. J., and Losos, J. B. 2013. Exceptional convergence on the macroevolutionary landscape in island lizard radiations. Science, 341.6143:292-295.

Maliet, O., F. Hartig, and H. Morlon. 2019. A model with many small shifts for estimating species-specific diversification rates. Nature Ecology and Evolution 3:1086–1092. Nature Publishing Group.

Maliet, O., and H. Morlon. 2021. Fast and accurate estimation of species-specific diversification rates using data augmentation. Systematic Biology, doi: 10.1093/sysbio/syab055. Cold Spring Harbor Laboratory.

Many Primates, Altschul, D. M., M. J. Beran, M. Bohn, J. Call, S. DeTroy, S. J. Duguid, C. L. Egelkamp, C. Fichtel, J. Fischer, M. Flessert, D. Hanus, D. B. M. Haun, L. M. Haux, R. A. Hernandez-Aguilar, E. Herrmann, L. M. Hopper, M. Joly, F. Kano, S. Keupp, A. P. Melis, A. M. Rodrigo, S. R. Ross, A. Sánchez-Amaro, Y. Sato, V. Schmitt, M. K. Schweinfurth, A. M. Seed, D. Taylor, C. J. Völter, E. Warren, and J. Watzek. 2019. Establishing an infrastructure for collaboration in primate cognition research. PLoS ONE 14:e0223675. Public Library of Science San Francisco, CA USA.

Matzke, N. J. 2013. Probabilistic historical biogeography: new models for founder-event speciation, imperfect detection, and fossils allow improved accuracy and model-testing. Frontiers of Biogeography.

Matzke, N. J. 2016. Stochastic mapping under biogeographical models. PhyloWiki BioGeoBEARS.

Melchionna, M., A. Mondanaro, C. Serio, S. Castiglione, M. Di Febbraro, L. Rook, J. A. F. Diniz-Filho, G. Manzi, A. Profico, G. Sansalone, and P. Raia. 2020. Macroevolutionary trends of brain mass in Primates. Biological Journal of the Linnean Society 129:14–25. Oxford University Press UK.

Merceron, G., J. Scott, R. S. Scott, D. Geraads, N. Spassov, and P. S. Ungar. 2009. Folivory or fruit/seed predation for mesopithecus, an earliest colobine from the late miocene of eurasia? Journal of human evolution 57:732–738. Elsevier.

Milham, M. P., L. Ai, B. Koo, T. Xu, C. Amiez, F. Balezeau, M. G. Baxter, E. L. A. Blezer, T. Brochier, A. Chen, and Others. 2018. An open resource for non-human primate imaging. Neuron 100:61–74. Elsevier.

Milton, K. 1981. Distribution patterns of tropical plant foods as an evolutionary stimulus to primate mental development. American Anthropologist 83:534–548. Wiley Online Library.

Minot, E. O. 1981. Effects of interspecific competition for food in breeding blue and great tits. The Journal of Animal Ecology 50:375. JSTOR.

Modolo, L., W. Salzburger, and R. D. Martin. 2005. Phylogeography of Barbary macaques (*Macaca sylvanus*) and the origin of the Gibraltar colony. Proceedings of the National Academy of Sciences of the United States of America 102:7392–7397. National Acad Sciences.

Morlon, H., E. Lewitus, F. Condamine, M. Manceau, J. Clavel, and J. Drury. 2016. RPANDA: An r package for macroevolutionary analyses on phylogenetic trees. Methods in Ecology and Evolution 7:589–597.

Mundry, R. 2014. Statistical issues and assumptions of phylogenetic generalized least squares. Pp. 131–153 *in* Modern phylogenetic comparative methods and their application in evolutionary biology. Springer.

Navarrete, A. F., E. L. A. Blezer, M. Pagnotta, E. S. M. De Viet, O. S. Todorov, P. Lindenfors, K. N. Laland, and S. M. Reader. 2018. Primate Brain Anatomy: New Volumetric MRI Measurements for Neuroanatomical Studies. Brain, Behavior and Evolution 91:109–117. Karger Publishers.

Navarrete, A. F., C. P. van Schaik, and K. Isler. 2011. Energetics and the evolution of human brain size. Nature 480:91–93. Nature Publishing Group.

Nuismer, S. L., and L. J. Harmon. 2015. Predicting rates of interspecific interaction from phylogenetic trees. Ecology Letters 18:17–27. Wiley Online Library.

Olupot, W., P. M. Waser, and C. A. Chapman. 1998. Fruit finding by mangabeys (*Lophocebus albigena*): Are monitoring of fig trees and use of sympatric frugivore calls possible strategies? International Journal of Primatology 19:339–353. Springer.

Pavoine, S., M. B. Bonsall, T. J. Davies, and S. Masi. 2019. Mammal extinctions and the increasing isolation of humans on the tree of life. Ecology and Evolution 9:914–924. Wiley Online Library.

Pearce, F., C. Carbone, G. Cowlishaw, and N. J. B. Isaac. 2013. Space-use scaling and home range overlap in primates. Proceedings of the Royal Society B: Biological Sciences 280:20122122. The Royal Society.

Pennell, M., J. Eastman, G. Slater, J. Brown, J. Uyeda, R. Fitzjohn, M. Alfaro, and L. Harmon. 2014. Geiger v2.0: An expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. Bioinformatics 30:2216–2218.

Pepperberg, I. M. 2002. Allospecific referential speech acquisition in grey parrots (*Psittacus erithacus*): Evidence for multiple levels of avian vocal imitation. Imitation in Animals and Artifacts 109–131. MIT Press.

Persson, T., G. A. Sauciuc, and E. A. Madsen. 2018. Spontaneous cross-species imitation in interactions between chimpanzees and zoo visitors. Primates 59:19–29. Springer.

Pigot, A. L., and J. A. Tobias. 2013. Species interactions constrain geographic range expansion over evolutionary time. Ecology Letters 16:330–338. Wiley Online Library.

Porter, L. M. 2001. Benefits of polyspecific associations for the Goeldi's monkey (*Callimico goeldii*). American Journal of Primatology 54:143–158. Wiley Online Library.

Powell, L. E., R. A. Barton, and S. E. Street. 2019. Maternal investment, life histories and the evolution of brain structure in primates. Proceedings of the Royal Society B: Biological Sciences 286:20191608. The Royal Society.

Powell, L. E., K. Isler, and R. A. Barton. 2017. Re-evaluating the link between brain size and behavioural ecology in primates. Proceedings of the Royal Society B: Biological Sciences 284:20171765. The Royal Society.

Price, T. D., and M. Kirkpatrick. 2009. Evolutionarily stable range limits set by interspecific competition. Proceedings of the Royal Society B: Biological Sciences 276:1429–1434. The Royal Society London.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rabosky, D. L., and I. J. Lovette. 2008. Density-dependent diversification in North American wood warblers. Proceedings of the Royal Society B: Biological Sciences 275:2363–2371. The Royal Society London.

Rafacz, M., and J. J. Templeton. 2003. Environmental unpredictability and the value of social information for foraging starlings. Ethology 109:951–960. Wiley Online Library.

Ramdarshan, A., G. Merceron, and L. Marivaux. 2012. Spatial and temporal ecological diversity amongst eocene primates of france: Evidence from teeth. American Journal of Physical Anthropology 147:201–216. Wiley Online Library.

Reader, S. M., and K. N. Laland. 2002. Social intelligence, innovation, and enhanced brain size in primates. Proceedings of the National Academy of Sciences of the United States of America 99:4436–4441. National Acad Sciences.

Revell, L. J. 2012. Phytools: An r package for phylogenetic comparative biology (*and other things*). Methods in Ecology and Evolution 3:217–223.

Robira, B., S. Benhamou, S. Masi, V. Llaurens, and L. Riotte-Lambert. 2021. Foraging efficiency in temporally predictable environments: Is a long-term temporal memory really advantageous? Royal Society Open Science. The Royal Society.

Rosati, A. G. 2017. Foraging cognition: Reviving the Ecological Intelligence Hypothesis. Trends in Cognitive Sciences 21:691–702. Elsevier.

Sayol, F., O. Lapiedra, S. Ducatez, and D. Sol. 2019. Larger brains spur species diversification in birds. Evolution 73:2085–2093. Wiley Online Library.

Schilder, B. M., H. M. Petry, and P. R. Hof. 2020. Evolutionary shifts dramatically reorganized the human hippocampal complex. Journal of Comparative Neurology 528:3143–3170. Wiley Online Library.

Schreier, B. M., A. H. Harcourt, S. A. Coppeto, and M. F. Somi. 2009. Interspecific competition and niche separation in primates: A global analysis. Biotropica 41:283–291. Wiley Online Library.

Shettleworth, S. J. 2009. Cognition, evolution, and behavior. Oxford university press.

Shultz, S., and R. I. M. Dunbar. 2006. Chimpanzee and felid diet composition is influenced by prey brain size. Biology Letters 2:505–508. The Royal Society London.

Slater, G., L. Harmon, D. Wegmann, P. Joyce, L. Revell, and M. Alfaro. 2012. Fitting models of continuous trait evolution to incompletely sampled comparative data using approximate bayesian computation. Evolution 66:752–762.

Smaers, J. B., R. S. Rothman, D. R. Hudson, A. M. Balanoff, B. Beatty, D. K. Dechmann, D. de Vries, J. C. Dunn, J. G. Fleagle, C. C. Gilbert, and others. 2021. The evolution of mammalian brain size. Science Advances 7:eabe2101. American Association for the Advancement of Science.

Sokolov, A. A., R. C. Miall, and R. B. Ivry. 2017. The cerebellum: Adaptive prediction for movement and cognition. Trends in Cognitive Sciences 21:313–332. Elsevier.

Sol, D., T. Székely, A. Liker, and L. Lefebvre. 2007. Big-brained birds survive better in nature. Proceedings of the Royal Society B: Biological Sciences 274:763–769. The Royal Society London.

Springer, M. S., R. W. Meredith, J. Gatesy, C. A. Emerling, J. Park, D. L. Rabosky, T. Stadler, C. Steiner, O. A. Ryder, J. E. Janečka, and others. 2012. Macroevolutionary dynamics and historical biogeography of primate diversification inferred from a species supermatrix. PloS one 7:e49521. Public Library of Science San Francisco, USA.

Todorov, O. S., V. Weisbecker, E. Gilissen, K. Zilles, and A. A. De Sousa. 2019. Primate hippocampus size and organization are predicted by sociality but not diet. Proceedings of the Royal Society B: Biological Sciences 286:20191712. The Royal Society.

Tomasello, M. 2019. The cultural origins of human cognition. Harvard university press.

Trapanese, C., B. Robira, G. Tonachella, S. Di Gristina, H. Meunier, and S. Masi. 2019. Where and what? Frugivory is associated with more efficient foraging in three semi-free ranging primate species. Royal Society Open Science 6:181722. The Royal Society.

van Schaik, C. P., and J. M. Burkart. 2011. Social learning and evolution: The cultural intelligence hypothesis. Philosophical Transactions of the Royal Society B: Biological Sciences 366:1008–1016. The Royal Society.

van Schaik, C. P., Z. Triki, R. Bshary, and S. A. Heldstab. 2021. A Farewell to the Encephalization Quotient: A New Brain Size Measure for Comparative Primate Cognition. Brain, Behavior and Evolution 96:1–12. Karger Publishers.

Weir, J. T., and S. Mursleen. 2013. Diversity-dependent cladogenesis and trait evolution in the adaptive
radiation of the auks (*Aves Alcidae*). Evolution 67:403–416. Wiley Online Library.

Whiten, A., and C. P. van Schaik. 2007. The evolution of animal 'cultures' and social intelligence.
Philosophical Transactions of the Royal Society B: Biological Sciences 362:603–620. The Royal Society
London.

Willems, E. P., B. Hellriegel, and C. P. van Schaik. 2013. The collective action problem in primate
territory economics. Proceedings of the Royal Society B: Biological Sciences 280:20130081. The Royal
Society.

Wilson, A. 1991. From molecular evolution to body and brain evolution. Pp. 331–340 *in* Perspectives on
cellular regulation: From bacteria to cancer. J. Wiley & Sons.

Wiltgen, B. J., R. A. M. Brown, L. E. Talton, and A. J. Silva. 2004. New circuits for old memories: The
role of the neocortex in consolidation. Neuron 44:101–108. Elsevier.

Wisniewski, A. L., G. T. Lloyd, and G. J. Slater. 2022. Extant species fail to estimate ancestral
geographical ranges at older nodes in primate phylogeny. Proceedings of the Royal Society B
289:20212535. The Royal Society.

923