Species sympatry shapes brain size evolution in Primates

Benjamin Robira[1,2,*], and Benoît Perez-Lamarque[3,4]

[1] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France.

[2] Eco-anthropologie (EA), Muséum National d'Histoire Naturelle, CNRS, Université de Paris, Musée de l'Homme, Paris, France.

[3] Institut de Biologie de l'École Normale Supérieure (IBENS), École Normale Supérieure, CNRS, INSERM, Université PSL, Paris, France.

[4] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France.

[*] Correspondence: Benjamin Robira <benjamin.robira@normalesup.org>

Abstract | The main hypotheses about the evolution of animal cognition emphasise the role of conspecifics. Yet, space is often simultaneously occupied by multiple species from the same ecological guild. These sympatric species can compete for food, which may thereby stimulate or hamper cognition. Considering brain size as a proxy for cognition, we tested whether species sympatry impacted the evolution of cognition in frugivorous primates. We first retraced the evolutionary history of sympatry between frugivorous primate lineages. We then fitted phylogenetic models of the evolution of the size of several brain areas in frugivorous primates, considering or not species sympatry. We found that the whole brain or brain areas used in immediate information processing were best fitted by models not considering sympatry. By contrast, models considering species sympatry best predicted the evolution of brain areas related to long-term memory of interactions with the social or ecological environment, with a decrease of their

size the higher the sympatry. We speculate that species sympatry, by generating intense food
depletion, leads to an over-complexification of resource spatio-temporality that counteracts the
benefits of high cognitive abilities and thereby induces lower brain area sizes. In addition, we
reported that species in sympatry diversify more slowly. This comparative study suggests that
species sympatry significantly contributes to shaping primate cognition and diversification.

Short title: Sympatry shapes primates' brain size

Keywords: Brain size - Cognition - Diversification - Frugivory - Primates - Sympatry

INTRODUCTION

Cognition evolution is shaped by the balance between socio-ecological drivers promoting cognitive abilities (González-Forero and Gardner 2018) and physiological and energetic constraints limiting them (Navarrete, Schaik, and Isler 2011). Primates are pivotal species for cognitive studies (Byrne 2000) because their cognition is thought to be promoted by interactions of individuals with conspecifics within the social unit (Byrne 2018; Dunbar and Shultz 2017), among generations (Wilson 1991; Whiten and Schaik 2007; Reader and Laland 2002; Herrmann et al. 2007; Tomasello 2019; Schaik and Burkart 2011), between social units (Ashton, Kennedy, and Radford 2020), or with the rest of their environment (Clutton-Brock and Harvey 1980; Milton 1981; Rosati 2017). However, space is often occupied by many primate species sharing the same diet. Because of competition for food between these species, both direct and indirect interactions between heterospecifics in sympatry are also likely to shape the evolution of their cognition.

Retracing the evolutionary history of cognitive abilities proves to be challenging because there is still no consensual measurement for cognition applicable across all species. Up to now, a raw approximation consists in considering the (relative) brain size as a proxy for cognitive abilities, with larger sizes considered equivalent to more advanced cognitive abilities (Benson-Amram et al. 2016). Although the relevance of this assumption is heavily limited within species, in part because of plasticity (Gonda, Herczeg, and Merilä 2013), this holds true when comparing different species (e.g., in primates, Reader and Laland 2002). Instead of considering the brain as a whole, the multifaceted aspect of animal cognition is then more precisely depicted by appreciating the mosaic nature of the brain (Barton and Harvey 2000). For instance, variations in the size of some specific brain areas have been robustly associated with variations in cognition related to the function of these areas (Healy and Rowe 2007). The brain is therefore a patchwork of areas cognitively specialised that may follow different evolutionary trajectories.

Because species sympatry might play on different aspects of the socio-ecological environment, the

brain areas might be differently affected by species sympatry. First, sympatric species from the

same dietary guild may show some dietary overlap. Thus, sympatry often leads to an increase in

food depletion of the shared resource compared with an environment with only one foraging

species (Minot 1981). As an indirect effect of depletion, sympatric species competing for the same

food resource may therefore complexify the pattern of resource distribution and availability in

space and time. This complexification may in turn affect the selective pressures upon brain areas

involved in the storing of spatio-temporal information, such as the Hippocampus (Burgess, Maguire,

and O'Keefe 2002, Hypothesis 1: memory is affected by sympatry). Second, all sympatric species

may enrich the landscape of visual, olfactory or acoustic cues usable to locate available food (e.g.,

Avarguès-Weber, Dawson, and Chittka 2013; Kashetsky, Avgar, and Dukas 2021). Consequently, it

may impact the selective pressures upon brain areas involved in processing more immediate

sensory information, such as the Main Olfactory Bulb (MOB), the Cerebellum (Koziol et al. 2014;

Sokolov, Miall, and Ivry 2017), and the Neocortex (Wiltgen et al. 2004) (Hypothesis 2: cue

processing is affected by sympatry). Besides indirect interaction(s) through foraging, cognition can

also be triggered by direct "social" interactions with other individuals (Byrne 2018; Dunbar and

Shultz 2017). The Striatum, a brain area stimulated during social interactions (Báez-Mendoza and

Schultz 2013), may therefore be affected by the increase of direct social interactions between

heterospecifics (Hypothesis 3: sociality is affected by sympatry).

Under these (non-exclusive) hypotheses, sympatry could stimulate or hamper cognition evolution.

Memory stands as a valuable tool to infer food availability and location when food is rare and

ephemeral but predictable (Milton 1981; Rosati 2017). Thus, having a better memory should be

advantageous under reasonable food depletion. In addition, competition for the shared resource

between species should promote anticipatory behaviour, hence high cognition, as expected for

within-species competition (Ashton, Kennedy, and Radford 2020). In this case, the size of the

Hippocampus (reflecting long-term memory abilities) should be larger the higher the sympatry

intensity (Prediction 1.1). On the other hand, intense depletion also increases environmental

unpredictability. In the case of a frugivore searching for fruit, for instance, the perceived synchrony

in fruit production between trees of the same or different species, used to infer food availability

(Janmaat et al. 2012), can be lowered by depletion, eventually limiting the benefits of memory

(Robira et al. 2021). Thus, with such a scenario and due to the energy constraints of maintaining a

large brain, the Hippocampus size could be smaller in highly sympatric species (Prediction 1.2).

Meanwhile, cues left out by heterospecifics and usable to locate available food might also add to

environmental ones already available. Hence, sympatry could be associated with larger sizes of the

MOB, the Cerebellum, or the Neocortex (Prediction 2). Finally, an increase in direct interactions

between species, such as with the formation of mixed-group species (Goodale et al. 2010), should

imply an upsurge of social stimuli leading to a larger size of the Striatum in sympatry (Prediction 3).

Here, we investigated whether species sympatry affected the evolution of cognition using

frugivorous primates as a study example. Frugivorous primates are an interesting group for such a

question because fruit is the archetype of a hard-to-find resource yet predictable (Janmaat et al.

2016), for which cognition considerably shapes the foraging strategy (Trapanese et al. 2019). To

infer the effect of species sympatry on cognition in frugivorous primates, we evaluated the support

for models of brain size evolution accounting or not for species sympatry, and investigated the

directionality of the selection induced by sympatry on brain size evolution. Finally, we tested for

correlative patterns between brain size or current sympatry and the species diversification in all

primates, to better understand the impact of cognition and interactions between primates on their

evolutionary success.

METHODS

Data processing, analyses, and plots were computed with R software (v.4.1.2, R Core Team 2020).

Data Collection

Phylogeny

We used a block of chronogram trees of the primate taxon of the 10kTrees project (downloaded on May 2021, version 3), as well as a consensus tree of 1000 trees for the subsequent phylogenetic analyses. The trees contain 301 primate species (Figure 2). Note that in all these analyses, we discarded *Homo sapiens* and *Macaca sylvanus*. The latter was discarded because of its complete geographic isolation and repeated intervention of human people in population maintenance (Modolo, Salzburger, and Martin 2005). A summary of available data per species is presented in Supplementary Material Figure S3.

Trait data

Data were pooled from previous literature surveys (see Supplementary Material "Data availability"). Brain data were obtained from DeCasien and Higham (2019) for the whole brain and all mentioned other areas (Cerebellum, Hippocampus, Main Olfactory Bulb (MOB), Neocortex, Striatum), Powell, Isler, and Barton (2017) and Powell, Barton, and Street (2019) for the whole brain, Cerebellum and Neocortex size, Todorov et al. (2019) for Hippocampus and Neocortex size, Grueter (2015) for the whole brain size and Navarrete et al. (2018) for the whole brain, Cerebellum, Hippocampus and Striatum size. They were freely-available in the main manuscript or supplementary materials. For each primate species, the percentage of frugivory and/or folivory was obtained based on a freely available dataset from DeCasien, Williams, and Higham (2017), Powell, Isler, and Barton (2017), and Willems, Hellriegel, and Schaik (2013). The availability of trait and distribution range for the 301 primate species represented in the primate phylogeny of the 10kTrees project is depicted in Supplementary Material Figure S3. From the global endocranial brain volume, we obtained the Encephalization Quotient (EQ, $N_{EQ,max}$ = 182) as follows (DeCasien, Williams, and Higham 2017)

$$EQ = 1.036 \times \text{Brain volume}/(0.085 \times \text{Body mass}^{0.775})$$

with the brain volume in cm^3, 1.036 g/cm^3 being the assumed homogeneous brain density, and the

body mass in g. EQ indicates whether the brain size ranges above (> 1) or below (< 1) expected

given the body mass. Body mass was obtained from DeCasien, Williams, and Higham (2017), Powell,

Isler, and Barton (2017), Grueter (2015), Pearce et al. (2013). The sub-parts of the brain were

chosen because they were involved in immediate sensory information processing (MOB, $N_{MOB,max}$ =

39), in movement and/or general information processing and retention (Neocortex, $N_{Neocortex,max}$

= 69, Wiltgen et al. 2004; Cerebellum, $N_{Cerebellum,max}$ = 70, Koziol et al. 2014; Sokolov, Miall, and

Ivry 2017), short-term working memory and long-term spatio-temporal memory (Hippocampus,

$N_{Hippocampus,max}$ = 63, Burgess, Maguire, and O'Keefe 2002). The Striatum ($N_{Striatum,max}$ = 63)

supports information processing during social interaction, reward assessment, planning or goal-

oriented behaviours (Báez-Mendoza and Schultz 2013; Johnson, Meer, and Redish 2007). To

investigate their evolutionary history, we used the ratio between their volume and body mass, so as

to maximize comparability. As such, the use of specific area sizes relative to the body mass and not

raw sizes depicts the evolution of cognitive abilities in terms of allocation rather than abilities per se

(but see discussion in Deaner, Nunn, and Schaik 2000).

Ranging Data

The current biogeographic range of each primate species was assessed using ranging maps

provided by the IUCN red list (IUCN 2021). Ranging data were available for 249 species among the

301 represented in the 10kTrees primate phylogeny.

Retracing past sympatry between primate species

Based on the biogeographic distribution of each extant primate species, we first reconstructed the

history of past sympatry between primate lineages. To do so, we followed Drury et al. (2018) and

first reconstructed the biogeographic history of each primate lineage to then retraced which pairs of

primate lineages were likely to be simultaneously present at the same place. Leaning on Kamilar

(2009), we considered that the biogeography of primates can be described by 12 discrete

biogeographic areas with highly similar community structures shaped by both the environment

geography and climatic correlates. These geographic areas, mapped using Google earth professional

(v7.3.3), are represented in Figure 1. One to multiple biogeographic areas were assigned to each

species as soon as 10 of their current distribution range overlapped on the surface with a given

biogeographic area. We also replicated these biogeographic assignations by using instead a larger

threshold of 30%. This upper threshold was chosen because a species could occupy as far as three

areas, Figure 1). Overlap of primate current range with biogeographic areas was calculated with the

"gIntersection" function from the *rgeos* package (Bivand and Rundel 2021) applied to Mercator-

projected data to get the overlapping contour, and the "area" function from the *geosphere* package

(Hijmans 2021), applied directly on unprojected longitudinal-latitudinal data for area size

calculation.

Given these 12 biogeographic areas, we retraced the biogeographic history of primates with the

BioGeoBEARS package (Matzke 2013), using the biogeographic stochastic mapping algorithm

(Matzke 2016). We fitted non-time-stratified dispersal-extinction-cladogenesis (DEC) models

specifically suiting analyses of range data since it accounts for spatially explicit processes of

cladogenetic and anagenetic events (see Matzke (2013) for further details on these events). We

fixed to three biogeographic areas the maximum number of areas that a lineage can simultaneously

occupy since it offers the possibility to occupy a complete mainland continent while keeping

computational time reasonable. DEC models were independently fitted when considering either a

10% or a 30% threshold of range overlaps. Finally, to account for the uncertainty in biogeographic

reconstructions, we sampled 10 histories of primate biogeographic ranges. We assumed that

primate lineages were in sympatry at a given time whenever the species occupied the same

biogeographic area.

Inferring past diets of primate lineages

Next, we retraced the evolutionary history of frugivorous lineages in primates. We first classified

extant species as either "frugivorous" or "folivorous" based on the availability of frugivorous rate

and folivorous rate, prioritizing frugivory over folivory. A species was classified as frugivorous if the

frugivory rate was at least above 20%. If this was not the case, or if the frugivory rate was

unavailable, a species could be classified as folivorous if the folivory rate was at least above 40%.

Otherwise, DeCasien, Williams, and Higham (2017) gave a binary classification of diet, species being

categorised as frugivorous or folivorous, partly based on anatomical criteria. Whenever the rate was

not available, we referred to this classification. In any other cases, the species was discarded. We

also replicated these diet assignments by considering a threshold of 40% for frugivory and 60% for

folivory.

Second, considering diet as a binary variable (frugivory versus folivory), we retraced the

evolutionary history of such discrete traits based on a continuous Markovian process (extended Mk

models) using a Bayesian inference (Bollback 2006), with the "simmap" function of the *phytools*

package (Revell 2012) and internally setting up the prior probability of trait, but with no prior on

the transition matrix. Ancestral diet reconstructions were performed using both combinations of

dietary thresholds (20/40% and 40/60%). To account for the uncertainty in the reconstructions, we

obtained 10 stochastic diet history timelines. The latter were used in combination with the history

of primate ranges to assess whether a frugivorous species was in sympatry with another

frugivorous species or not (i.e., we obtained reconstructions of the evolutionary history of sympatry

between frugivorous primate lineages).

Phylogenetic models

We assessed the effect of sympatry on primate brain evolution using two approaches. First, we used

phylogenetic models of trait evolution to assess the role of sympatry in the evolution of brain size.

Second, we investigated how sympatry has influenced brain size evolution (i.e., selection towards

smaller or larger brain sizes) by evaluating correlations between current levels of sympatry and

brain sizes, using linear modelling. Besides, we also checked for correlative patterns between

primate brain size and diversification rates to have insights into primate evolutionary success.

Phylogenetic models of trait evolution: does species sympatry shape brain size evolution?

(a)   Fitting models of trait evolution

We restricted the analyses to frugivorous species to test whether species sympatry has impacted

the evolution of cognition, depicted either by the whole brain (using the encephalic quotient, EQ), or

the size of the aforementioned specific brain areas relative to the whole-body mass (Figure 3). For

such a task, we fitted phylogenetic models of the evolution of the size of the different brain areas

independently (Drury et al. 2016). For models implying species sympatry, this was made possible

by combining the historical timeline of diet and biogeography evolution, so that we could retrace

the history of sympatry between frugivorous lineages. In practice, we obtained a series of

interaction matrices (i.e., lines and columns correspond to frugivorous species, and each cell

indicates whether a given species pair is in sympatry (value of 1) or not (value of 0)), along the

phylogenetic tree (see Drury et al. 2016). This was used to fit models that considered species

sympatry to model brain size evolution: the matching competition (MC) model (Nuismer and

Harmon 2015) and density-dependent models (DD_{lin} and DD_{exp}, Drury et al. 2016). Specifically,

these models expand classical models of stochastic evolution (Brownian Motion), by including an

additional variable related to current brain size of sympatric species (MC), or by considering

density-dependent evolutionary rate (DD models). Specifically, the matching competition model

(MC) considers the repulsion of traits of sympatric lineages from the same dietary guild due to

competition (character displacement), that is $z_i(t + dt) = z_i(t) + S\big(\mu(t) - z_i(t)\big)dt + \sigma dB_i$ where z

is the brain size of a species i at time t, μ is the mean value of the trait of sympatric species, S

reflects the strength of the effect of species sympatry and σdB_i is the drift with a constant

evolutionary rate σ (Drury et al. 2016). Here, S is constrained to be negative, which means that

sympatric species would tend to divergently evolve either lower, or higher, EQ or relative brain size.

Linear (DD_{lin}) or exponential (DD_{exp}) density-dependence (Drury et al. 2016; Weir and Mursleen

2013) means that the evolutionary rate, σ, of trait change, varies either positively or negatively as a

function f of the number of frugivorous sympatric lineages, such as

$$\sigma_l = f_{lin}(l) = \sigma_0 + rl$$

$$\sigma_l = f_{exp}(l) = \sigma_0 \exp(rl)$$

where σ_0 corresponds to the value of the initial ancestor, l indicates the number of lineages, r allows

for modelling the speed and direction of the dependency to lineage number ($r > 0$ leads to an

increase of trait changes, while $r < 0$ leads to a decline of the trait changes). We fitted models

considering species sympatry using the "fit_t_comp" function from the *RPANDA* package (Morlon et

al. 2016).

Depending on the brain area and the frugivory threshold we considered, the models were fitted on

different sample sizes: EQ: 148 to 182, Striatum: 56 to 63, MOB: 34 to 39, Neocortex: 61 to 69,

Hippocampus: 56 to 63, Cerebellum: 62 to 70 frugivorous species. For a given set of models (i.e.,

within a brain area), the sample was strictly identical, allowing within-set comparisons. Prior to

fitting, trait parameters were log-transformed to reach more symmetrical distributions.

We compared the support of models considering species sympatry to the support of simpler

models assuming no effect of species sympatry on the evolution of brain sizes: the Brownian Motion

(BM), the Ornstein-Uhlenbeck process (OU, a model with an optimum value, see Blomberg,

Rathnayake, and Moreau (2020) for a review), or the Early-Burst model (EB), for assessing a time-

dependence of the evolutionary rate, irrespective of the intensity of species sympatry (Blomberg,

Garland, and Ives 2003). These models without species sympatry were fitted using the

"fitContinuous" function from the *geiger* package (Slater et al. 2012; Pennell et al. 2014). All these

models were repeated 10 times, using 10 different combinations for the evolutionary history of

primate biogeography and diet. They were then compared within an information-theoretic

framework (Burnham and Anderson 2002), based on the weights of Akaike Information Criteria

corrected for small samples (AICc) when considering all six models (MC, DD_{lin}, DD_{exp}, BM, OU, EB).

The model weight depicts how well the model fits the observed data compared with the other tested

models.

(b)   Determining the effect of sympatry on brain sizes

If diversity-dependent models of traits evolution considering species sympatry can be used to

assess whether or not species sympatry has impacted the evolution of the brain size by increasing

or decreasing the tempo of trait evolution, they do not say anything about the directionality of the

effect (i.e., are brain sizes in frugivorous sympatric primates increasing or decreasing?). To

determine whether species sympatry positively or negatively affected the sizes of brain areas, we

independently fitted Gaussian Pagel's lambda phylogenetic regressions for each brain area of extant

frugivorous species. This model is a derivative of the Brownian Motion model, where the

phylogenetic variance-covariance matrix has all coefficients, but its diagonal ones, multiplied by

lambda: it thus relaxes the hypothesis of Brownian Motion since we included brain areas for which

the evolutionary history was best described by models considering sympatry (see Results). To fit

these models, we used a frequentist-based approach with the "phylolm" function from the *phylolm*

package (Ho and Ane 2014). We considered the least stringent frugivory assessment, with the

frugivory threshold fixed at 20% and the folivory threshold fixed at 40%. If due to data variability, a

species did not robustly fit into the categorical classification "frugivorous versus folivorous" (i.e.,

could be either of the two), it was considered as frugivorous nonetheless.

The response variable was the relative size of each brain area. Due to data variability, we took the

mean of the possible values given the different datasets and assessed the sensitivity using non-

averaged values (see Supplementary Material "Phylogenetic regressions: results, stability, and

assumption"). In this model, we used as covariates (i.e., continuous predictors) two explicit

measures of species sympatry intensity for each extant frugivorous species: (1) the number of

frugivorous sympatric species (square-rooted to reach symmetrical distribution) and (2) the

average percentage of overlapping current range (assessed based on IUCN data) with other

sympatric frugivorous species. For a given species A, sympatry with another species B was

considered when at least 10% of the range of species A overlaps with the range of species B. This

was done to reduce the noise induced by coarse identification of species range.

Eventually, it means that the results for each model represent the average of 10 (uncertainty on

diet/ranging evolution) x 10 (uncertainty in brain/diet rate data) x 2 (geographic overlap

threshold) x 2 (frugivory threshold) x 2 (folivory threshold) = 800 sub-models. We stopped

computations when the calculation of the likelihood was excessively long (> 1 week). The final

sample size thus was 730 models.

To sum up, when assessing the interplay between sympatry and the evolution of frugivorous

primates' brain architecture, we considered sympatry under different forms. To assess whether it

affected brain size evolution, sympatry was added to classical phylogenetic models of trait evolution

as an additional variable depicting the mean trait value of sympatric species (MC models), or as a

density-dependent term (i.e., the total number of sympatric lineages at a given time; in DD_{lin} and

DD_{exp} models). Then, to assess the directionality of the effect of sympatry on brain sizes, sympatry

was used as a tested predictor in phylogenetic linear regressions, under two forms: the number of

currently sympatric species, and the average range overlap with currently sympatric species.

Models of species diversification

Next, to investigate whether cognition and/or species sympatry have affected primate

diversification, we inferred how primates diversified over time and across lineages. Lineage-specific

net diversification rates (defined as speciation minus extinction rates) were estimated using an

updated version of the *ClaDS* algorithm (Maliet, Hartig, and Morlon 2019) boosted for

computational speed based on data augmentation techniques (Maliet and Morlon 2021).

Particularly, we used *ClaDS2*, the model with constant turnover (i.e., constant ratio between

extinction and speciation rates; see Supplementary Material "Primate diversification rate over time"

for further explanations). We extracted the mean diversification rates through time and the lineage-

specific diversification rate of each extant species.

We also fitted Gaussian Pagel's lambda phylogenetic regressions of the different relative brain sizes

against the net diversification rates, estimated for each extant species by the *ClaDS* algorithm.

Because assumptions for a frequentist-based approach were unmet, we used a Bayesian-based

approach. We used the "MCMCglmm" function of the *MCMCglmm* package (Hadfield 2010). Each

chain was based on a burn-in period of 5000 iterations, among a total of 50 000 iterations, and was

sampled every 50 iterations. We used the least informative priors. Fixed priors were let to default

(Gaussian distribution of mean 0 and variance 10^8). Again, we took the mean of the brain trait

values for the main model and assessed the sensitivity by re-running the model several times using

non-averaged values.

To determine whether species sympatry was associated with lower or larger diversification rates,

we fitted frequentist-based Gaussian Pagel's lambda phylogenetic regressions with the lineage-

specific diversification rate as the output variable, and used the two metrics for describing sympatry

(the number of frugivorous sympatric species and the average percentage of overlapping range with

other sympatric frugivorous species) as the tested variables, as in (a).

Details on the implementation, stability, and uncertainty of phylogenetic regressions are provided in

Supplementary Material (see "Phylogenetic regressions: results, stability, and assumption").

RESULTS

The database we gathered contained between 34 to 182 frugivorous primate species (depending on

the brain area considered). After pondering by whole-body mass, we observed ample variations in

brain area relative sizes. For instance, the lemuriformes, which are known to prioritize smell

compared with other primate species, have the largest relative MOB size (Lemuriformes: mean $\pm$ SE

= 0.23 $\pm$ 0.07, other: 0.12 $\pm$ 0.04, 3). Similarly, platyrrhini, and callitrichine primates in particular,

are known to form poly-specific associations (Heymann and Buchanan-Smith 2000). The latter

show the highest relative size of the Striatum (Platyrrhini: mean $\pm$ SE = 0.91 $\pm$ 0.07, other: 0.59 $\pm$

0.07, 3). In terms of the measures of sympatry, we observed that on average ($\pm$ SE), the considered

primate species had 52% ($\pm$ 2) of their range overlapping with other species. That ranged from 0%

of overlap (*Macaca nigra*), to 100% of overlap (*Cercopithecus pogonias, Alouatta pigra, Loris*

tardigradus, Hylobates moloch, Cercocebus galeritus, Presbytis melalophos, Semnopithecus entellus).

In terms of the distribution range, the considered primate species co-occurred on average with 6.38

($\pm$ 0.39) other primate species, ranging from 0 other species to 21.

To retrace the history of past species sympatry between frugivorous lineages, we first

reconstructed primate biogeographic history when considering 12 biogeographic areas (Figure 1,

Kamilar 2009) and their diet evolution. We then modelled the evolution of the size of the whole

brain (EQ), or regionalised areas (Neocortex, Cerebellum, MOB, Hippocampus, and Striatum) when

considering species sympatry or not. We found that models not considering species sympatry best

described the evolutionary history of the EQ, the Neocortex, and the Cerebellum (Figures 3 and 4),

two areas specifically involved in immediate sensory information processing (Wiltgen et al. 2004;

Koziol et al. 2014; Sokolov, Miall, and Ivry 2017), and also in memory consolidation for the

Neocortex (Wiltgen et al. 2004). The fact that these biggest areas are best described by the Ornstein-

Uhlenbeck process suggests a stabilization towards an optimal size, which may illustrate the trade-

off between costs and benefits of brain development (Isler and Schaik 2009). By contrast, density-

dependent models considering species sympatry (DD_{lin} and DD_{exp}) were best supported in the

foraging-related and social-related areas respectively: the Hippocampus, specialised in spatio-

temporal memory (Burgess, Maguire, and O'Keefe 2002) and the Striatum, involved in social

interactions (Báez-Mendoza and Schultz 2013). The fact that we inferred positive rates r of density-

dependence (Figure 4) suggested an acceleration of the evolutionary tempo of trait evolution

together with increased diversity of frugivorous sympatric lineages for the Hippocampus and the

Striatum. The MOB, the area involved in sensory abilities, also tended to be best fitted by models

considering sympatry as a whole. Yet, Brownian Motion (BM) was as likely as density-dependent or

MC models, preventing firm conclusions on whether sympatry affected or not MOB size evolution

(Figures 3 and 4), especially since this coincided with the most reduced sample size we had ($N_{species}$

= 34 to 39).

Next, we assessed whether species sympatry leads to "bigger" or "smaller" brain area sizes the

more sympatric species. To do so, we fitted phylogenetic regressions in extant frugivorous species

between the relative sizes of the different brain areas and two measures of sympatry (1) the

average percentage of overlapping range with other frugivorous sympatric species, and (2) the

number of such sympatric frugivorous species across their current entire distribution range. The

number of sympatric species never significantly influenced the relative brain sizes (Table 1).

Conversely, we found that the average percentage of overlapping range correlated with the relative

size of brain areas that were better fit with models considering sympatry: the Hippocampus and the

Striatum (Hippocampus: t = -1.94, p = 0.058; Striatum: t = -2.26, p = 0.028). The correlations were

all negative (Hippocampus: est. = -0.39, CI95% = [-0.76,-0.01]; Striatum: est. = -0.4, CI95% = [-0.77,-

0.04]), which means that higher range overlap between sympatric species associates with lower

relative size, insensitive to data and phylogenetic uncertainties (Appendix Table S1, Appendix

Figure S8, Appendix Table S1). Given the acceleration of the evolutionary tempo with species

sympatry ($r > 0$ in the density-dependent models), it suggests that compared with isolated species,

sympatric species are subject to a positive selection towards smaller brains, and not to a less intense

selection for advanced cognitive abilities.

Finally, we investigated the evolutionary consequences of cognition and species sympatry by

evaluating whether brain sizes and sympatry intensities correlated with the lineage-specific net

diversification rates of primates (defined as speciation minus extinction rates). Overall, species

diversification rates, estimated based on the primate molecular phylogeny, particularly boomed in

the early and late Miocene, around 25 and 11 Myr ago (Appendix Figure S4). When accounting for

phylogenetic dependence, no significant relationship between the net diversification rate and the

relative size of brain areas was found (Table 2, Appendix Figure S8; see robustness in

Supplementary Material Table S2). Although diversification was uncorrelated with brain size in

frugivorous primates, it was influenced by the sympatry context. In particular, phylogenetic

regressions highlighted a negative effect of the number of sympatric species on the diversification

rate (est. = -5.04e-03, CI95% = [-0.01,1.34e-04], t = 2.56e-03, p < 0.001, Table 3, Appendix Figure S8,

Appendix Table S3). In other words, the higher the number of sympatric species, the lower the

diversification rate.

DISCUSSION

Bigger brains are not necessarily better, as the size of the brain is subject to a compromise between

the energy it incurs, and the increase of fitness it allows. This is clearly emphasised by the fact that

the evolution of the biggest brain areas, the Cerebellum and the Neocortex, as well as the whole

brain (EQ), were best fitted by the Ornstein-Uhlenbeck process. This suggests a stabilisation

towards an optimal size resulting from an equilibrium between costs and benefits. Although

allometric and developmental constraints, as well as spatial proximity in the brain, can induce

correlation in the evolution of different brain areas (Gómez-Robles, Hopkins, and Sherwood 2014),

brain areas underpin different cognitive functions and can thus be under different, independent,

selective pressures (Barton and Harvey 2000). The functional regionalisation is for instance

evidenced here by the differences in relative sizes across lineages in the MOB, with larger sizes in

the lemuriformes that mostly rely on smell to forage. The differences in evolutionary trajectories are

highlighted by the variations in the best fit models of size evolution for the different brain areas. We

indeed show that sympatry is one factor that affects the selective regime under which only some

brain area evolves: although the brain as a whole was insensitive to species sympatry, the latter

nonetheless induced a change in the relative size of the Hippocampus and the Striatum. These areas

are involved in individual-based and social-based information processing, pinpointing that the two

components might be under strong selection in primates (DeCasien, Williams, and Higham 2017;

Powell, Isler, and Barton 2017; González-Forero and Gardner 2018).

Overall, the fact that the Hippocampus, particularly relevant to process and memorise spatio-

temporal information, is sensitive to sympatry, is consistent with the idea of an effect of sympatric

species on resource spatio-temporality (Hypothesis 1). Competition is generally the first-thought

mechanism to describe community structures (de Almeida Rocha et al. 2015) because it might affect

the environment in which species evolve. We show that a higher intensity of sympatry is actually

associated with smaller sizes of the Hippocampus (in accordance with Prediction 1.2). This suggests

that indirect competition for food might contribute to convoluting the environment, and such an

over-complexification of the resource spatio-temporality may render cognitive foraging not

advantageous anymore. As a result, it might even generate a selection for smaller brains.

By contrast, potential indirect facilitation between species due to "social" cues (Hypothesis 2), is

ruled out by the absence of an effect of sympatry on brain areas involved in immediate sensory

information processing (e.g., Cerebellum or Neocortex). This absence of effect can stem from two

possibilities. Either foragers do not exploit cues left out by sympatric heterospecifics. Otherwise, it

has been shown that foragers tend to use social information over environmental (i.e., personal)

information, in particular in non-perfectly predictable environments (Rafacz and Templeton 2003;

Dunlap et al. 2016). Thus, if environmental complexity increases too much, "social" cues provided by

heterospecifics might replace environmental ones. As such, stimulation intensity of the MOB, the

Cerebellum, or the Neocortex would somehow remain equivalent when in sympatry or not. Further

work should explicitly test for these possibilities.

As expected (Hypothesis 3), the Striatum size was relatively larger in callitrichines, particularly

known to form mixed-species groups (Heymann and Hsia 2015). Yet, overall, the Striatum size was

negatively affected by sympatry. This puzzle might take root in secondary, but key, functions

supported by the Striatum, namely reward expectation, goal-directed behaviour, and planning

abilities (Johnson, Meer, and Redish 2007). These three functions might as well be advantageous

when foraging. As for the Hippocampus, then, the increase in environment unpredictability could

diminish the benefits of these future-oriented skills.

Given the context-dependence of the direction of selection (towards bigger sizes when sympatry is

low, smaller sizes otherwise), there is no surprise that we do not observe a correlation between the

net diversification rate and the three brain areas affected by species sympatry. Surprisingly

however, we found no positive association between the net diversification rate and the EQ, the

Cerebellum or the Neocortex, which were insensitive to species sympatry. By contrast, a positive

association between brain size and diversification was also found in birds (Sayol et al. 2019) given

that bigger brains act as a buffer to environmental challenges (Sol et al. 2007). A visual inspection of

the regressions clearly evidenced a positive trend if not considering phylogeny (EQ and Neocortex,

Figure S6). Sudden encephalisation in primates is clearly associated with a limited number of

closely-related species (DeCasien, Williams, and Higham 2017; Melchionna et al. 2020). Thus, this

clearly limits the statistical power of our phylogenetically-corrected analyses, as we cannot decipher

whether larger brain size and faster species diversification result from a true biological link or

appeared simultaneously but independently. This means that, despite what we found here, a

positive association between brain size and species diversification remains a likely possibility (as

previously suggested in primates, Melchionna et al. 2020). Species sympatry, however, induced a

significant slowdown in primate diversification, a density-dependence trend frequently observed in

many tetrapod clades (Condamine, Rolland, and Morlon 2019). This frames coherently with a

competitive scenario, where the tempo of species diversification decreases when ecological niches

are progressively filled up (Rabosky and Lovette 2008). Species competing for resources are

thought to contribute to limiting competitors' range (Price and Kirkpatrick 2009), hence

constraining population size and diversification rate (Pigot and Tobias 2013).

CONCLUSION

The use of brain size as a proxy for cognition is a central debate with no optimal solution (see

grounded criticism from Deaner, Nunn, and Schaik 2000; Healy and Rowe 2007; Logan et al. 2018).

The current flourishment of consortia, allowing for much more detailed and standardised

anatomical measurements (e.g., in primates: Milham et al. 2018), or with standardised

behaviourally explicit comparisons (e.g., on captive, Altschul et al. 2019; or wild primates, Janmaat

et al. 2021), might alleviate biases stemming from brain size analysis, but this will take time to

generate large-enough datasets. In the meanwhile, brain size is a proxy much appreciated in

practice, because of its easy accessibility for a "large" number of species, while the multifaceted

aspect of cognition can simply be taken into account by considering the brain as a mosaic of singular

and independent regionalised areas that are cognitively specialised. Here, we showed that species

sympatry is an important factor shaping the evolutionary history of animals' brains, but the

proximate mechanisms at play remain to be elucidated. Finally, it is very likely that any hypothesis

on cognition evolution, generally discussed within species, could be broadened to a between-species

context: foraging facilitation between species does exist (Olupot, Waser, and Chapman 1998;

Havmøller et al. 2021), and so do polyspecific social associations (Porter 2001), as well as inter-

species territory defence (Drury, Cowen, and Grether 2020; Losin et al. 2016) or imitation and

copying (Persson, Sauciuc, and Madsen 2018; Pepperberg 2002). Similarly, prey-predator races

could shape selection on cognitive abilities (Shultz and Dunbar 2006). As Alice said "It's a great huge

game of chess that's being played—all over the world" (Carroll 1871, chap. II) and all individuals are

just pieces to play with or against, no matter the species.

ACKNOWLEDGEMENTS

We considerably value the help provided by J. Drury in making some scripts available functions in

the *RPANDA* package in *R* and helping us run them, and that of M-C. Quidoz for assistance in using

the CEFE cluster. We thank S. Benhamou and M. Clairbaux for discussions on spatial projections, M.

Quéroué, V. Lauret, A. Caizergues, and C. Teplistky for feedback on Bayesian computations, and H.

Morlon for discussions on models of trait evolution. Finally, this work could not have been possible

without prior data collection from the IUCN Red List (primate ranging), the 10kTrees project

(phylogenetic trees), and A. R. DeCasien and collaborators, L. E. Powell and collaborators, O. S.

Todorov and collaborators, E. P. Willems and collaborators, F. Pearce and collaborators, A. F.

Navarrete and collaborators, and C. C. Grueter who provided primate trait data we used as

(supplementary) material with their articles. Their indirect input is therefore tremendous. Both

authors were supported by a doctoral grant from the *École Normale Supérieure*, Paris.

AUTHORS' CONTRIBUTION

482 BR conceived the study, gathered, cleaned and analysed the data, drew the figures and wrote the

483 first version of the manuscript and subsequently revised it. BP-L implemented the ClaDS algorithm

484 for our data, helped with running other analyses, and revised the manuscript multiple times. The

485 authors declare having no conflict of interest. All authors gave final approval for publication and

486 agree to be held accountable for the work performed therein.

DATA ACCESSIBILITY

488 The data and codes that support the findings of this study are openly available at

489 https://github.com/benjaminrobira/Meta_analysis_cognition_primates.

 Table 1: Species sympatry correlates negatively with the size of some brain areas of extant frugivorous primate

492 species | Model estimates and significance of phylogenetic regressions to assess the relationship between relative

493 brain sizes and species sympatry. Est.=Estimate, CI2.5%=Lower border of the CI95%, CI97.5%=Upper border of

494 the CI95%, Sd=Standard deviation, t=Statistics t-value. The brain areas (as well as the associated sample sizes)

495 are indicated prior to each list of estimates. The transformations applied to variables are indicated between

496 parentheses (logarithm, log, or square-root, sqrt), as well as the ponderation by bodymass (/bodymass).

	Est.	CI2.5%	CI97.5%	Sd	t	p-value
EQ (log) (N=127)						
Intercept	-0.17	-0.53	0.22	0.20	-	-
% of overlapped range	0.02	-0.08	0.13	0.05	0.41	0.68
Number of sympatric frugivores (sqrt)	0.02	-0.02	0.05	0.02	1.03	0.31
Lambda	0.98	0.94	1.00			
Hippocampus (/bodymass, log) (N=50)						
Intercept	-0.92	-1.95	0.05	0.53	-	-
% of overlapped range	-0.39	-0.76	-0.01	0.20	-1.94	0.06
Number of sympatric frugivores (sqrt)	0.08	-0.06	0.20	0.07	1.21	0.23
Lambda	0.99	0.92	1.00			
Neocortex (/bodymass, log) (N=56)						
Intercept	2.07	1.31	2.86	0.41	-	-

% of overlapped range	-0.23	-0.54	0.11	0.16	-1.46	0.15
Number of sympatric frugivores (sqrt)	0.02	-0.08	0.13	0.05	0.48	0.63
Lambda	0.99	0.91	1.00			

Cerebellum (/bodymass, log) (N=57)

Intercept	0.60	-0.15	1.35	0.39	-	-
% of overlapped range	-0.08	-0.32	0.17	0.12	-0.7	0.49
Number of sympatric frugivores (sqrt)	-0.01	-0.1	0.07	0.04	-0.34	0.74
Lambda	1.00	0.96	1.00			

Striatum (/bodymass, log) (N=50)

Intercept	-0.36	-1.18	0.44	0.44	-	-
% of overlapped range	**-0.40**	**-0.77**	**-0.04**	**0.18**	**-2.26**	**0.03**
Number of sympatric frugivores (sqrt)	0.03	-0.08	0.15	0.06	0.61	0.54
Lambda	0.98	0.85	1.00			

MOB (/bodymass, log) (N=31)

Intercept	-2.76	-4.61	-0.93	1.00	-	-
% of overlapped range	-1.20	-2.65	0.35	0.80	-1.49	0.15
Number of sympatric frugivores (sqrt)	0.21	-0.18	0.56	0.19	1.12	0.27

Lambda   1.00   1e-07   1.00

Table 2: Relative brain sizes do not correlate with primate species diversification | Model estimates and

significance of Bayesian phylogenetic regressions to assess the correlation between the net diversification rates

and the relative brain sizes. Est.=Estimate, HDP2.5%=Lower border of the 95% Highest Posterior Density,

HDP97.5%=Upper border of the 95% Highest Posterior Density, Eff. samp.=Effective sample (adjusted for

autocorrelation). The brain areas (as well as the associated sample sizes) are indicated prior to each list of

estimates. The (log) indicates log-transformed variables, while the (/body mass) indicates variables pondered by

body mass.

	Est.	HDP2.5%	HDP97.5%	Eff. samp	pMCMC
Diversification EQ (N=148)					
Intercept	0.12	0.08	0.16	900.00	-
EQ (log)	0.02	-7.91e-03	0.05	789.25	0.15
Lambda	0.83	0.76	0.9		
Diversification Hippocampus (N=61)					
Intercept	0.13	0.09	0.18	900.00	-
Hippocampus (/bodymass, log)	9.10e-03	-9.48e-03	0.03	900.00	0.34
Lambda	0.73	0.6	0.85		
Diversification Neocortex (N=67)					
Intercept	0.1	0.04	0.17	991.53	-

Neocortex (/bodymass, log)	7.26e-03	-0.02	0.03	900.00	0.56
Lambda	0.74	0.6	0.86		
Diversification Cerebellum (N=68)					
Intercept	0.12	0.07	0.16	900.00	-
Cerebellum (/bodymass, log)	3.94e-03	-0.02	0.03	989.21	0.76
Lambda	0.74	0.6	0.86		
Diversification Striatum (N=61)					
Intercept	0.12	0.08	0.17	900.00	-
Striatum (/bodymass, log)	9.11e-03	-0.01	0.03	900.00	0.44
Lambda	0.73	0.59	0.85		
Diversification MOB (N=37)					
Intercept	0.11	0.05	0.17	900.00	-
MOB (/bodymass, log)	-4.79e-03	-0.02	0.01	900.00	0.59
Lambda	0.65	0.46	0.83		

506 **Table 3: Species sympatry slowdowns primate diversification | Model estimates and significance of phylogenetic**

507 **regressions to assess the correlation between diversification rate and species sympatry. Est.=Estimate,**

508 **CI2.5%=Lower border of the CI95%, CI97.5%=Upper border of the CI95%, Sd= Standard deviation, t= Statistics t-**

509 **value. The brain areas (as well as the associated sample sizes) are indicated prior to each list of estimates. The**

510 **transformation (logarithm or square-root) is indicated in parentheses by the abbreviation (log or sqrt).**

	Est.	CI2.5%	CI97.5%	Sd	t	p-value
Diversification (N=128)						
Intercept	0.15	0.10	0.2	0.03	-	-
% of overlapped range	-5.40e-03	-0.02	9.35e-03	8.14e-03	-0.66	0.51
Number of sympatric frugivores (sqrt)	**-5.04e-03**	**-0.01**	**1.34e-04**	**2.56e-03**	**-1.97**	**0.05**
Lambda	0.96	0.89	0.99			

Figure 1: Biogeographic areas used for reconstructing the history of sympatry in frugivorous primates represented on the Mercator projection of the world | Areas were defined as a combination of geographic and environmental criteria relative to the primate taxonomy following results from Kamilar (2009): (1) East Madagascar (2) West Madagascar (3) West Africa (4) Central Africa (5) East/South Africa (6) Central America (7) North South-America (8) South South-America (9) West Asia (10) Central/East Asia (11) South Asia (12) Asian peninsula and islands. Note that the north part of Africa and the south of Europe were discarded because *Macaca sylvanus* was not considered.

Figure 2: The intensity of species sympatry varies across the primate phylogeny | Primate phylogeny from the consensus tree of the 10kTrees project is depicted in the center, together with abbreviated species names. The corresponding non-abbreviated names can be found using Appendix Figure S3. Sympatric frugivorous (based on a frugivory threshold of 20% and folivory threshold of 40%) species are linked by light grey lines. The geographic areas occupied by a species are depicted by coloured rectangles. Presence was assessed given an overlap between the species range and the geographic area of 10%.

Figure 3: Variations in relative brain size areas among extant frugivorous primates | (Left) Circular plot of the relative sizes of the different brain areas. Colours indicate the rows for the different brain areas. The darker background emphasises when values are above average, while the lighter background emphasises when values are below average. The mean value (after scaling and based on one random sampling among possible values, but see Supplementary Material Figure S2 for visualization of measure variability) for the Encephalization Quotient (EQ) or relative size of brain areas, when available, is depicted by a plain circle for frugivorous species. The frugivorous threshold was fixed to 20% and the folivory threshold to 40%. (Right) The different studied brain areas (human brain as an illustration). In short, the MOB is involved in immediate olfactory information

processing, the Neocortex and the Cerebellum support working memory and memory consolidation of immediate sensory information processing (Wiltgen et al. 2004; Koziol et al. 2014; Sokolov, Miall, and Ivry 2017), and the Hippocampus supports a working memory and a long-term spatio-temporal memory (Burgess, Maguire, and O'Keefe 2002). The Striatum is involved in social information processing (Báez-Mendoza and Schultz 2013).

Figure 4: The evolution of the Hippocampus and Striatum in frugivorous primates are best fitted by models of trait evolution considering species sympatry | Plotted is the AICc weight, a measure of relative support for a given model, for models not considering species sympatry (BM, OU, EB) or considering species sympatry (MC, DD_{lin}, DD_{exp}). The points represent the average AICc weight obtained (when considering the six models from the same run), while the vertical bars indicate the standard deviation given all tested conditions (see Phylogenetic models of trait evolution: does species sympatry shape brain size evolution?).

REFERENCES

Altschul, Drew M., Michael J. Beran, Manuel Bohn, Josep Call, Sarah DeTroy, Shona J. Duguid, Crystal L. Egelkamp, et al. 2019. "Establishing an infrastructure for collaboration in primate cognition research." *PLoS ONE* 14 (10): e0223675. https://doi.org/10.1371/journal.pone.0223675.

Ashton, Benjamin J., Patrick Kennedy, and Andrew N. Radford. 2020. "Interactions with conspecific outsiders as drivers of cognitive evolution." *Nature Communications* 11 (1): 1–9. https://doi.org/10.1038/s41467-020-18780-3.

Avarguès-Weber, A., E. H. Dawson, and L. Chittka. 2013. "Mechanisms of social learning across species boundaries" 290 (1): 1–11. https://doi.org/10.1111/jzo.12015.

Báez-Mendoza, Raymundo, and Wolfram Schultz. 2013. "The role of the striatum in social behavior." *Frontiers in Neuroscience* 7 (7 DEC): 233. https://doi.org/10.3389/fnins.2013.00233.

Barton, Robert A., and Paul H. Harvey. 2000. "Mosaic evolution of brain structure in mammals." *Nature* 405 (6790): 1055–58. https://doi.org/https://doi.org/10.1038/35016580.

Benson-Amram, Sarah, Ben Dantzer, Gregory Stricker, Eli M. Swanson, and Kay E. Holekamp. 2016. "Brain size predicts problem-solving ability in mammalian carnivores." *Proceedings of the National Academy of Sciences of the United States of America* 113 (9): 2532–37. https://doi.org/10.1073/pnas.1505913113.

Bivand, Roger, and Colin Rundel. 2021. *Rgeos: Interface to Geometry Engine - Open Source ('GEOS')*. https://CRAN.R-project.org/package=rgeos.

Blomberg, Simone P., Theodore Garland, and Anthony R. Ives. 2003. "Testing for phylogenetic signal in comparative data: Behavioral traits are more labile." *Evolution* 57 (4): 717–45. https://doi.org/10.1111/j.0014-3820.2003.tb00285.x.

Blomberg, Simone P., Suren I. Rathnayake, and Cheyenne M. Moreau. 2020. "Beyond brownian

motion and the ornstein-uhlenbeck process: Stochastic diffusion models for the evolution of

quantitative characters." *American Naturalist* 195 (2): 145–65. https://doi.org/10.1086/706339.

Bollback, Jonathan P. 2006. "SIMMAP: Stochastic character mapping of discrete traits on

phylogenies." *BMC Bioinformatics* 7 (1): 1–7. https://doi.org/10.1186/1471-2105-7-88.

Burgess, Neil, Eleanor A. Maguire, and John O'Keefe. 2002. "The human hippocampus and spatial

and episodic memory." *Neuron* 35 (4): 625–41. https://doi.org/10.1016/S0896-6273(02)00830-9.

Burnham, Kenneth P., and David R. Anderson. 2002. *Model selection and multimodel inference: A*

practical Information-theoretic approach. Second. New York: Springer.

Byrne, Richard W. 2000. "Evolution of primate cognition." *Cognitive Science* 24 (3): 543–70.

https://doi.org/10.1207/s15516709cog2403_8.

———. 2018. *Machiavellian intelligence retrospective*. Vol. 132. 4. Oxford University Press Oxford.

https://doi.org/10.1037/com0000139.

Carroll, Lewis. 1871. *Through the looking glass*. Macmillan London U.K.

Clutton-Brock, T. H., and Paul H. Harvey. 1980. "Primates, brains and ecology." *Journal of Zoology*

190 (3): 309–23. https://doi.org/10.1111/j.1469-7998.1980.tb01430.x.

Condamine, Fabien L., Jonathan Rolland, and Hélène Morlon. 2019. "Assessing the causes of

diversification slowdowns: temperature-dependent and diversity-dependent models receive

equivalent support." *Ecology Letters* 22 (11): 1900–1912. https://doi.org/10.1111/ele.13382.

de Almeida Rocha, Juliana Monteiro, Míriam Plaza Pinto, Jean Philippe Boubli, and Carlos Eduardo

Viveiros Grelle. 2015. "The role of competition in structuring primate communities under different

productivity regimes in the amazon." *PLoS ONE* 10 (12): e0145699.

https://doi.org/10.1371/journal.pone.0145699.

Deaner, Robert O., Charles L. Nunn, and Carel P. van Schaik. 2000. "Comparative tests of primate

cognition: Different scaling methods produce different results." *Brain, Behavior and Evolution* 55

(1): 44–52. https://doi.org/10.1159/000006641.

DeCasien, Alex R., and James P. Higham. 2019. "Primate mosaic brain evolution reflects selection on

sensory and cognitive specialization." *Nature Ecology and Evolution* 3 (10): 1483–93.

https://doi.org/10.1038/s41559-019-0969-0.

DeCasien, Alex R., Scott A. Williams, and James P. Higham. 2017. "Primate brain size is predicted by

diet but not sociality." *Nature Ecology and Evolution* 1 (5): 1–7. https://doi.org/10.1038/s41559-017-

0112.

Drury, Jonathan P., Julien Clavel, Marc Manceau, and Hélène Morlon. 2016. "Estimating the effect of

competition on trait evolution using maximum likelihood inference." *Systematic Biology* 65 (4):

700–710. https://doi.org/10.1093/sysbio/syw020.

Drury, Jonathan P., Madeline C. Cowen, and Gregory F. Grether. 2020. "Competition and

hybridization drive interspecific territoriality in birds." *Proceedings of the National Academy of*

Sciences of the United States of America 117 (23): 12923–30. https://doi.org/10.1073/pnas.1921380117.

Drury, Jonathan P., Joseph A. Tobias, Kevin J. Burns, Nicholas A. Mason, Allison J. Shultz, and Hélène

Morlon. 2018. "Contrasting impacts of competition on ecological and social trait evolution in

songbirds." *PLoS Biology* 16 (1): e2003563. https://doi.org/10.1371/journal.pbio.2003563.

Dunbar, Robin I. M., and Susanne Shultz. 2017. "Why are there so many explanations for primate

brain evolution?" *Philosophical Transactions of the Royal Society B: Biological Sciences* 372 (1727):

20160244. https://doi.org/10.1098/rstb.2016.0244.

Dunlap, Aimee S., Matthew E. Nielsen, Anna Dornhaus, and Daniel R. Papaj. 2016. "Foraging bumble

bees weigh the reliability of personal and social information." *Current Biology* 26 (9): 1195–99.

https://doi.org/10.1016/j.cub.2016.03.009.

Gómez-Robles, Aida, William D. Hopkins, and Chet C. Sherwood. 2014. "Modular structure facilitates

mosaic evolution of the brain in chimpanzees and humans." *Nature Communications* 5 (1): 1–9.

https://doi.org/10.1038/ncomms5469.

Gonda, Abigé, Gábor Herczeg, and Juha Merilä. 2013. "Evolutionary ecology of intraspecific brain

size variation: A review." *Ecology and Evolution* 3 (8): 2751–64. https://doi.org/10.1002/ece3.627.

González-Forero, Mauricio, and Andy Gardner. 2018. "Inference of ecological and social drivers of

human brain-size evolution." *Nature* 557 (7706): 554–57. https://doi.org/10.1038/s41586-018-0127-x.

Goodale, Eben, Guy Beauchamp, Robert D. Magrath, James C. Nieh, and Graeme D. Ruxton. 2010.

"Interspecific information transfer influences animal community structure." *Trends in Ecology and*

Evolution 25 (6): 354–61. https://doi.org/10.1016/j.tree.2010.01.002.

Grueter, Cyril C. 2015. "Home range overlap as a driver of intelligence in primates." *American*

Journal of Primatology 77 (4): 418–24. https://doi.org/10.1002/ajp.22357.

Hadfield, Jarrod D. 2010. "MCMC Methods for Multi-Response Generalized Linear Mixed Models: The

MCMCglmm R Package." *Journal of Statistical Software* 33 (2): 1–22. https://www.jstatsoft.org/v33/i02/.

Havmøller, Linnea W., J. Carter Loftus, Rasmus W. Havmøller, Shauhin E. Alavi, Damien Caillaud,

Mark N. Grote, Ben T. Hirsch, Lucia L. Tórrez-Herrera, Roland Kays, and Margaret C. Crofoot. 2021.

"Arboreal monkeys facilitate foraging of terrestrial frugivores." *Biotropica* 53 (6): 1685–97.

https://doi.org/10.1111/btp.13017.

Healy, Susan D., and Candy Rowe. 2007. "A critique of comparative studies of brain size."

Proceedings of the Royal Society B: Biological Sciences 274 (1609): 453–64.

https://doi.org/10.1098/rspb.2006.3748.

Herrmann, Esther, Josep Call, María Victoria Hernández-Lloreda, Brian Hare, and Michael

Tomasello. 2007. "Humans have evolved specialized skills of social cognition: The cultural

intelligence hypothesis." *Science* 317 (5843): 1360–66. https://doi.org/10.1126/science.1146282.

Heymann, Eckhard W, and Hannah M. Buchanan-Smith. 2000. "The behavioural ecology of mixed-

species troops of callitrichine primates." *Biological Reviews* 75 (2): 169–90.

https://doi.org/10.1111/j.1469-185X.1999.tb00044.x.

Heymann, Eckhard W, and Shin S Hsia. 2015. "Unlike fellows–a review of primate–non-primate

associations." *Biological Reviews* 90 (1): 142–56.

Hijmans, Robert J. 2021. *Geosphere: Spherical Trigonometry*. https://CRAN.R-

project.org/package=geosphere.

Ho, Lam Si Tung, and Cecile Ane. 2014. "A Linear-Time Algorithm for Gaussian and Non-Gaussian

Trait Evolution Models." *Systematic Biology* 63: 397–408.

Isler, Karin, and Carel P. van Schaik. 2009. "The Expensive Brain: A framework for explaining

evolutionary changes in brain size." *Journal of Human Evolution* 57 (4): 392–400.

https://doi.org/10.1016/j.jhevol.2009.04.009.

IUCN. 2021. *The IUCN Red List of Threatened Species*. Downloaded on the 19/01/2021.

https://www.iucnredlist.org.

Janmaat, Karline R. L., Christophe Boesch, Richard Byrne, Colin A. Chapman, Zoro B. Goné Bi,

Josephine S. Head, Martha M. Robbins, Richard W. Wrangham, and Leo Polansky. 2016. "Spatio-

temporal complexity of chimpanzee food: How cognitive adaptations can counteract the ephemeral

nature of ripe fruit." *American Journal of Primatology* 78 (6): 626–45.
https://doi.org/10.1002/ajp.22527.

Janmaat, Karline R. L., Colin A. Chapman, R. Meijer, and Klaus Zuberbühler. 2012. "The use of fruiting
synchrony by foraging mangabey monkeys: A 'simple tool' to find fruit." *Animal Cognition* 15 (1):
83–96. https://doi.org/10.1007/s10071-011-0435-0.

Janmaat, Karline R. L., Miguel de Guinea, Julien Collet, Richard W. Byrne, Benjamin Robira, Emiel van
Loon, Haneul Jang, et al. 2021. "Using natural travel paths to infer and compare primate cognition in
the wild." *iScience* 24 (4): 102343. https://doi.org/10.1016/j.isci.2021.102343.

Johnson, Adam, Matthijs AA van der Meer, and A. David Redish. 2007. "Integrating hippocampus and
striatum in decision-making." *Current Opinion in Neurobiology* 17 (6): 692–97.
https://doi.org/10.1016/j.conb.2008.01.003.

Kamilar, Jason M. 2009. "Environmental and geographic correlates of the taxonomic structure of
primate communities." *American Journal of Physical Anthropology* 139 (3): 382–93.
https://doi.org/10.1002/ajpa.20993.

Kashetsky, Tovah, Tal Avgar, and Reuven Dukas. 2021. "The cognitive ecology of animal movement:
Evidence from birds and mammals." *Frontiers in Ecology and Evolution* 0 (September): 633.
https://doi.org/10.3389/FEVO.2021.724887.

Koziol, Leonard F., Deborah Budding, Nancy Andreasen, Stefano D'Arrigo, Sara Bulgheroni, Hiroshi
Imamizu, Masao Ito, et al. 2014. "Consensus paper: The cerebellum's role in movement and
cognition." *Cerebellum* 13 (1): 151–77. https://doi.org/10.1007/s12311-013-0511-x.

Logan, Corina J., Shahar Avin, Neeltje Boogert, Andrew Buskell, Fiona R. Cross, Adrian Currie, Sarah
Jelbert, et al. 2018. "Beyond brain size: Uncovering the neural correlates of behavioral and cognitive

specialization." *Comparative Cognition and Behavior Reviews* 13 (na): 55–90. https://doi.org/10.3819/CCBR.2018.130008.

Losin, Neil, Jonathan P. Drury, Kathryn S. Peiman, Chaya Storch, and Gregory F. Grether. 2016. "The ecological and evolutionary stability of interspecific territoriality." *Ecology Letters* 19 (3): 260–67. https://doi.org/10.1111/ele.12561.

Maliet, Odile, Florian Hartig, and Hélène Morlon. 2019. "A model with many small shifts for estimating species-specific diversification rates." *Nature Ecology and Evolution* 3 (7): 1086–92. https://doi.org/10.1038/s41559-019-0908-0.

Maliet, Odile, and Hélène Morlon. 2021. "Fast and accurate estimation of species-specific diversification rates using data augmentation." *Systematic Biology*. https://doi.org/10.1093/sysbio/syab055.

Matzke, Nicholas J. 2013. "Probabilistic historical biogeography: new models for founder-event speciation, imperfect detection, and fossils allow improved accuracy and model-testing." *Frontiers of Biogeography* 5 (4): na. https://doi.org/10.21425/f5fbg19694.

———. 2016. "Stochastic mapping under biogeographical models." *PhyloWiki BioGeoBEARS* na (na): na.

Melchionna, M., A. Mondanaro, C. Serio, S. Castiglione, M. Di Febbraro, L. Rook, J. A. F. Diniz-Filho, et al. 2020. "Macroevolutionary trends of brain mass in Primates." *Biological Journal of the Linnean Society* 129 (1): 14–25. https://doi.org/10.1093/biolinnean/blz161.

Milham, Michael P, Lei Ai, Bonhwang Koo, Ting Xu, Céline Amiez, Fabien Balezeau, Mark G Baxter, et al. 2018. "An open resource for non-human primate imaging." *Neuron* 100 (1): 61–74.

Milton, Katharine. 1981. "Distribution Patterns of Tropical Plant Foods as an Evolutionary Stimulus

to Primate Mental Development." *American Anthropologist* 83 (3): 534–48.

https://doi.org/10.1525/aa.1981.83.3.02a00020.

Minot, Edward O. 1981. "Effects of interspecific competition for food in breeding blue and great tits."

The Journal of Animal Ecology 50 (2): 375. https://doi.org/10.2307/4061.

Modolo, Lara, Walter Salzburger, and Robert D. Martin. 2005. "Phylogeography of Barbary

macaques (*Macaca sylvanus*) and the origin of the Gibraltar colony." *Proceedings of the National*

Academy of Sciences of the United States of America 102 (20): 7392–97.

https://doi.org/10.1073/pnas.0502186102.

Morlon, H, E Lewitus, FL Condamine, M Manceau, J Clavel, and J Drury. 2016. "RPANDA: An r

Package for Macroevolutionary Analyses on Phylogenetic Trees." *Methods in Ecology and Evolution*

7: 589–97. https://CRAN.R-project.org/package=RPANDA.

Navarrete, Ana F., Erwin L. A. Blezer, Murillo Pagnotta, Elizabeth S. M. De Viet, Orlin S. Todorov,

Patrik Lindenfors, Kevin N. Laland, and Simon M. Reader. 2018. "Primate Brain Anatomy: New

Volumetric MRI Measurements for Neuroanatomical Studies." *Brain, Behavior and Evolution* 91 (2):

109–17. https://doi.org/10.1159/000488136.

Navarrete, Ana F., Carel P. van Schaik, and Karin Isler. 2011. "Energetics and the evolution of human

brain size." *Nature* 480 (7375): 91–93. https://doi.org/10.1038/nature10629.

Nuismer, Scott L., and Luke J. Harmon. 2015. "Predicting rates of interspecific interaction from

phylogenetic trees." *Ecology Letters* 18 (1): 17–27. https://doi.org/10.1111/ele.12384.

Olupot, William, Peter M. Waser, and Colin A. Chapman. 1998. "Fruit finding by mangabeys

(*Lophocebus albigena*): Are monitoring of fig trees and use of sympatric frugivore calls possible

strategies?" *International Journal of Primatology* 19 (2): 339–53. https://doi.org/10.1023/A:1020391901832.

Pearce, Fiona, Chris Carbone, Guy Cowlishaw, and Nick J. B. Isaac. 2013. "Space-use scaling and home range overlap in primates." *Proceedings of the Royal Society B: Biological Sciences* 280 (1751): 20122122. https://doi.org/10.1098/rspb.2012.2122.

Pennell, MW, JM Eastman, GJ Slater, JW Brown, JC Uyeda, RG Fitzjohn, ME Alfaro, and LJ Harmon. 2014. "Geiger V2.0: An Expanded Suite of Methods for Fitting Macroevolutionary Models to Phylogenetic Trees." *Bioinformatics* 30: 2216–18.

Pepperberg, Irene M. 2002. "Allospecific referential speech acquisition in grey parrots (*Psittacus erithacus*): Evidence for multiple levels of avian vocal imitation." *Imitation in Animals and Artifacts*, 109–31.

Persson, Tomas, Gabriela Alina Sauciuc, and Elainie Alenkær Madsen. 2018. "Spontaneous cross-species imitation in interactions between chimpanzees and zoo visitors." *Primates* 59 (1): 19–29. https://doi.org/10.1007/s10329-017-0624-9.

Pigot, Alex L., and Joseph A. Tobias. 2013. "Species interactions constrain geographic range expansion over evolutionary time." *Ecology Letters* 16 (3): 330–38. https://doi.org/10.1111/ele.12043.

Porter, Leila M. 2001. "Benefits of polyspecific associations for the Goeldi's monkey (*Callimico goeldii*)." *American Journal of Primatology* 54 (3): 143–58. https://doi.org/10.1002/ajp.1019.

Powell, Lauren E., Robert A. Barton, and Sally E. Street. 2019. "Maternal investment, life histories and the evolution of brain structure in primates." *Proceedings of the Royal Society B: Biological Sciences* 286 (1911): 20191608. https://doi.org/10.1098/rspb.2019.1608.

Powell, Lauren E., Karin Isler, and Robert A. Barton. 2017. "Re-evaluating the link between brain

size and behavioural ecology in primates." *Proceedings of the Royal Society B: Biological Sciences* 284

(1865): 20171765. https://doi.org/10.1098/rspb.2017.1765.

Price, Trevor D., and Mark Kirkpatrick. 2009. "Evolutionarily stable range limits set by interspecific

competition." *Proceedings of the Royal Society B: Biological Sciences* 276 (1661): 1429–34.

https://doi.org/10.1098/rspb.2008.1199.

R Core Team. 2020. "R: A language and environment for statistical computing." Vienna, Austria: R

Foundation for Statistical Computing. http://www.r-project.org https://www.r-project.org/.

Rabosky, Daniel L., and Irby J. Lovette. 2008. "Density-dependent diversification in North American

wood warblers." *Proceedings of the Royal Society B: Biological Sciences* 275 (1649): 2363–71.

https://doi.org/10.1098/rspb.2008.0630.

Rafacz, Michelle, and Jennifer J. Templeton. 2003. "Environmental unpredictability and the value of

social information for foraging starlings." *Ethology* 109 (12): 951–60. https://doi.org/10.1046/j.0179-

1613.2003.00935.x.

Reader, Simon M., and Kevin N. Laland. 2002. "Social intelligence, innovation, and enhanced brain

size in primates." *Proceedings of the National Academy of Sciences of the United States of America* 99

(7): 4436–41. https://doi.org/10.1073/pnas.062041299.

Revell, Liam J. 2012. "Phytools: An r Package for Phylogenetic Comparative Biology (*and Other*

Things)." *Methods in Ecology and Evolution* 3: 217–23.

Robira, Benjamin, Simon Benhamou, Shelly Masi, Violaine Llaurens, and Louise Riotte-Lambert.

2021. "Foraging efficiency in temporally predictable environments: Is a long-term temporal memory

really advantageous?" *Royal Society Open Science*.

Rosati, Alexandra G. 2017. "Foraging cognition: Reviving the Ecological Intelligence Hypothesis."

Trends in Cognitive Sciences 21 (9): 691–702. https://doi.org/10.1016/j.tics.2017.05.011.

Sayol, Ferran, Oriol Lapiedra, Simon Ducatez, and Daniel Sol. 2019. "Larger brains spur species

diversification in birds." *Evolution* 73 (10): 2085–93. https://doi.org/10.1111/evo.13811.

Schaik, Carel P. van, and Judith M. Burkart. 2011. "Social learning and evolution: The cultural

intelligence hypothesis." *Philosophical Transactions of the Royal Society B: Biological Sciences* 366

(1567): 1008–16. https://doi.org/10.1098/rstb.2010.0304.

Shultz, Susanne, and Robin I. M. Dunbar. 2006. "Chimpanzee and felid diet composition is influenced

by prey brain size." *Biology Letters* 2 (4): 505–8. https://doi.org/10.1098/rsbl.2006.0519.

Slater, GJ, LJ Harmon, D Wegmann, P Joyce, LJ Revell, and ME Alfaro. 2012. "Fitting Models of

Continuous Trait Evolution to Incompletely Sampled Comparative Data Using Approximate

Bayesian Computation." *Evolution* 66: 752–62.

Sokolov, Arseny A., R. Chris Miall, and Richard B. Ivry. 2017. "The cerebellum: Adaptive prediction

for movement and cognition." *Trends in Cognitive Sciences* 21 (5): 313–32.

https://doi.org/10.1016/j.tics.2017.02.005.

Sol, Daniel, Tamás Székely, András Liker, and Louis Lefebvre. 2007. "Big-brained birds survive

better in nature." *Proceedings of the Royal Society B: Biological Sciences* 274 (1611): 763–69.

https://doi.org/10.1098/rspb.2006.3765.

Todorov, Orlin S., Vera Weisbecker, Emmanuel Gilissen, Karl Zilles, and Alexandra A. De Sousa. 2019.

"Primate hippocampus size and organization are predicted by sociality but not diet." *Proceedings of*

the Royal Society B: Biological Sciences 286 (1914): 20191712. https://doi.org/10.1098/rspb.2019.1712.

Tomasello, Michael. 2019. *The cultural origins of human cognition.* Harvard university press.

https://doi.org/10.2307/j.ctvjsf4jc.

Trapanese, Cinzia, Benjamin Robira, Giordana Tonachella, Silvia Di Gristina, Hélène Meunier, and

Shelly Masi. 2019. "Where and what? Frugivory is associated with more efficient foraging in three

semi-free ranging primate species." *Royal Society Open Science* 6 (5): 181722.

https://doi.org/10.1098/rsos.181722.

Weir, Jason T., and Sara Mursleen. 2013. "Diversity-dependent cladogenesis and trait evolution in

the adaptive radiation of the auks (*Aves: Alcidae*)." *Evolution* 67 (2): 403–16.

https://doi.org/10.1111/j.1558-5646.2012.01786.x.

Whiten, Andrew, and Carel P. van Schaik. 2007. "The evolution of animal 'cultures' and social

intelligence." *Philosophical Transactions of the Royal Society B: Biological Sciences* 362 (1480): 603–

20. https://doi.org/10.1098/rstb.2006.1998.

Willems, Erik P., Barbara Hellriegel, and Carel P. van Schaik. 2013. "The collective action problem in

primate territory economics." *Proceedings of the Royal Society B: Biological Sciences* 280 (1759):

20130081. https://doi.org/10.1098/rspb.2013.0081.

Wilson, AC. 1991. "From molecular evolution to body and brain evolution." In *Perspectives on*

Cellular Regulation: From Bacteria to Cancer, 331–40. J. Wiley & Sons.

Wiltgen, Brian J., Robert A. M. Brown, Lynn E. Talton, and Alcino J. Silva. 2004. "New circuits for old

memories: The role of the neocortex in consolidation." *Neuron* 44 (1): 101–8.

https://doi.org/10.1016/j.neuron.2004.09.015.