The synergistic effects of plants and nitrogen on microbial

hitchhiking

Zhibin Liu[1,7], Ziyuan Wang[1,7], Qini Xia[1], Qin Zhou[1], Xiaobo

Wu[1], Wenqing Kong[1], Wenyan Lei[1], Jiayi Zeng[1], Chao Liu[2,3],

Yongfeng Wang[4,5], Wei Chang[6], Zhi Li[6], Yi Yang[1], Liang

Yang[6*], Xiao Tan[2,3*]

[1] Key Laboratory of Bio-Resource and Eco-Environment of

Ministry of Education, College of Life Sciences, Sichuan

University, Chengdu 610064, Sichuan, PR China

[2] State Key Laboratory of Hydraulics and Mountain River

Engineering, Sichuan University, Chengdu 610065, Sichuan,

PR China

[3] College of Water Resource & Hydropower, Sichuan

University, Chengdu 610065, Sichuan, PR China

[4] Institute of Environment and Ecology, Institute of

Environmental Health and Ecological Security, Jiangsu

University, 301 Xuefu Road, Zhenjiang, 212013, PR China.

[5] Guangdong Provincial Key Laboratory of Silviculture,

Protection and Utilization, Guangdong Academy of Forestry,

233 Guangshan 1st Road, Guangzhou, PR China.

[6] Vegetable Germplasm Innovation and Variety Improvement

Key Laboratory of Sichuan Province/Horticulture Research

Institute, Sichuan Academy of Agricultural Sciences, Chengdu

610066, PR China

[7] These authors contributed equally to this work and should be

considered co-first authors.

* To whom correspondence should be addressed.

Contact: yangliang_saas@foxmail.com   (Liang   Yang),

xiaotan@scu.edu.cn (Xiao Tan)

The synergistic effects of plants and nitrogen on

microbial hitchhiking

Abstract

Microbial hitchhiking demonstrates that some nonmotile

microbes utilize trans-species motility to traverse their

environment; however, whether driving forces, such as plants

and nitrogen, affect microbial hitchhiking is not clear. In our

study, we explored the effects of plants and nitrogen fertilizer

on *Bacillus*- hitchhiking by setting filter membranes and

different nitrogen fertilizer concentration gradients. In the

experimental treatment, we added a filter membrane to the soil

to prevent hitchhiking. In the absence of plants, nitrogen alone

had little influence on motile bacteria and hitchhiking.

However, *Bacillus* contents were significantly impacted by the

nitrogen concentration when the plants were rooted, leading to

a great variation in cell motility function according to the

functional analysis in the soil microbial community. After

applying the filter membrane, there were no significant

differences in *Bacillus* contents, microbial community structure

or cell motility functional abundance, which illustrated that

hitchhiking impacted the microbial community. Our analysis of

co-occurrence between bulk soil motile bacteria (*Bacillus*) and

rhizosphere bacteria also confirmed this. The correlation

between bulk soil motile bacteria and the rhizosphere microbial

community was strong in the groups with suitable nitrogen

concentrations without filter membranes and was weak at all

nitrogen levels in the no-membrane treatments. Thus, we

concluded that plants and different nitrogen doses

synergistically altered the soil microbiome by hitchhiking,

whose effect depends on nitrogen.

Keywords: microbial hitchhiking; rhizosphere microbes;

nitrogen fertilizer

1. Introduction

Cell motility is responsible for diverse and interesting

survival behaviors(Kearns 2010; Lambert et al. 2019). Many

free-living motile microbes use motility machinery to navigate

their surroundings in search of optimum conditions(Lambert et

al. 2019). Motile microbes, which can be attracted or repelled

by specific stimuli on surfaces, can travel by creeping, gliding,

rolling, swarming, or twitching, among other motile behaviors.

In addition to the microbe motility machinery, a novel mode of

dispersal used by many microbes, hitchhiking on motile

microbes, was recently reported (Schäfer et al. 1998; Jarrell and

McBride 2008; Kearns 2010). Hitchhiking enables nonmotile

microbes to navigate their world by successfully using the

motility machinery of their motile partner (Muok et al. 2021).

Transport by hitchhiking occurring among bacteria is found in

soils, plant tissues, abiotic surfaces, and human tissues(Muok

and Briegel 2021; Muok et al. 2021). Various species of

bacteria comigrate across air gaps, which allows nonmotile

bacteria to access novel microhabitats(Warmink and van Elsas

2009).

A previous study showed that plants shape rhizosphere

microbiome structures by hitchhiking through motility of

Bacillus   subtilis,   which

thrives near plant roots and moves toward the root(Muok   and

Briegel 2021; Muok et al. 2021). Bacterial motility

has two modes of flagellar-mediated motility, swimming and s

warming, and can also move through sliding(Muok et al. 2021).

It was reported that *Bacillus* was able to transport Streptomyces

spores to plant tissues. However, previous studies on how

Bacillus hitchhiking affected the soil microbiome were carried

out in an ideal laboratory environment without simulating the

real soil ecological environment. At the same time, in addition

to Streptomyces, it was not clear whether the other bacteria

could be carried by *Bacillus*.

Some physical and chemical properties of soil, including

pH and water content, determine the composition of the

microbial community (Garrett et al. 2008; Tsagkari and Sloan

2018; Lambert et al. 2019). Nitrogen, by impacting physical

and chemical properties, acts directly as a key factor

influencing microbial community structure(Good and Beatty

2011; Yuan et al. 2015; Zhou et al. 2017; Kavamura et al. 2018;

Sun et al. 2020). A recent study showed that various kinds of

bacteria were involved in community interactions via a

community co-occurrence network analysis under different

nitrogen treatments, but no sources of the microbes responsible

for the changes in the microbial community were identified

(Hartman and Tringe 2019). Moreover, it did not reveal

whether hitchhiking played an important role in the changes in

the microbial community.

To simulate the natural soil environment and explore the

influence of nitrogen on microbial hitchhiking, we planted

spinach under wild conditions by applying different nitrogen

concentrations and set up a blank control group (no plants). In

addition, more kinds of bacteria can be identified to determine

whether they are related to hitchhiking. To explore the impact

of hitchhiking on the community, we set the filter group to

isolate bacterial movement. Then, high-throughput sequencing

of the obtained amplicons was performed to illustrate the

changes in the bulk soil and rhizosphere microbial communities

under different nitrogen concentrations and with or without

spinach planting. Furthermore, the mechanism responsible for

the effects of filter membrane applications near rhizosphere soil

exposed to various nitrogen concentrations and with or without

spinach planting on plant growth was investigated. We

assessed the current cooccurrence network and studied the

activation of the symbiotic mode under various nitrogen

concentrations and with or without *spinach* cultivation, and we

found that nitrogen concentration and with or without plants

influenced both motile and their associated nonmotile bacteria

together.

2 Materials and methods

2.1 Sites and experimental setup

To investigate the coupled effect of fertilization and microorganism on plant growth, a pot experiment was conducted in Shuangliu District, Chengdu, China (102°52'11''E, 30°33'21''N). The Shuangliu District belongs to subtropical humid monsoon climate zone with an annual average air temperature range of 15.6□ to 16.9□, an annual average precipitation range of 759.1 mm to 1,155.0 mm, and an annual average sunshine duration range of 825.0 h to 1,202.9 h. In this region, the soil is yellow loam and mostly slightly acidic.

The pot experiment was conducted from October 2019 to February 2020. Three nitrogen levels (no addition of extra nitrogen, addition of 20% extra nitrogen, addition of 40% extra nitrogen), two planting conditions (with/without plants), and two coating conditions (with/without filter membrane) were considered, resulted in 12 treatments. Each treatment has 3 replications. A plot (4×7.4 m in size) was selected from a greenhouse in the experimental site, consisting of three subplots (3×1.8 m in size). Each subplot has 12 sampling points, distributed in 3 rows and 4 columns (each column represents one treatment). Intervals with a width of 30 cm

between each sampling point and buffer zones with a width of

30 cm between the side row/column and the plot boundary

were set to reduce the disturbance (Supplementary Information).

The soil used in this experiment was taken from each sampling

point (diameter: 20 cm, depth: 20 cm), then air-dried, ground to

pass through a sieve of 2 mm, and mixed well (guarantee the

same soil texture in all treatments). Urea, nitro-compound

fertilizer, and potassium sulfate were applied as base fertilizers

at rates of 226.5 kg/ha, 400.6 kg/ha, and 120 kg/ha,

respectively, in accordance with local agricultural practices.

Afterwards, urea was added to achieve different nitrogen

concentrations: base fertilizer (H1), base fertilizer + 70.5 kg/ha

urea (H2), base fertilizer + 141 kg/ha urea (H3). The

well-mixed soil of each nitrogen concentration was then packed

into 12 fiber pots (diameter: 20 cm, height: 20 cm, 8 kg soil for

each pot, growth of spinach roots would be not limited in the

pot in such size) which allows mass and energy exchange

between the soil inside and outside of the pot. Taking H1 as an

example, half of the 12 pots (6 pots) were coated by 0.45 μm

filter membrane (Fig. 1B). Then, all the prepared pots were

carefully embedded into sampling points with no gaps between

pots and surrounding soils. The soil surface in pots was flush

with the ground. The spinach (Spinacia oleracea L.) seeds were

soaked in distilled water for 6 h and then sown into half of the

coated pots (3 pots) and half of the uncoated pots (3 pots). Each

column refers to a treatment in a subplot, details are shown in

Supplementary Information. Excessive seedlings were removed

when the third true leaf emerged, and one seedling was left in

each pot. Water was supplied in accordance with local water

management practices. The same operations were repeated for

the H2 and H3. All abbreviations in the text and corresponding

treatment are shown in Fig. 1C.

2.2 DNA extraction, PCR amplification, and Illumina

NovaSeq 6000 sequencing

Sample collection was conducted in February 2020.

Subplots (0.5 m × 0.5 m) were randomly selected from each

treatment site. Soil samples (0–15-cm depths) were drilled

using a soil sampler with a diameter of 3.5 cm. Each soil

sample was cut vertically in the middle, put into a sterile

self-sealing bag, and transported on ice. They were then stored

in a refrigerator at −80°C for microbial sequencing. Microbial

DNA was extracted using a HiPure Soil DNA Kit (or HiPure

Stool DNA Kit) (Magen, Guangzhou, China) in accordance

with the manufacturer's protocol. The 16S rDNA target regions

of the ribosomal RNA gene were amplified by PCR using the

following amplification protocol: 94°C for 2 min, followed by

30 cycles at 98°C for 10 s, 62°C for 30 s (except for 16S V4,

which was 55°C for 30 s), and 68°C for 30 s, with a final

extension at 68°C for 5 min. PCRs were performed in triplicate.

Each 50-µL mixture contained 5 µL of 10× KOD Buffer, 5 µL

of 2 mM dNTPs, 3 µL of 25 mM MgSO4, 1.5 µL of each

primer (10 µM), 1 µL of KOD polymerase, and 100 ng of

template DNA. The PCR reagents were purchased from

TOYOBO, Japan. Amplicons were extracted from 2% agarose

gels, purified using an AxyPrep DNA Gel Extraction Kit

(Axygen Biosciences, Union City, CA, USA) in accordance

with the manufacturer's instructions, and quantified using an

ABI StepOnePlus Real-Time PCR System (Life Technologies,

Foster City, CA, USA). Purified amplicons were pooled in

equimolar ratios and paired-end sequenced (PE250) on an

Illumina platform using standard protocols.

2.3 Community composition analysis

The representative OTU sequences were classified into

organisms by a naive Bayesian model using RDP classifier[15]

(version 2.2) based on SILVA[16] database (version 132), with

the confidence threshold value of 0.8 (Table S1, Table S2). The

stacked bar plot of the community composition was visualized

in R project ggplot2 package (version 2.2.1).

2.4 Statistical analyses

2.4.1 Alpha- and beta-diversity analyses

Chao1 and Sob indices were calculated using QIIME (Caporaso et al. 2010) (version 1.9.1). An OTU rarefaction curve, as well as rank abundance curves, were plotted in R project ggplot2 package (version 2.2.1, URL: http://CRAN. R-project. org/package= ggplot2, URL: http://CRAN.R-project.org/package=ggplot2). An alpha-index comparison between groups was calculated using Welch's t-tests in the R project Vegan package (version 2.5.3). Alpha-index comparisons among groups were conducted using Kruskal–Wallis H tests in the R project Vegan package (version 2.5.3).

The sequence alignment was performed using Muscle (Edgar 2004) (version 3.8.31), and a phylogenetic tree was constructed using FastTree(Price et al. 2010) (version 2.1). Then, weighted and unweighted unifrac distance matrices were generated using the GuniFrac package (Lozupone and Knight 2005) (version 1.0) in R project. Bray–Curtis distance matrices were calculated in the R project Vegan package (version 2.5.3). Multivariate statistical techniques, including a principal

coordinates analysis (PCoA) and the Bray–Curtis distances,

were generated in the R project Vegan package (version 2.5.3)

and plotted in R project ggplot2 package(version 2.2.1.).

Statistical analyses of Welch's t-tests, Kruskal-Wallis H tests,

and Anosim tests were calculated in the R project Vegan

package (version 2.5.3).

2.4.2 Functional predictions

The KEGG pathway analysis of the OTUs was inferred

using Tax4Fun (version 1.0)(Aßhauer et al. 2015). Tax4Fun

arranges all the samples, including SILVA species and

annotated OTUs. An absolute abundance table, which is a

program based on the SILVA species annotation information

database, selects OTU reference species and then combines the

OTU abundance table and species-gene nets. The output of

each sample is the relative abundance of the KEGG Orthology

pathway.

2.4.3 Co-occurrence analysis

Graphical representations were generated using Cytoscape

(Version 3.8.2). A bacterial co-occurrence analysis was

computed in the Vegan package of R 4.0.2. One network per

treatment combination was calculated to compare the effects of

different nitrogen concentrations. The input data for the visual

comparison were mean abundances over the three field

replicates per time point collapsed into taxonomic families. For

statistical evaluation, topological networks were computed for

each plot, and Cytoscape (Version 3.8.2) was used to generate

graphical representations of the resulting consensus networks.

The average degree, average clustering coefficient, and average

path length were computed and compared statistically using

distance and Pearson's correlation values to show the

co-occurrence networks of each bacterial class.

3. Results

3.1 Bulk and rhizosphere soil compositional differences

Major microbial classes in the bulk soil without plants

were *Gammaproteobacteria*, *Alphaprotebacteria*, and

Planctomycetacia (Fig. 2A, Table S1). The content of

Planctomycetacia increased with the nitrogen concentration,

and this phenomenon was blocked by the filter membrane. In

addition, more classes were influenced by nitrogen

(*Planctomycetacia*, *Verrucomicrobiae*, and *Actinobacteria*).

Nevertheless, the percentages of *Planctomycetacia*,

Verrucomicrobiae, and *Actinobacteria* did not change

significantly in the filter membrane treatment. Compared with

treatments without plants, *Bacteroidia* accounted for a larger

proportion of the community, while Planctomycetacia

accounted for a smaller proportion (Fig. 2B). In the rhizosphere

microbial community, *Gammaproteobacteria* abundance in H2

was greater than that in H1 and H3, irrespective of the

membrane (Fig. 2C).

In the soil microbial community without plants, there was

no significant difference in the content of *Bacillus* regardless of

the filter membrane (Fig. 2A). When there were plants in the

soil, there was a significant difference in the content of *Bacillus*

only if there was no filter membrane ($p < 0.05$ without filter

membrane and $p = 0.0714$ with; Fig. 2B). In the rhizosphere

microbial community, the content of *Bacillus* at different

nitrogen concentrations showed a similar trend as the soil

microbial community with the plants ($p < 0.05$ without filter

membranes and $p = 0.9286$ with; Fig. 2C).

3.2 Microbial alpha- and beta-diversity levels

The Sob indices were used to estimate and compare the

alpha-diversity of bacterial communities among different

nitrogen treatments (Table 1A, Table S3). In treatments with or

without the filter membrane, whether plants were added and the

bulk soil/rhizosphere bacterial species diversity was not

significantly different among the three different nitrogen

concentrations (P> 0.05; Table 1A, Table S4A). PCoA showed

a distinct clustering pattern for the microbiota in H1-NF/

H2-NF/ H3-NF(Fig. S1A), H1-S-NF/ H2-S-NF/ H3-S-NF (Fig.

3A) and H1-R-NF/H3-R-NF/H2-R-NF (Fig. 3B).

At the class level, for the bulk soil microbial community

with and without plants, the bacterial beta-diversity (P =

0.003**, Table 1B; P=0.026*, Table S4B) varied greatly with

nitrogen concentration. Nevertheless, there were no significant

differences in beta-diversity levels among the different nitrogen

concentration groups after membrane addition (P = 0.263,

Table 1B; P=0.2; Table S4B). Moreover, there was no

significant change in the beta diversity of the soil microbial

communities between groups with and without filter-membrane

(Fig. S1 B-D). A further analysis of the rhizosphere microbial

colonies showed the same results as the beta-diversity analysis

in the bulk soil microbial community with plants.

We discovered differences between the microbial

communities at the same nitrogen concentration with and

without the filter membrane treatment (Fig. 3C– E). In the bulk

soil, both alpha and beta diversity showed significant

differences between groups with and without filter membranes,

whether there were plants or not. In a comparison of H1-R-NF

and H1-R-F, although no significant difference in their

alpha-diversity levels was found, there was a significant

difference in their beta-diversity levels, which reflects a

significant difference in their microbial community structures.

When comparing H2-R-NF to H2-R-F and H3-R-NF to H3-R-F,

the same result was found. Thus, although there were great

effects of nitrogen on the bacterial community, the addition of

the membrane counteracted the effects.

3.3 Abundance of function

To determine the functional changes among the different

nitrogen treatments, we utilized and compared the relative

functional abundances of the different groups(Table S5). In the

absence of plants, nitrogen fertilizer had very little effect on

cell motility abundance (Fig. S2 A). However, in

H1-NF/H2-NF/H3-NF, the greatest functional abundance

difference occurred in the H2 group compared with the other

two treatments in both bulk and rhizosphere soils, and the

functions were weakened in the membrane-addition treatments

(Fig. 4A; Fig. S2 B). In particular, cell motility showed

significant differences, with H2-S/R-NF showing significantly

more cell motility than H1-S/R-NF and H3-S/R-NF ($P < 0.05$;

Fig. 4 C-D). However, the level of cell motility in the

membrane-addition group was not significantly different

among the different nitrogen treatments. In addition, this

phenomenon was not observed in soil microbial communities

regardless of whether plants were grown.

3.4 Co-occurrence networks between bulk soil and

rhizosphere soil microbial communities under different

nitrogen concentrations

We used Cytoscape (Version 3.8.2) to draw the

Co-occurrence networks and analyze the interactions of

microbial communities exposed to different nitrogen

concentrations. The Pearson's and distance correlation values

between classes were calculated based on their occurrences

across samples using the R project Vegan package. The

microorganism classes with high abundance values and high

correlations between nodes were selected, and an

Co-occurrence network diagram was constructed. Because

certain bacteria can survive only in soil at a certain nitrogen

concentration, we chose the top 19 most abundant groups for

the analysis. To establish the Co-occurrence network graph,

edges with Pearson's correlation coefficients greater than 0.7

were selected.

While the number of edges in the bacteria indicating an

association between bulk soil and rhizosphere microbial

community increased with the nitrogen concentration, the

numbers of edges under H2 and H3 were significantly greater

than those under H1, which was consistent with groups having

the filter-membrane treatment. In H2, without the

filter-membrane treatment, the relationships between the bulk

soil motile bacteria (*Bacillus*) and the rhizosphere

microorganisms were significantly stronger than those in the

other two groups (Fig. 5A–C). However, in groups with the

filter membrane treatment, the numbers of edges indicated that

associations with motile bacteria were significantly reduced in

all three treatments compared with groups without filter

membranes. Moreover, there were no differences among the

three treatments with the filter membrane (Fig. 5D–F).

3.5 Plant trait data

Without the 0.45-µm filter membrane, plant growth

differed under different nitrogen concentrations. The *spinach* in

H2 was larger than the *spinach* in H1 and H3, and the *spinach*

fresh weight in H2 was 23.3% and 14.6% higher than the

weights in H1 and H3, respectively. After filter membrane

addition, there was no significant difference in *spinach* growth

among the three nitrogen treatments. Moreover, *spinach* grown

without the filter membrane coating grew better than *spinach*

grown with the filter membrane coating. The fresh weights of

spinach grown without the filter membrane were 48.5%

(H1-NF versus H1-F), 56.2% (H2-NF versus H3-F), and 57.8%

(H3-NF versus H3-F) greater than those grown with the filter

membrane addition (Fig. 6).

4. Discussion

Microorganisms in the soil play significant roles in plant

development(Warmink and van Elsas 2009; Zhao et al. 2014;

Sheng et al. 2021; Sun et al. 2021). They are an essential

component of soil ecology and serve as an important

mechanism for soil regeneration. However, culture-independent

approaches have shown that the microbial diversity levels of

the soil and rhizosphere are highly underestimated(Chaparro et

al. 2014; Korenblum et al. 2020; Wang et al. 2020). Moreover,

the driving forces behind changes in microbial diversity are

unclear. Our research found that hitchhiking strongly affects

bacterial community structure, which is strongly regulated by

nitrogen concentration and plants.

4.1 Plant and nitrogen concentrations affect microbial

composition, especially *Bacillus*

From the analysis of microbial composition, *Bacillus* was

found to have no significant changes among the different

nitrogen concentrations in the groups without plants, which

showed that the content of *Bacillus* could not be affected by

nitrogen fertilizer. In groups with plants, when 20% nitrogen

fertilizer was added, the content of *Bacillus* was highest

compared with the other two groups, which suggested that

nitrogen fertilizer can affect the recruitment of *Bacillus*. Under

different nitrogen concentrations, the quality of root exudates,

such as terpenoid secretion, often changed, which was

previously believed to help root colonization of *Bacillus*.

Furthermore, this trend was prevented by the filter membrane,

which indicated that the filter membrane can prevent the

motility of *Bacillus*.

Similar to the contents of *Bacillus*, there was no significant

difference in beta diversity when plants were absent. With

plants, the bulk soil and rhizosphere microbial community

structure of the groups without filter membranes showed

significant beta-diversity levels. However, the addition of the

filter membrane eliminated the effects of different nitrogen

fertilizer treatments on the community structure. These results

suggest that nitrogen fertilizer and plants synergistically

regulate soil and rhizosphere microbial communities through

microbial migration. This is consistent with previous studies on

the driving forces of prokaryotic community change, which

concluded that soil physical and chemical properties affect

cellular dynamics and then microbial communities(Torsvik et

al. 2002; Mitchell and Kogure 2006).

4.2 Plant and nitrogen concentrations have important

effects on cell motility

If there were no plants in the soil, the functional abundance

of cell motility showed no great difference between different

nitrogen concentrations. This indicated that nitrogen had no

effect on cell motility alone. When plants are present, nitrogen

concentration has a greater effect on cell motility. Moreover,

the addition of the filter membrane weakened the variation in

the functional abundance of cell motility among different

nitrogen concentrations, which corresponds to the content of

Bacillus under different conditions. These results suggest that

nitrogen fertilizer and plants synergistically regulate microbial

motility. Especially when plants have no membrane, cell

motility and *Bacillus* of H2 are the highest compared with other

plants. Nitrogen can trigger genes related to motility in motile

bacteria and rewiring of the nitrogen regulation system(Zhou et

al. 2017; Kavamura et al. 2018; Sun et al. 2020; Karthikeyan et

al. 2021; Wang et al. 2021). At the same time, plant secretions

were different under different concentrations. Under

appropriate nitrogen concentrations, plant secretions might

have higher sensitivity to *Bacillus*(Yuan et al. 2015).

4.3 The effect of microbial hitchhiking was promoted by a

proper concentration of nitrogenous fertilizer

Studies using microscopy methods, motility assays, and

genetic approaches have shown that *Bacillus* transports

Streptomyces spores by direct attachment (hitchhiking) to the

bacterial flagella to utilize plant root exudate as a nutrient

source(Muok et al. 2021). Based on this, *Bacillus* was selected

to explore whether its interactions with other bacteria were

stronger. We discovered that more bacteria were impacted by

the motile *Bacillus* in H1, and this effect was also observed in

H3 but not in H2. In the membrane-addition treatment group,

there were fewer interactions between motile bacteria in the

rhizosphere and soil under the three different nitrogen

concentrations. Thus, the presence of a filter membrane

precluded the interactions between soil and rhizosphere

microorganisms mediated by motile bacteria. We concluded

that changes in nitrogen concentration affect the motility of and

hitchhiking onto motile bacteria.

The co-occurrence network, which provides insights into

ecological interactions among microbial taxa, shows that the

interactions between different microbial strains are the main

driving factors of population structure and dynamics because

microbes can co-occur or exclude each other, revealing that

different microorganism metabolites can have an inhibitory or

symbiotic impact. Co-occurrence network analyses have shown

that taxonomic shifts owing to the variation in nitrogen

availability can shape patterns of ecological interactions

regulating the structure, function, and potential resilience of

soil microbial communities (Schmidt et al. 2019; Han et al.

2020). However, these networks have focused only on the soil

or rhizosphere microbiome and did not take into account the

interactions between these two communities. Our cooccurrence

network analysis combining soil and rhizosphere microbes

showed that there were variations in the cooccurrence of bulk

soil and rhizosphere microbial species at varying nitrogen

concentrations.

4.4 Nitrogen influenced plant growth by hitchhiking

Morphological variations in plants may result from growth

in soil with different fertilizer concentrations (Sun et al. 2021).

Fertilization with an appropriate nitrogen content promotes

plant growth, while fertilization with a high nitrogen content

inhibits plant growth, which is consistent with our results (Liu

et al. 2014; Razaq et al. 2017; Elhanafi et al. 2019; Bindraban

et al. 2020; Bright et al. 2021). Previous studies have suggested

that root length and root surface area increased under

intermediate N levels and that root growth was reduced under

both higher and lower fertilization levels, affecting plant

morphology(Good and Beatty 2011; Chen et al. 2020). On the

other hand, nitrogen is functional in the construction of and

chlorophyll, which can influence plant growth and development

by affecting photosynthesis and the uptake of minerals(Grossart

et al. 2001; Razaq et al. 2017; Kavamura et al. 2018; Wang et

al. 2021). In addition to these reasons, in our study, we found

that in the treatment group with filter-membrane addition, there

was no significant difference in the growth of *spinach* with

different nitrogen concentrations. Through correlation analysis,

we found that the hitchhiking phenomenon was blocked by the

filter membrane; as a result, we presumed that nitrogen may

affect plant growth by affecting hitchhiking. In the rhizosphere,

compared with H1 and H3, there was an increase in functional

abundance encouraging plant growth (amino acid metabolism,

energy metabolism, and cofactor and vitamin metabolism) and

a decline in functional abundance (infectious disease)

discouraging plant growth in the H2 group without a filter

membrane. At suitable concentrations, the nonmotile bacteria

brought by the hitchhiking phenomena modified the

rhizosphere microbial environment via transport on favorable

bacterial species, resulting in improved plant growth. It can be

concluded that nitrogen fertilizer can promote plant growth by

promoting microbial hitchhiking.

5. Conclusion

Nitrogen alone has no effect on hitchhiking, but it will

synergistically affect the movement of sports bacteria with

plants, thus causing the carrying of sports bacteria. Moreover,

different concentrations of nitrogen fertilizer have different

effects on hitchhiking. The appropriate concentration will

greatly improve, while an excessive concentration will reduce

its effect. These studies provide guidance for rational

fertilization.

Acknowledgments

This project was funded by the National Natural Science

Foundation of China (NSFC Grant No. 31870240 and No.

51909175), the Key Research and Development Program of

Sichuan Province (No. 2019YFN0153) and the Modern

Agriculture Discipline Construction and Promotion Program of

Sichuan Academy of Agricultural Sciences (2021XKJS032).

We thank Mr. Mingyuan Wang for his guidance and advice

during the preparation of this manuscript.

Author contributions

LZB, YL, LC, YY and TX designed the experiment; CW, ZQ, KWQ, WXB, and LZ conducted the experiments; WZY and XQN analyzed the data; WZY, LWY, LXR and WYF wrote the manuscript.

Ethics approval

Not applicable.

Conflict of interest

The authors declare no competing interests.

Reference

Aßhauer KP, Wemheuer B, Daniel R, Meinicke P (2015) Tax4Fun: Predicting functional profiles from metagenomic 16S rRNA data. Bioinformatics 31:2882–2884. https://doi.org/10.1093/bioinformatics/btv287

Bindraban PS, Dimkpa CO, Pandey R (2020) Exploring phosphorus fertilizers and fertilization strategies for improved human and environmental health. Biology and Fertility of Soils 56:299–317. https://doi.org/10.1007/s00374-019-01430-2

Bright MBH, Diedhiou I, Bayala R, et al (2021) An overlooked local resource: Shrub-intercropping for food production, drought resistance and ecosystem restoration in the Sahel. Agriculture, Ecosystems & Environment 319:107523. https://doi.org/10.1016/j.agee.2021.107523

Caporaso JG, Kuczynski J, Stombaugh J, et al (2010) QIIME allows analysis of high-throughput community sequencing data. Nature Methods 7:335–336. https://doi.org/10.1038/nmeth.f.303

Chaparro JM, Badri D v., Vivanco JM (2014) Rhizosphere microbiome assemblage is affected by plant development. ISME Journal

8:790–803. https://doi.org/10.1038/ismej.2013.196

Chen J, Liu L, Wang Z, et al (2020) Nitrogen Fertilization Increases Root Growth and Coordinates the Root–Shoot Relationship in Cotton . Frontiers in Plant Science 11

Edgar RC (2004) MUSCLE: Multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32:1792–1797. https://doi.org/10.1093/nar/gkh340

Elhanafi L, Houhou M, Rais C, et al (2019) Impact of Excessive Nitrogen Fertilization on the Biochemical Quality, Phenolic Compounds, and Antioxidant Power of *Sesamum indicum* L Seeds. Journal of Food Quality 2019:9428092. https://doi.org/10.1155/2019/9428092

Garrett TR, Bhakoo M, Zhang Z (2008) Bacterial adhesion and biofilms on surfaces. Progress in Natural Science 18:1049–1056. https://doi.org/10.1016/J.PNSC.2008.04.001

Good AG, Beatty PH (2011) Fertilizing Nature: A Tragedy of Excess in the Commons. PLOS Biology 9:e1001124. https://doi.org/10.1371/journal.pbio.1001124

Grossart HP, Riemann L, Azam F (2001) Bacterial motility in the sea and its ecological implications. Aquatic Microbial Ecology 25:247–258. https://doi.org/10.3354/ame025247

Han Q, Ma Q, Chen Y, et al (2020) Variation in rhizosphere microbial communities and its association with the symbiotic efficiency of rhizobia in soybean. ISME Journal 14:1915–1928. https://doi.org/10.1038/s41396-020-0648-9

Hartman K, Tringe SG (2019) Interactions between plants and soil shaping the root microbiome under abiotic stress. Biochemical Journal 476:2705–2724. https://doi.org/10.1042/BCJ20180615

Jarrell KF, McBride MJ (2008) The surprisingly diverse ways that prokaryotes move. Nature Reviews Microbiology 6:466–476. https://doi.org/10.1038/nrmicro1900

Karthikeyan S, Orellana LH, Johnston ER, et al (2021) Metagenomic Characterization of Soil Microbial Communities in the Luquillo Experimental Forest (Puerto Rico) and Implications for Nitrogen Cycling. Appl Environ Microbiol 87:e0054621. https://doi.org/10.1128/AEM.00546-21

Kavamura VN, Hayat R, Clark IM, et al (2018) Inorganic Nitrogen Application Affects Both Taxonomical and Predicted Functional Structure of Wheat Rhizosphere Bacterial Communities . Frontiers

Kearns DB (2010) A field guide to bacterial swarming motility. Nature Reviews Microbiology 8:634–644. https://doi.org/10.1038/nrmicro2405

Korenblum E, Dong Y, Szymanski J, et al (2020) Rhizosphere microbiome mediates systemic root metabolite exudation by root-to-root signaling. Proc Natl Acad Sci U S A 117:3874–3883.

https://doi.org/10.1073/pnas.1912130117

Lambert BS, Fernandez VI, Stocker R (2019) Motility drives bacterial
encounter with particles responsible for carbon export throughout
the ocean. Limnology and Oceanography Letters 4:113–118.
https://doi.org/10.1002/lol2.10113

Liu CW, Sung Y, Chen BC, Lai HY (2014) Effects of nitrogen fertilizers
on the growth and nitrate content of lettuce (Lactuca sativa L.).
International Journal of Environmental Research and Public Health
11:4427–4440. https://doi.org/10.3390/ijerph110404427

Lozupone C, Knight R (2005) UniFrac: A new phylogenetic method for
comparing microbial communities. Applied and Environmental
Microbiology 71:8228–8235.
https://doi.org/10.1128/AEM.71.12.8228-8235.2005

Mitchell JG, Kogure K (2006) Bacterial motility: Links to the
environment and a driving force for microbial physics. FEMS
Microbiology Ecology 55:3–16.
https://doi.org/10.1111/j.1574-6941.2005.00003.x

Muok AR, Briegel A (2021) Intermicrobial Hitchhiking: How Nonmotile
Microbes Leverage Communal Motility. Trends in Microbiology
29:542–550. https://doi.org/10.1016/j.tim.2020.10.005

Muok AR, Claessen D, Briegel A (2021) Microbial hitchhiking: how
Streptomyces spores are transported by motile soil bacteria. The
ISME Journal 15:2591–2600.
https://doi.org/10.1038/s41396-021-00952-8

Price MN, Dehal PS, Arkin AP (2010) FastTree 2 - Approximately
maximum-likelihood trees for large alignments. PLoS ONE
5:e9490. https://doi.org/10.1371/journal.pone.0009490

Razaq M, Zhang P, Shen HL, Salahuddin (2017) Influence of nitrogen and
phosphorous on the growth and root morphology of Acer mono.
PLoS ONE 12:e0171321.
https://doi.org/10.1371/journal.pone.0171321

Schäfer A, Ustohal P, Harms H, et al (1998) Transport of bacteria in
unsaturated porous media. Journal of Contaminant Hydrology
33:149–169. https://doi.org/10.1016/S0169-7722(98)00069-2

Schmidt JE, Kent AD, Brisson VL, Gaudin ACM (2019) Agricultural
management and plant selection interactively affect rhizosphere
microbial community structure and nitrogen cycling. Microbiome
7:1–18. https://doi.org/10.1186/s40168-019-0756-9

Sheng Z, Zhu W, Yao H, et al (2021) Niche selection by soil bacterial
community of disturbed subalpine forests in western Sichuan.
Forests 12:505. https://doi.org/10.3390/f12040505

Sun B, Bai Z, Bao L, et al (2020) Bacillus biofertilizer mitigating
agricultural ammonia emission and shifting ssubtilisoil nitrogen
cycling microbiomes. Environment International 144:105989.

702 https://doi.org/10.1016/j.envint.2020.105989

703 Sun Y, Snow D, Walia H, Li X (2021) Transmission Routes of the
704  Microbiome and Resistome from Manure to Soil and Lettuce.
705  Environmental   Science   &   Technology.
706  https://doi.org/10.1021/acs.est.1c02985

707 Torsvik V, Øvreås L, Thingstad TF (2002) Prokaryotic diversity -
708  Magnitude, dynamics, and controlling factors. Science (1979)
709  296:1064–1066. https://doi.org/10.1126/science.1071698

710 Tsagkari E, Sloan WT (2018) The Role of the Motility of
711  Methylobacterium in Bacterial Interactions in Drinking Water.
712  Water  2018,  Vol  10,  Page  1386  10:1386.
713  https://doi.org/10.3390/W10101386

714 Wang X, Wang M, Xie X, et al (2020) An amplification-selection model
715  for quantified rhizosphere microbiota assembly. Science Bulletin
716  65:983–986. https://doi.org/10.1016/j.scib.2020.03.005

717 Wang Y, Liao S, Gai Y, et al (2021) Metagenomic Analysis Reveals
718  Microbial Community Structure and Metabolic Potential for
719  Nitrogen Acquisition in the Oligotrophic Surface Water of the
720  Indian  Ocean.  Frontiers  in  Microbiology  12:1–13.
721  https://doi.org/10.3389/fmicb.2021.518865

722 Warmink JA, van Elsas JD (2009) Migratory response of soil bacteria to
723  Lyophyllum sp. strain Karsten In soil microcosms. Applied and
724  Environmental  Microbiology  75:2820–2830.
725  https://doi.org/10.1128/AEM.02110-08

726 Yuan J, Zhang N, Huang Q, et al (2015) Organic acids from root exudates
727  of banana help root colonization of PGPR strain Bacillus
728  amyloliquefaciens  NJN-6.  Scientific  Reports  5:1–8.
729  https://doi.org/10.1038/srep13438

730 Zhao J, Ni T, Li Y, et al (2014) Responses of bacterial communities in
731  arable soils in a rice-wheat cropping system to different fertilizer
732  regimes  and  sampling  times.  PLoS  ONE  9:e85301.
733  https://doi.org/10.1371/journal.pone.0085301

734 Zhou J, Jiang X, Wei D, et al (2017) Consistent effects of nitrogen
735  fertilization on soil bacterial communities in black soils for two crop
736  seasons  in  China.  Scientific  Reports  7:1–10.
737  https://doi.org/10.1038/s41598-017-03539-6

Figure 1. (A) The plots had the following soil treatment conditions with three different nitrogen fertilizer concentrations (H1–3): natural, untreated soil (NF) with plants (H1-NF, H2-NF, and H3-NF), filter membrane (F) used to separate the rhizosphere soil from the outer bulk soil (H1-F, H2-F, and H3-F). There were two treatments for each area, a no spinach and no membrane addition treatment and a spinach and **(B)** 0.45-μm filter membrane addition. **(C)** All abbreviations in the text and corresponding treatment are shown in (C).

Figure 2. Comparison of bacterial community classification differences (class level) under different nitrogen treatments. (A, B, C) The relative abundances of Bacillus contents in soil without plants(A), without plants(B) and rhizosphere (C). H1, H2, and H3, normal, normal + 20%, and normal + 40% nitrogen fertilizer concentrations, respectively. S, bulk soil. R, spinach rhizosphere soil. F, filter membrane present, and NF, no filter membrane.

* p<0.05,** p<0.01

Figure 3. Beta-diversity levels of bulk soil and rhizosphere bacterial communities. (A) The PCoAs of soil and rhizosphere bacteria. Soil type is the main source of bacterial community changes, and each spot corresponds to different soil-type samples. (B) Significance test analysis of rhizosphere bacteria. The ends of the line represent the minimum and maximum values, and the line in the box is the middle value. H1, H2, and H3, normal, normal + 20%, and normal + 40% nitrogen fertilizer concentrations, respectively. S, spinach planted. R, spinach rhizosphere soil. F, filter membrane present, and NF, no filter membrane.

A

Carbohydrate Metabolism
Amino Acid Metabolism
Membrane Transport
Signal Transduction
Metabolism of Cofactors and Vitamins
Energy Metabolism
Nucleotide Metabolism
Translation
Replication and Repair
Xenobiotics Biodegradation and Metabolism
Lipid Metabolism
Metabolism of Terpenoids and Polyketides
Metabolism of Other Amino Acids
Glycan Biosynthesis and Metabolism
Cell Motility
Cell Growth and Death
Biosynthesis of Other Secondary Metabolites
Endocrine System
Cell Communication

1.97
-0.04
-2.06

H1-R-NF H2-R-NF H3-R-NF H1-R-F H2-R-F H3-R-F

B

	H1-NF	H1-F
	H2-NF	H2-F
	H3-NF	H3-F
P-value	0.258	0.070

C

	H1-S-NF	H1-S-F
	H2-S-NF	H2-S-F
	H3-S-NF	H3-S-F
P-value	0.031*	0.174

D

	H1-R-NF	H1-R-F
	H2-R-NF	H2-R-F
	H3-R-NF	H3-R-F
P-value	0.024*	0.386

Figure 4. The relative abundances of the top 20 classes of the most abundant rhizosphere bacteria plotted as a functional heatmap. H1, H2, and H3, normal, normal + 20%, and normal + 40% nitrogen fertilizer concentrations, respectively(A). Cell Motility was selected and marked red, and its test in bulk soil without plants (B), with plants (C) and rhizosphere(D) and different nitrogen concentration. R, spinach rhizosphere soil. F, filter membrane present, and NF, no filter membrane.

Figure 5. Microbial interaction networks of the different treatments. Different nitrogen treatments (A–C) without the filter membrane (NF) and (D–F) with the filter membrane (F). H1, H2, and H3, normal, normal + 20%, and normal + 40% nitrogen fertilizer concentrations, respectively. The interaction network of dominant microbiota at the class level (top 19) between bulk soil (Blue nodes) and rhizosphere (Red Nodes). The sizes of the nodes indicate the abundance levels of different bacterial classes, and the different colors indicate the corresponding location of the bacteria. The edge colors represent positive (green) and negative (orange) correlations. Only significant interactions are shown (r > 0.7). The motile Bacillus in the bulk soil are separated from other bacteria.

Figure 6. Comparison of spinach growth under different nitrogen concentrations (H1–3), with (F) or without (NF) the filter membrane. (A) Photos of mature spinach; (B) Statistical analysis of spinach fresh weights. Error bars are standard errors, and asterisks indicate significance at **$P < 0.01$. H1, H2, and H3, normal, normal + 20%, and normal + 40% nitrogen fertilizer concentrations, respectively.