The evolution of information transmission in mammalian brain networks

Alessandra Griffa[1,2,3], Mathieu Mach[2], Julien Dedelley[2], Daniel Gutierrez-Barragan[4],

Alessandro Gozzi[4], Gilles Allali[3,5], Joanes Grandjean[6,7], Dimitri Van De Ville[2,1], Enrico

Amico[2,1]

[1] Department of Radiology and Medical Informatics, University of Geneva, Geneva,

Switzerland

[2] Institute of Bioengineering, Center for Neuroprosthetics, École Polytechnique Fédérale De

Lausanne (EPFL), Geneva, Switzerland

[3] Leenaards Memory Centre, Lausanne University Hospital and University of Lausanne,

Lausanne, Switzerland

[4] Functional Neuroimaging Laboratory, Center for Neuroscience and Cognitive systems,

Istituto Italiano di Tecnologia, Rovereto, Italy

[5] Division of Cognitive and Motor Aging, Department of Neurology, Albert Einstein College of

Medicine, Yeshiva University, Bronx, NY, USA

[6] Department of Medical Imaging, Radboud University Medical Center, 6525 GA Nijmegen,

The Netherlands

[7] Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center,

6525 EN Nijmegen, The Netherlands

Corresponding authors

Alessandra Griffa - orcid.org/0000-0003-1923-1653

B1 0 265.042 (Campus Biotech Bâtiment B1)

Ch. des Mines 9

CH-1202 Genève

alessandra.griffa@gmail.com

Enrico Amico - orcid.org/0000-0001-6705-9689

H4 3 153.079 (Campus Biotech Bâtiment H4)

Ch. des Mines 9

CH-1202 Genève

enrico.amico@epfl.ch

Abstract

Brain communication, defined as information transmission through white-matter connections,

is at the foundation of the brain's computational capacities that virtually subtend all aspects

of behavior: from sensory perception shared across mammalian species, to complex

cognitive functions in humans. How did communication strategies in macroscale brain

networks adapted across evolution to accomplish increasingly complex functions? By

applying a novel approach to measure information transmission in mouse, macaque and

human brains, we found an evolutionary gradient from selective information processing,

where brain regions share information through single polysynaptic pathways, to parallel

information processing, where regions communicate through multiple parallel pathways. In

humans, parallel processing acts as a major connector between unimodal and transmodal

systems. Communication strategies are unique to individuals across different mammalian

species, pointing at the individual-level specificity of information routing architecture. Our

work provides compelling evidence that different communication strategies are tied to the

evolutionary complexity of mammalian brain networks.

Introduction

Understanding how brain function can be supported by patterns of neural signaling through its structural backbone is one of the enduring challenges of network and cognitive neuroscience[1]. The brain is effectively a complex system, a network of neural units interacting at multiple spatial and temporal scales through the white-matter wiring[2,3]. *Information transmission through structural connections*, which can be defined as *brain communication*[1], give rise to macroscale patterns of synchronous activity–or functional connectivity–between remote areas of the brain. Communication processes are at the foundation of the brain's computational capacities that virtually subtend all aspects of behavior, from sensory perception and motor functions shared across mammalian species, to complex human functions including higher-level cognition[4]. From an evolutionary perspective, high communication efficiency at minimal structural wiring cost has long been recognized as a fundamental attribute constraining the evolution of neural systems[5–7]. Yet, quantitative and comparative assessments of macroscale communication processes in brain networks of increasing evolutionary complexity are lacking[1,8].

Systems-level neuroscience has made different attempts to map brain communication as interrelated patterns of macroscale structural and functional brain connectivity, highlighting strikingly complex structure-function interdependencies[9]. Structurally connected region pairs tend to have stronger functional connectivity than disconnected pairs[10,11], suggesting the presence of monosynaptic interactions[12]. Nonetheless, direct structural connections alone are not able to explain most of the dynamic functional repertoire observed in a functioning brain[13]. Beyond monosynaptic interactions, functional connectivity between remote brain areas is likely to emerge from more complex, higher-order communication mechanisms that involve larger groups of neural elements and their structural interconnections, by polysynaptic (multi-step) routing of neural information[1,14,15]. Higher-order communication in neural systems is important from both neurocognitive and evolutionary perspectives. Functional connectivity patterns extending beyond pairwise-connected regions (i.e., not constrained by the direct structural connections underneath) are highly specific to individuals, reflect behavioral traits[16], and have been identified across different mammals[17]. Moreover, functional patterns untethered from structure are dominant in cortical areas that underwent larger evolutionary expansion across primates, suggesting a relation between local information transmission mechanisms and evolution[18–20].

Nonetheless, the information transmission mechanisms implemented in mammalian brain networks are, to date, largely unknown. A first hypothesis has been that shortest structural

paths are favored for neural communication as they allow more direct (faster, metabolically

less expensive) information transmission. In support of this hypothesis, brain networks of

several mammalian and simpler species have short structural path length[21,22] at the price of

a relatively high wiring cost[5]. This suggests that shortest paths contribute to efficient

communication in brain networks and have been selected throughout evolution despite their

high wiring cost. However, relying on path length as the sole measure of information routing

may be an oversimplification[14]. Shortest-path communication explains a limited portion of

functional connectivity[10] and excludes a large fraction of brain network connections and

near-optimal alternative pathways from the communication process[23]. Recent studies have

started to account for more complex mechanisms of information routing, such as parallel

communication (i.e., relay of information through multiple, parallel communication

pathways)[23–26] or convergent routing (i.e., neural signals' interaction through convergent

pathways)[27,28]. Indeed, in many real-world systems, information transmission unfolds through

numerous alternative pathways[29]. Nevertheless, we do not know what is the relative

contribution of single-pathway versus parallel-pathway communication in mammalian neural

systems.

Comparative neuroimaging provides instruments to understand the emergence of function

across evolution[21,30]. Evidence of similarities between neural systems in different species are

assumed to reflect common organizational principles and functions that may be

evolutionarily preserved. In contrast, regions showing the greatest changes between

humans and other species highlight neural changes that may account for features of

cognition unique to humans. It has been shown that the overall topology of the structural and

functional brain networks is preserved across evolution[21,22] despite large variations in brain

size and cortical expansion[20]. However, differences in local connectivity patterns and

functional dynamics exist[31–33]. Recent reports suggest that mammalian neural processing is

organized along multiple hierarchies from unimodal to transmodal regions, describing how

information from distinct neural populations are integrated and segregated across the

cortex[20,34]. Yet, a comprehensive account of how evolution has shaped cortical organization

requires a way to measure how brain communication mechanisms change *in vivo* across

species, while also taking into account the underlying structural architecture. Are macroscale

brain communication mechanisms preserved across mammalian species? Or contrarily, are

there distinct brain communication strategies relating to mammals' evolution? , In particular,

was there a shift from selective (single-pathway) to parallel processing across evolution to

support increasingly complex brain functions? These questions do not have trivial answers

and demand for new ways of assessing brain communication across different species.

Here we apply a novel approach, rooted in information theory, to measure polysynaptic
information transmission in macroscale brain networks. Taking advantage of structural and
functional connectivity information extracted from multimodal brain data (i.e., functional MRI,
diffusion MRI, tract tracing), we explore the intricate pathways of communication in the
mouse, monkey and human connectomes. We employ information-theoretical
principles[24,35,36] to identify the structural pathways selected for neural communication by
different neural systems, and measure the level of selective and parallel information
processing across the different species. We report a strong evolutionary gradient in the brain
communication dynamics of mammals, with predominant selective information routing in
lower mammalian species such as mice and macaques, morphing into more complex
communication patterns in human brains. Parallel communication strategies appear to have
acted as a major connector of unimodal (sensory, attentional) and transmodal (fronto-
parietal, default mode) areas in the human brain, possibly contributing to the evolution of
more complex cognitive functions in humans. Notably, we also found that brain
communication strategies are highly specific to individuals across the different mammalian
species. Our results link for the first time the complexity of macroscale brain communication
dynamics inferred from *in vivo* data to an evolutionary gradient across mammalian lineages.
These findings pave the way to a deeper understanding of how brain communication and its
relationship to function have evolved across species.

Results

We propose a new framework to identify communication pathways in brain networks and

investigate their evolution in three mammalian species of increasing phylogenetic

complexity: mice, monkeys, and humans (Fig. 1). These species represent distinct

mammalian lineages and include animal models (mice, monkeys) often used in translational

research. We aimed to formally test two general assumptions on brain communication

dynamics. First, due to the noisy nature of neural signaling, neural messages transmitted

through the structural brain network can keep at most the same amount of information

present at the source region[24,35]. This holds true for many communication systems where the

information content tends to decay as one moves away from the information source[25].

Second, in an information transmission process, messages are typically *relayed* through a

set of statistically independent steps[44]; i.e., neural messages do not contain memory of the

transmission process itself and communication happens in a Markovian fashion[37]. These two

assumptions –information decay and memoryless transmission– are formally summarized by

a fundamental principle of information theory, the data processing inequality (*DPI*)[36], which

we here apply to cross-species structural data and fMRI recordings (Online Methods). Given

a structural network representing the white matter wiring of the brain, we first identify sets of

short polysynaptic paths connecting each pair of brain regions (Fig. 1a). Next, we

quantitatively assess which and how many of those structural paths are effectively selected

for neural communication. To this aim, we quantify fMRI-derived mutual information

measures along the paths (Fig. 1b) to assess the *DPI* on those paths (Fig. 1c). From here, a

parallel communication score can be computed for every pair of brain regions, by counting

the number of paths that respect the *DPI* (Fig. 1d). Parallel communication scores portray a

spectrum of communication strategies from *selective information processing*, where brain

regions selectively exchange information through a single pathway, to *parallel information*

processing, where regions communicate through multiple, parallel pathways (Fig. 1e). We

assessed parallel communication scores at the individual and group levels, and summarized

them for distinct brain systems and single brain regions in comparison to appropriate null

models. Finally, we compared the distribution of selective and parallel information

processing across the three different mammalian species.

Data for this study consisted of open-source whole-brain structural connectivity matrices and

individual resting-state functional MRI recordings of 100 healthy human subjects, 9 macaque

monkeys, and 10 wild-type mice, all in their young adulthood (Online Methods,

Supplementary Table 1; see below for replication datasets). Group-representative structural

connectivity matrices with comparable number of brain regions were derived from diffusion

MRI and tract tracing data, and weighted by the Euclidean distance between connected

regions (Supplementary Figure 1). Individual-level mutual information matrices were
computed from fMRI recordings of comparable duration and temporal resolution across
species.
185

186
Figure 1 Identifying communication pathways in macroscale brain networks. (a) A
weighted and symmetric structural connectivity matrix summarizes the white matter wiring of
the brain for each species. For every pair of brain regions *(i, j)*, the 5 shortest structural paths
(light blue) connecting the two regions are identified using the *k*-Shortest path algorithm[23].
(b) For every subject (human participant or animal), the mutual information between region
pairs is computed from *z*-scored regional timecourses obtained from fMRI recordings. **(c)** By
analyzing the mutual information values along each structural path, the data processing
inequality (*DPI*) is used to assess whether the specific paths represent valid communication
channels between regions *i* and *j*. Left panel: two brain regions *i, j* are connected by a
structural path crossing regions x_1, x_2; green lines represent direct structural connections
(white matter fibers). Each region is associated with a neural activity-related timecourse; the
amount of information shared by two regions is quantified by their mutual information *I*
(darker and thicker arcs indicate stronger *I*). Right panel: a structural path (*i*, x_1, x_2, *j*) is
labeled as communication (relay) channel if the pairwise mutual information values do not
increase along the (undirected) path (first row, red shading); it is not a communication
channel otherwise (second row, gray shading: $I_{j,i} > I_{x2,i}$). **(d)** A parallel communication score
(*PCS*) is computed at the individual level (i.e., for every subject *n*) and for every pair of brain
regions *i, j* by counting the number of structural paths that serve as relay channels between
the two regions. **(e)** Parallel communication scores are investigated across mammalian
species, highlighting a spectrum of communication strategies from selective information

processing (light yellow; low *PCS*), to parallel information processing (dark brown; high
PCS).
209
210
211 **Parallel communication in brain networks follows an evolutionary gradient across**
212 **mammalian species**
213
Using our approach we found that, in mammalian brains, polysynaptic structural paths are
used to relay information in a 'Markovian'-specific, sequential processing fashion. For all the
three considered species, the whole-brain density of relay communication pathways (i.e., the
percentage of structural paths respecting the *DPI*) was higher than in a strict null model
preserving the structural connectivity architecture and the multivariate statistics of fMRI
timecourses (Online Methods; $p < 10^{-5}$ for all species). This held true when considering
either the first shortest path connecting region pairs (mean±standard deviation across
individuals: humans = 39.5±3.8%; macaques = 37.1±6.0%; mice = 37.1±2.3%), or longer
paths (humans = 33.0±5.6%; macaques = 29.5±5.7%; mice = 28.2±5.9%), showing that
relay communication is not limited to the shortest path only (Fig. 2a). Specifically, the
communication density level of the shortest and second shortest paths was comparable for
all the three species, but decayed right after. Next, we assessed the amount of parallel
communication between all brain region pairs. We found that, on average, the parallel
communication score (*PCS*) progressively increases from mice and macaques to humans
(median [5-, 95-percentile] across region pairs: mice = 1.30 [0.20, 2.50]; macaques = 1.33
[0.11, 2.56]; humans = 1.52 [0.11, 3.02];). This is particularly evident when considering the
long-tailed distribution of human brain network communication as compared to the animals
(Fig. 2b). The three species' *PCS* distributions were pairwise statistically different (two-
sample Kolmogorov-Smirnov tests human-macaque: $D_{4950,3321} = 0.157$, $p < 10^{-42}$; human-
mouse: $D_{4950,3003} = 0.171$, $p < 10^{-47}$; macaque-mouse: $D_{3321,3003} = 0.058$, $p < 10^{-4}$). These
results indicate that, as brain complexity increases across the phylogenetics tree, interareal
communication is increasingly subserved by parallel processing (Fig. 2).
236

Figure 2 Parallel communication follows an evolutionary gradient across mammalian species. Left: drawing of human, macaque, and mouse brains; each row in the figure corresponds to one species. **(a)** Box plots representing the percentage of short paths in individual brain networks used for relayed communication (i.e., respecting the *DPI*). Each colored dot represents an individual; gray dots represent species-specific null distributions obtained from permutation of mutual information values (Online Methods); circles and vertical bars indicate mean ± one standard deviation across individuals or randomizations. Paths are grouped according to the 1st up to the 5th shortest path between region pairs, showing that relay communication is not limited to the 1st shortest path only. **(b)** Group-average parallel communication score (*PCS*) matrices representing *PCS*s between every pair of brain regions, averaged across individuals. For each species, brain regions are organized according to meaningful functional circuits which are highlighted by black squares along the matrices' diagonals (Online Methods). On the right, the histograms of the average *PCS* scores across region pairs highlight an evolutionary gradient from mice, with lower *PCS*s and mainly selective information processing, to humans, with higher *PCS*s and presence of parallel communication. Median [5-, 95-percentile] *PCS* values for each species are reported atop each histogram.

Communication complexity is species-dependent and relates to the functional organization of mammalian brains

When evaluating the spatial localization of the relay communication pathways, we found that it followed the characteristics of each species' functional cortical architecture (Fig. 3). Specifically, in lower species the relay (mostly sequential) pathways mainly encompassed

unimodal/multimodal regions spanning the barrel, auditory and somatomotor cortices in
mice, and the visual, somatomotor and dorsal attention cortices in macaques (Fig. 3a). In
humans we found similar evidence of relay sequential processing in unimodal and
multimodal areas, but also a high concentration of parallel communication pathways in
transmodal regions including association cortices of the executive-control network and the
precuneus of the default mode network (Fig. 3a; note the different color scales across
species).

Next, we investigated the communication patterns at the level of region pairs within and
between different brain functional systems (Online Methods, Supplementary Fig. 3, 4). In
mice, the relay pathways mainly connected brain nodes belonging to unimodal systems,
including auditory, barrel, somatomotor and visual cortices (Fig. 3b). A similar distribution
was observed for relay pathways in macaque networks, mainly connecting visual with
somatomotor and dorsal attentional regions. However, a gradient transitioning towards
transmodal regions started to appear between the default mode network and attentional
systems (Fig. 3b). In humans, stronger (parallel) relay communication mainly connected
somatosensory and attention regions with executive-control and default mode systems,
forming *cross-modal parallel streams* between unimodal and transmodal regions (outside-
diagonal entries in Fig. 3b). Notably, these patterns were stable at the individual level. We
report in Fig. 3c the amount of relay communication (average *PCS*s) within unimodal
systems, within transmodal systems, and between unimodal and transmodal systems for
each subject. Within species, the amount of relay communication varied between systems,
with transmodal networks consistently presenting the lower amount of relay communication
(Kruskal-Wallis tests, $p < .05$) (Fig. 3c). Across species, relay communication within
unimodal systems decreased with increasing phylogenetic complexity (mice > macaques >
humans, $H(2) = 22.45$, $p = .000013$), while cross-modal communication between unimodal
and transmodal regions strongly increased (mice < macaques < humans, $H(2) = 46.44$, $p <
10^{-10}$), with humans presenting 30% larger *PCS*s than macaques (Supplementary Fig. 2).
Relay communication within transmodal systems was relatively stable across species ($H(2)
= 6.24$, $p = .044$). Taken together, these findings show that communication strategies are
highly heterogeneous across the brain network and are partially preserved across evolution.
However, phylogenetically older species demonstrate more developed relay communication
for lower-order processing between unimodal and multimodal regions. Conversely, the
human brain is characterized by stronger parallel communication that serves as the main
neural processing stream between unimodal and transmodal areas[34,38].

Figure 3 Relay communication strategies reflect the functional organization of mammalian brains. (a) Cortical distributions of relay communication, quantified as the average *PCS* of each brain region with the rest of the brain network (first row: human, *fsaverage6* cortical surface; second row: macaque, *F99 template*; third row: mouse, *ABI template*). For each species, the light yellow-to-brown colormap is scaled between the 5th and 95th percentiles of the cortical values. On the right, the average nodal communication scores per brain system are represented in the bar plots. **(b)** Average *PCS*s within and between brain systems, for humans, macaques and mice. Brain systems have been organized into unimodal/multimodal regions (upper-left black square) and transmodal regions (lower-right black square). **(c)** Average *PCS*s between unimodal systems, between transmodal systems, and between unimodal and transmodal systems (*cross*-modal communication) for individual subjects. In the box plots, each dot represents an individual; vertical bars indicate mean ± standard deviation; notch bars indicate median and 1st-3rd quartiles; shaded areas indicate 1st-99th percentiles. Kruskal Wallis *p*-values for within-species comparisons are reported. Control = executive-control; Dorsal attn. = dorsal attention; ventral attn. = ventral attention.

Relay communication patterns are unique to individuals

Our results revealed a link between relay information processing strategies and phylogenetic
complexity in the mammalian brain. Are the observed communication patterns specific to
individual subjects within single species? We addressed this question by exploring the
identifiability properties[39,40] of the parallel communication matrices reported in Fig. 2, across
the three different species. To this aim, the fMRI recording of each subject was split into two
sections of equal duration. From these, test and retest parallel communication matrices were
computed. Note that, at the individual level, the matrices' entries (*PCS*s) can take integer
values between 0 and 5, with 0 indicating no relay communication, 1 indicating perfectly
selective information processing, and 5 strongly parallel information processing. We
quantified the similarity between test and retest data as the percentage of brain regions'
pairs with exactly the same *PCS* (Jaccard similarity index). The individual identifiability
through relay communication patterns was then quantified as the success rate (SR), i.e., the
percentage of subjects whose identity was correctly predicted out of the total number of
subjects for each species[40]. We found that parallel communication scores allow to identify
individual mammals in all the three species, at a level that exceeds chance-level (humans:
SR = 87.0%, null = 0.9±1.0%; macaques: SR = 66.7%, null = 11.2±10.3%; mice: SR =
60.0%, null = 10.6±9.2%) (Fig. 4a,b). However, individual identifiability decreased from
humans, to macaque, to mice. Intriguingly, the major contribution to individual identifiability
was given by brain regions pairs that, on average, tends to communicate through multiple
parallel rather than selective pathways. When splitting region pairs into two groups ('*low-
PCS*', '*high-PCS*') according to group-average parallel communication scores, the success
rate obtained from *high-PCS* values was higher than the one obtained from *low-PCS* values
for both humans and macaques (*PCS* threshold = 1.3; humans: $SR_{low\text{-}PCS}$ = 73.0%, $SR_{high\text{-}PCS}$
= 85.0%; macaques: $SR_{low\text{-}PCS}$ = 44.4%, $SR_{high\text{-}PCS}$ = 55.5%; mice: $SR_{low\text{-}PCS}$ = 40.0%, $SR_{high\text{-}}$
$_{PCS}$ = 40.0%; see Supplementary Table 2 for alternative thresholds) (Fig. 4c,d). No
differences were found in mice. Taken together, these data suggest that, within the inherent
constraints of each species, individual subjects may implement distinct communication
strategies to relay neural information through the brain network, particularly when
considering higher-order communication mechanisms such as parallel communication.

349
Figure 4 Parallel communication is unique to individuals. Left: drawing of human,
macaque, and mouse brains; each row in the figure corresponds to one species. **(a)**
Identifiability matrices for the three species, reporting subjects' similarities between test
(rows) and retest (columns) parallel communication data. Test-retest similarity was
quantified with the Jaccard similarity index. **(b)** Box plots representing *self*-similarity (*Iself*,
diagonal entries of the identifiability matrix) and *others*-similarity (*Iothers*, out-diagonal
entries of the identifiability matrix) values. **(c)** *Self*- and *others*-similarity values when
considering only region pairs with *low* parallel communication scores (PCS ≤ 1.3 on
average). **(d)** *Self*- and *others*-similarity values when considering only region pairs with *high*
parallel communication scores (PCS > 1.3 on average). The success rate (SR) for subjects'
identification is reported for each pair of box plots. In the box plots, vertical bars indicate
mean ± standard deviation; notch bars indicate median and 1st-3rd quartiles; shaded areas
indicate 1st-99th percentiles.

Robustness, sensitivity and replication analyses

To ensure the validity of our results, we asked whether parallel communication scores and
their cross-species gradients could be explained by the different structural connectome
architectures and multivariate statistical properties of fMRI recordings alone. To this aim, we
constructed null distributions of parallel communication scores for each species, by randomly
shuffling the fMRI time series across brain regions and computing surrogate *PCS*s on the
original structural connectome architecture (*n* = 3000) (Online Methods, Supplementary Fig.
5). By *z*-scoring the real *PCS* scores with respect to the null distributions and applying

appropriate statistical thresholds, we show that our findings do not trivially derive from

structural connectivity and fMRI statistical properties alone. In particular, the gradient of

increasing parallel communication from mice to humans (Fig. 2, Supplementary Fig. 6, 7, 8)

and the cortical topographies of parallel communication density across species (Fig. 3,

Supplementary Fig. 9) remained significantly different from the null ones.

Next, we investigated whether results were sensitive to some methodological choices,

including number of brain regions (brain parcellation), fMRI time series length, and number

of subjects. We found that our results are robust to these factors. In humans, *PCS* scores

and their cortical topographies were comparable when subdividing the cortex into 100 or 200

regions of interest (Supplementary Fig. 10, 11). However, we observed lower *PCS* scores

when using a finer-grain parcellation with 400 regions (Supplementary Fig. 10). This is

expected since brain connectivity and structure-function relationship have been shown to

vary with the number of brain regions[41]. *PCS* scores tended to increase for longer fMRI time

series, but this effect did not impact inter-species differences nor *PCS* cortical topographies

(Supplementary Fig. 10, 11), and it could relate to the improved reliability of functional

connectivity estimation for longer scan lengths[42]. Whenever possible, data from the three

species were matched both in the number of brain regions and fMRI scan duration. The

effect of the number of subjects on *PCS* scores was minor (Supplementary Fig. 12).

Finally, we assessed the replicability of our findings by analyzing a total of six distinct

datasets (Online Methods). We found that parallel communication scores (Fig. 2,

Supplementary Fig. 12), their cortical topographies (Fig. 3, Supplementary Fig. 11, 13, 14),

and the gradient of parallel communication between unimodal and transmodal regions

across species (Fig. 3, Supplementary Fig. 15) were consistent when considering alternative

datasets. In particular, overall cortical topographies were unaltered when considering awake

or anesthetized macaques and mice (Supplementary Fig. 13, 14). However, awake animals

tended to have larger parallel communication between transmodal regions and between

transmodal and unimodal regions (*cross-modal* streams), but unchanged communication

between unimodal regions as compared to anesthetized animals (Supplementary Fig. 16).

These state-dependent differences were smaller than the ones observed between mice and

macaques, and humans. The amount of parallel communication in awake macaques and

mice was lower than the amount observed in humans, confirming the parallel communication

gradient along the evolutionary axis.

Discussion

How networked neural elements intercommunicate at the systems level, ultimately giving rise to brain function, stands as one of the most intriguing and unsolved questions of modern neurosciences. *In vivo* measurements of brain structure and activity are providing us with windows of opportunities for modeling communication in brain networks, across different animal species. We propose here to bring a piece to this puzzle, by investigating the link between communication strategies in large-scale brain networks, on the one side, and the evolutionary complexity of mammals' brain functions, on the other.

By introducing a novel approach to formally test and detect relay communication pathways in brain networks, we provide compelling evidence that this link exists, and that different communication strategies are tied to the evolutionary complexity of mammalian brain networks along two main organizational principles. The first principle is the increasing recruitment of parallel communication pathways, which evolves from predominantly selective information processing mechanisms in mice, to parallel information processing in humans. The second principle involves the development of cross-system communication through parallel pathways that connect together functionally specialized brain regions (i.e., somatomotor, visual) with transmodal ones (i.e., fronto-parietal, default mode).

Specialized, unimodal brain systems are organized as serial, hierarchical streams where raw sensory information is relayed through stepwise progressive circuits to guide attention and direct actions[20,34]. Consistent with this hierarchical polarity, we found that unimodal regions are mainly characterized by selective information processing through single pathways, as quantified by low parallel communication scores. This held true for all the investigated mammalian species, suggesting that unimodal selective information processing is phylogenetically preserved. On the other hand, transmodal regions present a more complex and less understood organization. Back in 1998, Mesulam hypothesized that "the flow of information for intermediary [transmodal] processing displays patterns consistent with parallel and re-entrant processing"[34]. Our findings consolidate this view by showing–in a data-driven and hypothesis-independent way–that information transmission between unimodal and transmodal regions evolved from selective to parallel streams across species with increasing cognitive abilities. Parallel communication could therefore represent a more complex form of information transmission beyond hierarchical processing, which might support integration of perceptual modalities into more complex textures of cognition.

What evolutionary mechanisms may have promoted a higher involvement of parallel

communication strategies in humans? According to the tethering hypothesis proposed by

Buckner and Krienen[20], the fast cortical expansion of transmodal regions in humans

compared to non-human primates has led to the untethering of these regions from

developmental anchor points. This process allowed human transmodal regions to develop

unique cytoarchitectonic[43] and connectional[31] fingerprints, unbound from the hierarchical

architecture of topographically distant unimodal systems[38]. The same process may have

also favored the development of new information transmission strategies (parallel

communication) to bridge hierarchical unimodal and distributed transmodal regions. Indeed,

in humans we observed the largest parallel communication scores in regions that underwent

the largest cortical expansion across evolution, including fronto-parietal association cortices

and precuneus[44]. In addition, parallel information transmission may be functional to specific

processing needs of unimodal-transmodal communication supporting cognition. Recent

computational studies suggest that brain regions with largest allometric scaling privilege

fidelity rather than compression of incoming signals from unimodal areas[24]. High-fidelity

information transmission may be achieved through parallel streaming of redundant signals,

expression of a more resilient communication process.

Our results show that parallel communication also contributes to the individual specificity of

communication strategies in brain networks. Selective and parallel information transmission

patterns allowed identifying subjects in a group with significant accuracy, across different

mammalian species. This indicates that the individual layout of relay communication

pathways constitutes an important fingerprint of brain organization, and that this fingerprint is

present even at lower phylogenetic levels (i.e., in mice). Brain regions that tend to

communicate through parallel rather than selective streams, including transmodal regions,

provided the largest contribution to subject identifiability. Consistently, fMRI activity of

association and default mode cortices displays larger inter-individual variability in human and

nonhuman primates compared to lower-order regions[40,45,46]. The role of transmodal cortices

in individual identifiability is consistent with their protracted neurodevelopment and function

in higher-order cognition, and it could partially explain the identifiability gradient observed

from humans to macaques, to mice, with mice displaying lower parallel communication

levels and lower identifiability. However, the identifiability gradients may also be explained by

a larger homogeneity among laboratory animals compared to human samples in terms of

genetic pedigree and environmental conditions.

Importantly, the evolutionary gradient from selective to parallel information processing and

the cortical topographies of parallel communication patterns were not explained by cross-

species differences of structural connectivity architecture, statistical properties of fMRI data,

or conscious (i.e., awake *vs* anesthetized) state. In keeping with previous studies[22], we

found that the overall distribution of short structural path lengths was similar between

species, with comparable amounts of 2-step, 3-step, and 4-step pathways. Relative cross-

species differences of parallel communication were unchanged when contrasting data with

respect to a strict, species-specific null model which preserves multivariate fMRI statistics.

When considering fMRI data from awake and anesthetized animals, we found similar cortical

distributions of parallel communication patterns, with unimodal regions dominated by

selective information processing. This finding is in line with the observation that resting state

networks are globally preserved in conscious and unconscious states[17,47]. However, we

found that the overall amount of communication tends to increase during wakefulness,

particularly when considering information transmission streams interconnecting transmodal

regions with the rest of the brain network. A shift of the communication regime toward more

abundant and (partially) parallelized polysynaptic information transmission may

mechanistically support functional integration, inter-network cross-talk, and rich functional

repertoires departing from the underlying monosynaptic connectivity constraints, which have

been repeatedly observed in awake primates and mice compared to the anesthetized

ones[17,47,48].

Several higher-order communication models have been proposed to explain integration of

information between multiple brain network elements[1]. Nonetheless, the exact polysynaptic

communication mechanisms underlying macroscale neural signaling remain unclear.

Intriguingly, brain communication models mostly rely on the assumption of memoryless

(Markovian) information transmission[49]. This hypothesis is pervasive in network

neuroscience[37] but has never been formally tested in the brain. Our work adds to the field in

two important ways. First, it introduces a new approach to extract relay communication

pathways from multimodal brain data, in a way that is agnostic to specific communication

models and grounded in fundamental information-theoretic principles. Secondly, it formally

probes the existence of memoryless information transmission in brain networks by

introducing an empirical way to assess deviations from Markovity through the data

processing inequality[35]. Our results show that Markovian communication is present in brain

networks across different mammalian species, is not limited to the shortest structural path,

but involves multiple and less optimal structural paths in a way that is species-dependent

and consistent with the evolutionary complexity of each investigated species.

Polysynaptic memoryless information transmission is a simple form of higher-order

communication. There is no reason to assume that macroscale neural communication is

limited to such a particular form. Brain network hierarchies may confer neural signals a
memory of the regions previously visited along a path, thus modifying neural communication
pathways in a context-dependent manner[49]. This process would result in non-Markovian
communication regimes. The brain may also implement complex multi-object interactions not
attributable to information transmission alone, such as synergistic or modulatory behaviors
between multiple brain regions[37,50]. Biologically, these communication strategies may shape
important features of the mammalian brain, such as cortical temporal hierarchies[51,52] or
receptive time windows for attentional processes[53]. This work only focuses on Markovian
information transmission and does not inform us about other complementary information
processing strategies. As such, absence of relay communication (i.e., violation of the data
processing inequality) may indicate absence of any communication between those particular
brain regions, or communication through more complex information encoding mechanisms.
Notwithstanding the evidence that selective and parallel Markovian pathways serve as
important information streams for multimodal integration between unimodal and transmodal
systems[34,38], we speculate that low parallel communication scores between transmodal
regions may indicate predominance of more complex communication regimes in these
areas. In addition, sensory input decoding within the highly clustered unimodal systems
(diagonal entries of the parallel communication matrices, Fig. 3b) may be supported by
synergistic processes within dense structural motifs[54]. How these macroscale
communication mechanisms may have adapted to changing environments over the evolution
of mammalian brains remains an exciting open field of research, to which the present work
adds a first foundation.

Acknowledgements

We thank Prof. Joaquín Goñi, Prof. Dante Mantini and Prof. Martijn Van den Heuvel for their
critical suggestions and helpful discussions; Dr. Kelly Shen for the help with the main
macaque dataset; Dr. Maria Giulia Preti and Dr. Raphaël Liégeois for the processing of
human data. Human data were provided by the Human Connectome Project, WU-Minn
Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657)
funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience
Research; and by the McDonnell Center for Systems Neuroscience at Washington
University. A.Gr. acknowledges funding from the Swiss National Science Foundation (Grant
No. 320030_173153). E.A. acknowledges financial support from the SNSF Ambizione
project "Fingerprinting the brain: network science to extract features of cognition, behavior
and dysfunction" (Grant No. PZ00P2_185716). A.Go. acknowledges funding from the
European Research Council (ERC, #DISCONN; no. 802371), the Brain and Behavior

Online Methods

Human data

We used Magnetic Resonance Imaging (MRI) data of the Human Connectome Project

(HCP), U100 dataset (HCP900 data release), which includes 100 unrelated healthy adults

('*h-HCP*' dataset, 64 males; mean age = 29.1 ± 3.7 years)[55]. Informed consent forms,

including consent to share de-identified data, were collected for all subjects (within the HCP)

and approved by the Washington University institutional review board. All methods were

carried out in accordance with relevant guidelines and regulations. MRI scans were

performed on a 3T Siemens Prisma scanner and included the following sequences:

Structural MRI: 3D Magnetization Prepared Rapid Acquisition with Gradient Echoes

(MPRAGE) T1-weighted, TR = 2400 ms, TE = 2.14 ms, TI = 1000 ms, flip angle = 8°, FOV =

224 × 224, voxel size = 0.7 mm isotropic. Diffusion-weighted MRI: spin-echo Echo-Planar

Imaging (EPI), TR = 5520 ms, TE = 89.5 ms, flip angle = 78°, FOV = 208 × 180, 3 shells of

b-value = 1000, 2000, 3000 s/mm^2 with 90 directions plus 6 b-value = 0 s/mm^2 acquisitions.

One session of 15 min resting-state functional MRI (fMRI): gradient-echo EPI, TR = 720 ms,

TE = 33.1 ms, flip angle = 52°, FOV = 208 × 180, voxel size = 2 mm isotropic, recorded with

two phase-encoding directions (right-left and left-right). HCP minimally preprocessed data[56]

were used for all acquisitions.

Group-level structural connectivity. A group-representative structural connectome between

100 cortical regions of interest (Schafer parcellation[57]) was obtained from the 100 unrelated

HCP subjects. Different cortical parcellation resolutions were explored in supplementary

analyses (200- and 400-region Schaefer parcellations[57]). Briefly, diffusion-weighted scans

were analyzed using MRtrix3[58], including the following steps: multi-shell multi-tissue

response function estimation; constrained spherical deconvolution; tractogram generation

with 10^7 output streamlines. The Schaefer cortical atlas was used to parcellate the cortex

into 100 (200, 400) regions and generate individual structural connectomes, from which a

group-representative structural connectome was computed. The binary architecture of the

group-representative connectome was obtained by including only the structural connections

retrieved in 100% of the subjects. This step is meant to minimize the number of false

positives in the group-representative network. The group-representative connectome was

then weighted by the Euclidean distance (in millimeters) between region pairs' centroids

(Supplementary Fig. 1a). This choice was motivated by the exigence of homogenizing

structural connections' weights across species (see also *Mapping relay communication*

pathways in brain networks).

Individual functional information. Resting-state fMRI data were pre-processed according to a

state-of-the-art pipeline[18] including: general linear model regression of nuisance signals

(removal of linear and quadratic trends; removal of motion regressors and their first

derivatives; removal of white matter and cerebrospinal fluid signals and their first

derivatives). 100 (200, 400) regional time series were obtained by averaging voxel-wise time

series across all voxels belonging to each region of interest. The mutual information between

region pairs was computed from the histograms of the z-scored time series, binned with a

step of 0.5. This bin size was chosen by comparing real and null mutual information values,

with null values obtained from multivariate gaussian data, and by assessing the

fingerprinting accuracy[39] of mutual information across bin sizes (Supplementary Fig. 17).

Only the first 800 time points (9.6 min) were considered for mutual information computation

for consistency with other species data (Supplementary Table 1; other time series lengths

were explored in supplementary analyses, Supplementary Fig. 10). Mutual information

matrices obtained from left-right and right-left phase-encoding acquisitions were averaged to

obtain a single 100x100 (200x200, 400x400) mutual information matrix per subject

(Supplementary Fig. 1c).

Replication datasets. Analyses were repeated considering sub-samples of the whole U100

dataset (Supplementary Fig. 12).

Macaque data

We used structural and functional monkey data from TheVirtualBrain project[59]. The fMRI

dataset included 9 adult male rhesus macaque monkeys (8 Macaca mulatta, 1 Macaca

fascicularis) aged between 4 and 8 years ('*q-TVB*' dataset). All methods were carried out in

accordance with relevant guidelines and regulations and have been previously described[59].

Briefly, animals were lightly anesthetized before their scanning session and anesthesia was

maintained using 1-1.5% isoflurane. The scanning was performed on a 7T Siemens

MAGNETOM head scanner included: Structural MRI: 3D MPRAGE T1-weighted sequence,

128 slices, voxel size = 0.5 mm isotropic. Diffusion-weighted MRI: EPI sequence, 24 slices,

b-value = 1000 s/mm^2, 64 directions, recorded with two opposite phase-encoding directions.

One session of 10 min resting-state functional MRI (fMRI): 2D multiband EPI sequence, TR

= 1000 ms, 42 slices, 1 X 1 X 1.1 mm^3 voxel size.

Group-level structural connectivity. We used the whole-brain macaque structural

connectome provided by TheVirtualBrain[59], which summarizes the brain connectivity

between 82 regions of interest (Regional Map parcellation of Kötter and Wanke[60]) and

includes inter-hemispheric connections. Briefly, the structural connectome was obtained by

optimizing tractography-derived structural connectivity matrices with respect to a reference

tracer-derived connectivity matrix and averaging across animals[59]. For cross-species

consistency reasons, we considered undirected structural connectivity information. That is, in

the final structural connectome, two regions are connected if at least one unidirectional

connection exists between the two regions. Structural connections were weighted by the

Euclidean distance (in millimeters) between region pairs' centroids (Supplementary Fig. 1a).

Individual functional information. Resting-state fMRI data were pre-processed by others, as

previously described[59]. Briefly, the processing pipeline included motion correction, high-pass

filtering, regression of white matter and cerebrospinal fluid signals, spatial normalization and

smoothing. Z-scored regional time series (Regional Map parcellation) including 600 time

points (10 min) were used to compute individual mutual information matrices (bin size = 0.5,

consistently with other species) (Supplementary Fig. 1c).

Replication dataset. Analyses were repeated on an independent dataset of 9 adult rhesus

macaque monkeys (Macaca mulatta) aged between 5 and 12 years scanned on a vertical

Bruker 4.7T primate dedicated scanner at Newcastle University[61] ('*q-NCS*' dataset). Raw

data were publicly available through the Primate Data Exchange (PRIME-DE) initiative[62] and

included the following MRI sequences: Structural MRI: Modified Driven Equilibrium Fourier

Transform (MDEFT) T1-weighted, TR = 2000 ms, TE = 3.75 ms, TI = 750 ms, voxel size =

0.6 x 0.6 x 0.62 mm^3. Two runs of 10.8 min resting-state fMRI: TR = 2600 ms, TE = 17 ms,

voxel size = 1.2 mm isotropic. All animals were scanned awake. MRI data preprocessing

included: T1-weighted volumes denoising[63], skull-stripping (FSL[64]), N4 bias field correction,

spatial normalization to the F99 template obtained from the SumDB database

(http://brainvis.wustl.edu/sumsdb/public_archive_index.html), and registration to fMRI native

space (ANTs[65]); fMRI volumes were coregistered (FSL[66]), corrected for nuisance signals

including 6 motion signals, average white matter and cerebrospinal fluid signals, and band-

pass filtered to the band 0.01-0.15 Hz. Z-scored regional time series (Regional Map

parcellation) of the two concatenated fMRI runs were used to compute individual mutual

information values (bin size = 0.5). The fMRI scans were concatenated to reach a number of

time points comparable with the other datasets (500 time points, 21.6 min).

Mouse data

We used open-source fMRI data of 10 male wild-type mice aged 6 months ('*m-AD3*'

dataset), available at https://openneuro.org/datasets/ds001890[67]. All methods were carried

out in accordance with relevant guidelines and regulations and have been previously
described[17,68]. Briefly, animals were anesthetized with 4% isoflurane before their scanning
session and maintained with 0.5% isoflurane and a 0.05 mg/kg/h medetomidine infusion[69].
The scanning was performed on a 11.75T Brucker BioSpin scanner and included: Structural
MRI: spin-echo turboRARE sequence, TR = 2750 ms, TE = 30 ms, FOV = 17 x 11 mm^2,
matrix dimension = 200 x 100 voxels, slice thickness = 0.35 mm. One session of 10 min
resting-state functional MRI (fMRI): gradient-echo EPI sequence, TR = 1000 ms, TE = 15
ms, matrix dimension = 90 x 60 voxels.

Group-level structural connectivity. A mouse structural connectome between 78 cortical
regions covering the isocortex, cortical subplate, and hippocampal formation, as defined in
the Allen Brain Atlas, was derived from published viral tracing data[70]. In more details, the
binary architecture of the structural connectome was assessed according to the following
steps: (i) we considered the right-hemisphere ipsilateral and contralateral connections
reported by Oh and colleagues[70]; (ii) we symmetrized the right-hemisphere ipsilateral
connections (i.e., we considered a connection between ipsilateral regions i and j to be
present if at least one of the two tracts (i→j), (j→i) was detected; (iii) we duplicated the
symmetrized ipsilateral connections to the left hemisphere (in absence of more detailed
information, we therefore assume equal intra-hemispheric connectivity in the right and left
hemispheres); (iv) we transposed the contralateral connections of the right hemisphere to
the left hemisphere; (v) to minimize false positives due to minor tissue segmentation
artifacts, we excluded connections with connectivity strength < $10^{-3.5}$, as suggested in[70],
where the connectivity strength was defined as the total volume of segmented pixels in the
target normalized by the injection site volume. The binary structural connectome was then
weighted by the Euclidean distance between region pairs' centroids obtained from the Allen
Brain Atlas (CCF v3, © 2004 Allen Institute for Brain Science. Allen Mouse Brain Atlas.
Available from: http://www.brain-map.org/) (Supplementary Figure 4).

Individual functional information. Resting-state fMRI data were pre-processed as previously
described[68]. Briefly, the processing pipeline included motion correction, automatic brain
masking, spatial smoothing (FWHM = 0.45 mm), high-pass filtering (0.01 Hz cut-off), and
automated nuisance removal based on independent component analysis. Z-scored regional
time series (78-region Allen Brain Atlas parcellation) including 600 time points (10 min) were
used to compute individual mutual information matrices (bin size = 0.5) (Supplementary Fig.
1c).

Replication datasets. Analyses were repeated on two independent datasets. The first one

included 51 male wild-type mice scanned at 3 months ('*m-CSD1*' dataset)[71]. MRI

acquisitions were performed on a 9.4T Brucker BioSpin system on anesthetized animals

(3.5% isoflurane, maintained with 0.5% isoflurane and a 0.05 mg/kg/h medetomidine

infusion) and included a 6-min resting-state fMRI recording: gradient-echo EPI sequence, TR

= 1000 ms, TE = 9.2 ms, flip angle = 90°, field of view = 20 x 17.5 mm^2, matrix size = 90 x 70

voxels, slice thickness = 0.5 mm. FMRI volumes were preprocessed using the same pipeline

as the *m-AD3* dataset. The average time series of the 78 cortical regions (360 time points, 6

min) were z-scored and used to compute individual mutual information matrices (bin size =

0.5). Analyses were repeated considering sub-samples of the whole *m-CSD1* dataset

(Supplementary Fig. 12).

The second dataset included 10 C57Bl6/J adult male mice ('*m-GG*' dataset, < 6 months old)

subject to surgery for headposts placement, MRI habituation and awake fMRI acquisition, as

previously described[17]. MRI acquisitions were performed at the IIT laboratory in Rovereto

(Italy) on a Bruker Biospin 7T scanner and included a 32-min resting-state fMRI recording:

single-shot EPI sequence, TR = 1000 ms, TE = 15 ms, flip angle = 60°, voxels size = 0.23 x

0.23 x 0.6 mm^3. fMRI preprocessing included exclusion of the first 2 min of recording, time

series despiking, motion correction, nuisance signals regression (average cerebrospinal fluid

and motion signals plus their temporal derivative and corresponding squared regressors),

data censoring (Framewise Displacement > 0.075 mm), band-pass filtering (0.01-0.1 Hz),

spatial smoothing (FWHM = 0.5 mm) and spatial normalization[17]. Average time series were

computed for 66 regions of interest, which represents a subset of the 78 Allen Brain Atlas

regions (data for bilateral regions CA1, CA2, CA3, dorsal and ventral endopiriform nucleus,

and frontal pole were not available). The first 600 time points (10 min) were used for the

computation of individual mutual information matrices (z-scored time series binning = 0.5).

Assignment of cortical regions to resting state networks

For the *human dataset*, each cortical region was assigned to one the seven resting state

networks (RSNs) defined by Yeo and colleagues and according to the Schaefer parcellation

[57,72]. For the *macaque dataset*, each cortical region was first associated with one or multiple

Brodmann areas according to the CoCoMac Regional Map of the macaque cortex[60,73–75].

Each Brodmann area was then assigned to one of the seven RSNs defined by Yeo and

colleagues[72] using a majority voting procedure and published atlases in MNI space[76]. Finally,

Regional Map regions of the macaque cortex were assigned to Yeo RSNs with a similar

majority voting procedure (Supplementary Fig. 3). For the *mouse dataset*, each cortical

region was assigned to one out of 6 RSN as identified by Zerbi and colleagues using

independent component analysis of resting-state fMRI data[77]. The assignment was done
through a majority voting procedure (Supplementary Fig. 4). Note that the default mode
network (DMN) has been consistently identified in humans[78], macaques[79] and mice[80,81],
suggesting a conservation of this network across mammalian species. In our mouse cortex
subdivision[77], the DMN includes bilateral hippocampal regions (CA1, CA2, CA3 hippocampal
fields, subiculum and dentate gyrus), and lateral (ectorhinal and temporal association areas)
and prefrontal (infralimbic, prelimbic and perirhinal areas) isocortices, while it excludes other
regions which have been reported by others, such as the retrosplenial cortex[82]. For all
species, RSNs were assigned to unimodal or transmodal systems according to established
cortical subdivisions[38].

Mapping information transmission pathways in brain networks
In this work we introduce a new approach to infer relay communication pathways from
multimodal neuroimaging data. The approach builds upon and extends an information
theoretical framework proposed in previous work[35], and aims at identifying polysynaptic
(multi-step) structural pathways selected for information transmission in macroscale brain
networks. Information theory is a branch of mathematics that studies the transmission of
information through communication systems[36] and has found several applications in
neuroscience[83,84]. It allows model-independent analysis of noisy data, such as the fMRI
ones.

Structural brain network and structural paths. Let's consider a structural brain network as an
undirected graph $G \equiv \{V, W\}$ formed by a set of N nodes $V = \{v_1, v_2, \ldots, v_N\}$ and a
connectivity matrix $W = [w_{i,j}]$, with $w_{i,j} > 0$ distance between directly connected region pairs
v_i, v_j and $w_{i,j} = \infty$ otherwise. In this work we assigned $w_{i,j}$ equal to the Euclidean distance
(in millimeters) between the centroids of regions v_i, v_j. This choice has two motivations.
First, the distance between region centroids can be easily computed across different
datasets, thus allowing to select homogeneous structural connectivity weights across
species. Second, this choice conceptually links information transmission in brain networks
with the sender-channel-receiver schematics proposed in electronic communication by
Shannon[35,85]. A path between a source node v_i and a target node v_j is a sequence of
pairwise connected and non-repeating nodes $\Omega_{i,j} = \{v_i, v_a, v_b, \ldots, v_j\}$. The shortest path $\Omega_{i,j}^{SP}$
between regions v_i, v_j is the path of minimal length (i.e., minimal Euclidean distance, in the
case of this work) connecting the two regions. The path length is computed as the sum of
edge weights along the path. In this work we identified the first $k = 5$ k-shortest paths

$\Omega_{i,j}^{k-SP}$ connecting each region pair v_i , v_j[86]. K-shortest path ensembles identify meaningful

trade-offs between efficiency and resiliency for putative communication processes in brain

networks[23]. The choice of k was dictated by the fact that, for $k = 5$, all edges of the structural

brain network participate in at least one k-shortest path[23].

Functional information along structural paths. Each node v_i is associated with a neural

activity-related fMRI time series X^i that can be interpreted as the realization of a discrete

random variable with probability mass function $p_i(x^i)$. The amount of shared information

between two random variables can be quantified as their mutual information $I(X^i, X^j) =$

$\sum_{x^i \in X^i} \sum_{x^j \in X^j} p_{i,j}(x^i, x^j) log_2 (p_{i,j}(x^i, x^j)/p_i(x^i)p_j(x^j))$, with $p_{i,j}(x^i, x^j)$ joint probability

distribution between X^i, X^j. The sequence of pairwise mutual information values along a

structural path $\Omega_{i,j}$ with respect to the source node i is defined as

$\Phi_{i,j} = \{I(X^i, X^a), I(X^i, X^b), \ldots, I(X^i, X^j)\}$. We estimated the fMRI time series probability mass

functions from the z-scored time series' histograms with appropriate binning. Different bin

sizes between 0.05 and 2.00 were explored and evaluated with respect to (i) corresponding

mutual information values for multivariate Gaussian processes $\aleph(0, I)$; (ii) individual

identifiability scores[39]. We selected the smallest bin size for which (i) the mutual information

values obtained from real data (*h-HCP* dataset) were larger than expected for a multivariate

Gaussian process $\aleph(0, I)$, and (ii) the individual identifiability score reached a maximum

plateau (Supplementary Fig. 17).

Data Processing Inequality (DPI). The *DPI*, a fundamental principle of information theory,

states that the amount of information available at a target node j about a source node i

cannot be increased through operations performed along the transmission path.

Mathematically, the DPI states that if $X^i - X^a - X^j$ is a Markov chain, then $I(X^i, X^a) \geq$

$I(X^i, X^j), I(X^a, X^j) \geq I(X^i, X^j)$, i.e., the mutual information does not increase along the

chain[36]. Note that the double inequality condition derives from the fact that a Markov chain

has no directionality information, i.e., if $X^i - X^a - X^j$ is a Markov chain, then $X^j - X^a - X^i$ is

also a Markov chain. The *DPI* can be extended to Markov chains of any length.

Conceptually, the *DPI* embeds two assumptions about the information transmission process:

the first one is that (neural) messages transmitted through the structural infrastructure (brain

network) can keep at most the same amount of information present at the source region

(*information decay*). The second one is that (neural) messages do not contain memory of the

transmission process itself and communication happens in a Markovian fashion

(*memoryless transmission*).

Identification of information transmission pathways in brain networks. We used the DPI to

test (deviation from) Markovian behavior. Each k-shortest structural path was labeled a relay

communication pathway if the *DPI*s along the paths were satisfied. Note that here we use

the wording *relay communication* in Shannon's sense. That is, we aim to characterize the

presence of memoryless information transmission processes, with information decay along

the path measured through mutual information values.

Parallel communication scores (PCSs). We define the parallel communication score $PCS_{i,j}^{n}$

between a pair of brain regions v_i , v_j as the number of k-shortest paths connecting the two

regions which respect the *DPI*, with n indicating the subject. Note that, given the choice of

$k = 5$, *PCS* scores can assume integer values between 0 and 5, and that $PCS_{i,j}^{n} = PCS_{j,i}^{n}$. A

PCS score equal to 0 is interpreted as absence of (Markovian) information transmission

between two regions; a *PCS* score equal to 1 is interpreted as presence of *selective*

information processing through a single information transmission pathway; PCS scores

larger than 1 are interpreted as presence of progressively increasing *parallel information*

processing with information transmission through multiple parallel pathways (Fig. 1). *PCS*

scores were computed for every pair of brain regions and every subject, for all investigated

datasets. Parallel communication information was summarized at the group-level by

computing a group-average parallel communication matrix PCS^{avg} for each dataset, and its

corresponding histogram (Fig. 2). In addition, node-average, RSN-average, and system-

average *PCS* scores were computed by averaging the parallel communication scores over

the corresponding region pairs (Fig. 3).

Null model. A null model was defined by randomly shuffling the raw fMRI time series across

brain regions while preserving the original structural connectivity information (Supplementary

Fig. 5). Note that with this randomization we are preserving the statistical properties of both

the original functional and structural data, since we are merely rearranging spatially fMRI

time series across the brain network. Parallel communication matrices were then computed

for each randomization following the above-described procedure. For each dataset, the

randomization was repeated 3000 times *per* subject, which allowed to build 3000 group-

average parallel communication matrices (Supplementary Fig. 6). Each region pair was

therefore associated with a null distribution of group-average *PCS* values including 3000

elements. To assess whether group-average *PCS* scores observed in real data could be

trivially explained by the structural connectivity architecture and the multivariate statistical

properties of fMRI data, which are both preserved in the null model, we adopted two

strategies. The first one consisted of *PCS* scores screening by z-scoring individual group-
average scores $PCS_{i,j}^{avg}$ with respect to the corresponding null distribution; z-scored where
thresholded at 1.96 (Supplementary Fig. 7). The second strategy consisted of analyzing the
PCS scores with false discovery rate (FDR)-corrected *p*-values < .05 (*FDR < .05*), with p-
values computed as the number of entries in the null distribution exceeding the real *PCS*
score (Supplementary Fig. 8).

Subject identifiability analysis
For each investigated dataset, fMRI time series were split into two parts of equal duration
and considered as test and retest data. From these, test and retest parallel communication
matrices were computed for each subject. An identifiability matrix summarizing test-retest
subjects' similarities was then obtained for each dataset. Diagonal entries of the identifiability
matrix represent subjects' *self*-similarity between test and retest data ('*Iself*'); outside-
diagonal entries represent inter-subject similarity ('*Iothers*') (Fig. 4)[39]. The similarity between
test and retest parallel communication matrices was assessed with the Jaccard index,
defined as the size of the intersection divided by the size of the union of two label sets. For
example, a Jaccard index equal to 0.3 indicates that 30% of brain region pairs have exactly
the same *PCS* score, which can take integer values between 0 and 5. The level of individual
identifiability was quantified with the success rate (SR) defined as the percentage of test
subjects whose identity was correctly predicted out of the total set of retest subjects[40]. The
subject identifiability analysis was repeated when considering only region pairs with, on
average, low (high) *PCS* scores for the computation of test-retest similarities. Different
thresholds defining low (high) *PCS* scores were explored (Supplementary Table 2).

Data and material availability
The data that support the findings of his study are available on the Human Connectome
Project platform (db.humanconnectome.org) for human data; OpenNeuro[59], Zenodo[59], and
INDI PRIMatE Data Exchange (fcon_1000.projects.nitrc.org/indi/indiPRIME.html) platforms
for macaque data; OpenNeuro[67], XNAT[71], and Mendeley[17] platforms for mouse data. The
derived brain matrices necessary to reproduce the main analyses of this study are available
on Zenodo. The code (in MATLAB) and sample brain matrices are available as maintained
version on A.Gr.'s GitHub repository (github.com/agriffa/BrainComm_mammalian_evolution).

References

1. Avena-Koenigsberger, A., Misic, B. & Sporns, O. Communication dynamics in complex brain networks. *Nat Rev Neurosci* **19**, 17–33 (2018).

2. Bullmore, E. & Sporns, O. Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience* **10**, 186–198 (2009).

3. Bassett, D. S. & Sporns, O. Network neuroscience. *Nat Neurosci* **20**, 353–364 (2017).

4. Medaglia, J. D., Lynall, M.-E. & Bassett, D. S. Cognitive Network Neuroscience. *Journal of Cognitive Neuroscience* **27**, 1471–1491 (2015).

5. Kaiser, M. & Hilgetag, C. C. Nonoptimal Component Placement, but Short Processing Paths, due to Long-Distance Projections in Neural Systems. *PLOS Computational Biology* **2**, e95 (2006).

6. Bullmore, E. & Sporns, O. The economy of brain network organization. *Nature Reviews Neuroscience* **13**, 336–349 (2012).

7. Chen, Y., Wang, S., Hilgetag, C. C. & Zhou, C. Features of spatial and functional segregation and integration of the primate connectome revealed by trade-off between wiring cost and efficiency. *PLOS Computational Biology* **13**, e1005776 (2017).

8. Lynn, C. W. & Bassett, D. S. The physics of brain network structure, function and control. *Nat Rev Phys* **1**, 318–332 (2019).

9. Mišić, B. *et al.* Network-Level Structure-Function Relationships in Human Neocortex. *Cereb Cortex* **26**, 3285–3296 (2016).

10. Goñi, J. *et al.* Resting-brain functional connectivity predicted by analytic measures of network communication. *PNAS* **111**, 833–838 (2014).

11. Hagmann, P. *et al.* Mapping the Structural Core of Human Cerebral Cortex. *PLOS Biology* **6**, e159 (2008).

12. Singer, W. Neuronal Synchrony: A Versatile Code for the Definition of Relations? *Neuron* **24**, 49–65 (1999).

13. Honey, C. J. *et al.* Predicting human resting-state functional connectivity from

structural connectivity. *PNAS* **106**, 2035–2040 (2009).

14.   Suárez, L. E., Markello, R. D., Betzel, R. F. & Misic, B. Linking Structure and

Function in Macroscale Brain Networks. *Trends in Cognitive Sciences* **24**, 302–315

(2020).

15.   Battiston, F. *et al.* The physics of higher-order interactions in complex systems. *Nat.*

Phys. **17**, 1093–1098 (2021).

16.   Griffa, A., Amico, E., Liégeois, R., Ville, D. V. D. & Preti, M. G. Brain structure-

function coupling provides signatures for task decoding and individual fingerprinting.

NeuroImage 118970 (2022) doi:10.1016/j.neuroimage.2022.118970.

17.   Gutierrez-Barragan, D. *et al.* Unique spatiotemporal fMRI dynamics in the awake

mouse brain. *Current Biology* **32**, 631-644.e6 (2022).

18.   Preti, M. G. & Van De Ville, D. Decoupling of brain function from structure reveals

regional behavioral specialization in humans. *Nature Communications* **10**, 4747 (2019).

19.   Baum, G. L. *et al.* Development of structure–function coupling in human brain

networks during youth. *PNAS* **117**, 771–778 (2020).

20.   Buckner, R. L. & Krienen, F. M. The evolution of distributed association networks in

the human brain. *Trends in Cognitive Sciences* **17**, 648–665 (2013).

21.   van den Heuvel, M. P., Bullmore, E. T. & Sporns, O. Comparative Connectomics.

Trends in Cognitive Sciences **20**, 345–361 (2016).

22.   Assaf, Y., Bouznach, A., Zomet, O., Marom, A. & Yovel, Y. Conservation of brain

connectivity and wiring across the mammalian class. *Nat Neurosci* **23**, 805–808 (2020).

23.   Avena-Koenigsberger, A. *et al.* Path ensembles and a tradeoff between

communication efficiency and resilience in the human connectome. *Brain Struct Funct*

222, 603–618 (2017).

24.   Zhou, D. *et al.* Efficient coding in the economics of human brain connectomics.

Network Neuroscience 1–40 (2022) doi:10.1162/netn_a_00223.

25.   Bettinardi, R. G. *et al.* How structure sculpts function: Unveiling the contribution of

anatomical   connectivity to the brain's spontaneous correlation structure. *Chaos* **27**,

047409 (2017).

26.   Goñi, J. *et al.* Exploring the Morphospace of Communication Efficiency in Complex

Networks. *PLOS ONE* **8**, e58070 (2013).

27.   Hao, Y. & Graham, D. Creative destruction: Sparse activity emerges on the mammal

connectome under a simulated communication strategy with collisions and

redundancy. *Network Neuroscience* **4**, 1055–1071 (2020).

28.   Mišić, B. *et al.* Cooperative and Competitive Spreading Dynamics on the Human

Connectome. *Neuron* **86**, 1518–1529 (2015).

29.   Estrada, E. & Hatano, N. Communicability in complex networks. *Phys. Rev. E* **77**,

036111 (2008).

30.   Friedrich, P. *et al.* Imaging evolution of the primate brain: the next frontier?

NeuroImage **228**, 117685 (2021).

31.   Ardesch, D. J. *et al.* Evolutionary expansion of connectivity between multimodal

association areas in the human brain compared with chimpanzees. *PNAS* **116**, 7101–

7106 (2019).

32.   Xu, T. *et al.* Cross-species functional alignment reveals evolutionary hierarchy within

the connectome. *NeuroImage* **223**, 117346 (2020).

33.   Krubitzer, L. The Magnificent Compromise: Cortical Field Evolution in Mammals.

Neuron **56**, 201–208 (2007).

34.   Mesulam, M. M. From sensation to cognition. *Brain* **121**, 1013–1052 (1998).

35.   Amico, E. *et al.* Toward an information theoretical description of communication in

brain networks. *Network Neuroscience* 1–20 (2021) doi:10.1162/netn_a_00185.

36.   Cover, T. M. & Thomas, J. A. ELEMENTS OF INFORMATION THEORY. 774 (2006).

37.   Lambiotte, R., Rosvall, M. & Scholtes, I. From networks to optimal higher-order

models of complex systems. *Nature Physics* **15**, 313–320 (2019).

38.   Margulies, D. S. *et al.* Situating the default-mode network along a principal gradient

of macroscale cortical organization. *PNAS* **113**, 12574–12579 (2016).

39.   Amico, E. & Goñi, J. The quest for identifiability in human functional connectomes.

Sci Rep **8**, 1–14 (2018).

40.   Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using

patterns of brain connectivity. *Nature Neuroscience* **18**, 1664–1671 (2015).

41.   Messé, A. Parcellation influence on the connectivity-based structure–function

relationship in the human brain. *Human Brain Mapping* **41**, 1167–1180 (2020).

42.   Birn, R. M. *et al.* The effect of scan length on the reliability of resting-state fMRI

connectivity estimates. *NeuroImage* **83**, 550–558 (2013).

43.   Paquola, C. *et al.* Microstructural and functional gradients are increasingly

dissociated in transmodal cortices. *PLOS Biology* **17**, e3000284 (2019).

44.   Bruner, E., Preuss, T. M., Chen, X. & Rilling, J. K. Evidence for expansion of the

precuneus in human evolution. *Brain Struct Funct* **222**, 1053–1060 (2017).

45.   Xu, T. *et al.* Interindividual Variability of Functional Connectivity in Awake and

Anesthetized Rhesus Macaque Monkeys. *Biological Psychiatry: Cognitive Neuroscience*

and Neuroimaging **4**, 543–553 (2019).

46.   Mueller, S. *et al.* Individual Variability in Functional Connectivity Architecture of the

Human Brain. *Neuron* **77**, 586–595 (2013).

47.   Barttfeld, P. *et al.* Signature of consciousness in the dynamics of resting-state brain

activity. *Proceedings of the National Academy of Sciences* **112**, 887–892 (2015).

48.   Luppi, A. I. *et al.* Consciousness-specific dynamic interactions of brain integration

and functional diversity. *Nature Communications* **10**, (2019).

49.   Vézquez-Rodríguez, B., Liu, Z.-Q., Hagmann, P. & Misic, B. Signal propagation via

cortical hierarchies. *Network Neuroscience* **4**, 1072–1090 (2020).

50.   Battiston, F. *et al.* Networks beyond pairwise interactions: Structure and dynamics.

Physics Reports **874**, 1–92 (2020).

51.   Honey, C. J. *et al.* Slow Cortical Dynamics and the Accumulation of Information over

Long Timescales. *Neuron* **76**, 423–434 (2012).

52.   Murray, J. D. *et al.* A hierarchy of intrinsic timescales across primate cortex. *Nat*

Neurosci **17**, 1661–1663 (2014).

53.   Fries, P., Reynolds, J. H., Rorie, A. E. & Desimone, R. Modulation of Oscillatory

Neuronal Synchronization by Selective Visual Attention. *Science* **291**, 1560–1563 (2001).

54.   Reimann, M. W. *et al.* Cliques of Neurons Bound into Cavities Provide a Missing Link

between Structure and Function. *Front. Comput. Neurosci.* **11**, (2017).

55.   Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An overview.

NeuroImage **80**, 62–79 (2013).

56.   Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human

Connectome Project. *NeuroImage* **80**, 105–124 (2013).

57.   Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from

Intrinsic Functional Connectivity MRI. *Cereb Cortex* **28**, 3095–3114 (2018).

58.   Tournier, J.-D. *et al.* MRtrix3: A fast, flexible and open software framework for

medical image processing and visualisation. *NeuroImage* **202**, 116137 (2019).

59.   Shen, K. *et al.* A macaque connectome for large-scale network simulations in

TheVirtualBrain. *Sci Data* **6**, 123 (2019).

60.   Kötter, R. & Wanke, E. Mapping brains without coordinates. *Philosophical

Transactions of the Royal Society B: Biological Sciences* **360**, 751–766 (2005).

61.   Baumann, S. *et al.* Orthogonal representation of sound dimensions in the primate

midbrain. *Nat Neurosci* **14**, 423–425 (2011).

62.   Milham, M. P. *et al.* An Open Resource for Non-human Primate Imaging. *Neuron*

100, 61-74.e2 (2018).

63.   Manjón, J. V., Coupé, P., Martí-Bonmatí, L., Collins, D. L. & Robles, M. Adaptive

non-local means denoising of MR images with spatially varying noise levels. *Journal of

Magnetic Resonance Imaging* **31**, 192–203 (2010).

64.   Smith, S. M. Fast robust automated brain extraction. *Human Brain Mapping* **17**, 143–

155 (2002).

65.   Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance

in brain image registration. *NeuroImage* **54**, 2033–2044 (2011).

66.   Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved Optimization for the

Robust and Accurate Linear Registration and Motion Correction of Brain Images.

NeuroImage **17**, 825–841 (2002).

67.   Mandino, F., Yeow Ling Yun, Gigg, J., Olivo, M. C. & Joanes Grandjean.

Mouse_rest_3xTG. (2019) doi:10.18112/OPENNEURO.DS001890.V1.0.1.

68.   Mandino, F., Yeow, L. Y., Gigg, J., Olivo, M. & Grandjean, J. Preserved functional

networks in a hydrocephalic mouse. *Matters* **5**, e201905000001 (2019).

69.   Grandjean, J., Schroeter, A., Batata, I. & Rudin, M. Optimization of anesthesia

protocol for resting-state fMRI in mice based on differential effects of anesthetics on

functional connectivity patterns. *NeuroImage* **102**, 838–847 (2014).

70.   Oh, S. W. *et al.* A mesoscale connectome of the mouse brain. *Nature* **508**, 207–214

(2014).

71.   Grandjean, J. *et al.* Chronic psychosocial stress in mice leads to changes in brain

functional connectivity and metabolite levels comparable to human depression.

NeuroImage **142**, 544–552 (2016).

72.   Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic

functional connectivity. *J Neurophysiol* **106**, 1125–1165 (2011).

73.   Kötter, R. *et al.* Advanced database methodology for the Collation of Connectivity

data on the Macaque brain (CoCoMac). *Philosophical Transactions of the Royal Society*

of London. Series B: Biological Sciences **356**, 1159–1186 (2001).

74.   Kötter, R. Online retrieval, processing, and visualization of primate connectivity data

from the CoCoMac Database. *Neuroinform* **2**, 127–144 (2004).

75.   Stephan, K. E., Zilles, K. & Kötter, R. Coordinate–independent mapping of structural

and functional data by objective relational transformation (ORT). *Philosophical*

Transactions of the Royal Society of London. Series B: Biological Sciences **355**, 37–54

(2000).

76.   Pijnenburg, R. *et al.* Myelo- and cytoarchitectonic microstructural and functional

human cortical atlases reconstructed in common MRI space. *NeuroImage* **239**, 118274

(2021).

77.   Zerbi, V., Grandjean, J., Rudin, M. & Wenderoth, N. Mapping the mouse brain with

1048   rs-fMRI: An optimized pipeline for functional network identification. *NeuroImage* **123**, 11–

1049   21 (2015).

1050   78.   Greicius, M. D., Krasnow, B., Reiss, A. L. & Menon, V. Functional connectivity in the

1051   resting brain: A network analysis of the default mode hypothesis. *PNAS* **100**, 253–258

1052   (2003).

1053   79.   Vincent, J. L. *et al.* Intrinsic functional architecture in the anaesthetized monkey

1054   brain. *Nature* **447**, 83–86 (2007).

1055   80.   Gozzi, A. & Schwarz, A. J. Large-scale functional connectivity networks in the rodent

1056   brain. *NeuroImage* **127**, 496–509 (2016).

1057   81.   Grandjean, J. *et al.* Common functional networks in the mouse brain revealed by

1058   multi-centre resting-state fMRI analysis. *NeuroImage* **205**, 116278 (2020).

1059   82.   Whitesell, J. D. *et al.* Regional, Layer, and Cell-Type-Specific Connectivity of the

1060   Mouse Default Mode Network. *Neuron* **109**, 545-559.e8 (2021).

1061   83.   Timme, N. M. & Lapish, C. A Tutorial for Information Theory in Neuroscience. *eNeuro*

1062   **5**, (2018).

1063   84.   Dimitrov, A. G., Lazar, A. A. & Victor, J. D. Information theory in neuroscience. *J*

1064   *Comput Neurosci* **30**, 1–5 (2011).

1065   85.   Shannon, C. E. A Mathematical Theory of Communication. 53.

1066   86.   Yen, J. Y. Finding the K Shortest Loopless Paths in a Network. *Management Science*

1067   **17**, 712–716 (1971).

1068