The effect of sociality on competitive interactions among birds

Ilias Berberi[1], Eliot T. Miller[2], Roslyn Dakin[1]

[1]Department of Biology
Carleton University
1125 Colonel By Drive
Ottawa, Canada

[2]Cornell Lab of Ornithology
159 Sapsucker Woods Road
Ithaca, New York

To whom correspondence should be addressed:
Ilias Berberi
iliasberberi@gmail.com
Roslyn Dakin
roslyn.dakin@gmail.com

Keywords: competition, network, foraging, flocking, citizen science, behavioural ecology

ABSTRACT

Sociality can provide many benefits, including increased foraging success, reproductive opportunities, and defence against predation. How does sociality influence the dominance hierarchies of ecological competitors? Here, we address this question using a large dataset of competitive interactions among birds foraging at backyard feeders across North America, representing a network of over 88,000 interactions among 196 bird species. We quantify sociality as the number of conspecifics observed together, and show that this measure of group size varies across bird species. More social species are less likely to dominate similarly sized opponents, suggesting that the evolution of social clustering is associated with reduced competitive ability overall. At the same time, we find that more social species also gain a greater competitive advantage from the presence of their conspecifics. We show that within-species competition occurs more often than expected in a null model, with the most social species exhibiting the greatest proportion of conflict originating from conspecifics. Overall, these results demonstrate that sociality influences the outcome of competition in ecological networks. Species that evolve greater sociality exhibit decreased competitive ability as individuals, but increased competitive ability in groups.

36   **INTRODUCTION**

Animal sociality can provide many ecological and evolutionary advantages, including increased

38   foraging success [1,2], more frequent mating opportunities [3,4], and improved survival through

dilution and confusion effects on predators [5,6]. However, sociality may also incur costs under

40   certain circumstance, such as conflict among conspecific group members that share a similar

niche [7,8]. The causes and consequences of animal sociality have historically been of interest in

42   evolutionary biology, with several different hypotheses for benefits that may explain the

evolution of social grouping. One hypothesized benefit of sociality is that social coalitions may

44   provide an advantage during conflict with competitors [9,10]. While both inter- and intraspecific

dominance is often enhanced by body size [11,12] and age/experience [13], there may also be

46   opportunities for subordinate individuals to gain an advantage over dominant competitors

through social coalitions [9,14]. During foraging, animals face competition from both

48   conspecifics and heterospecifics, but the influence of social grouping on the structure of these

competitive systems has not yet been described. Our aim in this study is to evaluate how

50   sociality influences competition in communities where interactions occur both within and

between species.

52   One system which could measure such effects is the competition among seed-eating birds

at backyard bird feeders. In many temperate systems, survival in winter depends on acquiring

54   adequate food resources [15,16]. Moreover, the use of artificial food sources (e.g., bird feeders)

during winter foraging has been identified as having important evolutionary consequences in

56   birds [17,18], including high rates of competitive interactions among and within species foraging

at feeder locations [19,20]. In the context of sociality, group foraging can provide more

58   opportunities for resource acquisition than foraging independently due to the shared vigilant

effort [21,22] (but see [23]), social information gathering [24,25], and improved rates of

60   competition success over solitary foragers [26]. Individuals may be particularly likely to adopt

social foraging strategies in the winter when resources are scarce.

62   Here, we investigate how sociality influences competition using interactions among birds

that visit backyard feeders. Our general predictions are that more social species gain a benefit in

64   interspecific conflict, at a cost of increased rates of intraspecific conflict. We tested these

predictions using data from Project FeederWatch (PFW), a surveying program with a network of

66   tens of thousands of observers across North America who monitor the presence and interactions

of hundreds of bird species throughout winter (Figure 1). This project has yielded a rich dataset

68   indicating who competes with whom, and how often (Figure 1D). Previous research using PFW

behavioural observations has revealed that species body mass is the primary predictor of

70   competitive dominance within communities of winter resident birds across the North American

continent [12]. Owing to the large number of interactions observed and variation in competitive

72   outcomes (Figure 1D), the PFW program provides an ideal system to test how variation in

sociality influences competitive interactions after accounting for the effects of body size.

74   We quantify variation in social behaviour in the PFW database using conspecific group

size (i.e., the number of individuals of a given bird species observed at a given a location at the

76   same time; Figure 1A-C) [27]. It is important to note that sociality also has other dimensions

beyond the simple tempo-spatial aggregation studied here, such as the stability of group

78   membership, the types of relationships between group members, and other emergent properties

of group organization [28,29]. Nevertheless, conspecific groupings – individuals found in the

80   same place at the same time – represent the most fundamental dimension of sociality that we

expect to influence competitive outcomes across species.

82   As a first step in this study, we determine whether winter sociality differs across bird

species by quantifying the repeatability of group size (Figure 1A). We also estimate the

84   phylogenetic signal of average group size, which describes the extent to which this trait is more

similar among closely related species (Figure 1B). Second, we evaluate whether average group

86   size (as a species trait) is associated with species differences in competitive dominance against

similarly sized opponents (Figure 1D). We expect that larger social group size has coevolved

88   with greater interspecific dominance. Third, for each species studied, we test whether the

presence of conspecifics affects the outcome of interspecific dominance interactions (hereafter,

90   we refer to this as a "conspecific effect"). After estimating the conspecific effect for each

species, we test whether more social species exhibit a more positive conspecific effect. Finally,

92   to evaluate the hypothesized costs of social grouping, we consider the frequency of conflict with

conspecifics (Figure 1D), and compare the observed rates of within-group conflict to those

94   expected in a null model. If large group sizes lead to increased within-group conflict, we expect

more social species to experience a greater proportion of conflict originating from conspecifics.

MATERIALS AND METHODS

98   *(a) Data sources*

We used a dataset collected by Project FeederWatch (PFW), a population monitoring program

100   with over 9 million observations of bird species observed at winter feeding locations in North

America during the months of November through April, 2015-2020. Participants were invited to

102   monitor their chosen feeding location during weekly two-day observation periods throughout the

winter. Participants recorded which species they observed and the maximum number of co-

104   occurring individuals that they observed simultaneously at the feeding location for each species.

Additionally, as a measure of observation effort, each participant indicated whether they

106   performed morning and/or afternoon monitoring sessions during each day (up to a maximum

value of four monitoring sessions per two-day observation period). For the purpose of the present

108   study, we sampled species with at least 100 displacement interactions in PFW as the focal

species for our further comparative analyses (n = 68 species). All data analyses were performed

110   in R (version 4.0.3) [30].

112   *(b) Quantifying species differences in sociality*

As our measure of sociality, we used group size, defined as the maximum number of individuals

114   of a given species observed simultaneously during a given monitoring session (Figure 1A). To

assess whether focal species differ in their average winter sociality, we analyzed the repeatability

116   of this metric using the "rptR" package [31]. Hence, repeatability is defined as the proportion of

total variation in group size that is attributed to differences among species. A high value of

118   repeatability in this analysis indicates that most of the variation in group size is due to

differences among species, whereas a low value (near 0) indicates that species differences in this

120   trait are not detected. Because the size of the dataset was so large (n = 8,575,748 observations of

the 68 focal species), we used a bootstrapping approach to estimate repeatability. For each

122   bootstrap iteration, we randomly selected a subset of data with 1,000 observed group sizes for

each of the 68 focal species (randomly chosen from the full dataset available for each species).

124   We used this subset to compute conditional repeatability, adjusted by observation effort, using

the Poisson distribution and log link as appropriate for these count data. This procedure was

126   repeated 100 times to derive a distribution of repeatability estimates for species differences in

average group size.

128   Upon finding that average group size differs among species, we estimated each focal

species' average group size as our measure of sociality for further comparative analyses of

130   species traits. To do this, we fit a regression model for each focal species where the response

variable was the observed group size (log-transformed; n = 3,200 – 551,026 observations,

132   depending on the species), with a fixed effect of observation effort. From the fitted models, we

defined each species' sociality as its predicted average group size on a log scale given an effort

134   of one monitoring session (Figure 1B). We also estimated Pagel's lambda for species average

group size [32]. This value ranges between 0 and 1, where values closer to 1 indicate that closely

136   related species have similar trait values, and values near 0 indicate that trait variation is

independent of phylogenetic relatedness.

138   As an additional check on our metric of group size, we collected information on the

natural history each focal species during winter from species accounts in the Birds of the World

140   online resource [33]. We used these descriptive accounts to classify each focal species into one

of three categories: often flocking (clear descriptions of winter flocking and/or group behaviour),

142   sometimes flocking (indicators of occasional winter flocking and/or participation as a satellite

member), and rarely flocking (described as independent in winter).

(c) Quantifying species dominance traits

146   We studied dominance interactions by using a dataset of displacement events collected by

participants during a subset of the PFW observation periods. Dominance events were defined as

148   instances when one species displaced another at a feeding location. Participants recorded 88,988

displacement interactions among 196 bird species, with most of these observations (71%)

150   collected in winters of 2017-18 and 2018-19. Note that although we filtered non-focal species

from our comparative analyses, we retained the interactions they had with focal species when

152   characterizing focal species traits.

        For each focal species, we quantified two species dominance traits: a species' average

154   competitive dominance and its 'conspecific effect'. Competitive dominance was defined as the

        probability that a focal species would be the displacer in a dominance interaction against a same-

156   sized (body mass) heterospecific opponent. To estimate competitive dominance, we fit a

        binomial mixed-effects model for each focal species where the response variable was the

158   outcome of their displacement interactions (i.e., whether the focal species was the displacer, or

        was displaced). These models were fit using the "lme4" package [34] and included two fixed

160   effects: the focal species' observed group size during that monitoring session (Figure S1), and

        the relative body mass of the opponent species . Relative body mass was defined as the

162   difference in log-transformed species body masses (opponent species body mass – focal species

        body mass). To account for repeated measures at the same spatial location, the models also

164   included region as a random effect, defined by latitude and longitude coordinates rounded to the

        nearest degree. The average sample size for these models was 1,606 displacement interactions

166   per focal species (ranging 83 – 10,583).

        We used these same models to calculate the conspecific effect for each species, defined

168   as the average effect (positive or negative) of nearby conspecifics on the outcome of between-

        species competition. As illustrated in Figure 2B, C: the conspecific effect was calculated as the

170   expected increase in the probability of the focal species being the displacer over a heterospecific

        opponent, given a focal species group size of three, as compared to a focal species group size of

172   one. Note that we chose a group size of three because such smaller group sizes are far more

        commonly observed. The resulting metric, the conspecific effect, can take values from –1 to 1,

174 with negative values indicating that a species' dominance decreases in the presence of

conspecifics, and positive values indicating that a species' dominance increases in the presence

176 of conspecifics. A conspecific effect value of 0 would indicate that the presence of conspecifics

does not affect a species' dominance. We consider analyses of this trait to be conservative

178 because the PFW observers reported only one group size per species per monitoring session (the

maximum group size they observed). As a result, our estimates for the conspecific effect hinge

180 on the assumption that species group sizes are relatively stable at a given feeder within a short

span of time (e.g., [35–37]). Alternatively, if species group sizes are highly variable over the

182 span of a few hours, it would reduce our ability to detect conspecific effects.

184 *(d) Dominance traits in relation to sociality*

To test whether species dominance traits are related to species differences in sociality, we used

186 two regression models. The first model related a species' competitive dominance to group size,

and the second model related a species' conspecific effect to group size (n = 68 focal species). In

188 both models, the main predictor of interest was species average group size, and body mass was

included as an additional predictor. Models were fit using the "brms" package [38] with a

190 Gaussian error distribution. We ran these models both with and without accounting for the

phylogenetic relationships among focal species in the random effects structure. If the

192 phylogenetically adjusted model estimated a weaker effect size for a given predictor than the

simpler model, it would indicate that the relationship is driven by differences among broader

194 lineages. We ran models for 10,000 iterations using two chains and verified that Rhat values

were all between 1.00 and 1.01, indicating model convergence.

(e) Within-species conflict

198   Competitive interactions with conspecifics are among the most common (Figure 1D). To test

whether within-species conflict occurs more often than expected, we compared the observed

200   frequencies of within-species displacement interactions in PFW with expected values based on a

null model that was spatially constrained by region. The null model was computed as follows:

202   within each region (defined by latitude and longitude rounded to the nearest degree), we first

determined the proportion of total interactors for each species. Then, we calculated the expected

204   number of conspecific interactions for each species, under the assumption that interaction rates

are determined by the composition of interactors in that region. We took the square of a given

206   species' regional proportion as the expected probability of a conspecific interaction for that

species and region. We then multiplied this probability by the number of interactions in the

208   region to obtain an expected count. Finally, we took the sum of these expected conspecific

interaction counts across all regions for a given focal species. When analyzing the expected and

210   observed within-species interactions, we log-transformed both variables to normalize residuals.

To test whether a species' social behaviour predicts within-species conflict, we fit a

212   regression model where the response variable was the proportion of competitive displacement

interactions that were conspecific (one value per focal species). The predictors in this analysis

214   included three species traits: average group size, the conspecific effect, and body mass. This

model was fit using the Bayesian brms package as described above, and we examined the model

216   with and without accounting for the phylogenetic relationships among focal species.

218   **RESULTS**

(a) Species differ in the degree of sociality

220   There is substantial variation both among and within species in average group size, as indicated

the repeatability for this trait ($R_{\text{link-scale}}$ = 0.36, 95% CI = [0.29, 0.44]; Figure 1A, n = 68 focal

222   species). Species average group size also has a strong phylogenetic signal (lambda = 0.99, p <

0.001), indicating that closely related species share a very similar degree of sociality (Figure 1B).

224   Species average group size also corresponds to broad descriptions from natural history accounts

(Figure 1C). Notably, our metric captures group size variation both within and among species.

(b) Estimates of the conspecific effect

228   The conspecific effect varied widely across species, with most estimates centred on 0 (average

conspecific effect of -0.10, 95% CI = [-0.37, 0.17]; Figure S1). We detected a significant positive

230   conspecific effect in only one species, the pine siskin (*Carduelis pinus*), indicating that pine

siskins gain a large increase in competitive dominance when more group members are present

232   (0.12, 95% CI = [0.08, 0.17]; Figure 2A, B). We detected significant negative conspecific effects

in 14 species, including the pine warbler (*Dendroica pinus*; -0.27 [-0.44, -0.10]), the white-

234   winged dove (*Zenaida asiatica*; -0.16 [-0.32, -0.05]; Figure 2A, C), and the eastern bluebird

(*Sialia sialis*; -0.13 [-0.21, -0.04]). These species are less dominant when more conspecific group

236   members are present.

238   *(c) Social species are poor competitors, but gain a competitive advantage in groups*

Contrary to our initial expectation, we found that more social species were less likely to

240   dominate other similarly-sized opponent species (Figure 3A, Table S2). The conclusions of this

analysis were consistent when accounting for phylogeny (Table S2). More social species also

242 had more positive conspecific effects (Figure 3B, Table S3). This effect remained strong when accounting for phylogenetic nonindependence (Table S3).

(d) Social species have elevated within-group conflict

246 In 62 of the 68 focal species, displacement interactions with conspecifics occurred more often than expected in the null model (Figure 4A). As expected, more social species receive a greater

248 proportion of conflict originating from conspecifics (Figure 4B, S2, Table S4). This association was weaker when accounting for phylogenetic relationships, indicating that it is primarily driven

250 by differences among lineages within our sample (Table S4).

252 **DISCUSSION**

Here, we examine the effect of social clustering on competition among birds across the North

American continent. We find that species vary in their average group size, and that sociality during winter foraging has been highly conserved in avian evolution. Contrary to our initial

prediction, we find that more social species are also weaker competitors when interacting with similarly sized heterospecific opponents. This result suggests that there may be an evolutionary

trade-off whereby the evolution of greater sociality is correlated with reduced individual competitive ability against other opponent species. Our results also show that more social species

gain a greater dominance advantage in the presence of their conspecifics, whereas solitary species are less dominant in the presence of conspecifics. Collectively, these results suggest that

more social species have reduced competitive ability as individuals, but enhanced competitive ability in groups.

These results indicate that selection for pro-social behaviours may be associated with a

reduction in traits such as aggression that would provide an advantage to an individual in

266   heterospecific competition [39]. As species evolve to become more social, they may also acquire

behavioral plasticity in aggression, boldness, or vigilance behaviour that may facilitate greater

268   dominance over heterospecific opponents when in the presence of one or more conspecifics

[40,41]. It is also possible that opponents respond differently to social species depending on their

270   group size (e.g., because larger groups generate a more intimidating visual and/or acoustic

signal). To test these hypotheses, experiments are needed to measure behavioural outcomes

272   under controlled conditions where the social environment can also be carefully manipulated.

      Our results examining within-species conflict support the hypothesis that the evolution of

274   sociality leads to greater conflict within-groups. In the presence of many conspecifics,

individuals may interact more often, especially if they are collecting similar resources [7,8].

276   Members of a group may experience increased rates of aggressive interactions at times of food

scarcity [42,43]. One way to reduce competition pressures between conspecifics may be through

278   the participation in a mixed-species flock. Mixed-species flocks have been found to provide

members with greater advantages than conspecific coalitions, including increased foraging rates

280   and reduce vigilance effort [1]. While some evidence suggests that members of mixed-species

flocks are more similar than expected [44], they undoubtedly vary more in their ecological needs

282   than same-species flocks and therefore may demonstrate a reduced level of within-group conflict

than a conspecific coalition [45–47]. A key question for future research is whether mixed-species

284   flocking reduces the frequency of within-group conflict.

      It is important to note that while we define sociality as the aggregation of individuals in

286   space and time, there are other dimensions of sociality that are also expected to play a role in

influencing competition. For example, members of a temporary flock with short-term social

288 bonds may experience a weaker influence of conspecifics than of members in a structured

dominance hierarchy with long-term, persistent bonds [12,48–50]. This raises interesting

290 questions that can only be addressed by tracking individuals within groups, such as: How does

the strength of social bonds and the organization of a group influence a species' competitive

292 ability? Do such effects on competition change with temporal variation in species social

organization? Within a species, how does the strength of a coalitionary relationship affect the

294 dominance of one group over others?

Previous studies have examined the benefits of coalitionary behaviour on within-species

296 conflict [9,14,51]. Our results here demonstrate how sociality can also influence the dynamics of

ecological competition, and how a species' tendency to form social groups is associated with the

298 evolution of reduced competitive ability as individuals. These findings provided valuable insight

into the fundamental effects of grouping behaviour on competitive interactions during foraging.

300 Further questions remain as to the effects of specific social bonds and the emergent properties of

groups on competitive interactions.

FIGURE LEGENDS

Figure 1. North American bird species differ in winter sociality and competitive dominance. A) Example of daily average group sizes (± SE) for seven bird species during winter 2017-18, with average group size indicated by colour of the line. B) Phylogeny of 68 focal bird species with average group size shown by the size and colour of the outer bars. Note that the colour gradient throughout this figure follows the legend in (A). C) Species average group size in relation to descriptions from natural history accounts. Horizontal lines indicate medians. D) Network illustrating dominance interactions among a subset of the most common species. Each row represents a species, and the grey connections show frequencies of competitive interactions. Horizontal connections represent competitive interactions with conspecifics. Species nodes are ordered vertically (top to bottom) from the most to least dominant, and coloured by average group size. Labels represent species common names (see Table S1 for details).

Figure 2. Predictors of competitive dominance against other species. A) Partial residual plot showing the predicted probability of competitive dominance as a function of group size for two species models, the pine siskin (green circles) and white-winged dove (orange triangles). B-C) These fitted models were used to calculate the conspecific effect by determining the change in the probability of being the displacer when comparing a group size of three with a solitary condition. D) Caterpillar plot showing estimates of the conspecific effect for each focal species (± 95% confidence intervals). Species shown in grey have a confidence interval that includes zero, those shown in red have a significant negative conspecific effect, and the species shown in blue has a significant positive conspecific effect.

Figure 3. Social species are poor competitors as individuals, but gain a competitive advantage in groups. Partial residual plots show the analysis of two species traits: (A) competitive dominance, and (B) the conspecific effect, in relation to average group size (n = 68 focal species). Partial residuals on the y-axis account for other predictors in the analysis. The lines show the fitted model.

Figure 4. Conflict with conspecifics occurs more often than expected, especially in more social lineages. (A) Observed and expected counts of conspecific displacement interactions (n = 68 focal species). The red dotted line indicates a 1:1 line where the observed frequency matches the expected value in the null model; the black line is the line of best fit for the observed and expected values. (B) Partial residual plot showing the analysis of conspecific conflict. The line indicates the fitted model.

DATA ACCESSIBILITY

340 All data and R scripts are available at: https://figshare.com/s/1cd6708a27a541029409
The repository will be made public when the final version of the study is published.

AUTHOR CONTRIBUTIONS

344 IB and RD designed the study with input from ETM. ETM developed the behaviour interactions
methodology in Project FeederWatch. IB and RD analyzed the data and wrote the manuscript.

346 All authors edited the manuscript.

348 COMPETING INTERESTS
We have no competing interests.

FUNDING

352 Supported by an NSERC Discovery Grant to RD and Carleton University.

354 ACKNOWLEDGEMENTS
We would like to thank the Cornell Lab of Ornithology and Birds Canada for developing and

356 managing Project FeederWatch and sharing the results. We especially thank the thousands of
volunteers who have contributed their observations to Project FeederWatch.

References

1. Sridhar H, Beauchamp G, Shanker K. 2009 Why do birds participate in mixed-species
362 foraging flocks? A large-scale synthesis. *Anim. Behav.* **78**, 337–347.
(doi:10.1016/j.anbehav.2009.05.008)

364 2. Beauchamp G. 1998 The effect of group size on mean food intake rate in birds. *Biol. Rev.*
73, 449–472. (doi:10.1111/j.1469-185X.1998.tb00179.x)

366 3. McDonald DB, Potts WK. 1994 Cooperative Display and Relatedness Among Males in a
Lek-Mating Bird. *Science.* **266**, 1030–1032. (doi:10.1126/science.7973654)

368 4. Oh KP, Badyaev A V. 2010 Structure of social networks in a passerine bird:
Consequences for sexual selection and the evolution of mating strategies. *Am. Nat.* **176**.
370 (doi:10.1086/655216)

5. Sorato E, Gullett PR, Griffith SC, Russell AF. 2012 Effects of predation risk on foraging
372 behaviour and group size: Adaptations in a social cooperative species. *Anim. Behav.* **84**,
823–834. (doi:10.1016/j.anbehav.2012.07.003)

374 6. Cresswell W. 1994 Flocking is an effective anti-predation strategy in redshanks, *Tringa
totanus* . *Anim. Behav.* **47**, 433–442. (doi:10.1006/anbe.1994.1057)

376 7. Furness RW, Birkhead TR. 1984 Seabird colony distributions suggest competition for
food supplies during the breeding season. *Nature* **311**, 655–656. (doi:10.1038/311655a0)

378 8. Forero MG, Tella JL, Hobson KA, Bertellotti M, Blanco G. 2002 Conspecific Food
Competition Explains Variability in Colony Size: A Test in Magellanic Penguins. *Ecology*
380 **83**, 3466. (doi:10.2307/3072095)

9. Packer C, Pusey AE. 1982 Cooperation and competition within coalitions of male lions:
382 kin selection or game theory? *Nature* **296**, 740–742. (doi:10.1038/296740a0)

10. Lamprecht J. 1978 The Relationship between Food Competition and Foraging Group Size
384 in some Larger Carnivores. *Z. Tierpsychol.* **46**, 337–343. (doi:10.1111/j.1439-
0310.1978.tb01455.x)

386 11. Bribiesca R, Herrera-Alsina L, Ruiz-Sanchez E, Sánchez-González LA, Schondube JE.
2019 Body mass as a supertrait linked to abundance and behavioral dominance in
388 hummingbirds: A phylogenetic approach. *Ecol. Evol.* **9**, 1623–1637.
(doi:10.1002/ece3.4785)

390 12. Miller ET, Bonter DN, Eldermire C, Freeman BG, Greig EI, Harmon LJ, Lisle C,
Hochachka WM. 2017 Fighting over food unites the birds of North America in a
392 continental dominance hierarchy. *Behav. Ecol.* **28**, 1454–1463.
(doi:10.1093/beheco/arx108)

394 13. Šárová R, Špinka M, Stěhulová I, Ceacero F, Šimečková M, Kotrba R. 2013 Pay respect
to the elders: Age, more than body mass, determines dominance in female beef cattle.
396 *Anim. Behav.* **86**, 1315–1323. (doi:10.1016/j.anbehav.2013.10.002)

14. Zabel CJ, Glickman SE, Frank LG, Woodmansee KB, Keppel G. 1992 Coalition
398 formation in a colony of prepubertal spotted hyenas. In *Coalitions and Alliances in
Humans and other Animals* (eds AH Harcourt, FBM de Waal), pp. 113–136. Oxford:
400 Oxford University Press.

15. Lahti K, Orell M, Rytkonen S, Koivula K. 1998 Time and Food Dependence in Willow
402 Tit Winter Survival. *Ecology* **79**, 2904. (doi:10.2307/176525)

16. Jansson C, Ekman J, von Brömssen A, von Bromssen A. 1981 Winter Mortality and Food
404 Supply in Tits Parus spp. *Oikos* **37**, 313. (doi:10.2307/3544122)

17. Brittingham MC, Temple SA. 1988 Impacts of Supplemental Feeding on Survival Rates of Black-Capped Chickadees. *Ecology* **69**, 581–589. (doi:10.2307/1941007)

18. Plummer KE, Bearhop S, Leech DI, Chamberlain DE, Blount JD. 2013 Winter food provisioning reduces future breeding performance in a wild bird. *Sci. Rep.* **3**, 2002. (doi:10.1038/srep02002)

19. Kalinoski R. 1975 Intra- and Interspecific Aggression in House Finches and House Sparrows. *Condor* **77**, 375. (doi:10.2307/1366086)

20. Wojczulanis-Jakubas K, Kulpińska M, Minias P. 2015 Who bullies whom at a garden feeder? Interspecific agonistic interactions of small passerines during a cold winter. *J. Ethol.* **33**, 159–163. (doi:10.1007/s10164-015-0424-x)

21. Dias RI. 2006 Effects of position and flock size on vigilance and foraging behaviour of the scaled dove Columbina squammata. *Behav. Processes* **73**, 248–252. (doi:10.1016/j.beproc.2006.06.002)

22. Roberts G. 1996 Why individual vigilance declines as group size increases. *Anim. Behav.* **51**, 1077–1086. (doi:10.1006/anbe.1996.0109)

23. Beauchamp G. 2008 What is the magnitude of the group-size effect on vigilance? *Behav. Ecol.* **19**, 1361–1368. (doi:10.1093/beheco/arn096)

24. Bijleveld AI, van Gils JA, Jouta J, Piersma T. 2015 Benefits of foraging in small groups: An experimental study on public information use in red knots Calidris canutus. *Behav. Processes* **117**, 74–81. (doi:10.1016/j.beproc.2014.09.003)

25. Valone TJ. 2007 From eavesdropping on performance to copying the behavior of others: A review of public information use. *Behav. Ecol. Sociobiol.* **62**, 1–14. (doi:10.1007/s00265-007-0439-6)

26. Gompper ME. 1996 Sociality and asociality in white-nosed coatis (*Nasua narica*): foraging costs and benefits. *Behav. Ecol.* **7**, 254–263. (doi:10.1093/beheco/7.3.254)

27. Krause J, Ruxton GD. 2002 *Living in Groups*. Oxford: Oxford University Press.

28. Grueter CC *et al.* 2020 Multilevel Organisation of Animal Sociality. *Trends Ecol. Evol.* **35**, 834–847. (doi:10.1016/j.tree.2020.05.003)

29. Papageorgiou D, Farine DR. 2021 Multilevel Societies in Birds. *Trends Ecol. Evol.* **36**, 15–17. (doi:10.1016/j.tree.2020.10.008)

30. R Core Team. 2019 R: A Language and Environment for Statistical Computing.

31. Stoffel MA, Nakagawa S, Schielzeth H. 2017 rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* **8**, 1639–1644. (doi:10.1111/2041-210X.12797)

32. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884. (doi:10.1038/44766)

33. Billerman SM, Keeney BK, Rodewald PG, Schulenberg TS, editors. 2020 *Birds of the world*. Ithaca, NY, USA: Cornell Laboratory of Ornithology. See https://birdsoftheworld.org/bow/home.

34. Bates D, Mächler M, Bolker B, Walker S. 2014 Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

35. Balph MH. 1979 Flock Stability in Relation to Social Dominance and Agonistic Behavior in Wintering Dark-Eyed Juncos. *Auk* **96**, 714–722. (doi:10.1093/auk/96.4.714)

36. Odum EP. 1942 Annual Cycle of the Black-Capped Chickadee: 3. *Auk* **59**, 499–531. (doi:10.2307/4079461)

37. Stouffer PC, Caccamise DF. 1991 Roosting and diurnal movements of radio-tagged

American crows. *Wilson Bull.* **103**, 387–400.

452 38. Bürkner P-C. 2017 brms : An R Package for Bayesian Multilevel Models Using Stan. *J. Stat. Softw.* **80**, 1–28. (doi:10.18637/jss.v080.i01)

454 39. Magurran AE, Seghers BH. 1991 Variation in Schooling and Aggression Amongst Guppy (Poecilia Reticulata) Populations in Trinidad. *Behaviour* **118**, 214–234.
456 (doi:10.1163/156853991X00292)

40. Webster MM, Ward AJW, Hart PJB. 2007 Boldness is influenced by social context in
458 threespine sticklebacks (Gasterosteus aculeatus). *Behaviour* **144**, 351–371. (doi:10.1163/156853907780425721)

460 41. Humfeld SC, Marshall VT, Bee MA. 2009 Context-dependent plasticity of aggressive signalling in a dynamic social environment. *Anim. Behav.* **78**, 915–924.
462 (doi:10.1016/j.anbehav.2009.06.028)

42. Symons PEK. 1968 Increase in Aggression and in Strength of the Social Hierarchy among
464 Juvenile Atlantic Salmon Deprived of Food. *J. Fish. Res. Board Canada* **25**, 2387–2401. (doi:10.1139/f68-207)

466 43. Collie J, Granela O, Brown EB, Keene AC. 2020 Aggression Is Induced by Resource Limitation in the Monarch Caterpillar. *iScience* **23**, 101791.
468 (doi:10.1016/j.isci.2020.101791)

44. Sridhar H *et al.* 2012 Positive Relationships between Association Strength and Phenotypic
Similarity Characterize the Assembly of Mixed-Species Bird Flocks Worldwide. *Am. Nat.* **180**, 777–790. (doi:10.1086/668012)

45. Metcalfe NB, Furness RW. 1987 Aggression in shorebirds in relation to flock density and composition. *Ibis (Lond. 1859).* **129**, 553–563. (doi:10.1111/j.1474-919X.1987.tb08243.x)

46. Sasvári L. 1992 Great tits benefit from feeding in mixed-species flocks: a field experiment. *Anim. Behav.* **43**, 289–296. (doi:10.1016/S0003-3472(05)80224-6)

47. Hino T. 2000 Intraspecific differences in benefits from feeding in mixed-species flocks. *J. Avian Biol.* **31**, 441–446. (doi:10.1034/j.1600-048X.2000.310402.x)

48. Myers JP. 1983 Space, time and the pattern of individual associations in a group-living species: Sanderlings have no friends. *Behav. Ecol. Sociobiol.* **12**, 129–134.
(doi:10.1007/BF00343203)

49. Conklin JR, Colwell MA. 2008 Individual associations in a wintering shorebird
482 population: Do Dunlin have friends? *J. F. Ornithol.* **79**, 32–40. (doi:10.1111/j.1557-9263.2008.00143.x)

484 50. Tibbetts EA, Pardo-Sanchez J, Weise C. 2022 The establishment and maintenance of dominance hierarchies. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **377**, 20200450.
486 (doi:10.1098/rstb.2020.0450)

51. Markham AC, Alberts SC, Altmann J. 2012 Intergroup conflict: Ecological predictors of
488 winning and consequences of defeat in a wild primate population. *Anim. Behav.* **84**, 399–403. (doi:10.1016/j.anbehav.2012.05.009)

Figure 1

Figure 2

Figure 3

Figure 4