The effect of sociality on competitive interactions among birds

Ilias Berberi[1], Eliot T. Miller[2], Roslyn Dakin[1]

[1]Department of Biology
Carleton University
1125 Colonel By Drive
Ottawa, Canada

[2]Cornell Lab of Ornithology
159 Sapsucker Woods Road
Ithaca, New York

To whom correspondence should be addressed:
Ilias Berberi
iliasberberi@gmail.com
Roslyn Dakin
roslyn.dakin@gmail.com

Keywords: competition, network, foraging, flocking, citizen science, behavioural ecology

ABSTRACT

Sociality can provide many benefits, including increased foraging success, reproductive opportunities, and defence against predation. How does sociality influence the dominance hierarchies of ecological competitors? Here, we address this question using a large citizen science dataset of competitive interactions among birds foraging at backyard feeders, representing a network of over 55,000 interactions among 68 common species. We first show that species differ in average group size (the number of conspecifics observed together) as a fundamental measure of sociality. When analyzing heterospecific competition, we find that sociality is inversely related to dominance. On average, a single individual from a solitary species is more likely to displace a size-matched opponent than a single individual from a social species. Yet, we find that social species gain an increase in their competitive advantage when in the presence of their conspecifics, which may occur as a result of dynamics within their groups. Finally, we show that more social species have relatively fewer dominance interactions with heterospecifics, and more with conspecifics. Overall, these results demonstrate that sociality can influence competition in ecological networks. More social species have decreased competitive ability as individuals, but they may gain competitive ability in groups.

INTRODUCTION

Animal sociality can provide many ecological and evolutionary advantages, including increased foraging success [1,2], more frequent mating opportunities [3,4], and improved survival through dilution and confusion effects on predators [5,6]. However, sociality may also incur costs under certain circumstance, such as conflict among conspecific group members that share a similar niche [7,8]. The causes and consequences of animal sociality have historically been of interest in

evolutionary biology, with several different hypotheses for benefits that may explain the

44   evolution of social grouping. One hypothesized benefit of sociality is that social coalitions may

provide an advantage during conflict with competitors [9,10]. While both inter- and intraspecific

46   dominance is often enhanced by body size [11,12] and age/experience [13], there may also be

opportunities for subordinate individuals to gain an advantage over dominant competitors

48   through social coalitions [9,14]. During foraging, animals face competition from both

conspecifics and heterospecifics, but the influence of social grouping on the structure of these

50   competitive systems has not yet been described. Our aim in this study is to evaluate how

sociality influences competition in communities where interactions occur both within and

52   between species.

One system that provides an opportunity to study this is the network of competitive

54   interactions among birds at backyard bird feeders. In many temperate systems, survival in winter

depends on acquiring adequate food resources [15,16]. Moreover, the use of artificial food

56   sources (e.g., bird feeders) during winter foraging has been identified as having important

evolutionary consequences in birds [17,18], including high rates of competitive interactions

58   among and within species foraging at feeder locations [19,20]. These levels of competition may

depend on the rate of resource depletion, where regularly depleted feeders may generate higher

60   levels of resource competition than feeders that are continually replenished [21]. In the context

of sociality, group foraging can provide more opportunities for resource acquisition than

62   foraging independently due to shared vigilance [22,23] (but see [24]), social information

gathering [25,26], and improved dominance over solitary foragers [27]. Individuals in many

64   species have been found to adopt social foraging strategies in the winter when resources are

scarce [28]. Additionally, sociality in the winter has been associated with reproductive

66   advantages in subsequent mating seasons [29,30].

        Here, we investigate how sociality influences competition using interactions among birds

68   that visit backyard feeders. Our predictions at the outset of this study were that more social

      species may gain a benefit in interspecific conflict, at a cost of increased rates of intraspecific

70   conflict. We tested these predictions using data from Project FeederWatch (PFW), a surveying

      program with a network of tens of thousands of observers across North America who monitor the

72   presence and interactions of hundreds of bird species throughout winter (Figure 1). This project

      has yielded a rich dataset indicating who competes with whom, and how often (Figure 1D).

74   Previous research using PFW observations has revealed that species body mass is the primary

      predictor of competitive dominance within winter bird communities across North America

76   [12,31]. Owing to the large number of interactions observed and variation in competitive

      outcomes (Figure 1D), the PFW citizen science program provides an ideal system to test how

78   variation in sociality might influence competitive interactions after accounting for the effects of

      body size.

80   Here, we define species differences in winter sociality using a simple metric of

      conspecific group size (i.e., the number of individuals of a given bird species observed at a given

82   location at the same time in the PFW database; Figure 1A-C) [32]. It is important to note that

      sociality also has other dimensions beyond the simple tempo-spatial aggregation studied here,

84   such as the stability of group membership, the types of relationships between group members,

      and other emergent properties of group organization [33,34]. Nevertheless, conspecific

86   groupings – individuals found in the same place at the same time – represent the most

fundamental dimension of sociality that we expect to influence competitive outcomes across

88   species.

As a first step in this study, we established that winter sociality differs across bird species

90   by quantifying the repeatability of observed group sizes (Figure 1A). We also estimated the

phylogenetic signal of species-average group size, which describes the extent to which this trait

92   is more similar among closely related species (Figure 1B). Next, we analyzed the outcome of

>55,000 heterospecific dominance interactions (a subset of these is shown in Figure 1D). Our

94   aim was to evaluate whether average group size (as a species-level trait) is associated with

species differences in dominance against size-matched opponents, and whether the presence of

96   conspecifics would positively or negatively impact each species' competitive ability (hereafter,

we refer to this impact as the "conspecific effect"). We expected that more social species would

98   have greater dominance over size-matched opponents, and a more positive conspecific effect.

Finally, we analyzed the frequencies of conspecific and heterospecific interactions (Figure 1D),

100   to determine whether sociality is associated with variation in the most common sources of

conflict.

MATERIALS AND METHODS

104   *(a) Data sources*

We used a dataset collected by Project FeederWatch (PFW), an extensive, comprehensive and

106   ongoing monitoring program with over 9 million observations of birds at winter feeding

locations in North America during the months of November through April. For this study, we

108   collected PFW data from the period November 2015 – April 2020. Participants were invited to

monitor their chosen feeding location during weekly two-day observation periods throughout the

110   winter. Participants recorded which species they observed and the maximum number of co-

occurring individuals that they observed simultaneously at the feeding location for each species.

112   Some participants also recorded displacement interactions, used in our analysis of competition

described in section (c) of the Methods below. Additionally, as a measure of observation effort,

114   each participant indicated the number of two-hour monitoring sessions performed (up to a

maximum value of four monitoring sessions per two-day observation period). Collected PFW

116   data routinely undergo a data validation protocol (highlighted in [35]) to reduce biases in species

reporting and identification often associated with citizen science [36]. For the purpose of the

118   present study, we focused our analyses on a set of the most common species that had at least 100

displacement interactions (n = 68 species, Table S2). All data analyses were performed in R

120   (version 4.0.3) [37]. A detailed flow chart illustrating our main analysis methods is provided in

the supplement (Figure S1).

(b) Quantifying species differences in sociality

124   As our measure of sociality, we used group size, defined as the maximum number of individuals

of a given species observed simultaneously during a given monitoring session (see examples in

126   Figure 1A). To assess whether focal species differ in their average winter sociality, we analyzed

the repeatability of observed group size values (e.g., Figure 1A), [38]defined as the proportion of

128   total variation in observed group sizes that is attributed to differences among species. A high

value of repeatability in this analysis indicates that most of the variation in group size is due to

130   differences among species, whereas a low value (near 0) indicates that species differences in this

trait are not detected. Because group sizes change with the onset of spring, we included dates

132   from November 1 – February 28 (note that conclusions are unchanged when including a broader

span of dates). Because the size of the dataset was so large (n > 6.1 million observations of the

134   68 focal species), we used a bootstrapping approach to estimate repeatability. For each bootstrap

iteration, we randomly selected a subset of data with 1,000 observed group sizes for each of the

136   68 focal species (randomly chosen from the full dataset available for each species). We used this

subset to compute conditional repeatability of observed group size using the "rptR" package

138   [38], adjusted by observation effort as a fixed effect, and accounting for region (defined by

latitude and longitude to the nearest degree), using the Poisson distribution and log link for count

140   data. This procedure was repeated 100 times to derive a distribution of repeatability estimates for

species differences in average group size.

142   Upon finding that observed group size differs among species, we computed estimates of

species-average group size, which we use in further comparative analyses of our 68 focal species

144   (see Figure S1 for details). To do this, we fit a regression model for each focal species where the

response variable was the log-transformed observed group size (n = 2,257 – 414,471

146   observations, depending on the species), with a fixed effect of observation effort, and a random

effect of region. From each fitted model, we defined species-average group size as the predicted

148   group size given an effort of one monitoring session (Figure 1B; Table S2). We also estimated

Pagel's lambda for the 68 species-average group size values [39]. Lambda values close to 1

150   would indicate that closely related species have similar species-average group sizes, whereas

lambda values near 0 would indicate that variation in species-average group size is independent

152   of phylogenetic relatedness.

          As an additional check on our metric of species-average group size, we collected

154   information on the natural history each focal species during winter from species accounts in the

Birds of the World online resource [40]. We used these descriptive accounts to classify each

156 focal species into one of three categories: often flocking (clear descriptions of winter flocking

and/or group behaviour), sometimes flocking (indicators of occasional winter flocking and/or

158 participation as a satellite member), and rarely flocking (described as independent in winter).

(c) Modelling heterospecific competition

We modelled heterospecific dominance interactions by using a dataset of displacement events,

162 defined as instances when one individual purposefully takes over, or dislodges, another bird from

the feeding location (see supplement for further details). PFW participants recorded 88,988

164 displacement interactions among 196 bird species in winters of 2016-17, 2017-18, 2018-19 and

2019-20, with most of these observations (71%) collected in 2017-18 and 2018-19. In this

166 analysis, we used displacement interactions that involved two different species, wherein at least

one of the two interactors belonged to a focal species (n = 55,483 interactions). It is important to

168 note that these interactions involved one individual bird displacing another, and did not include

collective events such as mobbing (see supplement for details).

170 We used a binomial mixed-effects regression model in the "lme4" package [41] to model

the outcome of these heterospecific interactions. The aim of this model was to evaluate whether

172 average group size (as a species-level trait) is associated with species differences in dominance

against size-matched opponents, and whether the presence of conspecifics impacts competitive

174 ability differently across species. The model had random effects for the identity of the "primary"

species, and the identity of its opponent species. These identities were assigned randomly in a

176 bootstrapping procedure, with details below. We included an additional random effect of region.

The response variable was the outcome of the displacement interaction (i.e., whether the primary

178 species was dominant in the displacement interaction). The model included fixed effects of the

primary species' average body mass (log-transformed), the primary species' average group size

180   (log-transformed), its relative group size during that particular observation session (expressed

relative to its species average group size), and the relative body mass of its opponent (defined as

182   the difference in log-transformed species-average body masses, opponent – primary). In this

binomial model, the slope parameter for the primary species' average group size captures the

184   association between species-level differences in sociality and species-level differences in

dominance (we also describe further comparative analyses of this relationship in section (d)

186   below). The slope parameter for relative group size captures the "conspecific effect". A positive

estimate for this slope parameter would indicate that the presence of additional conspecifics can

188   boost competitive ability, whereas a negative slope would indicate that the presence of

conspecifics can hinder competitive ability. To test whether the conspecific effect is species-

190   specific, we included a random slope for the relative group size parameter. This allows the

conspecific effect to vary among species. We then compared the fit of the model with this

192   random slope to a simpler model that lacked the random (species-specific) slope using AIC and

the likelihood ratio test. Stronger support for the random slope model would indicate that the

194   conspecific effect varies significantly among species.

   To account for the fact that both species in a heterospecific interaction may be considered

196   as primary, we bootstrapped this analysis by repeating the procedure described above 100 times.

In each bootstrap iteration, the primary and opponent positions for each interaction were

198   randomly assigned. Hence, the bootstrapping procedure ensured that each displacement

interaction was only represented once, such that the analysis did not pseudoreplicate the data.

   (d) Comparative analyses of species dominance in relation to sociality

202 To test our main hypotheses about the relationships between dominance and winter sociality, we

used the bootstrapped model fits to quantify two further species traits for comparative analysis:

204 the 'competitive dominance' and 'conspecific effect' of each focal species. Competitive

dominance was defined as the predicted probability that a single focal individual (group size of

206 1) would be the displacer in a dominance interaction against a same-sized (equal body mass)

heterospecific opponent. We extracted these predictions from the 100 bootstrapped models

208 described in Methods section (c) above, and took the average for each species as our measure of

its competitive dominance when alone and facing size-matched opponents (see flow chart in

210 Figure S1 for details).

   To examine variation in the conspecific effect, we extracted the random (species-specific)

212 slopes for relative group size. Negative slope values indicate that a species' dominance decreases

in the presence of conspecifics, whereas positive slope values indicate that a species' dominance

214 increases in the presence of conspecifics. A value of 0 would indicate that the presence of

conspecifics does not affect a given species' dominance. As above, we extracted these values

216 from the 100 bootstrapped models in Methods section (c), and took the average for each species

as its 'conspecific effect'. We consider analyses of this trait to be conservative because the PFW

218 observers reported only one group size per species per monitoring session (the maximum group

size they observed). As a result, our estimates for the conspecific effect hinge on the assumption

220 that species group sizes are relatively stable at a given feeder within a short span of time (e.g.,

[42–44]). Alternatively, if species group sizes are highly variable over the span of a few hours, it

222 would reduce our ability to detect conspecific effects.

   To test the relations between species dominance and sociality, we ran two comparative

224 analyses using the "brms" package [45] (n = 68 focal species). The response variables were

species' competitive dominance, and the conspecific effect, respectively, modelled with

226   Gaussian error distributions. The predictor of interest was species-average group size (log

transformed). We ran these models both with and without accounting for the phylogenetic

228   relationships among species in the random effects structure. If the phylogenetically adjusted

model estimated a weaker effect size for a given predictor than the simpler model, it would

230   indicate that the relationship is driven by differences among broader lineages. We ran models for

10,000 iterations using two chains and verified that Rhat values were all between 1.00 and 1.05,

232   indicating model convergence. We verified that residuals for Gaussian regression models were

normally distributed with no heteroscedasticity. To check whether our conclusions were robust

234   to error in the y-values (Figure 2B,E) we repeated these comparative analyses 100 times using

the full distribution of y-values (repeating the analyses once for each unique set of bootstrapped

236   y-values). We found that the conclusions of the comparative analyses were consistent each time.

238   *(e) Sources of conflict*

Sociality may also influence the occurrence of competition (as well as its outcome). To compare

240   the typical sources of conflict for each species with those expected by chance, we used a null

model that was spatially constrained by region. The null model was computed as follows: within

242   each i^{th} region (defined by latitude and longitude rounded to the nearest degree), we first

determined the proportion of total interactors for each j^{th} focal species, p_{ij}. Then, we calculated

244   the expected number of conspecific and heterospecific interactions for each species, under the

assumption that interaction rates are determined by the composition of interactors in that region.

246   We took p_{ij}^2 as the expected probability of a conspecific interaction, and $2 \times p_{ij} \times (1 - p_{ij})$ as the

expected probability of a heterospecific interaction for each species-region. We then multiplied

248 these probabilities by the number of interactions in the region to obtain expected counts. Finally,

we took the sum of the expected counts across all regions to get the total expected frequencies

250 for each focal species and compared these null expectations with observed values.

To test whether a species' social behaviour predicts variation in the sources of conflict,

252 we fit a regression model where the response variable was the observed proportion of

interactions against heterospecific opponents (one value per focal species). The predictor in this

254 analysis was the species-average group size. This model was fit using the Bayesian brms

package as described above, and we examined the model with and without accounting for the

256 phylogenetic relationships among focal species. We verified that residuals for this model were

normally distributed with no heteroscedasticity.

RESULTS

260 *(a) Species differ in the degree of sociality*

Observed group size varies both within and among species (Figure 1A), as indicated by

262 significant repeatability ($R_{\text{link-scale}}$ = 0.37, range 0.36-0.37, bootstrapped analysis of n = 68,000

observed group size values). Species-average group size has a very strong phylogenetic signal

264 (lambda = 0.99, p < 0.001, n = 68, one value per focal species), indicating that closely related

species share a very similar degree of sociality (Figure 1B). Species-average group size values

266 also correspond to broad descriptions from natural history accounts (Figure 1C).

268 *(b) Species dominance in relation to sociality*

In the bootstrapped model of all interactions, we found a strong negative association between

270 heterospecific dominance and species-average sociality (Figure 2A). Consistent with this result,

comparative analysis confirmed that competitive dominance is negatively related to sociality

272   (Figure 2B, Table S3). Hence, a single individual from a more social species is much less likely

to dominate size-matched opponents, as compared to a single individual from a less social

274   species. This conclusion is upheld when accounting for phylogeny (Table S3).

We found strong support for species differences in the 'conspecific effect' (bootstrap

276   average of $\Delta AIC = 40.2$ in favour of random slope models; all LRT p-values < 0.006). Some

species, like the pine siskin (*Carduelis pinus*), gain a large increase in competitive ability when

278   more conspecifics are present (Figure 2C). Other species, like the northern cardinal (*Cardinalis*

cardinalis), become less dominant when more conspecifics are present (Figure 2D). We found

280   that the conspecific effect tends to be more positive in more social species (Figure 2E, Table S4).

Thus, more social species gain a greater boost in dominance from the presence of conspecifics,

282   as compared to less social species. This association, while weak, was upheld when accounting

for phylogeny (Table S4).

(c) Sources of conflict

286   Nearly all species experience less conflict with heterospecifics, and more conflict with

conspecifics, than expected in a null model (Figure 3A-B). When examining species differences

288   in the sources of conflict, we find that more social species have proportionately less conflict with

heterospecifics (and hence, a greater proportion of their conflict is with conspecifics; Figure 3C,

290   Table S5). This association was weaker when accounting for phylogeny, indicating that it is

primarily driven by differences among lineages within our sample (Table S5).

DISCUSSION

294   Here, we examine the effect of social clustering on competition among winter resident birds. We

find that species vary in group sizes, and that certain phylogenetic clades are more social than

296   others. Contrary to our initial prediction, we find that individuals from more social species are

also weaker competitors when facing size-matched heterospecific opponents (Figure 2B). This

298   result suggests that there may be an evolutionary trade-off whereby the evolution of greater

sociality is associated with reduced competitive ability of an individual against other species

300   [31]. Our results also show that more social species tend to experience an increase in competitive

ability when in the presence of their conspecifics (Figure 2E). Collectively, these results suggest

302   that more social species have reduced competitive ability as individuals, but gain a boost in

competitive ability in groups.

304   We can hypothesize several potential mechanisms that may drive the patterns in Figure 2.

First, the strong decrease in competitive dominance observed in more social species (Figure 2B)

306   may be due to a loss of aggressive or antagonistic behaviours against heterospecifics during the

evolution of sociality. It is also possible that social species simply invest less into competition

308   with other species over food, for example if it is more beneficial for social species to limit their

interactions to conspecific competitors, or if social species are less motivated to compete because

310   they can rely on other strategies for finding food (e.g., social information transfer) [46,47]. To

test these hypotheses, further studies are required to test behavioural differences under conditions

312   where motivation and experience can be controlled.

        We can also propose several hypotheses that may explain variation in the conspecific

314   effect, and the fact that it is generally more positive in more social species (Figure 2E). One

possibility is that the conspecific effect may be driven intrinsically, as a consequence of other

316   species-level traits. For example, social species may have increased plasticity in aggression,

boldness, and/or vigilance that allows them to adjust their agonistic behaviour depending on the

318   context [48,49]. The evolution of sociality has also been shown to be associated with enhanced

cognition [50] and collective intelligence [51], which could also underly this effect. Yet another

320   possibility is that the conspecific effect could be a by-product of within-species dominance

hierarchies in more social species. For example, when a social flock visits a feeder, access may

322   be biased toward the most dominant and aggressive individual(s) in the group, who may in turn

be more likely to prevail against heterospecific competitors. The conspecific effect may also be

324   driven extrinsically by opponent responses (e.g., if larger groups of a more social species are

more coordinated in ways that generate more intimidating visual and/or acoustic signals). These

326   intrinsic and extrinsic explanations for the conspecific effect are not mutually exclusive. Testing

causal hypotheses will require careful experimentation.

328   Our results examining sources of conflict suggest that the evolution of sociality leads to

less frequent conflict with heterospecific competitors, and more frequent conflict among

330   conspecifics (Figure 3). In the presence of many conspecifics, individuals may interact more

often, especially if they are collecting similar resources [7,8]. Members of a group may

332   experience increased rates of aggressive interactions at times of food scarcity [52,53]. One way

to reduce competition pressures between conspecifics may be through participation in mixed-

334   species flocks. Mixed-species flocks have been found to provide members with greater

advantages than conspecific coalitions, including increased foraging rates and reduce vigilance

336   effort [1]. While some evidence suggests that members of mixed-species flocks are more similar

than expected [54], they undoubtedly vary more in their ecological needs than same-species

338   flocks and therefore may demonstrate a reduced level of within-group conflict than a conspecific

coalition [55–57]. A key question for future research is whether mixed-species flocking can

340   reduce the frequency of within-group conflict.

        Our results raise interesting questions for future research on other ecological factors. For

342   example, low resource availability is expected to generate higher levels of competition [21]. It

would be interesting to test whether environmental harshness influences the magnitude of the

344   conspecific effect. It is possible that social species experience even stronger conspecific effects

when resources are more limited. Additionally, weather conditions such as precipitation and

346   temperature may constrain foraging behaviour and may affect the costs and benefits of

competitive interactions as well as social clustering behaviour [58]. Further studies are thus

348   required to investigate how these environmental challenges may affect the structure of

competitive dominance networks.

350   It is important to note that while we define sociality as the aggregation of individuals in

space and time, there are other dimensions of sociality that are also expected to play a role in

352   influencing competition. For example, members of a temporary flock with short-term social

bonds may experience a weaker influence of conspecifics than members in a structured

354   dominance hierarchy with long-term, persistent bonds [12,59–61]. This raises interesting

questions for future work that can be addressed by tracking specific individuals within groups.

356   For example, how does the strength of social bonds and the organization of a group influence a

species' competitive ability? Do such effects on competition change with temporal variation in

358   species social organization? Within a species, how does the strength of a coalitionary

relationship affect the dominance of one group over others?

360   Previous studies have examined the benefits of coalitionary behaviour on within-species

conflict [9,14,62]. Our results here demonstrate how sociality can also influence the dynamics of

362   ecological competition, and how a species' tendency to form social groups can be associated

with the evolution of reduced competitive ability as individuals. These findings provide valuable

364   insight into the fundamental effects of grouping behaviour on competitive interactions during

foraging. Further questions remain as to the effects of specific social bonds and the emergent

366   properties of groups on competitive interactions, along with the effects of additional influential

variables, such as cognitive ability, resource abundance, and environmental conditions on the

relationships between sociality and competition.

FIGURE LEGENDS

Figure 1. Bird species differ in winter sociality and competitive dominance. A) Example of observed group sizes (daily averages ± SE) for seven bird species during winter 2017-18, with
species identity indicated by the line type. These species were chosen for visualization purposes to represent a wide range of avian families. B) Phylogeny of 68 focal bird species with species-
average group size shown by the size and colour of the outer bars. Note that the colour gradient throughout this figure follows the legend shown in (B). C) Species-average group size in relation
to descriptions from natural history accounts. Horizontal lines indicate medians. D) Competitive network illustrating dominance interactions among a subset of common species, chosen for
visualization purposes, with at least 2,000 displacement interactions. Each row represents a species, and the thickness of grey connecting lines indicates the frequency of competitive
interactions. Horizontal connections represent competitive interactions with conspecifics. Species nodes are ordered vertically (top to bottom) from the most to least dominant and
coloured by average group size. Labels represent short-form common names (see Table S1 for details).
386
       Figure 2. Relationship between sociality and heterospecific dominance. A) There is a strong
negative association between species-average sociality and heterospecific dominance. This density plot shows the distribution of effect sizes from the bootstrapped models of heterospecific
interactions in PFW. B) Scatterplot showing each species' competitive dominance (the predicted probability of a single individual winning a heterospecific competition) in relation to species-
average group size (n = 68 focal species). C-D) Examples illustrating how species differ in the relationship between dominance and the presence of conspecifics. Panels C-D show partial
residual plots for two species: the pine siskin, where larger group sizes are associated with increased competitive ability, and the northern cardinal, where larger group sizes are associated
with reduced competitive ability. E) Scatterplot showing the conspecific effect in relation to species-average group size (n = 68 focal species). In B and E, the y-values show bootstrapped
averages (error bars represent 2.5th and 97.5th percentiles), and the grey lines show model fits based on 100 bootstrapped estimates (see methods for details).
400
       Figure 3. Conflict with heterospecifics occurs less often than expected, especially in more
402   **social lineages.** Observed frequencies of (A) heterospecific and (B) conspecific dominance interactions in relation to null expected values for 68 focal species. Along the red dotted 1:1
404   lines, the observed frequency equals the expected value. (C) Scatterplot showing the proportion of conflict with heterospecifics in relation to species-average group size. Note that the line of
406   best fit does not account for phylogeny (see Table S5 for additional details).

DATA ACCESSIBILITY

408 All data and R scripts are available at: https://figshare.com/s/1cd6708a27a541029409
The repository will be made public when the final version of the study is published.
410

AUTHOR CONTRIBUTIONS

412 IB and RD designed the study with input from ETM. ETM developed the behavioural
interactions methodology in Project FeederWatch. IB and RD analyzed the data and wrote the
414 manuscript. All authors edited the manuscript.

COMPETING INTERESTS

416 We have no competing interests.
418

FUNDING

420 Supported by an NSERC Discovery Grant to RD and Carleton University.

ACKNOWLEDGEMENTS

422 We would like to thank the Cornell Lab of Ornithology and Birds Canada for developing and
424 managing Project FeederWatch and sharing the results. We especially thank the thousands of
volunteers who have contributed their observations to Project FeederWatch, and we thank the
426 peer reviewers for helpful feedback on this study.

References

1. Sridhar H, Beauchamp G, Shanker K. 2009 Why do birds participate in mixed-species foraging flocks? A large-scale synthesis. *Anim Behav* **78**, 337–347. (doi:10.1016/j.anbehav.2009.05.008)

2. Beauchamp G. 1998 The effect of group size on mean food intake rate in birds. *Biological Reviews* **73**, 449–472. (doi:10.1111/j.1469-185X.1998.tb00179.x)

3. McDonald DB, Potts WK. 1994 Cooperative display and relatedness among males in a lek-mating bird. *Science* **266**, 1030–1032. (doi:10.1126/science.7973654)

4. Oh KP, Badyaev A V. 2010 Structure of social networks in a passerine bird: Consequences for sexual selection and the evolution of mating strategies. *American Naturalist* **176**. (doi:10.1086/655216)

5. Sorato E, Gullett PR, Griffith SC, Russell AF. 2012 Effects of predation risk on foraging behaviour and group size: Adaptations in a social cooperative species. *Anim Behav* **84**, 823–834. (doi:10.1016/j.anbehav.2012.07.003)

6. Cresswell W. 1994 Flocking is an effective anti-predation strategy in redshanks, *Tringa totanus*. *Anim Behav*. **47**, 433–442. (doi:10.1006/anbe.1994.1057)

7. Furness RW, Birkhead TR. 1984 Seabird colony distributions suggest competition for food supplies during the breeding season. *Nature* **311**, 655–656. (doi:10.1038/311655a0)

8. Forero MG, Tella JL, Hobson KA, Bertellotti M, Blanco G. 2002 Conspecific food competition explains variability in colony size: A test in Magellanic penguins. *Ecology* **83**, 3466. (doi:10.2307/3072095)

9. Packer C, Pusey AE. 1982 Cooperation and competition within coalitions of male lions: kin selection or game theory? *Nature* **296**, 740–742. (doi:10.1038/296740a0)

10. Lamprecht J. 1978 The relationship between food competition and foraging group size in some larger carnivores. *Z Tierpsychol* **46**, 337–343. (doi:10.1111/j.1439-0310.1978.tb01455.x)

11. Bribiesca R, Herrera-Alsina L, Ruiz-Sanchez E, Sánchez-González LA, Schondube JE. 2019 Body mass as a supertrait linked to abundance and behavioral dominance in hummingbirds: A phylogenetic approach. *Ecol Evol* **9**, 1623–1637. (doi:10.1002/ece3.4785)

12. Miller ET, Bonter DN, Eldermire C, Freeman BG, Greig EI, Harmon LJ, Lisle C, Hochachka WM. 2017 Fighting over food unites the birds of North America in a continental dominance hierarchy. *Behavioral Ecology* **28**, 1454–1463. (doi:10.1093/beheco/arx108)

13. Šárová R, Špinka M, Stěhulová I, Ceacero F, Šimečková M, Kotrba R. 2013 Pay respect to the elders: Age, more than body mass, determines dominance in female beef cattle. *Anim Behav* **86**, 1315–1323. (doi:10.1016/j.anbehav.2013.10.002)

14. Zabel CJ, Glickman SE, Frank LG, Woodmansee KB, Keppel G. 1992 Coalition formation in a colony of prepubertal spotted hyenas. In *Coalitions and Alliances in Humans and other Animals* (eds AH Harcourt, FBM de Waal), pp. 113–136. Oxford: Oxford University Press.

15. Lahti K, Orell M, Rytkonen S, Koivula K. 1998 Time and food dependence in willow tit winter survival. *Ecology* **79**, 2904. (doi:10.2307/176525)

16. Jansson C, Ekman J, von Brömssen A, von Bromssen A. 1981 Winter mortality and food supply in tits Parus spp. *Oikos* **37**, 313. (doi:10.2307/3544122)

474   17.   Brittingham MC, Temple SA. 1988 Impacts of supplemental feeding on survival rates of
                black-capped chickadees. *Ecology* **69**, 581–589. (doi:10.2307/1941007)
476   18.   Plummer KE, Bearhop S, Leech DI, Chamberlain DE, Blount JD. 2013 Winter food
                provisioning reduces future breeding performance in a wild bird. *Sci Rep* **3**, 2002.
478   (doi:10.1038/srep02002)
        19.   Kalinoski R. 1975 Intra- and interspecific aggression in house finches and house sparrows.
480   *Condor* **77**, 375. (doi:10.2307/1366086)
        20.   Wojczulanis-Jakubas K, Kulpińska M, Minias P. 2015 Who bullies whom at a garden
482   feeder? Interspecific agonistic interactions of small passerines during a cold winter. *J
                Ethol* **33**, 159–163. (doi:10.1007/s10164-015-0424-x)
484   21.   Carpenter FL. 1987 Food abundance and territoriality: To defend or not to defend? *Integr
                Comp Biol* **27**, 387–399. (doi:10.1093/icb/27.2.387)
486   22.   Dias RI. 2006 Effects of position and flock size on vigilance and foraging behaviour of the
                scaled dove *Columbina squammata*. *Behavioural Processes* **73**, 248–252.
488   (doi:10.1016/j.beproc.2006.06.002)
        23.   Roberts G. 1996 Why individual vigilance declines as group size increases. *Anim Behav*
490   **51**, 1077–1086. (doi:10.1006/anbe.1996.0109)
        24.   Beauchamp G. 2008 What is the magnitude of the group-size effect on vigilance?
492   *Behavioral Ecology* **19**, 1361–1368. (doi:10.1093/beheco/arn096)
        25.   Bijleveld AI, van Gils JA, Jouta J, Piersma T. 2015 Benefits of foraging in small groups:
494   An experimental study on public information use in red knots Calidris canutus.
                Behavioural Processes **117**, 74–81. (doi:10.1016/j.beproc.2014.09.003)
496   26.   Valone TJ. 2007 From eavesdropping on performance to copying the behavior of others:
                A review of public information use. *Behav Ecol Sociobiol* **62**, 1–14. (doi:10.1007/s00265-
498   007-0439-6)
        27.   Gompper ME. 1996 Sociality and asociality in white-nosed coatis (*Nasua narica*):
500   foraging costs and benefits. *Behavioral Ecology* **7**, 254–263. (doi:10.1093/beheco/7.3.254)
        28.   Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015 Interspecific social networks
502   promote information transmission in wild songbirds. *Proceedings of the Royal Society B:
                Biological Sciences* **282**. (doi:10.1098/rspb.2014.2804)
504   29.   Beck KB, Farine DR, Kempenaers B. 2020 Winter associations predict social and extra-
                pair mating patterns in a wild songbird. *Proceedings of the Royal Society B: Biological
506   Sciences* **287**, 20192606. (doi:10.1098/rspb.2019.2606)
        30.   Firth JA, Sheldon BC. 2016 Social carry-over effects underpin trans-seasonally linked
508   structure in a wild bird population. *Ecol Lett* **19**, 1324–1332. (doi:10.1111/ele.12669)
        31.   Leighton GM, Lamour D, Malcolm K, Miller ET. 2022 Both morphological and
510   behavioral traits predict interspecific social dominance in birds. *J Ornithol*
                (doi:10.1007/s10336-022-02022-y)
512   32.   Krause J, Ruxton GD. 2002 *Living in Groups*. Oxford: Oxford University Press.
        33.   Grueter CC *et al.* 2020 Multilevel organisation of animal sociality. *Trends Ecol Evol* **35**,
514   834–847. (doi:10.1016/j.tree.2020.05.003)
        34.   Papageorgiou D, Farine DR. 2021 Multilevel societies in birds. *Trends Ecol Evol* **36**, 15–
516   17. (doi:10.1016/j.tree.2020.10.008)
        35.   Bonter DN, Cooper CB. 2012 Data validation in citizen science: a case study from Project
518   FeederWatch. *Front Ecol Environ* **10**, 305–307. (doi:10.1890/110273)

36. Balázs B, Mooney P, Nováková E, Bastin L, Jokar Arsanjani J. 2021 Data quality in
520   citizen science. In *The Science of Citizen Science*, pp. 139–157. Cham: Springer
International Publishing. (doi:10.1007/978-3-030-58278-4_8)

522 37. R Core Team. 2019 R: A language and environment for statistical computing.

38. Stoffel MA, Nakagawa S, Schielzeth H. 2017 rptR: Repeatability estimation and variance
524   decomposition by generalized linear mixed-effects models. *Methods Ecol Evol* **8**, 1639–
1644. (doi:10.1111/2041-210X.12797)

526 39. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–
884. (doi:10.1038/44766)

528 40. Billerman SM, Keeney BK, Rodewald PG, Schulenberg TS, editors. 2020 *Birds of the
world*. Ithaca, NY, USA: Cornell Laboratory of Ornithology.

530 41. Bates D, Mächler M, Bolker B, Walker S. 2014 Fitting linear mixed-effects models using
lme4. *J Stat Softw* **67**, 1–48. (doi:10.18637/jss.v067.i01)

532 42. Balph MH. 1979 Flock stability in relation to social dominance and agonistic behavior in
wintering Dark-eyed juncos. *Auk* **96**, 714–722. (doi:10.1093/auk/96.4.714)

534 43. Odum EP. 1942 Annual cycle of the Black-capped chickadee: 3. *Auk* **59**, 499–531.
(doi:10.2307/4079461)

536 44. Stouffer PC, Caccamise DF. 1991 Roosting and diurnal movements of radio-tagged
American crows. *Wilson Bulletin* **103**, 387–400.

538 45. Bürkner P-C. 2017 brms: An R package for Bayesian multilevel models using Stan. *J Stat
Softw* **80**, 1–28. (doi:10.18637/jss.v080.i01)

540 46. Duboscq J, Romano V, MacIntosh A, Sueur C. 2016 Social information transmission in
animals: Lessons from studies of diffusion. *Front Psychol*. **7**.
542   (doi:10.3389/fpsyg.2016.01147)

47. Chapman MR, Kramer DL. 1996 Guarded resources: The effect of intruder number on the
544   tactics and success of defenders and intruders. *Anim Behav* **52**, 83–94.
(doi:10.1006/anbe.1996.0154)

546 48. Webster MM, Ward AJW, Hart PJB. 2007 Boldness is influenced by social context in
threespine sticklebacks (Gasterosteus aculeatus). *Behaviour* **144**, 351–371.
548   (doi:10.1163/156853907780425721)

49. Humfeld SC, Marshall VT, Bee MA. 2009 Context-dependent plasticity of aggressive
550   signalling in a dynamic social environment. *Anim Behav* **78**, 915–924.
(doi:10.1016/j.anbehav.2009.06.028)

552 50. Ashton BJ, Ridley AR, Edwards EK, Thornton A. 2018 Cognitive performance is linked
to group size and affects fitness in Australian magpies. *Nature* **554**, 364–367.
554   (doi:10.1038/nature25503)

51. Galton F. 1907 Vox Populi. *Nature* **75**, 450–451. (doi:10.1038/075450a0)

556 52. Symons PEK. 1968 Increase in aggression and in strength of the social hierarchy among
juvenile Atlantic salmon deprived of food. *J. Fish. Res. Board Can.* **25**, 2387–2401.
558   (doi:10.1139/f68-207)

53. Collie J, Granela O, Brown EB, Keene AC. 2020 Aggression is induced by resource
560   limitation in the monarch caterpillar. *iScience* **23**, 101791.
(doi:10.1016/j.isci.2020.101791)

562 54. Sridhar H *et al.* 2012 Positive relationships between association strength and phenotypic
similarity characterize the assembly of mixed-species bird flocks worldwide. *Am Nat* **180**,
564   777–790. (doi:10.1086/668012)

55. Metcalfe NB, Furness RW. 1987 Aggression in shorebirds in relation to flock density and composition. *Ibis* **129**, 553–563. (doi:10.1111/j.1474-919X.1987.tb08243.x)

56. Sasvári L. 1992 Great tits benefit from feeding in mixed-species flocks: a field experiment. *Anim Behav* **43**, 289–296. (doi:10.1016/S0003-3472(05)80224-6)

57. Hino T. 2000 Intraspecific differences in benefits from feeding in mixed-species flocks. *J Avian Biol* **31**, 441–446. (doi:10.1034/j.1600-048X.2000.310402.x)

58. Kao AB, Hund AK, Santos FP, Young JG, Bhat D, Garland J, Oomen RA, McCreery HF. 2020 Changes in group size during resource shifts reveal drivers of sociality across the tree of life. *bioRxiv* (doi:10.1101/2020.03.17.994343)

59. Tibbetts EA, Pardo-Sanchez J, Weise C. 2022 The establishment and maintenance of dominance hierarchies. *Philos Trans R Soc Lond B Biol Sci* **377**, 20200450. (doi:10.1098/rstb.2020.0450)

60. Myers JP. 1983 Space, time and the pattern of individual associations in a group-living species: Sanderlings have no friends. *Behav Ecol Sociobiol* **12**, 129–134. (doi:10.1007/BF00343203)

61. Conklin JR, Colwell MA. 2008 Individual associations in a wintering shorebird population: Do Dunlin have friends? *J Field Ornithol* **79**, 32–40. (doi:10.1111/j.1557-9263.2008.00143.x)

62. Markham AC, Alberts SC, Altmann J. 2012 Intergroup conflict: Ecological predictors of winning and consequences of defeat in a wild primate population. *Anim Behav* **84**, 399–403. (doi:10.1016/j.anbehav.2012.05.009)

Figure 1

Figure 2

Figure 3