NO BRANCH LEFT BEHIND: TRACKING TERRESTRIAL BIODIVERSITY FROM A PHYLOGENETIC COMPLETENESS PERSPECTIVE

Jesús N. Pinto-Ledezma[1]*, Sandra Díaz [2,3] , Benjamin S. Halpern[4,5], Colin Khoury[6,7],

Jeannine Cavender-Bares[1]*

[1]Department of Ecology, Evolution and Behavior, University of Minnesota, 1479 Gortner Ave, Saint Paul, MN, 55108, USA

[2]Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto Multidisciplinario de Biología Vegetal (IMBIV), CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina

[3]Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba, Argentina

[4]National Center for Ecological Analysis and Synthesis, University of California, Santa Barbara, CA, 93101, USA

[5]Bren School of Environmental Science and Management, University of California, Santa Barbara, CA, 93106, USA

[6]San Diego Botanic Garden, 230 Quail Gardens Dr., Encinitas, CA 92024, USA

[7]International Center for Tropical Agriculture (CIAT), Km 17, Recta Cali-Palmira, Apartado Aéreo 6713, 763537, Cali, Colombia

*Corresponding Author: jpintole@umn.edu, cavender@umn.edu

*Equal contribution and shared first authorship

Running Header: No branch left behind

Type of article: Concepts & Questions

24   **Number of words:** Abstract (153), Main text (3717)

25   **Number of references:** 39

26   **Number of figures:** 3

27   **Number of tables:** 1

Abstract

Biodiversity, as we see it today, ultimately is the outcome of millions of years of evolution; however, biodiversity in its multiple dimensions is changing rapidly due to increasing human domination of Earth. Here, we present the "phylogenetic completeness" (PC) a concept and methodology that intends to safeguard Earth's evolutionary heritage by maintaining all branches of the tree of life. We performed a global evaluation of the PC approach using data from five major terrestrial clades and compared the results to an approach in which species are conserved or lost randomly. We demonstrate that under PC, for a given number of species extinctions, it is possible to maximize the protection of evolutionary innovations in every clade. The PC approach is flexible and can be used to conduct a phylogenetic audit of biodiversity under different conservation scenarios. The PC approach complements existing conservation efforts and is linked to the post-2020 Convention of Biodiversity targets.

Introduction

Over more than 3.5 billion years of life on Earth, evolution has generated and honed a vast array of innovations represented by the diversity of genomes and forms across the tree of life. Contemporary species collectively represent the genetic assets that contribute to the functioning of the current biosphere, and these functions in turn serve as the foundation of nature's contributions to people (NCP) (Díaz *et al.* 2019). Put another way, species embody evolutionary innovations that represent complex and unique approaches to life on Earth. These innovations not only support current NCP, but are also necessary for future benefits to humanity, including those not yet discovered.

Phylogenetic trees depict the hierarchy of life in which species are nested in larger and larger clades, each descended from a more distant common ancestor (Figure 1). They provide information on the breadth and variation of innovations evolution has generated and can be used to inform approaches to species conservation with the goal of minimizing extinction of evolutionary innovations (Faith 2002; Mace *et al.* 2003; Diniz-Filho *et al.* 2013; Larkin *et al.* 2016). Close relatives typically have a high proportion of shared genetics since they arose from a common ancestor at some point in the comparatively recent past, and thus share many of the same innovations.

A wide range of phylogenetic metrics exist that capture diversity across the tree of life (Tucker *et al.* 2017) and are relevant to discerning how much variation is captured under different conservation scenarios. Most evolutionary-based metrics of diversity are applied to community or regional phylogenies and represent different amounts of shared ancestry between taxa (sometimes accounting for other factors, such as probability of extinction). The canonical

example is phylogenetic diversity (PD, *sensu* (Faith 1992)), which, for a subset of species, is

defined as a sum of all branch lengths required to connect those species or the total amount of

evolutionary history contained by those species (Faith 1992, 2002). Since its introduction in

1992, PD has been used in ecological and conservation studies as a measure of biodiversity that

captures information about the number of accumulated features in the branches of the tree of life

(Scherson and Faith 2018).

In our era of rapid biodiversity loss (Tilman *et al.* 2017; IPBES 2019; Díaz *et al.* 2019)

keeping all species—the tips of the tree of life—is not realistic, and indeed is practically

unfeasible given recent extinction rates (Crozier 1997; Rounsevell *et al.* 2020). However, it may

still be possible to conserve most of the branches of the tree of life, assuming it is possible to

meaningfully define those branches (Figure 1). This concept is the principle behind phylogenetic

completeness, a concept and methodology we propose here that aims to preserve Earth's

evolutionary heritage (*sensu* Mooers *et al.* 2005) by maintaining all branches across the tree of

life. The approach aims to facilitate a phylogenetic audit of biodiversity. The central goal is to

track all of life's variation that has arisen over the course of evolution, and still remains, in both

early-diverged lineages and recently-derived lineages in each of the major living clades and

across the entire tree of life. Phylogenetic completeness (PC) and phylogenetic diversity (PD)

both aim to maintain as many branches as possible from the tree of life, recognizing it is not

possible to retain all species. But given the hierarchical nature of phylogenetic trees, how a

lineage is circumscribed—or what constitutes a "branch" that should be protected—can

influence the outcome (Vane-Wright *et al.* 1991; Diniz-Filho *et al.* 2013; Scherson and Faith

2018). PC differs from PD in that it slices across the tree at a given point in evolutionary history

and defines a set of branches based on this cut off (Figure 1), using a phylogenetic accounting

framework to identify and audit all distinctive branches in the tree of life at local, regional or

global scales, rather than only maximizing phylogenetic breadth.

Here we conducted a series of analyses to explore the implications for conservation

depending on the depth in the tree of life where a "branch" is defined. The goal was to develop

and apply a framework for conservation of species that minimizes hemorrhaging of Earth's

evolutionary assets, given a fixed level of species loss that is assumed to be unavoidable. We

then compared the loss of PD under conservation scenarios in which the set of species targeted

for conservation was based on an "informed" phylogenetic completeness approach or in which

species conserved or lost to extinction occurred randomly.

Methods

Phylogenetic completeness (PC) approach

We developed an approach in which a phylogeny is iteratively sliced at different periods of time

(T_N) until a specified finish time (T_F). For example, if a phylogeny is sliced every $T_N = 2$ million

years until $T_F = 50$ million years, a total of 25 slice points is obtained (see also Figure 1). These

slice points are then used to drop all but one terminal tips—or operational taxonomic units

(OTU)—from the phylogeny; with this approach, we ensure that at least one descendant OTU of

each lineage at a specific time (T_N) is kept. In other words, by keeping at least one OTU from

each lineage in the tree of life, we aim to maximize the preservation of the deepest evolutionary

history. At each slice point (T_N) we additionally calculated the number of species (T_{SP}) and

phylogenetic diversity (T_{PD}) as the simple sum of branch lengths at the specific slice point (T_N).

Empirical assessment

The empirical assessment focused on five major terrestrial clades (seed plants, amphibians,

squamates, birds, and mammals). Description of the data and software used can be found in the

WebPanel 1.

We tested the reliability of the PC approach by slicing each phylogeny every $T_N = 100$

Ky until $T_F = 100$ My and calculated the T_{SP} and T_{PD} at every slice point. These metrics were

then used to identify change points in the phylogenetic diversity over the 100 My time period.

Change points were evaluated using Bayesian Multiple Changing Points (MCP) regressions. The

first changing point plus its credible intervals (CIs) identified by the MCP analysis were used as

cutoff thresholds to estimate the number and identities of OTUs to be kept. This procedure

allowed us to identify different change points or cutoff thresholds in the phylogenetic diversity

over 100 My for each clade separately and consequently prevent us establishing a fixed arbitrary

cutoff threshold (e.g., setting a changing point at 2 My as cutoff threshold) for all clades.

We compared the diversity in each clade for a PC conservation scenario in which species

were managed to maintain all phylogenetic branches to random losses (RAND). In other words,

we removed OTUs at random until the identified cutoff threshold for each clade separately. This

procedure was repeated 1000 times, and at each step the T_{SP} and T_{PD} were estimated.

Finally, using the OTUs identities from both the PC and the random loss scenarios we

mapped the phylogenetic diversity of seed plants and terrestrial vertebrates globally. These maps

were used to estimate the difference (ΔPD) between the observed PD (PD_{OBS}) and the expected

PD (PD_{EXP}) under either the phylogenetic completeness (PD_{PC}) and the random loss approach

(PD_{RAND}).

134
$$\Delta PD = \frac{PD_{EXP} - PD_{OBS}}{PD_{OBS}} \; x \; 100$$

These maps represent the proportional difference between the observed (PD_{OBS}) and

expected (PD_{PC} or PD_{RAND}) phylogenetic diversity, where negative values suggest that a grid cell

will lose a proportion of its PD according to a specific conservation scenario, e.g., under the PC

scenario. Note that the lower CI from the Bayesian MCP regressions were used as variable cutoff

thresholds for each clade for mapping purposes.

Results

Phylogenetic completeness

Bayesian MCP models revealed variable cutoff thresholds for each clade (WebFigure 1;

WebTable 1). Based on these thresholds, losses of species ranging from 1.34% to 18.11% are

estimated to occur in each of the major clades (seed plants, amphibians, squamates, birds, and

mammals) (WebTable 1) while still safeguarding between 97.27% to 99.97% of the phylogenetic

diversity—i.e., of the evolutionary history of each clade (WebTable 1; Figure 2). If the lower

credible interval (LCI or the 2.5% quantile of the posterior distribution) of our Bayesian model

estimates were used to define the phylogenetic branches to be conserved (Figure 2; WebTable 1),

a higher number of species and branches in the tree of life would be safeguarded (Table 1).

These analyses demonstrate that if conservation efforts are focused on maintaining

defined branches of the tree of life it is possible to maximize the accumulated evolutionary

innovations that are safeguarded across all clades even when individual species go extinct.

Figure 2 shows the comparison between the estimates of PD under both phylogenetic

completeness (PC) and random loss (RAND) scenarios. Under PC, a higher number of branches

(Table 1) and greater evolutionary history in each clade is preserved for a given number of

species extinctions (Figure 2; WebFigure 2).

Spatial patterns of ΔPD under PC and RAND scenarios (Figure 3; WebFigure 3), show

how conservation informed by PC safeguards a greater proportion of evolutionary history even

with the same number of species extinctions (WebFigure 2). For example, for seed plants in

tropical regions across the world, conservation informed by PC resulted in PD loss below 10%,

whereas the Rand scenario resulted in 10-20 % of PD loss. Extinction patterns of terrestrial

biodiversity under PC and Rand scenarios at the biome level (WebFigure 4) also show greater

preservation of accumulated evolutionary innovations when conservation is targeted to maintain

branches of the tree of life. Nevertheless, we find that Tundra and Taiga biomes are susceptible

to high losses in PD, especially for seed plants, amphibians, and mammals, even under PC

scenarios. Indeed, further statistical analyses revealed small to no evidence in favor of the

phylogenetic completeness over random loss (evidence ratio < 5) in these biomes (WebTable 2).

In contrast, tropical biomes (for both forest and grasslands) show limited losses in PD for the

same threshold values used to define branches, as in Tundra and Taiga biomes (WebFigure 4;

WebTable 2). These results indicate that for a given number of species extinctions, tropical

biomes will lose fewer branches of the tree of life and are thus less susceptible to loss of

evolutionary history.

Discussion

Our "phylogenetic completeness" framework for informing biodiversity conservation focuses on

maintaining the accumulated evolutionary innovations across the tree of life with the intent of

leaving no branch behind. We introduce a rigorous approach for defining branches across clades

of terrestrial organisms to ascertain where in the tree of life there is high evolutionary

redundancy and where a single species may represent an entire branch. In doing so, we establish

phylogenetic branches as units of conservation priority rather than specific species or OTUs. By

defining these branches and the species contained within them, the phylogenetic completeness

approach provides critical information on branches at risk of extinction where there is low

redundancy as well as flexibility in which species could be targeted for conservation in cases of

high redundancy (Mooers *et al.* 2005).

The approach is particularly useful in developing priorities for *ex situ*, as well as for *in*

situ in protected areas, by tracking which branches of the tree of life are currently safeguarded

and by identifying the branches that are at highest risk—those not currently protected or with the

least amount of their range protected. The Kunming-Montreal Global Biodiversity Framework,

just adopted by the Convention on Biological Diversity aims to reduce the extinction rate and

risk of all species by tenfold by 2050 (Goal A) (CBD 2022), but does not explicitly include the

tree of life. Yet calls by the international scientific community to set ambitious goals for

biodiversity recognize that not all extinctions have equal consequences and that phylogenetic

dimensions of biodiversity need to be recognized among the criteria for implementation (Díaz *et*

al. 2020). The PC approach is consistent with the overall the CBD agenda because, at a given

overall extinction rate, it can help identify what species or clades will represent disproportionally

large or small losses of evolutionary history. As well as having self-evident intrinsic value,

evolutionary history is considered a major basis of "maintenance of options", one of the major

categories of nature's contributions to people recognized by the intergovernmental biodiversity

science-policy interface (IPBES 2019).

Increasing human domination of Earth and its ecosystems is rapidly changing

biodiversity patterns and negatively impacting the biosphere's capacity to provide essential

contributions to humanity (Tilman *et al.* 2017; Díaz *et al.* 2019). Safeguarding all remaining

biodiversity, although desirable, is unrealistic on the basis of virtually all projections (Pimm *et*

al. 2014; Tilman *et al.* 2017; Scherson and Faith 2018); the footprint of humanity is currently too

large to completely avoid further extinctions. Scientists have recognized the challenge of

developing logical conservation solutions given the complexity of stakeholders, managers, and

indirect actors as a 'wicked problem' that has no straightforward solution (Vane-Wright *et al.*

1991; Crozier 1997; Mace 2014). Focusing on the conservation of evolutionary history has been

hailed as an integrative way to safeguard most of the biodiversity and its functions (Faith 1992;

Mooers 2007; Mishler *et al.* 2014; Larkin *et al.* 2016). For example, a recent study by Molina-

Venegas and collaborators (Molina-Venegas *et al.* 2021) (Molina-Venegas *et al.* 2021) found

strong evidence that plant evolutionary history is tightly linked to multiple plant use categories

and therefore to human well-being (Molina-Venegas *et al.* 2021). These findings, among others,

support the idea that conserving evolutionary history is critical for future human well-being

(Forest *et al.* 2007; Molina-Venegas *et al.* 2021).

We demonstrate that under the PC framework, the loss of evolutionary history—

measured as the total branch lengths in the tree of life—can be minimized (Figure 2, 3). Our

results show that the loss of evolutionary heritage for five major terrestrial clades is lower

compared to phylogenetically random extinctions (Figure 2, WebFigure 2). In addition, although

current species extinctions are non-random (Purvis *et al.* 2000), evidence suggests that the

impact of random extinctions erases more evolutionary history on clades with imbalanced

phylogenetic trees (Heard and Mooers 2002) as the ones assessed in this study.

By assessing the evolutionary history loss at biome level (WebFigure 4) we found that

PC does not outperform the conservation scenario of random extinction in biomes at high

latitudes (e.g., Tundra and Taiga), in particular for the seed plants, amphibians, and squamates

(WebTable 2). At the scale of the biome, the loss of evolutionary history under the PC

framework seems to have acted selectively—extinctions are concentrated in specific taxa—

causing those depauperate lineages in these biomes to become even more depauperate. Indeed, at

high latitudes an individual species frequently represents an entire phylogenetic branch while at

low latitudes (tropics), a branch is likely to contain many species. This pattern is largely the

consequence of more recent divergence times and higher speciation rates in the tropics.

However, the spatial scale (grain size) must also be considered; for example, one hectare in a

tropical forest can hold ~650 tree species, which is more than all tree species that occur at high

latitudes (Coley and Kursar 2014). Despite this high diversity, tropical forests are usually hyper-

dominated by a fraction of species (~1.4% of about 16,000 tree species estimated for the

Amazonian Forest are considered as hyper-dominant) that are specialists to their habitats and

have large geographical ranges (ter Steege *et al.* 2013). The less abundant species or poorly

known species with small geographical ranges are potentially threatened.

Although it is beyond the scope of this article to conduct, PC audits at local and regional

scales could help to identify which species may be prioritized to prevent losing branches of the

tree of life at local and regional scales. To demonstrate scaling our approach to local sites, we

assessed a plant community-level phylogeny from the Cedar Creek Ecosystem Science Reserve

(Pinto-Ledezma *et al.* 2020). This analysis reveals which taxa are unique and which taxa present

evolutionary redundancy, i.e., red branches in WebFigure 5. Moreover, using plant community

composition (N = 987 plots of 40 x 40 m) from the National Ecological Observatory Network

(NEON) we found that under scenarios where extinctions are predicted to occur all branches are

preserved according to our phylogenetic accounting framework. Our results demonstrate that

phylogenetic diversity is nearly equivalent to that calculated from observed data where no

extinctions are assumed to have occurred (WebFigure 6). This additional analysis reveals that,

despite variation in the number of species lost (WebFigure 6A), focusing on branches as units of

conservation aids in maintaining the evolutionary history of local communities.

Multiple approaches have been proposed to assess changes in biodiversity focusing on

"hotspot" areas (spatial prioritization), or taxa (taxonomic prioritization) for conservation

purposes (Margules and Sarkar 2007). These approaches rely on the use of metrics that capture

different dimensions of biodiversity, e.g., metrics that capture evolutionary changes among a set

of taxa (Rodrigues *et al.* 2005; Margules and Sarkar 2007) or the variation in form and function

of taxa within communities (Díaz and Cabido 2001; Petchey and Gaston 2006). Despite their

usefulness for assessing the state and the fate of biodiversity, most of these metrics, if not all, are

sensitive to information (in)completeness. Missing information can result in misleading metric

calculations and inappropriate interpretations of spatial or taxonomic comparisons (Isaac 2004;

Rodrigues *et al.* 2011; Diniz-Filho *et al.* 2013; Weedop *et al.* 2019). The PC framework

introduced here represents a complementary approach to counting numbers of species or

comparing levels of phylogenetic diversity to assess biodiversity under alternative conservation

scenarios. It provides an accounting framework that prioritizes conservation of branches of the

tree of life rather than individual taxa (Table 1; Figure 3). It also allows the identification of

areas susceptible to high losses of evolutionary heritage (Figure 3; WebFigure 3), which can be

used as baseline information for spatial prioritization, providing a broader context for local

decision making (Mace *et al.* 2003; Mishler *et al.* 2014; Chaplin-Kramer *et al.* 2022; Silvestro *et*

al. 2022).

Moreover, given that the PC framework focuses on the branches of the tree of life, in the

context of spatial prioritization, our approach my help existing spatial phylogenetic

approaches—e.g., categorical analysis of neo- and paleo-endemism (Mishler *et al.* 2014;

Thornhill *et al.* 2016), phylogenetic endemism (Rosauer *et al.* 2009)—to deal with the issue of

missing data. To illustrate this point, if we consider protecting at least one descendant taxon from

a specific node in the phylogeny, this taxon contains information—genetic, traits, functions—

that captures most of the evolutionary history of the lineage, i.e., the evolutionary history of the

taxon plus its common ancestor (see Figure 1; WebFigure 5). If species within the branch have

not yet been identified or are not readily observed, the branch itself is still preserved, with the

caveat that phylogenetic information remains imperfect—i.e., Darwinian shortfall in biodiversity

and conservation (Diniz-Filho *et al.* 2013). By focusing on branches of the tree of life as an

additional conservation criterion beyond those prioritized in other ways, it is possible to reduce

the impact of missing data and taxonomic inflation on evolutionary diversity estimations and

conservation planning (Isaac 2004; Rodrigues *et al.* 2011; Diniz-Filho *et al.* 2013; Allen and

Mishler 2022).

Finally, we are aware that our empirical evaluation was based on incomplete data, but it

is the best phylogenetic and geographical data available to date. Although our estimations are

robust, more research may be needed to fully understand the potential of the PC framework in

biodiversity conservation. In addition, this study is limited to terrestrial biodiversity; thus,

evaluations of aquatic biodiversity are required. Future studies focusing on aquatic systems or

terrestrial biodiversity not considered here (e.g., Hexapoda) may help to refine the PC's potential

in biodiversity assessment and conservation. These studies should also investigate the relative

impact of branch selection, i.e., "the agony of choice," or which branch should be prioritized and

its conservation implications. In other words, where should we focus our efforts to minimize the

loss of evolutionary heritage. Lastly, we emphasize that phylogenetic completeness is not a

metric of phylogenetic diversity but an accounting framework that identifies targets (unique

branches) for eco-evolutionary or conservation questions in a phylogenetic context. Once the

unique branches are identified, existing metrics can be used (WebFigure 6) to estimate the

impact of lineage loss on biodiversity estimations, conservation scenarios or ecological research.

Moving forward

Phylogenetic completeness provides a means to account for phylogenetic branches and facilitates

the safeguarding of evolutionary heritage, adding to but not replacing other priorities for

conservation. For example, the approach will highlight when a single species is the lone member

of a branch, increasing its conservation priority. The spatial scale of evaluation is important to

consider, and application at local scales in coordination with regional scales is critical. It would

be dangerous to consider that maintaining only one species in a branch is adequate at large

scales, mainly because some species were extirpated locally and currently occur in a fraction of

their original geographic distribution (e.g., Hyacinth macaw, African lion). However, an

informed and coordinated conservation may allow individual protected areas to focus on

conservation of species that maintain all branches locally, emphasizing particular species within

a branch based on other conservation criteria. An auditing procedure with appropriate tools

would enable coordination at regional scales to ensure that most or all species in a particular

branch are being managed. By focusing on branches of the tree of life as an additional

conservation criterion beyond those prioritized in other ways, we will be less likely to overlook

the importance of management for taxa that alone represent an evolutionary branch (Mace *et al.*

2003; Mace 2014; Allen and Mishler 2022).

Conclusion

In our era of rapid global change and rapid biodiversity loss, conservation efforts must reckon with the reality that we will not succeed in saving all species on Earth. We outline an approach for conservation, which we call phylogenetic completeness, that focuses on saving the accumulated innovations that have evolved in Earth's biota by counting individual branches in the tree of life as units of conservation priority. The approach benefits from detailed information of the tree of life that is only now sufficiently resolved to be applicable to all of life on Earth. The approach complements other conservation efforts and is directly relevant to the targets of the Kunming-Montreal Global Biodiversity Framework established by the Convention on Biological Diversity.

Acknowledgements

Support to this project was provided by the National Science Foundation (NSF) through the Macrosystems Biology & NEON-Enabled Science program (DEB-2017843). Further support was provided by the NSF BII ASCEND (DBI-2021898). The work is part of the FITBITs working group hosted by the National Center for Ecological Analysis and Synthesis.

Statement of Authorship

All authors contributed intellectually to the manuscript. J.C.-B. and J.N.P.-L. conceived of and framed the manuscript. J.N.P.-L. performed all the statistical and spatial analyses with input from J.C.-B. All authors edited the manuscript.

Open Research

Data Availability Statement

All data used in this manuscript are publicly available. Main sources are provided in the

WebPanel 1. R functions and examples for data analyses are publicly available at

https://github.com/jesusNPL/FITBITs.

References

Allen MF and Mishler BD. 2022. A Phylogenetic Approach to Conservation: Biodiversity and

349   Ecosystem Functioning for a Changing Globe. In: Swartz B, Mishler BD (Eds).

350   Speciesism in Biology and Culture. Cham: Springer International Publishing.

351   CBD. 2022. Kunming-Montreal Global biodiversity framework: Draft decision submitted by the

352   President. Convention on Biological Diversity CBD/COP/15/L.25. Conference of the

353   Parties to the Convention on Biological Diversity. Fifteenth meeting – Part II Montreal,

354   Canada, 7-19 December 2022.

355   Chaplin-Kramer R, Brauman KA, Cavender-Bares J, *et al.* 2022. Conservation needs to integrate

356   knowledge across scales. *Nat Ecol Evol* **6**: 118–9.

357   Coley PD and Kursar TA. 2014. On Tropical Forests and Their Pests. *Science* **343**: 35–6.

358   Crozier RH. 1997. Preserving the Information Content of Species: Genetic Diversity, Phylogeny,

359   and Conservation Worth. *Annu Rev Ecol Syst* **28**: 243–68.

360   Díaz S, Settele J, Brondízio ES, *et al.* 2019. Pervasive human-driven decline of life on Earth

361   points to the need for transformative change. *Science* **366**: eaax3100.

Díaz S, Zafra-Calvo N, Purvis A, *et al.* 2020. Set ambitious goals for biodiversity and

sustainability. *Science* **370**: 411–3.

Diniz-Filho JAF, Loyola RD, Raia P, *et al.* 2013. Darwinian shortfalls in biodiversity

conservation. *Trends in Ecology & Evolution* **28**: 689–95.

Díaz S and Cabido M. 2001. Vive la différence: plant functional diversity matters to ecosystem

processes. *Trends in Ecology & Evolution* **16**: 646–55.

Faith DP. 1992. Conservation evaluation and phylogenetic diversity. *Biological Conservation*

61: 1–10.

Faith DP. 2002. Quantifying Biodiversity: a Phylogenetic Perspective. *Conservation Biology* **16**:

248–52.

Forest F, Grenyer R, Rouget M, *et al.* 2007. Preserving the evolutionary potential of floras in

biodiversity hotspots. *Nature* **445**: 757–60.

Heard SB and Mooers AØ. 2002. Signatures of Random and Selective Mass Extinctions in

Phylogenetic Tree Balance. *Systematic Biology* **51**: 889–97.

IPBES. 2019. Summary for policymakers of the global assessment report on biodiversity and

ecosystem services (pp. XIV-LXI). Díaz S, Settele J, Brondízio ES, Ngo HT, Guèze M,

Agard J, Arneth A, Balvanera P, Brauman KA, Butchart SHM, Chan KMA, Garibaldi

LA, Ichii K, Liu J, Subramanian SM, Midgley GF, Miloslavich P, Molnár Z, Obura D,

Pfaff A, Polasky S, Purvis A, Razzaque J, Reyers B, Roy Chowdhury R, Shin YJ,

Visseren-Hamakers IJ, Willis KJ, and Zayas CN (eds). IPBES secretariat, Bonn,

Germany.

Isaac N. 2004. Taxonomic inflation: its influence on macroecology and conservation. *Trends in*

Ecology & Evolution **19**: 464–9.

Larkin DJ, Jacobi SK, Hipp AL, and Kramer AT. 2016. Keeping All the PIECES:

Phylogenetically Informed Ex Situ Conservation of Endangered Species. *PLoS ONE* **11**:

e0156973.

Mace GM. 2014. Whose conservation? *Science* **345**: 1558–60.

Mace GM, Gittleman JL, and Purvis A. 2003. Preserving the Tree of Life. *Science* **300**: 1707–9.

Margules CR and Sarkar S. 2007. Systematic conservation planning. Cambridge ; New York:

Cambridge University Press.

Mishler BD, Knerr N, González-Orozco CE, *et al.* 2014. Phylogenetic measures of biodiversity

and neo- and paleo-endemism in Australian Acacia. *Nat Commun* **5**: 4473.

Molina-Venegas R, Rodríguez MÁ, Pardo-de-Santayana M, *et al.* 2021. Maximum levels of

global phylogenetic diversity efficiently capture plant services for humankind. *Nat Ecol*

Evol **5**: 583–8.

Mooers AØ. 2007. The diversity of biodiversity. *Nature* **445**: 717–8.

Mooers AØ, Heard SB, and Chrostowski E. 2005. Evolutionary heritage as a metric for

conservation. In: Purvis A, Gittleman JL, Brooks T (Eds). Phylogeny and Conservation.

Cambridge University Press.

Petchey OL and Gaston KJ. 2006. Functional diversity: back to basics and looking forward. *Ecol*

Letters **9**: 741–58.

Pimm SL, Jenkins CN, Abell R, *et al.* 2014. The biodiversity of species and their rates of

extinction, distribution, and protection. *Science* **344**: 1246752.

Pinto-Ledezma JN, Villalobos F, Reich PB, *et al.* 2020. Testing Darwin's naturalization

conundrum based on taxonomic, phylogenetic, and functional dimensions of vascular

plants. *Ecol Monogr* **90**.

Purvis A, Agapow P-M, Gittleman JL, and Mace GM. 2000. Nonrandom Extinction and the Loss

of Evolutionary History. *Science* **288**: 328–30.

Rodrigues ASL, Brooks TM, and Gaston KJ. 2005. Integrating phylogenetic diversity in the

selection of priority areas for conservation: does it make a difference? In: Purvis A,

Gittleman JL, Brooks T (Eds). Phylogeny and Conservation. Cambridge University

Press.

Rodrigues ASL, Grenyer R, Baillie JEM, *et al.* 2011. Complete, accurate, mammalian

phylogenies aid conservation planning, but not much. *Phil Trans R Soc B* **366**: 2652–60.

Rosauer D, Laffan SW, Crisp MD, *et al.* 2009. Phylogenetic endemism: a new approach for

identifying geographical concentrations of evolutionary history. *Molecular Ecology* **18**:

4061–72.

Rounsevell MDA, Harfoot M, Harrison PA, *et al.* 2020. A biodiversity target based on species

extinctions. *Science* **368**: 1193–5.

Scherson RA and Faith DP (Eds). 2018. Phylogenetic Diversity: Applications and Challenges in
Biodiversity Science. Cham: Springer International Publishing.

Silvestro D, Goria S, Sterner T, and Antonelli A. 2022. Improving biodiversity protection
through artificial intelligence. *Nat Sustain* **5**: 415–24.

Steege H ter, Pitman NCA, Sabatier D, *et al.* 2013. Hyperdominance in the Amazonian Tree
Flora. *Science* **342**: 1243092.

Thornhill AH, Mishler BD, Knerr NJ, *et al.* 2016. Continental-scale spatial phylogenetics of
Australian angiosperms provides insights into ecology, evolution and conservation.
Journal of Biogeography **43**: 2085–98.

Tilman D, Clark M, Williams DR, *et al.* 2017. Future threats to biodiversity and pathways to
their prevention. *Nature* **546**: 73–81.

Tucker CM, Cadotte MW, Carvalho SB, *et al.* 2017. A guide to phylogenetic metrics for
conservation, community ecology and macroecology: A guide to phylogenetic metrics for
ecology. *Biol Rev* **92**: 698–715.

Vane-Wright RI, Humphries CJ, and Williams PH. 1991. What to protect?—Systematics and the
agony of choice. *Biological Conservation* **55**: 235–54.

Weedop KB, Mooers AØ, Tucker CM, and Pearse WD. 2019. The effect of phylogenetic
uncertainty and imputation on EDGE Scores. *Anim Conserv* **22**: 527–36.

Tables

Table 1. Number of branches (NB) conserved under the scenarios of phylogenetic completeness (PC) and random loss (Rand) conservation scenarios. The number of branches conserved under the PC approach is higher than the random loss scenario for the same number of species except for birds and mammals. Percentages of NB under PC and Rand scenarios are displayed within brackets. Number of branches within a clade assuming no extinction is displayed for reference. The threshold number of species under either the PC or Rand scenarios corresponds to the number of species at the lower CI identified using the Bayesian MCP regressions.

Clade	Observed N species	Threshold N species	Threshold (My)	NB all species	NB-PC	NB-Rand
Seed plants	353185	312540	1.76	438863	397974 [90.68%]	391851 [89.29%]
Amphibians	7238	5927	4.43	14474	12614 [87.15%]	11852 [81.88%]
Squamata	9755	8141	2.61	19508	17186 [88.10%]	16280 [83.45%]
Birds	9993	9859	0.33	19984	19716 [98.66%]	19716 [98.66%]
Mammals	5911	5802	0.31	11820	11602 [98.16%]	11602 [98.16%]

Note: the observed number of species (second column) corresponds to the number of species sampled in each phylogenetic tree and might not represent each clade's true number of species.

Figure legends

Figure 1. Schematic diagram of the phylogenetic relationships of species within a lineage of

organisms, showing the hierarchical nesting of hypothetical species. The number of species that

need to be conserved in order to prevent the extinction of any branches in the tree depends on the

depth in the phylogeny where we define the branch. Dashed lines represent different ages (T_N),

or depths in the phylogeny, used to define branches—the corresponding number of species

preserved (T_{SP}), if one species per branch is conserved, is colored the same as the dashed line.

For example, compare purple circles (fewer branches and species with a deep slice at T_{N3}) to red

ones (more branches and species with a more recent slice at T_{N1}). The deeper we "slice" the

phylogeny, the fewer the species need to be saved in order to preserve all branches of the

phylogeny at that slice. Black circles represent species at T_{N0}, i.e., the current species, assuming

no extinction. Open circles filled with "x" indicate species that went extinct at a specific T_N.

Figure 2. Difference between the remaining phylogenetic diversity under phylogenetic

completeness (red) and random loss (blue) scenarios for five major terrestrial clades (seed plants,

amphibians, squamates, birds, and mammals). The X-axis was log-transformed for plotting

purposes. In all cases, the phylogenetic diversity is higher for the phylogenetic completeness

approach.

Figure 3. Mapped phylogenetic diversity for seed plants (A-C) and birds (B-D) globally under

scenarios of phylogenetic completeness and random loss. Legends indicate the proportional loss

of phylogenetic diversity (ΔPD). Blue tones indicate that more branches of the tree of life have

been preserved and red tones that more branches have been lost. Comparing the two scenarios,

globally more branches of the tree of life are conserved in the phylogenetic completeness

scenario for the same number of species extinctions. Maps for amphibians, squamates, and

mammals can be found in the supplementary material (WebFigure 3).

Figures

484   Figure 1

488   Figure 2

492   Figure 3

494