1

No branch left behind: tracking terrestrial biodiversity from a phylogenetic completeness

perspective

Jesús N. Pinto-Ledezma[1]*, Sandra Díaz [2,3] , Benjamin S. Halpern[4,5], Colin Khoury[6,7],

Jeannine Cavender-Bares[1]*

*Equal contribution and shared first authorship

1 Department of Ecology, Evolution and Behavior, University of Minnesota, 1479 Gortner Ave,

Saint Paul, MN, 55108, USA

2 Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto Multidisciplinario de

Biología Vegetal (IMBIV), CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina

3 Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba,

Argentina

4 National Center for Ecological Analysis and Synthesis, University of California, Santa

Barbara, CA, 93101, USA

5 Bren School of Environmental Science and Management, University of California, Santa

Barbara, CA, 93106, USA

6 San Diego Botanic Garden, 230 Quail Gardens Dr., Encinitas, CA 92024, USA

7 International Center for Tropical Agriculture (CIAT), Km 17, Recta Cali-Palmira, Apartado

Aéreo 6713, 763537, Cali, Colombia

Abstract

Biodiversity, as we see it today, ultimately is the outcome of millions of years of evolution;

however, biodiversity in its multiple dimensions is changing rapidly due to increasing human

domination of Earth. Here, we present the "phylogenetic completeness" (PC) a concept and

methodology that intends to safeguard Earth's evolutionary assets that have arisen across the tree

of life. We performed a global evaluation of the PC approach using data from five major

terrestrial clades and compared the results to an approach in which species are conserved or lost

randomly. We demonstrate that under PC, it is possible to maximize the protection of greater

evolutionary assets of each clade for a given number of species extinctions. The PC approach is

flexible and can be used to assess biodiversity under different conservation scenarios. The PC

approach complements existing conservation efforts and is linked to the post-2020 Convention of

Biodiversity targets.

Introduction

Over more than 3.5 billion years of life on Earth, evolution has generated and honed a vast array

of innovations represented by the diversity of forms and genomes across the tree of life.

Contemporary species collectively represent the genetic assets that contribute to the functioning

of the current biosphere, and these functions in turn serve as the foundation of Nature's

Contributions to People (NCP) (Díaz *et al.* 2019). Put another way, species embody evolutionary

innovations that represent complex and unique approaches to life on Earth. These innovations

not only support current NCP, but are also necessary for future benefits to humanity, including

those not yet discovered.

Phylogenetic trees depict the hierarchy of life in which species are nested in larger and

larger clades, each descended from a more distant common ancestor (Figure 1). They provide

information on the breadth and variation of innovations evolution has generated and can be used

to inform approaches to species conservation with the goal of minimizing extinction of

evolutionary innovations (Faith 2002; Larkin *et al.* 2016). Close relatives typically have a high

proportion of shared genetics since they arose from a common ancestor at some point in the

comparatively recent past, and thus share many of the same innovations.

A wide range of phylogenetic metrics exist that capture diversity across the tree of life

(Tucker *et al.* 2017) and are relevant to discerning how much variation is captured under

different conservation scenarios. Most metrics of phylogenetic diversity are applied to

phylogenetic trees and represent a sum over phylogenetic branch lengths (possibly after

augmentation to account for other factors, such as probability of extinction). The canonical

example is phylogenetic diversity (PD, *sensu* (Faith 1992)), which, for a subset of species, is

defined as a sum of all branch lengths required to connect those species (Faith 1992, 2002).

In our era of rapid biodiversity loss (Tilman *et al.* 2017; Díaz *et al.* 2019) (Tilman *et al.*

2017; Díaz *et al.* 2019), keeping all species—the tips of the tree of life—is not realistic, and

indeed is impossible given recent extinction rates (Rounsevell *et al.* 2020). However, it may still

be possible to conserve all branches of the tree of life, depending on how deeply we define the

branches. This concept is the principle behind phylogenetic completeness, a concept and

methodology we propose here that aims to preserve Earth's evolutionary assets that have arisen

across the entire tree of life. Phylogenetic completeness (PC) and phylogenetic diversity (PD)

both aim to maintain as many branches as possible from the tree of life, recognizing it is not

possible to retain all species. But given the hierarchical nature of phylogenetic trees, how a

lineage is circumscribed—or what constitutes a "branch" that should be protected—can

influence the outcome. PC differs from PD in that it slices across the tree at a given point in

evolutionary history and defines a set of branches based on this cut off (Figure 1), using an

accounting framework, rather than merely maximizing phylogenetic breadth.

Here we conducted a series of analyses to explore the implications for conservation

depending on the depth in the tree of life where a "branch" is defined. The goal was to develop

and apply a framework for conservation of species that minimizes hemorrhaging of Earth's

evolutionary assets, given a fixed level of species loss that is assumed to be unavoidable. We

then compared the loss of PD under conservation scenarios in which the set of species targeted

for conservation was based on an "informed" phylogenetic completeness approach or in which

species conserved or lost to extinction occurred randomly.

Methods

Phylogenetic completeness (PC) approach

We developed an approach in which a phylogeny is iteratively sliced at different periods of time

(T_N) until a specified finish time (T_F). For example, if a phylogeny is sliced every $T_N = 2$ million

years until $T_F = 50$ million years, a total of 25 slice points is obtained (see also Figure 1). These

slice points are then used to drop all but one terminal tips—or operational taxonomic units

(OTU)—from the phylogeny; with this approach, we ensure that at least one OTU of each

lineage at a specific time (T_N) is kept. In other words, by keeping at least one OTU from each

lineage in the tree of life, we aim to maximize the preservation of the deepest evolutionary

history. At each slice point (T_N) we additionally calculated the number of species (T_{SP}) and

phylogenetic diversity (T_{PD}) as the simple sum of branch lengths at the specific slice point (T_N).

Empirical assessment

The empirical assessment was focused on five major terrestrial clades (seed plants, amphibians,

squamates, birds, and mammals). Description of the data can be found in the WebPanel 1.

We tested the reliability of the PC approach by slicing each phylogeny every $T_N = 100$

Ky until $T_F = 100$ My and calculated the T_{SP} and T_{PD} at every slice point. These metrics were

then used to identify change points in the phylogenetic diversity over the 100 My time period.

Change points were evaluated using Bayesian Multiple Changing Points (MCP) regressions. The

first changing point plus its credible intervals (CIs) identified by the MCP analysis were used as

cutoff thresholds to estimate the number and identities of OTUs to be kept. This procedure

allowed us to identify different change points or cutoff thresholds in the phylogenetic diversity

over 100 My for each clade separately and consequently prevent us establishing a fixed arbitrary

cutoff threshold (e.g., setting a changing point at 2 My as cutoff threshold) for all clades.

We compared the diversity in each clade for a PC conservation scenario, in which species

were managed to maintain all phylogenetic branches to random losses (RDM). In other words,

we removed OTUs at random until the identified cutoff threshold for each clade separately. This

procedure was repeated 1000 times, and at each step the T_{SP} and T_{PD} were estimated.

Finally, using the OTUs identities from both the PC and the random loss scenarios we

mapped the phylogenetic diversity of seed plants and terrestrial vertebrates globally. These maps

were used to estimate the difference (ΔPD) between the observed PD (PD_{OBS}) and the expected

PD (PD_{EXP}) under either the phylogenetic completeness (PD_{PC}) and the random loss approach

(PD_{RDM}).

$$\Delta PD = \frac{PD_{EXP} - PD_{OBS}}{PD_{OBS}} \; x \; 100$$

These maps represent the proportional difference between the observed (PD_{OBS}) and

expected (PD_{PC} or PD_{RDM}) phylogenetic diversity, where negative values suggest that a grid cell

will lose a proportion of its PD according to a specific conservation scenario, e.g., under the PC

scenario. Note that the lower CI from the Bayesian MCP regressions were used as variable cutoff

thresholds for each clade for mapping purposes.

Protected areas assessment

We were also interested in evaluating the role of protected areas (PAs) in protecting the

evolutionary history of terrestrial biodiversity. To do so, we overlayed the PAs with the

vertebrate and seed plants PAMs to obtain the presence-absence of species within the PAs

(PAM_{PA}) globally (WebPanel 1). These PAM_{PA} were then used to estimate the number of

species and phylogenetic diversity within each PA. Using this information, we estimate and map

the ΔPD between the observed PD (PD_{OBS}) and the PD within PAs (PD_{PA}). The resulting maps

show how much evolutionary history are currently protected within the PAs.

Results

Phylogenetic completeness

Bayesian MCP models revealed variable cutoff thresholds for each clade (Figure 2; WebTable

1). Based on these thresholds, losses of species ranging from 1.34% to 18.11% are estimated to

occur in each of the major clades (seed plants, amphibians, squamates, birds, and mammals)

(WebTable 1) while still safeguarding between 97.27% to 99.97% of the phylogenetic

diversity—i.e., of the evolutionary history of each clade (WebTable 1; Figure 2). If the lower

credible interval (LCI) of our Bayesian model estimates were used to define the phylogenetic

branches to be conserved (Figure 2; WebTable 1), a higher number of species and branches in

the tree of life would be safeguarded (Table 1).

These analyses demonstrate that if conservation efforts are focused on maintaining

defined branches of the tree of life it is possible to maximize the accumulated evolutionary

innovations that are safeguarded across all clades even when individual species go extinct.

Figure 3 shows the comparison between the estimates of PD under both phylogenetic

completeness (PC) and random loss (RDM) scenarios. Under PC, a higher number of branches

(Table 1) and greater evolutionary history in each clade is preserved for a given number of

species extinctions (Figure 3).

Spatial patterns of ΔPD under PC and RDM scenarios (Figure 4; WebFigure 1), show

how conservation informed by PC safeguards a greater proportion of evolutionary history even

with the same number of species extinctions. For example, for seed plants in tropical regions

across the world, conservation informed by PC resulted in PD loss below 10%, whereas the

RDM scenario resulted in 10-20 % of PD loss. Extinction patterns of terrestrial biodiversity

under PC and RDM scenarios at the biome level (WebFigure 2) also show greater preservation

of accumulated evolutionary innovations when conservation is targeted to maintain branches of

the tree of life. Nevertheless, we find that Tundra and Taiga biomes are susceptible to high losses

in PD, especially for seed plants, birds, and mammals, even under PC scenarios. In contrast,

tropical biomes (for both forest and grasslands) show limited losses in PD for the same threshold

values used to define branches, as in Tundra and Taiga biomes (WebFigure 2). These results

indicate that for the a given number of species extinctions, tropical biomes will lose fewer

branches of the tree of life and are thus less susceptible to loss of evolutionary history.

Protected areas assessment

The currently implemented PAs across the world cover an area of approximately 26,775,820

km—~18% of the land surface. Collectively, ~97% of the species in our dataset overlap their

ranges with the system of PAs globally (Table 2). Although the proportion of likely protected

species is high, a large proportion of the Earth's land surface, and thus most of the ranges of

most of these species, is not protected (Figure 4; WebFigure 3) and is threatened by land use

change and other human activities.

Discussion

Our "phylogenetic completeness" framework for informing biodiversity conservation focuses on

maintaining the accumulated evolutionary innovations across the tree of life with the intent of

leaving no branch behind. We introduce a rigorous approach for defining branches across clades

of terrestrial organisms to ascertain where in the tree of life there is high evolutionary

redundancy and where a single species may represent an entire branch. In doing so, we establish

phylogenetic branches as units of conservation priority. By defining these branches and the

species contained within them, the phylogenetic completeness approach provides critical

information on branches at risk of extinction where there is low redundancy as well as flexibility

in which species can targeted for conservation in cases of high redundancy. The approach is

particularly useful in developing conservation priorities in relation to protected areas by tracking

which branches of the tree of life are currently safeguarded and by identifying the branches that

are at highest risk—those not currently protected or with the least amount of their range

protected. This approach is consistent with the Convention on Biodiversity draft Milestone A.2,

that *the increase in the extinction rate is halted or reversed, and the extinction risk is reduced by*

at least 10 per cent, with a decrease in the proportion of species that are threatened, and the

abundance and distribution of populations of species is enhanced or at least maintained. Yet it

also captures elements of diversity not delineated by the CBD by accounting for the breadth of

evolved variation in plant and vertebrate-animal life.

Increasing human domination of Earth and its ecosystems is rapidly changing

biodiversity patterns and negatively impacting the capacity of ecosystems to provide goods and

services to humanity (Tilman *et al.* 2017; Díaz *et al.* 2019). Safeguarding all remaining

biodiversity, although desirable, is unrealistic on the basis of virtually all projections (Pimm *et*

al. 2014; Urban 2015; Tilman *et al.* 2017); the footprint of humanity is currently too large to

completely avoid further extinctions. Scientists have recognized the challenge of developing

logical conservation solutions given the complexity of stakeholders, managers, and indirect

actors as a 'wicked problem' that has no straightforward solution (Vane-Wright *et al.* 1991;

DeFries and Nagendra 2017). Focusing on the conservation of evolutionary history has been

hailed as an integrative way to safeguard most of the biodiversity and its functions (Faith 1992;

Mooers 2007). For example, a recent study by Molina-Venegas and collaborators (Molina-

Venegas *et al.* 2021) (Molina-Venegas *et al.* 2021) found strong evidence that plant evolutionary

history is tightly linked to multiple plant use categories and therefore to human well-being

(Molina-Venegas *et al.* 2021). These findings, among others, support the idea that conserving

evolutionary history is critical for future human well-being (Forest *et al.* 2007; Molina-Venegas

et al. 2021).

Multiple approaches have been proposed to assess changes in biodiversity focusing on

"hotspot" areas (spatial prioritization), or taxa (taxonomic prioritization) for conservation

purposes (Margules and Sarkar 2007). These approaches rely on the use of metrics that capture

different dimensions of biodiversity, e.g., metrics that capture evolutionary changes among a set

of taxa (Margules and Sarkar 2007) or the variation in form and function of taxa within

communities (Díaz and Cabido 2001; Petchey and Gaston 2006). Despite their usefulness for

assessing the state and the fate of biodiversity, most of these metrics, if not all, are sensitive to

information completeness. Missing information can result in misleading metric calculations and

inappropriate interpretations of spatial or taxonomic comparisons (e.g., Weedop *et al.* 2019). The

PC framework introduced here represents a complementary approach to counting numbers of

species or comparing levels of phylogenetic diversity to assess biodiversity under alternative

conservation scenarios. It provides an accounting framework that prioritizes conservation of

branches of the tree of life rather than individual species (Table 1; Figure 2). It also allows the

identification of areas susceptible to high losses of evolutionary assets (Figure 4; WebFigure 1),

which can be used as baseline information for spatial prioritization, providing a broader context

for local decision making (Chaplin-Kramer *et al.* 2022). Moreover, in the context of spatial

prioritization, the PC framework may be less susceptible to missing data given that it focuses on

the branches of the tree of life. To illustrate this point, if we consider protecting at least one

descendant taxon from a specific node in the phylogeny, this taxon contains genetic information

that captures most of the evolutionary history of the entire branch (see Figure 1). If species

within the branch have not yet been identified or are not readily observed, the branch itself is still

preserved, with the caveat that phylogenetic information remains imperfect.

Substantial efforts have been invested to prevent the extinction of biodiversity. In

particular, the establishment, expansion, and enhancement of protected areas (PAs) have

received considerable attention due to their critical role in protecting Earth's biodiversity and

preventing the erosion of its benefits to humans (Naughton-Treves *et al.* 2005; Watson *et al.*

2014). Despite important progress in establishing terrestrial PAs around the world (Table 2),

several critical areas are still unprotected, leaving many branches of the tree of life vulnerable to

extinction (Figure 4; WebFigure 3). Note that in our evaluation we used the species presence

within current PAs to estimate which species are less vulnerable to extinction for the purposes of

deciphering which branches of the tree of life are most at risk. We thus assume that species not

present in PAs are the most vulnerable to extinction and in greatest need of conservation or

assistance.

At high latitudes (tundra) an individual species frequently represents an entire

phylogenetic branch while at low latitudes (tropics), a branch is likely to contain many species.

This pattern is largely the consequence of more recent divergence times and higher speciation

rates in the tropics. However, the spatial scale (grain size) must also be considered, for example,

one hectare in a tropical forest can hold ~650 tree species more than all tree species that occur at

high latitudes (Coley and Kursar 2014). Despite this high diversity, tropical forests are usually

hyperdominated by a fraction of species (~1.4% of about 16,000 tree species estimated for the

Amazonian Forest are considered as hyperdominant) that are specialists to their habitats and

have large geographical ranges (ter Steege *et al.* 2013). The less abundant species or poorly

known species with small geographical ranges are potentially threatened. Although it is beyond

the scope of this article, PC evaluations at local and regional scales could help to identify which

species may be prioritized to prevent losing branches of the tree of life at local and regional

scales.

Furthermore, the combined effects of climate and land-use changes are likely to limit the

role of PAs in safeguarding biodiversity (Hoffmann *et al.* 2019; Asamoah *et al.* 2021). About

one-quarter of the PAs globally are projected to suffer rapid climate change, with small PAs

being the most impacted (Hoffmann *et al.* 2019; Asamoah *et al.* 2021). Advancing projections of

combined limits to species tolerance to climate change and migration capability without

assistance could be oriented towards understanding which branches of the tree of life currently

protected are most likely to be lost. Doing so, could help inform costly efforts to establish

corridors for movement within human-dominated landscapes (Gibson *et al.* 2011; Wintle *et al.*

2019) and assisted migration plans to target the protection of species that maintain all the

branches of the tree of life.

Conclusion

In our era of rapid global change and rapid biodiversity loss, conservation efforts must reckon

with the reality that we will not succeed in saving all species on Earth. We outline an approach

for conservation, which we call phylogenetic completeness, that focuses on saving the

accumulated innovations that have evolved in Earth's biota by counting individual branches in

the tree of life as units of conservation priority. The approach benefits from detailed information

of the tree of life that is only now sufficiently resolved to be applicable to all of life on Earth.

The approach complements other conservation efforts and is directly relevant to the targets of the

post-2020 Convention of Biodiversity.

Acknowledgements

Support to this project was provided by the National Science Foundation (NSF) through the

Macrosystems Biology & NEON-Enabled Science program (DEB 2017843). Further support

was provided by the NSF BII ASCEND (DBI-2021898). The work is part of the FITBITs

working group hosted by the National Center for Ecological Analysis and Synthesis.

Statement of Authorship

All authors contributed intellectually to the manuscript. J.C.-B. and J.N.P.-L. conceived of and

framed the manuscript. J.N.P.-L. performed all the statistical and spatial analyses with input from

J.C.-B. All authors edited the manuscript.

Open Research

Data Availability Statement

All data used in this manuscript are publicly available. Main sources are provided in the

WebPanel 1. R functions and examples for data analyses are publicly available at

https://github.com/jesusNPL/FITBITs.

References

Asamoah EF, Beaumont LJ, and Maina JM. 2021. Climate and land-use changes reduce the

benefits of terrestrial protected areas. *Nat Clim Chang* **11**: 1105–10.

Chaplin-Kramer R, Brauman KA, Cavender-Bares J, *et al.* 2022. Conservation needs to integrate

knowledge across scales. *Nat Ecol Evol* **6**: 118–9.

Coley PD and Kursar TA. 2014. On Tropical Forests and Their Pests. *Science* **343**: 35–6.

DeFries R and Nagendra H. 2017. Ecosystem management as a wicked problem. *Science* **356**:

265–70.

Díaz S, Settele J, Brondízio ES, *et al.* 2019. Pervasive human-driven decline of life on Earth

points to the need for transformative change. *Science* **366**: eaax3100.

Díaz S and Cabido M. 2001. Vive la différence: plant functional diversity matters to ecosystem

processes. *Trends in Ecology & Evolution* **16**: 646–55.

Faith DP. 1992. Conservation evaluation and phylogenetic diversity. *Biological Conservation*

61: 1–10.

Faith DP. 2002. Quantifying Biodiversity: a Phylogenetic Perspective. *Conservation Biology* **16**:

248–52.

Forest F, Grenyer R, Rouget M, *et al.* 2007. Preserving the evolutionary potential of floras in

biodiversity hotspots. *Nature* **445**: 757–60.

Gibson L, Lee TM, Koh LP, *et al.* 2011. Primary forests are irreplaceable for sustaining tropical

biodiversity. *Nature* **478**: 378–81.

Hoffmann S, Irl SDH, and Beierkuhnlein C. 2019. Predicted climate shifts within terrestrial

protected areas worldwide. *Nat Commun* **10**: 4787.

Larkin DJ, Jacobi SK, Hipp AL, and Kramer AT. 2016. Keeping All the PIECES:

Phylogenetically Informed Ex Situ Conservation of Endangered Species (N Cellinese,

Ed). *PLoS ONE* **11**: e0156973.

Margules CR and Sarkar S. 2007. Systematic conservation planning. Cambridge ; New York:

Cambridge University Press.

Molina-Venegas R, Rodríguez MÁ, Pardo-de-Santayana M, *et al.* 2021. Maximum levels of

global phylogenetic diversity efficiently capture plant services for humankind. *Nat Ecol*

Evol **5**: 583–8.

Mooers AØ. 2007. The diversity of biodiversity. *Nature* **445**: 717–8.

Naughton-Treves L, Holland MB, and Brandon K. 2005. THE ROLE OF PROTECTED AREAS

IN CONSERVING BIODIVERSITY AND SUSTAINING LOCAL LIVELIHOODS.

Annu Rev Environ Resour **30**: 219–52.

Petchey OL and Gaston KJ. 2006. Functional diversity: back to basics and looking forward. *Ecol*

Letters **9**: 741–58.

Pimm SL, Jenkins CN, Abell R, *et al.* 2014. The biodiversity of species and their rates of

extinction, distribution, and protection. *Science* **344**: 1246752.

Rounsevell MDA, Harfoot M, Harrison PA, *et al.* 2020. A biodiversity target based on species

extinctions. *Science* **368**: 1193–5.

Steege H ter, Pitman NCA, Sabatier D, *et al.* 2013. Hyperdominance in the Amazonian Tree

Flora. *Science* **342**: 1243092.

Tilman D, Clark M, Williams DR, *et al.* 2017. Future threats to biodiversity and pathways to

their prevention. *Nature* **546**: 73–81.

Tucker CM, Cadotte MW, Carvalho SB, *et al.* 2017. A guide to phylogenetic metrics for

conservation, community ecology and macroecology: A guide to phylogenetic metrics for

ecology. *Biol Rev* **92**: 698–715.

Urban MC. 2015. Accelerating extinction risk from climate change. *Science* **348**: 571–3.

Vane-Wright RI, Humphries CJ, and Williams PH. 1991. What to protect?—Systematics and the

agony of choice. *Biological Conservation* **55**: 235–54.

Watson JEM, Dudley N, Segan DB, and Hockings M. 2014. The performance and potential of

protected areas. *Nature* **515**: 67–73.

Weedop KB, Mooers AØ, Tucker CM, and Pearse WD. 2019. The effect of phylogenetic

uncertainty and imputation on EDGE Scores. *Anim Conserv* **22**: 527–36.

Wintle BA, Kujala H, Whitehead A, *et al.* 2019. Global synthesis of conservation studies reveals

the importance of small habitat patches for biodiversity. *Proc Natl Acad Sci USA* **116**:

909–14.

Tables

Table 1. Number of branches (NB) conserved under the scenarios of phylogenetic completeness

(PC) and random loss (RDM) conservation scenarios. The number of branches conserved under

the phylogenetic completeness approach is higher than the random loss scenario for the same

number of species except for birds and mammals. Percentages of NB under PC and RDM

scenarios are displayed within brackets. Number of branches for all species within a clade

assuming no extinction is displayed for reference. The number of species under PC/RDM

correspond to the number of species at the minimum threshold (lower credible interval)

identified using the Bayesian MCP regressions. Note that the observed number of species

(second column) corresponds to the number of species sampled in each phylogenetic tree and

might not represent each clade's true number of species.

Clade	Observed N species	PC/RDM N species	Threshold (My)	NB all species	NB-PC	NB-RDM
Seed plants	353185	312540	1.76	438863	397974 [90.68%]	391851 [89.29%]
Amphibians	7238	5927	4.43	14474	12614 [87.15%]	11852 [81.88%]
Squamata	9755	8141	2.61	19508	17186 [88.10%]	16280 [83.45%]
Birds	9993	9859	0.33	19984	19716 [98.66%]	19716 [98.66%]
Mammals	5911	5802	0.31	11820	11602 [98.16%]	11602 [98.16%]

Table 2. Summary of terrestrial biodiversity within protected areas. Table summarizes the

number and percentage of species currently protected in protected areas. Our estimations of

%protected and % of loss are based on the number of species available in the geographical

ranges and phylogenetic datasets (see methods). Note that the number of seed plant species with

available geographic ranges is 207,146, so the estimations of %protected and % of loss may be

underestimated. The observed number of species (second column) corresponds to the number of

species sampled in each phylogenetic tree and might not represent each clade's true number of

species.

Clade	Observed N species	WPA N species	%Protected	%Loss
Seed plants	353,185	191,830	92.61	7.39
Amphibians	7,238	6,630	91.60	8.40
Squamata	9,755	8,872	90.95	9.05
Birds	9,993	9,710	97.17	2.83
Mammals	5,911	5,661	95.77	4.23

Figure legends

Figure 1. Schematic diagram of the phylogenetic relationships of species within a lineage of organisms, showing the hierarchical nesting of hypothetical species. The number of species that need to be conserved in order to prevent the extinction of any branches in the tree depends on the depth in the phylogeny we define the branch. Dashed lines represent different ages (T_N) or depths in the phylogeny used to define branches—the corresponding number of species preserved (T_{SP}), if one species per branch is conserved, is colored the same as the dashed line. For example, purple circles (fewer branches and species with a deep slice at T_{N3}) to red ones (more branches and species with a more recent slice at T_{N1}). The deeper we "slice" the phylogeny, the fewer the species need to be saved in order to preserve all branches of the phylogeny at that slice. Black circles represent species at T_{N0}, i.e., the current species, assuming no extinction. Open circles filled with "x" indicate species that went extinct at a specific T_N.

Figure 2. Changes in the phylogenetic diversity and number of species over 100 million years for terrestrial biodiversity, including seed plants (A), woody plants (B), amphibians (C), squamates (D), birds (E), and mammals (F). Black dashed vertical lines represent the first changing point identified by the Bayesian Multiple Changing Points (MCP) regressions. Black dotted vertical lines represent the 95 credible intervals. Red long-dashed horizontal lines represent the observed number of species for each clade. Red dashed horizontal lines are the number of species expected under the first changing point. Red dotted horizontal lines represent the expected number of species under the lower bound (or credible interval) of the first changing point. See WebTable 1

for a comprehensive numerical summary of the changing points. Silhouettes obtained from

Phylopic (http://phylopic.org).

Figure 3. Difference between the remaining phylogenetic diversity under phylogenetic

completeness (red) and random loss (green) scenarios. X-axis was log-transformed for plotting

purposes. In all cases, the phylogenetic diversity is higher for the phylogenetic completeness

approach.

Figure 4. Mapped phylogenetic diversity for seed plants (A-B) and birds (D-E) globally under

scenarios of phylogenetic completeness and random loss. Legends indicate the proportional loss

of phylogenetic diversity (ΔPD). Blue tones indicate that more branches of the tree of life have

been preserved and red tones that more branches have been lost. Comparing the two scenarios,

globally more branches of the tree of life are conserved in the phylogenetic completeness

scenario for the same number of vertebrate species extinctions. Bottom panels (C and F) show

the differences in phylogenetic diversity (ΔPD) in 1º grid cells between the species currently

estimated to occur in those cells and the fraction of those estimated to occur in protected areas

globally. Red colors represent greater ΔPD, meaning more branches of the tree of life are not

currently protected. Continental China is shaded gray given that protected area information is

currently unavailable in the WDPA for this country. Maps for amphibians, squamates, and

mammals can be found in the supplementary material.

Figures

Figure 1

423   Figure 2

429   Figure 3

Phylogenetic diversity (Log)

Figure 4