Protection of high-frequency low-intensity pulsed electric fields and brain-derived neurotrophic factor for SH-SY5Y cells against hydrogen peroxide-induced cell damage

Guan-Bo Lin[1,2], Wei-Ting Chen[1,2], Yu-Yi Kuo[1,2], You-Ming Chen[2,3], Hsu-Hsiang Liu[2,3], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author

E-mail: cychao@phys.ntu.edu.tw (CYC)

Abstract

Neurodegenerative diseases have been a major threat to public health worldwide nowadays. In particular, Alzheimer's disease is the most common type causing dementia which has remained an incurable disease, despite improvement of the symptoms via some therapies. The pathogenic mechanism of Alzheimer's disease is deemed to be related to the aggregation of beta-amyloid peptide (Aβ), a major cause for amyloid plaques, resulting in neuronal death via the generation of reactive oxygen species (ROS). Scientists have regarded oxidative stress as a potential treatment target in the development of novel remedies. In the study, human neural cell line SH-SY5Y was used in probing the effect of combining non-invasive high-frequency low-intensity pulsed electric field (H-LIPEF) and brain-derived neurotrophic factor (BDNF) in protection against hydrogen peroxide (H_2O_2)-induced neuron damage. Our result finds that the combination approach has intensified the neuroprotective effect significantly, perhaps due to H-LIPEF and BDNF synergistically increasing the expression level of the phosphorylated epidermal growth factor receptor (p-EGFR), which induces the survival-related mitogen-activated protein kinases (MAPK) proteins. The study confirmed the activation of extracellular signal-regulated kinase (ERK), a member of MAPK family, and the downstream pro-survival and antioxidant proteins as the mechanism underlying neuron protection. These findings have highlighted the potential of H-LIPEF combining with BDNF in the treatment of neurodegenerative diseases. Furthermore, the application of BDNF-mimetic drugs and non-invasive H-LIPEF to patients is a promising approach worthy of further research, and this points to strategies and mechanisms for selecting drugs or compounds to cooperate with electric fields in the treatment of neurodegenerative disorders.

Introduction

In line with the aging of global population, neurodegenerative diseases (NDDs) have emerged as a mainstream medical issue, not only causing discomfort and even death for patients but also disrupting the life of their families. Among these brain diseases, Alzheimer's disease is the most common type and a major cause of dementia in aged people. Alzheimer's pathogenic mechanism is believed to be related to the aggregation of β-amyloid (Aβ), which leads to amyloid plaques, causing neuronal death via the generation of reactive oxygen species (ROS) [1]. Another well-known hypothesis is phosphorylation of tau protein, which forms neurofibrillary tangles in nerve cell, damaging nerve cells and leading to progressive brain degeneration and deterioration of cognitive function [2]. Although Alzheimer's disease has been discovered for many years, there has yet to be an effective treatment, except a few therapies to relieve its symptoms. Hence, there has been a global scramble for the development of effective Alzheimer's disease therapy.

Neurotrophins are potential therapeutic agents for treating neurodegenerative disorder of the central nervous system. Among these neurotrophins, brain-derived neurotrophic factor (BDNF) is the most abundant neurotrophin in the human brain, which has a remarkable capability to repair brain damage. With proven effect in promoting neuronal survival, synaptic plasticity and neurogenesis, BDNF can resist brain damage caused by oxidative inflammation-related factors [3]. Some studies have found that BDNF produces neuroprotective effects against ROS-induced cell death, even with the potential for treating NDDs [4-6]. However, it should be noted that reduction in BDNF concentration can cause symptoms such as forgetfulness and poor learning ability, and many studies have blamed insufficient BDNF contents for depression and even Alzheimer's disease [7-9]. In addition, there has been evidence

suggesting that Aβ can inhibit the proteolytic conversion of pro-BDNF to BDNF,

decreasing BDNF level [10]. Therefore, stimulation of BDNF secretion or use of

exogenous BDNF is a promising approach in the treatment of Alzheimer's disease.

Many studies have proposed treatments capable of increasing the production of

endogenous BDNF, thereby achieving nerves protection [11-13]. Meanwhile, the use

of exogenous BDNF is also a direct and effective approach in dealing with the

abnormal secretion of endogenous BDNF caused by the disease, as some studies have

confirmed the significant neuroprotection effect of exogenous BDNF [14-16].

However, a major problem is the short half-life of BDNF, which increases the

difficulty of many therapeutic applications [17]. Therefore, how to improve BDNF

efficiency will be an important issue.

In addition to traditional drug medication, physical therapy has gradually

emerged as a mainstream approach in many diseases, such as electric current

stimulation with proven value in treating muscle injury, wound, cancer, and neuronal

diseases [18-21]. Many studies have pointed out that external electric stimulation can

stimulate proteins to change their structure, affect their distribution, and even

influence relevant biochemical pathways [22-24]. On the other hand, it is worth

noting that epidermal growth factor receptor (EGFR) is a transmembrane protein

deemed to be relevant to cell survival pathways. Especially, EGFR signalling

abnormalities are believed to be associated with certain diseases, such as

atherosclerosis, Parkinson's disease, and Alzheimer's disease [25, 26]. Furthermore, it

has been reported that the conventional electric current stimulation can affect cell

migration via EGFR and enhance the expressions of downstream survival-related

proteins, such as mitogen-activated protein kinases (MAPK) and phosphatidylinositol

3-kinase (PI3K) protein families in some medical fields [27].

To date, some issues remain unsolved for the application of electric current

stimulation in many diseases. As the electric current stimulations are applied via contact or even invasive method, electric current would pass through tissues directly, causing the Joule effect [28]. It is known that invasive treatments are at higher risk and difficult in execution for some tissues, especially brain. Therefore, the development of non-invasive and non-contact electrotherapy is an inevitable trend. Recently, our team has developed a unique non-invasive and non-contact low-intensity pulsed electric field (LIPEF) stimulation for biomedical application. The pulsed electric field (PEF) can be considered as a composition of multiple sinusoidal subcomponents with different frequencies and intensities, which may interact with molecules, proteins, or organelles to induce specific bioeffect at the same time. Consequently, PEF stimulation can facilitate significantly the development of new therapies to meet different needs of different diseases, with great potential for applications in various medical fields. First, our team has demonstrated, for the first time, the capability of LIPEF to combine with epigallocatechin-3-gallate (EGCG) in inhibiting the pancreatic cancer cell line PANC-1 [29]. Second, low-dose curcumin, along with LIPEF, also has good anticancer effect, underscoring LIPEF's capability to work with various chemical molecules [30]. Third, ultrasound can be added to the combination treatment, further boosting the anticancer effect [31]. Moreover, our team has found another effective anticancer triple therapy, combining LIPEF, chlorogenic acid (CGA), and thermal cycling-hyperthermia technique [32]. Besides, LIPEF can join hands with fucoidan to modulate rho-associated protein kinase (ROCK) and the downstream Akt protein, thereby protecting motor neurons NSC-34 against damage by oxidative stress [33]. On the other hand, in human neuronal SH-SY5Y cells under hydrogen peroxide (H_2O_2) stress, we found that high-frequency LIPEF (H-LIPEF) can modulate and activate the extracellular signal-regulated kinase (ERK) pathway, thereby protecting neuronal cells against oxidative stress [34]. In

sum, in addition to safety, LIPEF (or H-LIPEF) alone or in combination with

chemical molecules and/or other physical stimuli has significant potential for a wide

range of applications.

In this paper, our team first combined the features of BDNF and H-LIPEF

stimulation to study the effect of combination treatment on neuroprotection against

the oxidative insult caused by H_2O_2 to SH-SY5Y human neural cells. The result

showed that the combination of BDNF and H-LIPEF significantly inhibited the

H_2O_2-induced cytotoxicity in SH-SY5Y cells. Examination of the underlying

mechanism also indicated that the addition of BDNF is beneficial via regulation of

ERK, a member of MAPK family proteins, a possible reason for the enhanced

neuroprotective effect of combination BDNF and H-LIPEF treatment. Moreover, the

combination treatment was found to significantly upregulate the expression of

phosphorylated EGFR (p-EGFR) protein. These findings suggest that the combination

of neurotrophic-related chemical molecules and H-LIPEF promises to be an effective

therapy for neurodegenerative diseases.

Materials and methods

Experimental setup for exposure of the cells to non-contact

H-LIPEF

The H-LIPEF stimulation device described in our previous study [34] was used for

exposure of the SY-SY5Y cells to electric field. In brief, the cells were placed

between two copper flat and parallel electrodes. The pulsed electric signal was

produced from a function generator and amplified by a voltage amplifier. In the study,

consecutive pulses with the electric field 10 V/cm, frequency 200 Hz, and pulse

duration 2 ms were applied across the electrodes for the treatment. Cells treated with

continuous exposure of non-contact H-LIPEF were kept in a humidified incubator of

5% CO_2 and 95% air at 37°C.

Cell culture

The SH-SY5Y cells were purchased from American Type Culture Collection (ATCC,

Manassas, VA, USA) and cultured in MEM/F-12 mixture (HyClone; Cytiva,

Marlborough, MA, USA) supplemented with 10% fetal bovine serum (FBS)

(HyClone; Cytiva), 100 units/ml penicillin (Gibco; Thermo Fisher Scientific, Inc.,

Waltham, MA, USA), 100 mg/ml streptomycin

(Gibco; Thermo Fisher Scientific, Inc.), 1 mM sodium pyruvate (Sigma-Aldrich;

Merck KGaA, Darmstadt, Germany), and 0.1 mM non-essential amino acids

(Gibco; Thermo Fisher Scientific, Inc.). Cells were maintained in a humidified

incubator composed of 5% CO_2 and 95% air at 37°C, and harvested with 0.05%

trypsin-0.5 mM EDTA solution (Gibco; Thermo Fisher Scientific, Inc.) for in vitro

experiments.

H-LIPEF and BDNF treatment

169   BDNF (Alomone Labs, Jerusalem, Israel) was dissolved in distilled water as a stock

170   solution and stored at -20°C. SH-SY5Y cells were pretreated with various

171   concentrations of BDNF and then subjected to H-LIPEF treatment for 4 h. Subsequently,

172   cells were challenged with H_2O_2 and put back again to the H-LIPEF stimulation device

173   for further treatment.

Cell viability assay

Viability of SH-SY5Y cells after treatments with or without H-LIPEF was assessed by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) assay. Briefly, the medium was replaced with MTT solution (0.5 mg/mL in SH-SY5Y culture medium) and incubated at 37°C for 4 h. During this period, the ability of mitochondrial dehydrogenases to reduce the MTT to formazan crystal is an indicator of cellular viability. Following MTT reaction, formazan dissolution was performed by using 10% sodium dodecyl sulfate (SDS) in 0.01M HCl. The optical density of each well was then determined at 570 nm subtracting the background at 690 nm using Multiskan GO spectrophotometer (Thermo Scientific, Hudson, NH, USA). The cell viability was calculated based on the intensity of formazan, and expressed as percentage of the non-treated control.

Western blot analysis

After the treatment of H_2O_2, cells were washed with phosphate buffered saline (PBS) and then lysed on ice for 30 m in lysis buffer (EMD Millipore, Billerica, MA, USA) containing fresh protease and phosphatase inhibitor cocktail (EMD Millipore). The lysates were centrifuged at 23,000 × g for 1 h at 4°C, and the supernatant was collected and used for the determination of protein concentration. Equal amounts of protein extracts were loaded in the 10% SDS-PAGE wells and transferred to polyvinylidene difluoride (PVDF) membranes (EMD Millipore). The blots were incubated overnight with primary antibodies at 4°C, followed by incubation with horseradish peroxidase-coupled secondary antibodies (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA, USA) at room temperature for 1 h. In this study, the specific primary antibodies against p-EGFR, phosphorylated ERK (p-ERK),

nuclear factor erythroid 2-related factor 2 (Nrf2), phosphorylated cAMP response

element binding protein (p-CREB), B-cell lymphoma 2 (Bcl-2), Bcl-2-associated X

protein (Bax) (Cell Signaling Technology, Inc., Danvers, MA, USA), and GAPDH

(GeneTex, Inc., Irvine, CA, USA) were used. All the antibodies were diluted at the

optimal concentration according to the manufacturer's instructions. Finally, protein

bands were visualized with an enhanced chemiluminescence substrate (Advansta, Inc.,

Menlo Park, CA, USA) and detected with the Amersham Imager 600 imaging system

(GE Healthcare Life Sciences, Chicago, IL, USA). The images were analyzed with

Image Lab software (Bio-Rad Laboratories, Inc., Hercules, CA, USA). For

normalization of proteins, GAPDH was used as an internal control.

ROS level detection

After the treatments, the SH-SY5Y cells were collected and washed with PBS. The

fluorescent dye dihydroethidium (DHE) (Sigma-Aldrich; Merck KGaA) was used to

detect the intracellular ROS levels. Cells were resuspended in PBS and incubated with

5 µM DHE dye for 20 min at 37°C in the dark. The fluorescence intensity was

measured by FACSCanto II system (BD Biosciences, San Jose, CA, USA) in the PE

channel. The ROS levels were expressed as mean fluorescence intensity for

comparison.

Statistical analysis

221 The results were presented as mean ± standard deviation and performed in triplicate.

222 Differences of statistical significance were determined by a one-way analysis of

223 variance (ANOVA), followed by Tukey's post-hoc test. P-value < 0.05 was

224 considered to indicate a statistically significant difference. Analyses were carried out

225   using OriginPro 2015 software (OriginLab).

Results

In vitro-applied H-LIPEF

To avoid the harmful effects caused by conduction current and Joule effect, the

culture dishes were placed in the non-invasive H-LIPEF device as described

previously [34]. The H-LIPEF stimulation was performed in a non-contact manner

with the apparatus consisting of two parallel conductive electrodes mounted on the

insulating acrylic frame (Fig 1A). Therefore, the H-LIPEF device can provide the

electric field stimulation in a non-invasive manner, so that the cells were not in

contact with the electrodes directly. The pulsed electric signal was from a function

generator and amplified by a voltage amplifier that made pulsed electric signal

tunable. The applied electric signal was characterized by a pulse train waveform (Fig

1B) with various field strengths and frequencies for different applications. In this

study, we used the electric field strength of 10 V/cm at 200 Hz, which was adopted in

our previous study to be the optimal H-LIPEF condition for neuroprotection on

SH-SY5Y cell line [34].

Fig 1. The experimental setup of H-LIPEF. (A) Image of the H-LIPEF stimulation

device. The SH-SY5Y cells were placed in the H-LIPEF stimulation device

constructed of two parallel electrodes. The green arrows shown here indicate the

schematic representation of non-invasive and non-contact H-LIPEF exposure on cells.

(B) Schematic representation of the pulse train waveform applied across the

electrodes.

Effect of the H-LIPEF and BDNF on H_2O_2-induced cytotoxicity in SH-SY5Y cells

Because oxidative stress plays a key role in the etiology of neurodegenerative diseases, reducing oxidative stress is often a top priority for neuroprotection. As shown in Fig 2A, administration of H_2O_2 on SH-SY5Y cells for 24 h decreased the cell viability in a concentration-dependent manner. Treatment of SH-SY5Y cells with 500 μM H_2O_2 for 24 h reduced cell viability to about 50% compared to the untreated control group. In order to understand whether the H-LIPEF combined with BDNF could attenuate the H_2O_2-induced neuron cell damage, we first applied various concentrations of BDNF (0 to 100 ng/ml) in SH-SY5Y cells and see their neuroprotection effect. As shown in Fig 2B, we found a slight protective effect of these concentrations of BDNF against H_2O_2-induced cell oxidation. On the other hand, in the combination treatment, SH-SY5Y cells were pretreated with H-LIPEF and BDNF for 4 h, and then exposed to 500 μM H_2O_2 in the continuous administration of H-LIPEF for another 24 h. In present work, we applied the most effective H-LIPEF parameters (200 Hz, 10 V/cm, and 2 ms pulse duration) previously used by our team for neuroprotection study [34] to human neural cells SH-SY5Y. As shown in Fig 2C, the cell viability decreased to about half of the control group after treating SH-SY5Y cells with 500 μM H_2O_2 for 24 h. When cells were treated by combination treatment with H-LIPEF and BDNF, the cell viability was significantly rescued. With the help of H-LIPEF stimulation, administration of 25 ng/ml and 50 ng/ml BDNF on SH-SY5Y cells increased cell viability to approximately 80% and 90%, respectively. Our results showed that, while BDNF alone had a slight protective effect on SH-SY5Y cells, the combination of H-LIPEF and BDNF further enhanced the neuroprotective effect significantly. Our goal is to amplify the therapeutic effect and reduce the dosage of drug via H-LIPEF

stimulation. Therefore, based on these data, the H-LIPEF (200 Hz, 10 V/cm, and 2 ms

pulse duration) parameters in combination with the concentration of 25 ng/ml BDNF

were chosen to be used in all the following experiments.

Fig 2. Effect of H-LIPEF and/or BDNF on H_2O_2-induced cytotoxicity in

SH-SY5Y cells. (A) The concentration response of SH-SY5Y cells treated with

different concentrations of H_2O_2 and the cell viability was measured by MTT assay at

24 h after the H_2O_2 treatment. (B) SH-SY5Y cells were pretreated with different

concentrations of BDNF for 4 h and challenged with 500 μM H_2O_2 for 24 h and then

the cell viability was measured by MTT assay. (C) SH-SY5Y cells administered by 25

and 50 ng/ml BDNF were subsequently treated with or without H-LIPEF for 4 h and

then challenged with 500 μM H_2O_2, respectively. The cell viability was also measured

by MTT assay at 24 h after the H_2O_2 treatment. Data represent the mean ± standard

deviation (n=3). The concentration response curve of SH-SY5Y cells was statistically

analyzed using one-way ANOVA with Tukey's post hoc test (**$P < 0.01$ and ***$P <$

0.001).

Combination of H-LIPEF and BDNF attenuates

H_2O_2-induced ROS generation

To the best of our knowledge, the damage of ROS accumulation to nerves is known to

be the main cause of many brain diseases, such as Alzheimer's disease and

Parkinson's disease. In view of this, we studied the effect of combination treatment

with H-LIPEF and BDNF on the regulation of ROS level by flow cytometry

measurements (Fig 3A). As shown in Fig 3B, the quantification results show that the

ROS level was increased to about 200% of the control level after exposure to 500 μM

H_2O_2 for 24 h. Compared to the H_2O_2 alone group, H-LIPEF treatment significantly

ameliorated the H_2O_2-induced increase in ROS level, while the combination of

H-LIPEF and BDNF further enhanced the effect. This result indicates that the

combination treatment (H-LIPEF+BDNF) is more effective in reducing the oxidative

stress of SH-SY5Y cells.

Fig 3. Effect of H-LIPEF alone or in combination with BDNF on H_2O_2-induced

ROS generation in SH-SY5Y cells. (A) ROS level was measured by flow cytometry

with DHE fluorescent dye. (B) Quantification of the ROS levels after H_2O_2,

H-LIPEF+H_2O_2, and H-LIPEF+BDNF+H_2O_2 treatment. Data represent the mean ±

standard deviation (n=4). Comparisons of the effect of H_2O_2, H-LIPEF+H_2O_2, and

H-LIPEF+BDNF+H_2O_2 on the ROS generation were analyzed using one-way

ANOVA with Tukey's post hoc test (**$P < 0.01$ and ***$P < 0.001$).

Role of the EGFR pathways in the neuroprotective effect of

the H-LIPEF and BDNF

BDNF is an important neurotrophin that regulates the survival of existing

neurons and promotes the growth and differentiation of new neurons. It has been

previously reported that BDNF can activate EGFR proteins and cause intracellular

signalling pathways [35]. Consequently, to further identify whether the protective

effect of the H-LIPEF and BDNF on SH-SY5Y cells was related to the EGFR

pathway, we examined the protein expression of p-EGFR on human neuroblastoma

SH-SY5Y cells. As shown in Fig 4, when SH-SY5Y cells were treated with H_2O_2

alone, the expression of p-EGFR was slightly decreased. In contrast, the H-LIPEF

treatment was found to induce a significant recovery of p-EGFR protein level in the

cells under H_2O_2 oxidative stress. Meanwhile, the result finds that the combination

treatment of BDNF and H-LIPEF can further increase the expression level of p-EGFR

protein significantly. The study suggests that H-LIPEF could have an interaction with

EGFR, and that binding of BDNF to EGFR would further enhance the

neuroprotective effect induced by H-LIPEF on H_2O_2-treated SH-SY5Y cells.

Subsequently, we studied the downstream signalling pathways that could participate

in the neuroprotective mechanism of H-LIPEF and BDNF.

Fig 4. Effect of H-LIPEF alone or in combination with BDNF on expressions of

p-EGFR protein in SH-SY5Y cells. (A) Western blot analysis of p-EGFR protein

expressions. (B) Quantification of p-EGFR protein expression level after H_2O_2,

H-LIPEF+H_2O_2, and H-LIPEF+BDNF+H_2O_2 treatment. H_2O_2 and BDNF

concentrations in treatments were 500 μM and 25 ng/ml, respectively. GAPDH was

used as the loading control, and the relative expression level of each group was

compared with the untreated control group. Data represent the mean ± standard

deviation (n=3). Comparisons of the effect of H_2O_2, H-LIPEF+H_2O_2, and

H-LIPEF+BDNF+H_2O_2 on p-EGFR protein levels were analyzed using one-way

ANOVA with Tukey's post hoc test (**$P < 0.01$ and ***$P < 0.001$).

Activation of the ERK pathway in the neuroprotective effect

of the H-LIPEF and BDNF

It is well known that EGFR phosphorylation regulates the expression of its

downstream proteins such as MAPK family. ERK protein is one of the MAPK family

and has been known to be widely associated with cell survival. Upon phosphorylation,

activated ERK proteins enter the nucleus, where they can phosphorylate certain

transcription factors, thereby regulating the expression of specific genes that

contribute to cell survival [36]. As shown in Fig 5, in comparison to the untreated

control cells, the expression level of p-ERK protein was decreased when cells were

exposed to 500 μM H_2O_2. Notably, the level of p-ERK was significantly restored in

the cells treated with H-LIPEF stimulation. Furthermore, the result found that p-ERK

expression was further elevated significantly when SH-SY5Y cells were co-treated by

the combination of BDNF and H-LIPEF. Therefore, the study indicates that the

neuroprotection effect produced by the combination of BDNF and H-LIPEF could be

due to the activation of ERK pathway, thus inducing a significant recovery of p-ERK

level in the cells under H_2O_2 oxidative stress.

Fig 5. Effect of H-LIPEF alone or in combination with BDNF on expressions of

p-ERK protein in SH-SY5Y cells. (A) Western blot analysis of p-ERK protein

expressions. (B) Quantification of p-ERK protein expression level after H_2O_2,

H-LIPEF+H_2O_2, and H-LIPEF+BDNF+H_2O_2 treatment. H_2O_2 and BDNF

concentrations in treatments were 500 μM and 25 ng/ml, respectively. GAPDH was

used as the loading control, and relative expression level of each group was compared

with the untreated control group. Data represent the mean ± standard deviation (n=3).

Comparisons of the effect of H_2O_2, H-LIPEF+ H_2O_2, and H-LIPEF+BDNF+H_2O_2 on

p-ERK protein levels were analyzed using one-way ANOVA with Tukey's post hoc

test (***$P < 0.001$).

Effect of combination of H-LIPEF and BDNF on CREB, Nrf2, Bcl-2,

and Bax in H_2O_2-treated SH-SY5Y cells

Our previous research pointed out that the protective effect of H-LIPEF on SH-SY5Y

cells was through activation of ERK pathway [34]. In present study, we examined the

effect of the H-LIPEF combined with BDNF on SH-SY5Y cells under H_2O_2 oxidative

stress. The ERK downstream proteins should be measured to determine whether the

combined treatment of H-LIPEF and BDNF can further enhance the ERK signalling

pathway. In the following, we further examined the expression levels of p-CREB,

Nrf2, Bcl-2, and Bax proteins using western blot analysis. As a transcription factor

that regulates the expression of pro-survival proteins, CREB plays an important role

in neuronal survival [37]. As shown in Fig 6A, the protein expression of p-CREB in

SH-SY5Y cells was decreased when cells were exposed to 500 μM H_2O_2. Meanwhile,

we found that the expression of p-CREB was restored significantly in SH-SY5Y cells

treated with H-LIPEF stimulation. In addition, the combination of BDNF and

H-LIPEF was found to further enhance the expression of p-CREB in H_2O_2-treated

cells significantly. Next, Nrf2 is a key transcription factor that regulates the

expression of antioxidant proteins participating in the protection of oxidative damage

triggered by injury and inflammation [38]. As shown in Fig 6B, the level of Nrf2 was

also decreased in H_2O_2-treated SH-SY5Y cells compared to the untreated cells. In

contrast with the cells exposed to 500 μM H_2O_2, the expression level of Nrf2 in the

cells treated with H-LIPEF was restored significantly. Moreover, the result found that

the combined treatment of BDNF and H-LIPEF further increased the expression of

Nrf2 in H_2O_2-treated cells significantly.

Furthermore, it is known that both phosphorylated CREB and Nrf2 proteins can

regulate the expression of Bcl-2 family proteins [39, 40], and the Bcl-2/Bax ratio

determines whether the neurons undergo survival or death after being stimulated [33,

34, 41]. Therefore, the expressions of Bcl-2 and Bax proteins were also measured in

this study. As shown in Fig 6C, in comparison to the untreated control cells, the level

of Bcl-2/Bax ratio of SH-SY5Y cells was found to decrease when cells were exposed

to 500 μM H_2O_2. On the other hand, for SH-SY5Y cells treated with the H-LIPEF, the

expression levels of Bcl-2 and Bax were increased and decreased respectively, leading

to a higher elevation of the Bcl-2/Bax ratio compared to the H_2O_2-treated cells.

Moreover, it was observed that the combined treatment of BDNF and H-LIPEF

further enhanced the ratio of Bcl-2/Bax significantly in the cells under H_2O_2 oxidative

stress. The above data indicate that the combination treatment not only can greatly

increase the expressions of the phosphorylated CREB and Nrf2 proteins, but also can

remarkably enhance the ratio of Bcl-2/Bax for better neuroprotection effect, compared

to the single H-LIPEF treatment. Consequently, the result suggests that the

combination of the BDNF and H-LIPEF synergistically promotes the cell survival

against H_2O_2 oxidative stress via ERK signalling pathway.

Fig 6. Effect of H-LIPEF alone or in combination with BDNF on survival-related

proteins in SH-SY5Y cells. Western blot analysis of p-CREB (A), Nrf2 (B), Bcl-2

and Bax (C) protein expressions. H_2O_2 and BDNF concentrations in treatments were

500 μM and 25 ng/ml, respectively. The expression levels of p-CREB and Nrf2 were

normalized to GAPDH and each relative expression level was compared with control.

Data represent the mean ± standard deviation (n=4). Comparisons of the effect of

H_2O_2, H-LIPEF+H_2O_2, and H-LIPEF+BDNF+H_2O_2 on different protein levels were

analyzed using one-way ANOVA with Tukey's post hoc test (**$P < 0.01$ and ***$P <$

0.001).

Discussion

The study aimed to investigate the effect of combining BDNF and H-LIPEF in

protecting SH-SY5Y neural cells against oxidative stress. In the study, BDNF was

combined with the non-invasive and non-contact H-LIPEF with 200 Hz frequency

and 10 V/cm intensity, enhancing the survival rate of SH-SY5Y cells under oxidative

stress significantly. As one of the most abundant proteins in human brain, BDNF's

importance cannot be overstated. Many studies have pointed out that low-level of

BDNF can cause cognitive impairment in neurodegenerative diseases, such as

Alzheimer's disease, which has affected an increasing number of people worldwide

[42-44]. Some studies on low endogenous BDNF expression caused by injury or

disease have found that employment of exogenous BDNF is effective in protecting

neuron cells or brain tissues [14, 15, 45]. In the study, with H_2O_2 as oxidative stress in

a cellular model, MTT data show that the synergy of H-LIPEF and BDNF

significantly increases the viability of SH-SY5Y cells under H_2O_2 stress, plus

significant improvement in stress-induced apoptosis and suppression of elevated ROS

level caused by H_2O_2.

As for the neuroprotective mechanism, our previous study showed that the

non-invasive H-LIPEF stimulation, within a specific range of electric field intensity

and frequency, can effectively increase the survival rate of SH-SY5Y cells under

H_2O_2 stress to achieve neuroprotection effect [34]. The main mechanism is attributed

to the ability of electric field to increase the expression of survival-related proteins

such as p-ERK and p-CREB, and thus improve the viability of SH-SY5Y cells. In this

work, we used H-LIPEF together with BDNF drug to protect SH-SY5Y cells against

H_2O_2-induced cell injury. This study employed the combination treatment using the

electric field with the optimal intensity and frequency identified in our previous study

[34] and tried to pinpoint more upstream sources of survival signals in SH-SY5Y cells

for neuroprotection.

In recent years, people have gradually come to know the importance of EGFR

protein in the nervous system, as some studies have found the correlation between

neurodegeneration and reduced EGFR expression [25]. Increased activation of EGFR

may be conducive to neuroprotection. It is known that EGFR has to be bound with a

specific ligand such as epidermal growth factor (EGF) for activation and enhancement

of the expression of its downstream MAPK family proteins [46]. Moreover, recent

literature has found that EGFR can also be triggered by BDNF to activate its

downstream signalling pathways [35]. In addition, it has been reported that electric

current stimulation alone could trigger a ligand-independent activation of EGFR and

even cell migration [47]. Since activation of EGFR is thought to be associated with

cell survival, this study aims to investigate whether the combination BDNF and

H-LIPEF treatment can synergistically activate EGFR to intensify the ability of

SH-SY5Y cells to cope with oxidative stress.

While most electrical stimulation applications called for direct contact of

electrode plate with tissues and cells, the study employed the non-invasive and

non-contact H-LIPEF to stimulate SH-SY5Y cells, demonstrating that H-LIPEF alone

can significantly enhance p-EGFR protein expression in SH-SY5Y cells under H_2O_2

oxidative stress (Fig 4). Moreover, a further significant increase in the expression of

p-EGFR was observed in the presence of H-LIPEF in combination with BDNF. As

shown in Fig 5, the data also demonstrated that H-LIPEF combined with BDNF

significantly boosting the expression of p-ERK, on top of the modulation of other

proteins, such as Nrf2 and p-CREB. It is known that Nrf2 is a cellular transcription

factor that regulates the expression of various antioxidant proteins. The combination

H-LIPEF and BDNF treatment can effectively activate Nrf2 protein to protect neurons

via the antioxidant pathways. The other protein, CREB, is also a transcription factor

activated by the phosphorylation from ERK, which can trigger an increase in Bcl-2

protein expression. Experimental results also show that the combination of H-LIPEF

and BDNF can create a synergy effect raising the Bcl-2/Bax ratio and exerting a

pro-survival effect on SH-SY5Y cells.

Recently, the importance of EGFR in neuroprotection has been on the rise, which

may be a noteworthy trend for the development of neurodegenerative drugs [25].

Besides, the study demonstrates the potential of combining H-LIPEF and BDNF in

activating EGFR and related downstream survival proteins, leading to higher survival,

which is an approach that may augment drug efficacy and reduce drug dosage. This

suggests that the non-invasive H-LIPEF treatment can synergize with BDNF drug to

achieve desirable effect at low drug dosage, minimizing side effects. On the other

hand, it should be mentioned that conventional electric stimulation was conducted

mainly via contact or invasive manner using direct currents, limiting its applications

significantly, especially in some organs, such as brain. Therefore, it is believed that

the development of non-contact, non-invasive electric field therapy will be inevitable,

which can also be further applied in conjunction with drugs for various therapeutic

purposes. Previously, our team has achieved synergistic inhibitory effects on cancer

cells by combining such electric fields with herbal extracts such as EGCG, CGA, and

curcumin, and has also accomplished satisfactory results in repairing mouse motor

neuron cell NSC-34, demonstrating the versatility of non-contact, non-invasive

electric field stimulations [29-33]. In application, the electric field's strength and

frequency would vary among different cellular tissues, depending on accompanying

drugs, tissue structure, and protein response, which needs further research.

In summary, the study demonstrates that H-LIPEF can work together with BDNF,

creating a synergy effect in protecting human neuronal cells SH-SY5Y against

antioxidant stress. The underlying mechanism is believed to involve activation of

EGFR and ERK pathway to regulate ROS level, thereby enhancing cell survival. The

study also suggests further research on the combination of H-LIPEF and

BDNF-mimetic drugs for treating neurodegenerative brain diseases effectively.

Funding

This work was supported by research grants from Ministry of Science and Technology (MOST 110-2112-M-002-004, MOST 109-2112-M-002-004, and MOST 108-2112-M-002-016 to CYC) and Ministry of Education (MOE 106R880708 to CYC) of the Republic of China. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors have declared that no competing interests exist.

Acknowledgments

The authors would like to acknowledge the service provided by Technology Commons in College of Life Science, National Taiwan University for use of flow cytometry system.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Guan-Bo Lin, Chih-Yu Chao.

Formal analysis: Guan-Bo Lin, Wei-Ting Chen, Yu-Yi Kuo, You-Ming Chen, Hsu-Hsiang Liu, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Guan-Bo Lin, Wei-Ting Chen, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Guan-Bo Lin.

Writing – original draft: Guan-Bo Lin, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References

1. Chen Z, Zhong C. Oxidative stress in Alzheimer's disease. Neurosci Bull. 2014;
30(2):271-81.
2. Spillantini MG, Goedert M. Tau protein pathology in neurodegenerative diseases.
Trends Neurosci. 1998; 21(10):428-33.
3. Bathina S, Das UN. Brain-derived neurotrophic factor and its clinical
implications. Arch Med Sci. 2015; 11(6):1164-78.
4. Chen SD, Wu CL, Hwang WC, Yang DI. More insight into BDNF against
neurodegeneration: anti-apoptosis, anti-oxidation, and suppression of autophagy.
Int J Mol Sci. 2017; 18(3):545.
5. Habtemariam S. The brain-derived neurotrophic factor in neuronal plasticity and
neuroregeneration: new pharmacological concepts for old and new drugs. Neural
Regen Res. 2018; 13(6):983-4.
6. Palasz E, Wysocka A, Gasiorowska A, Chalimoniuk M, Niewiadomski W,
Niewiadomska G. BDNF as a promising therapeutic agent in Parkinson's disease.
Int J Mol Sci. 2020; 21(3):1170.
7. Ng TKS, Ho CSH, Tam WWS, Kua EH, Ho RC. Decreased serum brain-derived
neurotrophic factor (BDNF) levels in patients with Alzheimer's disease (AD): a
systematic review and meta-analysis. Int J Mol Sci. 2019; 20(2):257.
8. Lee BH, Kim YK. Reduced platelet BDNF level in patients with major

555  depression. Prog Neuropsychopharmacol Biol Psychiatry. 2009; 33(5):849-53.

556 9. Laske C, Stransky E, Leyhe T, Eschweiler G, Wittorf A, Richartz E, et al.
557  Stage-dependent BDNF serum concentrations in Alzheimer's disease. J Neural
558  Transm. 2006; 113(9):1217-24.

559 10. Zheng Z, Sabirzhanov B, Keifer J. Oligomeric amyloid-β inhibits the proteolytic
560  conversion of brain-derived neurotrophic factor (BDNF), AMPA receptor
561  trafficking, and classical conditioning. J Biol Chem. 2010; 285(45):34708-17.

562 11. Wang R, Holsinger RMD. Exercise-induced brain-derived neurotrophic factor
563  expression: Therapeutic implications for Alzheimer's dementia. Ageing Res Rev.
564  2018; 48:109-21.

565 12. Hoppe JB, Coradini K, Frozza RL, Oliveira CM, Meneghetti AB, Bernardi A, et
566  al. Free and nanoencapsulated curcumin suppress β-amyloid-induced cognitive
567  impairments in rats: involvement of BDNF and Akt/GSK-3β signaling pathway.
568  Neurobiol Learn Mem. 2013; 106:134-44.

569 13. Uddin MS, Al Mamun A, Kabir MT, Ahmad J, Jeandet P, Sarwar MS, et al.
570  Neuroprotective role of polyphenols against oxidative stress-mediated
571  neurodegeneration. Eur J Pharmacol. 2020; 886:173412.

572 14. Xu AH, Yang Y, Sun YX, Zhang CD. Exogenous brain-derived neurotrophic
573  factor attenuates cognitive impairment induced by okadaic acid in a rat model of
574  Alzheimer's disease. Neural Regen Res. 2018; 13(12):2173-81.

575 15. Arancibia S, Silhol M, Moulière F, Meffre J, Höllinger I, Maurice T, et
576  al.Protective effect of BDNF against beta-amyloid induced neurotoxicity in vitro
577  and in vivo in rats. Neurobiol Dis. 2008; 31(3):316-26.

578 16. Tapia-Arancibia L, Aliaga E, Silhol M, Arancibia S. New insights into brain
579  BDNF function in normal aging and Alzheimer disease. Brain Res Rev. 2008;
580  59(1):201-20.

581 17. Poduslo JF, Curran GL. Permeability at the blood-brain and blood-nerve barriers
582  of the neurotrophic factors: NGF, CNTF, NT-3, BDNF. Brain Res Mol Brain Res.
583  1996; 36(2):280-6.

584 18. Ainsworth L, Budelier K, Clinesmith M, Fiedler A, Landstrom R, Leeper B, et al.
585  Transcutaneous electrical nerve stimulation (TENS) reduces chronic hyperalgesia
586  induced by muscle inflammation. Pain. 2006; 120(1-2):182-7.

587 19. Wagstaff PG, Buijs M, van den Bos W, de Bruin DM, Zondervan PJ, de la
588  Rosette JJ, et al. Irreversible electroporation: state of the art. Onco Targets Ther.
589  2016; 9:2437-46.

590 20. Perlmutter JS, Mink JW. Deep brain stimulation. Annu Rev Neurosci. 2006;
591  29:229-57.

592 21. Thakral G, LaFontaine J, Najafi B, Talal TK, Kim P, Lavery LA. Electrical

stimulation to accelerate wound healing. Diabet Foot Ankle. 2013; 4(1):22081.

22. Lamprea MR, Cardenas FP, Vianna DM, Castilho VM, Cruz-Morales SE, Brandão ML. The distribution of fos immunoreactivity in rat brain following freezing and escape responses elicited by electrical stimulation of the inferior colliculus. Brain Res. 2002; 950(1-2):186-94.

23. Hekstra DR, White KI, Socolich MA, Henning RW, Šrajer V, Ranganathan R. Electric-field-stimulated protein mechanics. Nature. 2016; 540(7633):400-5.

24. Kawamura K, Kano Y. Electrical stimulation induces neurite outgrowth in PC12m3 cells via the p38 mitogen-activated protein kinase pathway. Neurosci Lett. 2019; 698:81-4.

25. Romano R, Bucci C. Role of EGFR in the Nervous System. Cells. 2020; 9(8):1887.

26. Dreux AC, Lamb DJ, Modjtahedi H, Ferns GA. The epidermal growth factor receptors and their family of ligands: their putative role in atherogenesis. Atherosclerosis. 2006; 186(1):38-53.

27. Chen C, Bai X, Ding Y, Lee IS. Electrical stimulation as a novel tool for regulating cell behavior in tissue engineering. Biomater Res. 2019; 23(1):1-12.

28. Novac BM, Banakhr FA, Smith IR, Pecastaing L, Ruscassie R, De Ferron AS, et al. Demonstration of a novel pulsed electric field technique generating neither conduction currents nor Joule effects. IEEE Trans Plasma Sci. 2014; 42(1):216-28.

29. Hsieh CH, Lu CH, Chen WT, Ma BL, Chao CY. Application of non-invasive low strength pulsed electric field to EGCG treatment synergistically enhanced the inhibition effect on PANC-1 cells. PLoS One. 2017; 12(11):e0188885.

30. Lu CH, Lin SH, Hsieh CH, Chen WT, Chao CY. Enhanced anticancer effects of low-dose curcumin with non-invasive pulsed electric field on PANC-1 cells. Onco Targets Ther. 2018; 11:4723-32.

31. Hsieh CH, Lu CH, Kuo YY, Chen WT, Chao CY. Studies on the non-invasive anticancer remedy of the triple combination of epigallocatechin gallate, pulsed electric field, and ultrasound PLoS One. 2018; 13(8):e0201920.

32. Lu CH, Kuo YY, Lin GB, Chen WT, Chao CY. Application of non-invasive low-intensity pulsed electric field with thermal cycling-hyperthermia for synergistically enhanced anticancer effect of chlorogenic acid on PANC-1 cells. PLoS One. 2020; 15(1):e0222126.

33. Hsieh CH, Lu CH, Kuo YY, Lin GB, Chao CY. The protective effect of non-invasive low intensity pulsed electric field and fucoidan in preventing oxidative stress-induced motor neuron death via ROCK/Akt pathway. PLoS One. 2019; 14(3):e0214100.

34. Chen WT, Lin GB, Kuo YY, Hsieh CH, Lu CH, Sun YK, et al. Effect of
high-frequency low-intensity pulsed electric field on protecting SH-SY5Y cells
against hydrogen peroxide and β-amyloid-induced cell injury via ERK pathway.
PLoS One. 2021; 16(4):e0250491.

35. Qiu L, Zhou C, Sun Y, Di W, Scheffler E, Healey S, et al. Crosstalk between
EGFR and TrkB enhances ovarian cancer cell migration and proliferation. Int J
Oncol. 2006; 29(4):1003-11.

36. Almeida RD, Manadas BJ, Melo CV, Gomes JR, Mendes CS, Grãos MM, et al.
Neuroprotection by BDNF against glutamate-induced apoptotic cell death is
mediated by ERK and PI3-kinase pathways. Cell Death Differ. 2005;
12(10):1329-43.

37. Sakamoto K, Karelina K, Obrietan K. CREB: a multifaceted regulator of
neuronal plasticity and protection. J Neurochem. 2011; 116(1):1-9.

38. Ma Q. Role of nrf2 in oxidative stress and toxicity. Annu Rev Pharmacol Toxicol.
2013; 53(1):401-26.

39. Walton M, Woodgate AM, Muravlev A, Xu R, During MJ, Dragunow M. CREB
phosphorylation promotes nerve cell survival. J Neurochem. 1999;
73(5):1836-42.

40. Niture SK, Jaiswal AK. Nrf2 protein up-regulates antiapoptotic protein Bcl-2 and
prevents cellular apoptosis. J Biol Chem. 2012; 287(13):9873-86.

41. Korsmeyer SJ, Shutter JR, Veis DJ, Merry DE, Oltvai ZN. Bcl-2/Bax: a rheostat
that regulates an anti-oxidant pathway and cell death. Semin Cancer Biol. 1993;
4(6):327-32.

42. Shimada H, Makizako H, Doi T, Yoshida D, Tsutsumimoto K, Anan Y, et al. A
large, cross-sectional observational study of serum BDNF, cognitive function,
and mild cognitive impairment in the elderly. Front Aging Neurosci. 2014; 6:69.

43. Siuda J, Patalong-Ogiewa M, Żmuda W, Targosz-Gajniak M, Niewiadomska E,
Matuszek I, et al. Cognitive impairment and BDNF serum levels. Neurol
Neurochir Pol. 2017; 51(1):24-32.

44. Weinstein G, Beiser AS, Choi SH, Preis SR, Chen TC, Vorgas D, et al. Serum
brain-derived neurotrophic factor and the risk for dementia: the Framingham
Heart Study. JAMA Neurol. 2014; 71(1):55-61.

45. Zhang L, Fang Y, Lian Y, Chen Y, Wu T, Zheng Y, et al. Brain-derived
neurotrophic factor ameliorates learning deficits in a rat model of Alzheimer's
disease induced by aβ1-42. PLoS One. 2015; 10(4):e0122415.

46. Wells A. EGF receptor. Int J Biochem Cell Biol. 1999; 31(6):637-43.

47. Wolf-Goldberg T, Barbul A, Ben-Dov N, Korenstein R. Low electric fields induce
ligand-independent activation of EGF receptor and ERK via electrochemical

elevation of H(+) and ROS concentrations. Biochim Biophys Acta. 2013;
1833(6):1396-408.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6