1 Rapid evolution in response to climate-change-induced drought and its
2 demographic and genetic controls
3

4 **John W. Benning***
5 University of Wyoming
6 Department of Botany
7 Laramie, WY, USA
8 *jwbenning@gmail.com
9
10 **Alexai Faulkner**
11 University of Minnesota
12 Department of Plant and Microbial Biology
13 Saint Paul, MN, USA
14
15 **David A. Moeller**
16 University of Minnesota
17 Department of Plant and Microbial Biology
18 Saint Paul, MN, USA

Abstract

Populations often vary in their evolutionary responses to a shared environmental
perturbation. A key hurdle in building more predictive models of rapid evolution is
understanding this variation – why do some populations and traits evolve while others do not?
We combined long-term demographic and environmental data, estimates of quantitative genetic
variance components, a resurrection experiment, and individual-based evolutionary simulations
to gain mechanistic insights into contrasting evolutionary responses to a severe multiyear
drought. We examined five traits in two populations of a native California plant, *Clarkia*
xantiana, at three timepoints over 7 years. Earlier flowering phenology evolved in only one of
the two populations, though both populations experienced similar drought severity and
demographic declines and were estimated to have considerable additive genetic variance for
flowering phenology. Pairing demographic and experimental data with evolutionary simulations
suggested that while seed banks in both populations likely constrained evolutionary responses, a
stronger seed bank in the non-evolving population resulted in evolutionary stasis. Gene flow
through time via germ banks may be an important, underappreciated control on rapid evolution
in response to extreme environmental perturbations.
35
36 **Keywords**: adaptation; evolutionary rescue; *Clarkia xantiana*; seed bank; genetic constraints

Introduction

Extreme environmental perturbations offer excellent opportunities to examine evolution in natural populations on ecological timescales [1]. Rapid evolution in response to strong episodes of environmental change has been documented in many taxa, including lizards [2,3], finches [4], and plants [5]. Beyond offering fundamental insights into the evolutionary process, this and related work speak to the potential for natural populations to evolve in response to anthropogenic forcings like climate change, and to the importance of adaptation for the persistence of populations [6–8]. However, it remains poorly understood as to why rapid evolution often fails to occur in some populations despite the same environmental episode resulting in evolution in other nearby populations.

Resurrection experiments are increasingly used to directly examine evolutionary change before and after an environmental event predicted to exert strong selection on populations [5,9]. The power of resurrection experiments lies in their ability to precisely quantify short-term evolutionary responses, rather than only making predictions of the trajectory of evolution, while controlling for confounding environmental effects on phenotypes. In short, the resurrection approach entails rearing different generations of a population contemporaneously in a common environment to directly quantify phenotypic evolution. Though only a modest number of resurrection studies have been published to date [10], such experiments with plants have sometimes demonstrated rapid evolution in response to periods of drought [5,11,12].

Just as important as these positive results, however, are instances in which resurrection experiments do *not* find evidence of rapid evolution. Populations of a species can vary in their responses to the same climatic fluctuation [12–14], and even extreme climatic anomalies can result in no discernible phenotypic evolution of putatively important traits [15]. What underlies

this variation in populations' evolutionary responses to extreme events? While the resurrection

approach is a powerful method to determine if phenotypic evolution has occurred, it does not

typically provide insight into the demographic and genetic controls on evolutionary change.

A more comprehensive understanding of the causes and consequences of rapid evolution

can be gained by coupling evolutionary analyses with long-term demographic and quantitative

genetic investigations. Sharp declines in population size due to environmental perturbations will

reduce the efficacy of selection and increase the influence of stochastic processes like genetic

drift [16,17]. Such demographic, environmental, or genetic stochasticity may stymie adaptation,

potentially causing population extinction before 'evolutionary rescue' can occur [18–20]. Models

and experiments have further shown that evolutionary rescue is less probable when the

environment changes suddenly compared to when the change is gradual [21]. As such, the rate

and severity of demographic decline can provide insight into instances where some populations

fail to evolve in response to an extreme event that drove rapid evolution in other conspecific

populations.

For organisms with dormancy, long-term outcomes of selection can also be affected by

the presence of a germ bank (seed or egg bank). Populations that often experience significant

environmental fluctuations may evolve mechanisms that confer dormancy as a bet-hedging

strategy [22]. For annual plants, seed banks thus result in overlapping, instead of discrete

generations. While germ banks may buffer populations from extinction [23,24], they may also

constrain evolutionary responses to environmental change due to gene flow among generations

[25,26]. This "temporal migration" can slow the rate of adaptive evolution [27,28]. Alleles

favored prior to a selective event will be 'reintroduced' to the population from the germ bank

during and after the event. If those same alleles are selected against during the event, this gene

flow can retard adaptation [but see 29]. Moreover, the environmental and demographic history of

populations just prior to an extreme event likely affects the magnitude of gene flow from the

past. For example, high fecundity of generations immediately prior to an extreme environmental

event would increase input to the seed bank. Therefore, historical demographic data can also

provide insight into the influence of germ banks on population responses to selection.

Strong environmental perturbations typically exert selection on traits, but the same

perturbation may still result in selection regimes that vary among sites [5]. The response to any

selection pressure will depend upon the presence of sufficient additive genetic variance for traits

underlying fitness variation [30,31]. A population's response to selection will be directly

proportional to a given trait's narrow-sense heritability, $h^2 = V_A/V_p$ [32]. In the context of

resurrection experiments, measuring the additive genetic variance of ecologically-important traits

may help us understand why traits do or do not evolve in different populations exposed to similar

selective regimes. While tests of rapid evolution are increasingly common, few studies have

simultaneously examined quantitative genetic variation and its potential role in facilitating or

hindering adaptation [but see 5].

Furthermore, resurrection experiments usually examine cohorts from before and at the

end of an extreme event [but see 33]. A key question is how long evolutionary changes that

occur in response to these events persist beyond them. Including later cohorts in a resurrection

experiment allows us to assess the temporal stability of evolutionary responses and the extent to

which dormancy might delay responses to selection. When paired with associated demographic

data from natural populations, such a design gives inference to how rapid evolution (or its

absence) influences a populations' longer-term phenotypic and demographic trajectory.

The Southwest of the United States is in the midst of the most severe multi-decadal

drought (i.e., megadrought) in recorded history (at least since 800 C.E.) [34,35]. Recent analyses

have shown that anthropogenic climate change has been a key driver of this drought and

accounts for an estimated 42% of the soil moisture anomaly [35,36]. We examined evolutionary

responses to the most severe episode of this climate anomaly, which began in the latter half of

2011 and ended in late 2015 in our study area. Our work focused on a well-studied annual plant,

Clarkia xantiana ssp. *xantiana*, which is endemic to Southern California, U.S.A. We used a

resurrection experiment to test whether traits likely to mediate drought adaptation exhibited rapid

evolution and whether genetic or demographic constraints modulated responses.

We monitored environmental and demographic variation over 12 years (2006 – 2017),

spanning a period from six years prior to drought until two years after the drought ended. Those

data provided insight into the severity and rate of environmental and demographic change, as

well as the magnitude of input to the seed bank in years just prior to the drought episode. To test

for rapid evolution in response to drought, we used a resurrection experiment with pedigreed

individuals from three time points — prior to the prolonged drought (2011), at the end of the

drought (2015), and two years later, after more average precipitation resumed (2017). This latter

sample allowed us to examine whether any evolutionary responses persisted, increased, or

evolved back toward the original population mean. We measured a suite of traits that prior

studies have shown to confer drought escape and/or avoidance, and estimated additive genetic

variance for each of these traits in each population. Lastly, we leveraged these demographic and

experimental data in the construction and analysis of an individual-based evolutionary simulation

model to test the hypothesis that a seed bank influenced the evolutionary responses of these

populations to extreme drought.

Material and Methods

Study System

Clarkia xantiana ssp. *xantiana* A. Gray (Onagraceae) is a predominantly outcrossing

winter annual native to the southern Sierra Nevada foothills and Transverse Ranges of

California, USA [37]. In this Mediterranean climate, the bulk of precipitation occurs during

winter and early spring. Plants germinate November - March during the rainy season, begin

flowering in May, and set seed in late June. In this study, we focus on two populations, KYE and

S22. KYE occupies oak woodland habitat on granite-derived soils characteristic of the more

mesic, western portion of *C. x. xantiana*'s distribution (35.6240674°, -118.5156798°). S22 is

located near the subspecies' eastern geographic range limit and occupies a more xeric, higher

elevation site in pine woodland on metasedimentary-derived soils (35.83996°, -118.450386°).

These metasedimentary-derived soils occur along the Kern Canyon Fault, a ca. 150km fault that

parallels the Kern River through the Southern Sierra [38,39]. [Site identifiers are 57x (KYE) and

22x (S22) for consistency with previous and future *C. x. xantiana* studies).] Though it was long

assumed that *Clarkia* species had little if any seed storage [e.g., 40], recent work has shown that

C. x. xantiana populations can harbor substantial seed banks [41,42].

Climatic and demographic data

We obtained growing season precipitation data (November - June) for years 2006-2017

from long-term weather monitoring stations (HOBO Onset) at each site. Demographic data were

collected as part of a long-term study of 36 populations across the range of *C. x. xantiana* in

years 2006-2017 (detailed in Supp. Mat. A). Using a population projection model approach, we

combined estimates of seed input in each year with estimates of seed bank vital rates (survival

and germination) for KYE and S22 from [42] to estimate the age distribution of germinants in

each population in each year (assuming a maximum seed age of 3 years). Siegmund et al. (2022)

estimated these vital rates for the two populations in our study as: seed survival rate (KYE =

0.66; S22 = 0.61), germination rate (KYE = 0.13; S22 = 0.27). The multi-year field experiments

used to estimate these vital rates are detailed in [42] and [41].

Seed collection and refresher generation

We used seeds collected as part of the aforementioned long-term demographic study on

C. x. xantiana. Seeds were collected from a haphazard sampling of plants (dozens to hundreds of

plants, depending on field conditions) across the spatial extent of sites KYE and S22 in late June

of years 2011 (pre-drought), 2015 (end of drought), and 2017 (after two years of more average

precipitation following the drought). All seeds from a given dam constitute a maternal family.

Because *C. x. xantiana* is a highly outcrossing species [43] with multiple paternity (DM, unpubl.

data), each maternal family sampled from the field was a collection of full and half-sibs. Seeds

were stored at room temperature in plastic boxes containing silica desiccant, with maternal

families each in their own coin envelope. To standardize maternal environmental effects, we

grew plants from each year together in a greenhouse for one "refresher" generation to produce

seeds for the resurrection experiment [10].

For each of the three generations of each population (six "cohorts"), we grew 1-5 plants

from each of 20 haphazardly selected maternal families (N = 219 plants total) in a fully

randomized design in the greenhouse in spring 2018. Most (66%) of maternal families were

represented by two plants; representation of families varied due to unequal germination. Within

each cohort, half the plants were randomly assigned as sires and each sire was mated to two

unique dams (with all plants serving as dams) to produce a pedigreed population for the

measurement round of the experiment. We also grew individuals from a third site, Mill Creek,

during this generation prior to the resurrection experiment, but dropped this site from the

subsequent experiment for feasibility; data from the Mill Creek refresher generation did not

suggest evolution of any of the three measured traits (growth rate, days to flowering, SLA; Fig.

S4).

Measurement generation

With the seeds produced from crosses, we grew the six offspring cohorts in a fully

randomized design in the greenhouse in spring 2019 to assess phenotypic changes across

generations for each population. Of the original 20 maternal families planted for each year cohort

in the refresher generation, 16-20 were represented (as sire and/or dam) in this pedigreed

population of offspring. We sowed six seeds for each of 22-28 dams from each refresher cohort

in plug trays with germination mix in the growth chamber and scored germination for 33 days.

There were 83-122 germinants per cohort (Table S2); germination rates ranged from 0.61 to 0.75

(Table S1). Up to four seedlings per dam (haphazardly chosen) were transplanted into individual

656 mL Deepots (Stuewe & Sons, Tangent, OR) filled with a 1:1 mix of sand and potting soil.

Pots were arranged in a completely randomized design in the greenhouse on a 16/8 hr light

schedule and watered as needed. Final sample sizes varied due to unequal seed availability and

germination among cohorts ($n = 65 - 97$ individuals per year cohort for post-germination traits;

Table S2).

Trait measurements

We measured five traits that have previously been shown to be related to drought escape

and/or avoidance: days to germination, growth rate, days to flowering, specific leaf area (SLA),

and leaf succulence. Faster germination, growth, or flowering phenology may allow plants to

take advantage of the relatively mesic early growing season and complete their life cycle before

the onset of drier conditions [5,11,44], though delayed germination may also be selected for in

arid environments [e.g., 45]. Lower SLA and increased leaf succulence can increase water use

efficiency, and evolution in these physiological traits may represent drought avoidance

adaptations [44,46]. We also measured two fitness proxies: total number of flowers produced and

shoot (aboveground) biomass.

Days to germination was the number of days elapsed between sowing and cotyledon

emergence. Growth rate was measured as the number of leaves produced per day, measured for

plants 41-43 days post-germination (i.e., growth rate was calculated as the number of leaves at

measurement divided by the plant age in days at measurement). One fully-expanded leaf was

collected from each plant 64-75 days post-germination; we recorded fresh and dry weight for that

leaf and used ImageJ [47] to measure leaf area from a photo of the fresh leaf. SLA was

calculated as $mm^2\,mg^{-1}$; leaf succulence was calculated as leaf wet weight / leaf dry weight.

Days to flowering was measured as the days elapsed between seed sowing and the opening of a

plant's first flower. Measuring flowering time in such a way, as opposed to the days elapsed

between germination and flowering, incorporates variability in time to germination and is more

directly relevant to phenological timing as measured in wild populations; both traits showed a

similar pattern across years (Table S1). Total number of flowers produced was measured when

all flowers had opened on a plant. Shoot biomass was collected at the end of the experiment and

plants were dried before weighing. Two pairs of traits were strongly correlated [number of

flowers produced and shoot biomass (r = 0.72); SLA and leaf succulence (r = 0.81); Fig. S5]. We

report on number of flowers produced and specific leaf area below, as flowers produced is a

more direct proxy for plant fitness than shoot biomass, and SLA is a more commonly-reported

trait than leaf succulence. We include descriptive statistics for shoot biomass and leaf succulence

in Table S1. Our analyses of invisible fraction bias [sensu 48] indicated that any bias in

estimation of trait means due to unequal seed survival among genotypes was likely minimal

(Supp. Mat. B).

Statistical analyses for resurrection experiment

224 All statistical analyses were conducted in R version 4.1.2 [49]. Data were organized and

225 summarized using the tidyr [50] and dplyr [51] packages and plotted using ggplot2 [52]. All data

226 and code necessary to reproduce these analyses are uploaded with the submission and will be

227 archived at FigShare upon acceptance.

Phenotypic evolution

We used linear mixed models [package lme4 [53]] to test for differences among year

cohorts in the measured phenotypic traits (days to germination, growth rate, flowering date,

SLA, and flower number). We analyzed populations separately because our main interest was in

differences among year cohorts, not populations, and we had limited power to detect interactions

between population and year. All models took the general form of *response ~ year + transplant*

age + (1|sire/dam). Transplant age (days elapsed between germination and transplanting from

germination tray to full pot) was included as a covariate to account for differing seedling ages

when transplanted to pots. Sire, and dam nested within sire, were included as random effects to

account for non-independence amongst related individuals. For days to germination, dam

average seed weight was included as a covariate and transplant age was not. Only seeds that

germinated were included in days to germination analyses to avoid any confounding of

dormancy with seed viability. Days to germination and SLA were log transformed prior to

analysis to better meet assumptions of homoscedasticity and normality of model residuals. We

tested whether inclusion of terms improved model fit with Wald λ^2 tests using Type II SS

[Anova.mermod() function in the car package [54]]. If year was a significant predictor at

$\alpha = 0.05$, we followed up with pairwise Tukey contrasts between year cohorts using the

emmeans package [55]. We computed estimated marginal means for each year cohort using

emmeans.

Genetic variance components

We estimated quantitative genetic variance components for all traits in both populations

using an animal model implemented in the MCMCglmm package [56]. In short, the animal

model approach comprises a linear mixed effects model with an individual's breeding (i.e.,

additive genetic) value modeled as a random effect [57,58]. A pedigree of the population

provides an expectation of how breeding values should covary between individuals, and thus

allows for an estimate of a trait's additive genetic variance in that population. MCMCglmm uses

a Markov Chain Monte Carlo approach in a Bayesian framework to approximate the posterior

distribution of quantitative genetic variance components (analyses detailed in Supp. mat. C).

Following [59] we obtained estimates of additive genetic variance (V_A), residual variance (V_R),

and the variance explained by fixed effects (V_F) from the model, and calculated $h^2 = V_A / (V_A +$

$V_R + V_F$); we report the mean of the posterior distribution of h^2 as our estimate. We considered a

trait to exhibit significant heritability if its 95% credible interval for h^2 did not touch 0.001 (in

MCMCglmm, variance estimates must be greater than zero). MCMCglmm diagnostic plots and

posterior distribution of model parameters are included in SI: MCMCglmm.

Simulation model

We used an individual-based, genomically-explicit evolutionary simulation model, paired

with Approximate Bayesian Computation, to test the hypothesis that seed bank dynamics

influenced evolutionary patterns in our populations. We focused this investigation on one trait,

flowering phenology, which displayed contrasting evolutionary responses in our two populations

(Results). Using the SLiM modeling framework [60], we built a model tracking the demography

and evolution of a population of diploid, hermaphroditic individuals that experience an

environmental perturbation. Individual female fitness (here, the probability of surviving to set

seed) is determined by the deviation of an individual's phenotype (here, flowering phenology,

which is under stabilizing selection and controlled by multiple QTL) from the optimal

phenotype; the optimal phenotype is assumed to change based on current environmental

conditions. We focus on female fitness (i.e., siring success is not affected by an individual's

phenotype) because earlier work in *C. x. xantiana* has demonstrated strong selection on

flowering phenology via female fitness in arid conditions [61] but we have no such tests of

selection via male fitness. After a long burn-in period with a relatively stable environment (and

thus phenotypic optimum), the model environment and population size are explicitly tied to our

observed 12 year data set. There is an environmental perturbation (e.g., drought) in year 7 that

changes the phenotypic optimum, and modeled population sizes in each year are proportional to

the observed census sizes of natural populations in each year. Full model details are in Supp.

Mat. D.

We used Approximate Bayesian Computation (ABC) to 1) test the hypothesis that seed

banks influenced evolutionary trajectories in our populations, 2) test the hypothesis that the

phenotypic optimum returned to its pre-drought state after drought, and 3) estimate seed bank

demographic (seed survival and germination) and evolutionary (change in phenotypic optimum

during drought) parameters in both populations. ABC is a statistical method used to estimate

model parameters and choose among a set of candidate models when likelihood calculations are

intractable [62]. First, many simulations are run where values of the parameters of interest are

randomly drawn from their defined prior distributions (Table 1). Then, summary statistics are

calculated for each simulation run and are used to compare simulation outcomes with observed

data, in order to compare models and estimate parameters. Our three summary statistics were: 1)

the difference in mean phenotype between 2011 to 2015 (which we measured in the resurrection

experiment), 2) the difference in mean phenotype between 2015 and 2017 (measured in the

resurrection experiment), and 3) the ratio of survival rates between the drought (averaging 2012-

2015) and pre-drought (averaging 2006-2011) periods (measured in the field). The observed

values of the first two summary statistics were calculated using the estimated marginal means

from the linear mixed effects models described above in *Statistical analyses*. For the observed

value of the third summary statistic, because survival in the model is only affected by the focal

trait but in the field is likely affected by multiple traits, we assumed that one third of the decrease

in survival during drought in natural populations was due to the focal trait (flowering

phenology).

For each of our two populations, we explored four models (Table 1). Models were run

separately for each population because model demography was explicitly tied to observed census

sizes in each population (see above). In the models without seed banks (Models A and B), seed

survival was set to 0.0 and seed germination was set to 1.0, while the change in optimum was

allowed to vary between -20.0 and 0.0 (Table 1). For models with seed banks (Models C and D),

seed survival and germination rates were also allowed to vary between 0.05 and 0.75, which

encompasses the range of values for these parameters estimated across 20 *C. x. xantiana*

populations in [42] (range of posterior mode for germination = [0.11, 0.27]; seed survival [0.45,

0.72]). We ran 20,000 simulations of each model for model comparison, then ran an additional

20,000 simulations of the best model to aid in parameter estimation. We used the `abc` package

[63] in R to compare models and estimate parameters (*Supp. Mat. D*). Graphical posterior

predictive checks were performed to compare observed summary statistics in each population to

the distribution of summary statistics resulting from 500 simulations run using parameter values

drawn from the estimated parameter posterior distributions.

Table 1. Description and parameter prior distributions for the four candidate models.

	Description	**Parameters**		
		Seed survival rate	**Seed germination rate**	**Change in optimum**
Model A	No seed bank, with the trait optimum changing at the beginning of the drought, then returning to the historical optimum after drought	0.0	1.0	U[-20.0, 0.0]
Model B	No seed bank, with the trait optimum changing at the beginning of the drought and staying the same post-drought	0.0	1.0	U[-20.0, 0.0]
Model C	With seed bank, with the trait optimum changing at the beginning of the drought, then	U[0.05, 0.75]	U[0.05, 0.75]	U[-20.0, 0.0]

	returning to the historical optimum after drought			
Model D	With seed bank, with the trait optimum changing at the beginning of the drought and staying the same post-drought	U[0.05, 0.75]	U[0.05, 0.75]	U[-20.0, 0.0]

To estimate the individual effects of aboveground demography (seed input), seed bank

vital rates (survival and germination), and selection environment (change in phenotypic optimum

during drought) on the evolutionary stasis of KYE relative to S22 (Results), we compared the

posterior predictive checks described above to additional batches of simulations (500 simulation

each) as follows:

1. Aboveground demographic effect: seed bank (survival and germination) and optimum

324   change parameters estimated for KYE, but with demographic history of S22

325   2. Seed bank effect: optimum change estimated for KYE, with KYE demographic

326   history, but with seed bank parameters estimated for S22

327   3. Selection environment effect: seed bank parameters estimated for KYE, with KYE

328   demographic history, but with optimum change parameter estimated for S22

Results

Climate and demography

Both sites experienced a period of substantially reduced precipitation in years 2012-2015

(Fig 1). For KYE, growing season precipitation was reduced by 45% compared to 2006-2011;

for S22, precipitation was reduced 57%. Precipitation increased in 2016 and 2017 at both sites,

with particularly high precipitation in 2017.

Overall, KYE, which is nearer the range center, had higher density and total population

size than S22, which is located near the range edge (average density across 12 years: 4.03 and

1.36 fruiting plants per m^2, respectively; average population size: 282,704 and 9,166 fruiting

plants). Reduced precipitation during the 2012-2015 drought period had strong demographic

effects at both sites (Fig. 1). At both sites, survival rates of plants in permanent plots decreased

over 85 percent during drought relative to pre-drought averages (KYE survival rates: pre-drought

= 0.34, drought = 0.05; S22 pre-drought = 0.35, drought = 0.02). At KYE, fruiting plant

estimates during this drought period were 94% lower than pre-drought estimates (0.5 and 7.7

fruiting plants per m^2, respectively), though the pre-drought average was largely influenced by

high abundances in 2010 and 2011 (Fig. 1). At S22, plant densities during the drought period

were 83% lower than pre-drought estimates (0.4 and 2.3 fruiting plants per m^2, respectively). In

the two years post-drought, fruiting plants remained few at KYE. At S22, the number of fruiting

plants somewhat rebounded in 2017.

Seed input prior to the drought was roughly twice as high at KYE relative to S22 (yearly

mean of 1,198 and 646 seeds per m^2, respectively). Seed input was especially high at KYE in the

two years preceding the drought, 2010-2011. Seed input was similar at KYE and S22 during the

2012-2015 drought period (92 and 76 seeds per m^2, respectively; Fig. 1). In the two years post-

drought, patterns of seed input mirrored patterns of plant density, with input remaining low at

KYE but trending upward at S22 in 2017. The decreased germination and increased seed

survival rates of KYE (estimated in [42]) along with high seed input resulted in KYE having a

more substantial seed bank than S22, with estimated mean germinant age ca. 20% higher than

S22 (1.27 vs. 1.04 mean germinant age, respectively; Fig. 1). In both populations, low seed input

358 during drought resulted in increased germinant age during that period due to older seeds

359 comprising a larger proportion of the seed bank (Figs. 1, S8).

Figure 1. Environmental and demographic trends in two populations (KYE, S22) of the California annual plant *Clarkia xantiana ssp. xantiana* from 2006-2017. Dotted vertical lines mark cohorts collected for the resurrection experiment. Precipitation represents the cumulative growing season precipitation, in mm, for a given year (i.e., precipitation for year 2011 was the cumulative precipitation from November 2010 – June 2011). Blue horizontal bars in top panels mark the mean precipitation across three periods – pre-drought, during drought, and post-drought, with the drought period (2012-2015) highlighted in gold. Germinants, fruiting plants, and seed input are per m^2. Survival rate is based on censuses of germinants and fruiting plants in permanent plots in each population; no survival data are presented for 2014 because zero germinants were recorded in both populations. Mean germinant age in each year is estimated based on seed input in years prior and seed bank vital rates from [42].

Phenotypic evolution

For S22, days to flowering differed among year cohorts ($P = 0.004$; Fig. 2; Table S3). Days to flowering decreased 1.6 days from 2011 to 2015, and another 1.2 days from 2015 to 2017. Pairwise Tukey tests identified the 2011 / 2017 contrast in days to flowering as significant (69.4 vs. 66.4 days, respectively; $P = 0.006$). Phenotypic variance in days to flowering also decreased across year cohorts, from 28.6 in 2011, to 21.3 in 2015, to 14.3 in 2017 (Table S2). There was also a trend in S22, albeit not significant, for growth rate to increase over this timeline, which may have contributed to earlier flowering time. There was no statistically significant evidence of phenotypic evolution of other traits for S22, nor any traits in the KYE population.

Figure 2. Measured phenotypes across the three sampling years for populations KYE and S22. Each panel shows estimated marginal means with 95% CI's estimated from the model *trait ~ year + transplant age + (1|sire/dam)*. Days to germination was measured as days elapsed between sowing and emergence; growth rate was measured as leaves produced per day; days to flowering was measured as days elapsed between sowing and opening of the first flower; SLA was calculated as mm^2 mg^{-1}; flower number was measured as the total number of flowers an individual produced in its lifetime. Raw data are shown in Fig. S6.

Additive genetic variance

There was evidence of significant additive genetic variance ( ) for days to germination, flowering date, and flower number in both populations (Fig. 3; Table S4) The h^2 estimates for these traits ranged from 0.32 – 0.59. There was also evidence of significant for growth rate in the S22 population, but not in the KYE population. We did not find evidence that for SLA was significantly different from zero in either population.

Figure 3. Estimates of narrow-sense heritability (h^2) for traits, with 95% credible intervals. Estimates of additive genetic variance for each trait are in Table S4.

Simulation model

For both populations, the model with the highest posterior probability was Model D (population with a seed bank, with post-drought phenotypic optimum equal to optimum during

drought) (Table S5). Posterior probability of Model D was much higher than all other models for

KYE (Bayes factors all > 12; Table S5). For S22, the contrast of Model D with the second best

model, Model B (population *without* a seed bank, with post-drought phenotypic optimum equal

to optimum during drought), was weaker (Bayes factor = 2.3), indicating less support for an

influential seed bank in that population compared to KYE. Overall, without a seed bank,

populations experiencing a strong demographic decline due to large changes in phenotypic

optima saw rapid phenotypic evolution toward the new optimum (Fig. S9a). However, with a

seed bank, even large changes in optima usually resulted in only modest phenotypic evolution

like that observed in our populations. These constrained evolutionary responses were caused by

gene flow from prior generations via the seed bank.

We can concisely describe the strength of a population's seed bank as a combination of

the seed germination and survival rates, $\phi = (1 - g)s$. Thus, ϕ ("strength of seed bank" in Fig.

4) describes the expected proportion of seeds available for germination in year t that will be

available for germination in year $t + 1$, with larger values of ϕ indicating a stronger seed bank.

The strength of the seed bank at KYE was estimated to be more than five-fold higher than that of

S22 (0.47 vs. 0.07) due to a higher seed survival rate and a lower germination rate (Fig. 4a).

KYE was also estimated to have experienced a 15% smaller change in optimum during drought

compared to S22 (-6.1 vs. -7.2 days; Fig. 4a). Observed evolutionary and demographic responses

of both populations fell within the distributions of posterior predictive checks (Fig. 4b).

KYE's mating population had a consistently higher mean age than S22's because of its

stronger seed bank (Fig. 4c). However, mean age for both populations increased during the

drought period due to low fecundity and increased representation of older generations. The

increased gene flow from earlier generations in KYE led to the mean phenotype of its mating

pool responding much more slowly to the environmental perturbation than S22 (Figs. 4c, S9b).

Further simulation runs indicated that the difference in evolutionary responses was

overwhelmingly due to differences in seed bank vital rates, as opposed to differences in

aboveground demography or selective environments (Fig. 4d). Simulations run with the selective

environment and aboveground demography of KYE, but with the seed bank rates of S22,

produced evolutionary change nearly equal to that observed in simulations run with full S22

parameters.

Figure 4. Results of evolutionary simulations and ABC analyses. (a) Estimated posterior
distributions for model parameters (germination rate, seed survival rate, change in trait optimum
during drought). Seedbank strength (   ) is a joint parameter defined as   , where *g* is
germination rate and *s* is seed survival rate. Red vertical lines mark the posterior means. (b)
Posterior predictive checks for the three summary statistics used in ABC analyses (change in trait
mean 2015-2011, change in trait mean 2017-2015, and ratio of survival during drought to
survival pre-drought). Blue vertical lines mark observed summary statistics from empirical data.
(c) Mean age and phenotype of mating pools in both populations over the 12 focal years during
the simulation. Dashed vertical line marks the year when the trait optimum changed. (d)
Distributions of the summary statistic   (change in trait mean 2015-2011) under
varying model scenarios. Colored vertical lines mark posterior means for each scenario.

Discussion

Extreme environmental perturbations offer an opportunity to examine rapid evolution in

natural populations. We used a resurrection experiment to test whether native plant populations

rapidly evolved in response to a multi-year drought despite severe demographic declines. In one

population (S22), earlier flowering evolved by the end of the drought and continued to evolve in

the same direction after the drought ended. However, no other traits diverged at S22, and all

traits exhibited stasis in the second population, KYE. We detected substantial additive genetic

variance for multiple traits in both populations, including flowering phenology, suggesting that

responses to selection were not constrained by limited genetic variation. Evolutionary

simulations integrated with our demographic and experimental data suggested that seed banks

constrained evolutionary responses in both populations. In KYE, the seed bank had a stronger

effect resulting in evolutionary stasis; whereas in S22, the seed bank had a weaker effect, which

facilitated closer tracking of changing phenotypic optima.

Why does rapid evolution occur in some populations but not others? One hypothesis is

that a lack of quantitative genetic variation limits adaptive responses even when selection is

intense. However, we found substantial V_A and high heritability for flowering phenology in both

populations. In fact, KYE contained twice the additive genetic variance in flowering time

compared to S22. Thus, these results are inconsistent with the hypothesis that genetic variation

prevented rapid evolution. Other studies have similarly found that rapid evolution occurred in

some but not all studied populations [12–14]. However, estimates of V_A are rarely quantified in

this context and thus the importance of genetic constraints is poorly understood [but see 5]. It is

also worth noting that phenotypic variance in flowering phenology in S22 decreased 50% over

our study, consistent with a strong episode of selection. Thus, the erosion of genetic variation

due to this drought could constrain future responses to selection on phenology.

A more likely explanation for the lack of phenological evolution in KYE is that gene

flow from the seed bank reintroduced alleles that were maladaptive during the drought.

Simulations suggested that decreased germination and increased seed survival rates resulted in a

seed bank at KYE that had a stronger effect on responses to selection compared to S22. The

recruitment and mating of older individuals from the seed bank essentially caused maladaptive

gene flow through time (Fig. 4c). Similarly, a third population grown during the refresher

generation showed no significant evolutionary response to the drought (Fig. S4) and was

estimated in [42] to have seed bank vital rates similar to KYE (low germination, high seed

survival). It would be fruitful for future resurrection studies to explicitly sample populations

across a range of seed bank vital rates to further dissect seed bank effects on adaptation to

climate-change-induced selective episodes. Additionally, seed banks can serve to buffer

populations from extinction during times of environmental stress [64,65], and the interplay

between these two contrasting consequences of seed banks for populations is an area ripe for

study [22,28].

Of course, "all models are wrong" [66], and our simulation models do not perfectly

reflect reality for these natural populations. But they do demonstrate that, for a population with

substantial genetic variance for a trait under selection during an environmental perturbation with

severe demographic consequences, like our populations during the megadrought, selection is

expected to effectively drive rapid phenotypic evolution toward new trait optima. Seed banks can

greatly constrain this evolution and produce results in line with our own empirical data, with

parameter estimates that follow patterns of empirically estimated seed vital rates. However, the

hypotheses tested above are not exhaustive, and it is worth exploring alternative explanations for

the lack of evolutionary change in phenology at KYE. Trait plasticity could allow a population to

weather an environmental perturbation with little or no evolutionary change [67]. However,

plastic responses in flowering time were nearly identical for our two populations (Fig. S10),

suggesting that greater plasticity in KYE cannot explain its evolutionary stasis. Spatial gene flow

among populations could potentially oppose allele frequency changes favored by selection.

However, gene flow is unlikely to be strong enough over this short time window to prevent

evolutionary change, especially given that our populations are isolated from others (>1

kilometer) and lack adaptations for long-distance dispersal. It is also possible that advanced

phenology simply did not confer fitness benefits during drought at KYE, though this would be

contrary to both general trends in plants [44] and the well-documented relationship between

early flowering time and adaptation to aridity in *C. xantiana* and close relatives [37,45,61,68–

70]. It is also interesting that, in contrast to expectations, empirical and simulation results

suggested that phenotypic optima did not return to their pre-drought level after the drought

ended. This insight is consistent with the observation that population demography did not

rebound after drought as one might expect (Fig. 1). Although the causes of this trend are unclear,

it could be related to the timing of precipitation [e.g., most precipitation fell early in the season,

which seems to promote germination but not survival in *C. x. xantiana* (DM, JB, pers. obs.)].

523   The evolution of earlier flowering at S22 is consistent with post-drought resurrection

studies in *Brassica rapa* [5,33], and likely reflects a strategy of drought escape [reviewed in

44,71,72]. Past work on *C. x. xantiana* has shown that Q_{ST} for flowering phenology is more than

twice as high ($Q_{ST} = 0.7$) than Q_{ST} for five other ecologically-important traits [68]. This result

suggests that flowering phenology is often the target of spatially-variable selection and readily

evolves, as has been frequently observed in many systems [5,44,73,74]. Earlier flowering may

evolve more readily than other traits under selection if its genetic architecture is comparatively

simple [e.g., 71], which can quicken responses to selection relative to more highly polygenic

traits. The evolutionary lability of flowering phenology could also be due to its direct tie to

assortative mating [75] — early flowering individuals tend to mate with other early flowering

individuals. Theoretical work has shown that when there is directional selection on flowering

time, positive assortative mating can increase genetic variation and the rate of phenotypic

evolution compared to scenarios with random mating [76,77].

A population's response to environmental change will be determined by the interplay of

demography, genetics, selection, and stochastic processes. Here, we have shown that flowering

phenology rapidly evolved during and after a severe drought in one population. Whereas, in a

second population we observed no sign of rapid evolution despite similar environmental stress,

demographic decline, and even greater additive genetic variance in phenology. In both

populations, our results are consistent with the hypothesis that gene flow through time via seed

banks either slowed or prevented rapid evolution. Our results are inconsistent with the

hypotheses that plasticity or limited genetic variation constrained responses to selection. While

rapid evolution in response to environmental perturbations has been repeatedly demonstrated,

few studies have tackled the problem of why evolutionary stasis is so commonly observed across

studies and systems. Our results emphasize that studies of adaptation in plant populations

experiencing environmental change should consider both above and belowground processes

affecting population demography and fitness. As we seek to build more predictive models of the

evolutionary process, the synthesis of demographic, environmental, and quantitative genetic data

will be invaluable for understanding where and when rapid evolution is likely to occur.

Acknowledgments

We thank Monica Geber and Vince Eckhart for their assistance with collection of seeds, demographic and environmental data, and for their comments on a previous version of this manuscript. We thank Eric Bakken, Ryan Allen, Hannah Littel, Samantha Sorg, and Adam Kostanecki for assistance with experiments. Conversations with Ruth Shaw and Seema Sheth were invaluable in the design of the experiments.

References

1. Grant PR, Grant BR, Huey RB, Johnson MTJ, Knoll AH, Schmitt J. 2017 Evolution caused

by extreme events. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **372**.
(doi:10.1098/rstb.2016.0146)

2.   Donihue CM, Herrel A, Fabre A-C, Kamath A, Geneva AJ, Schoener TW, Kolbe JJ, Losos
JB. 2018 Hurricane-induced selection on the morphology of an island lizard. *Nature* **560**,
88–91.

3.   Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV. 2017
Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole
lizard. *Science* **357**, 495–498.

4.   Grant PR, Grant BR. 2002 Unpredictable evolution in a 30-year study of Darwin's finches.
Science **296**, 707–711.

5.   Franks SJ, Sim S, Weis AE. 2007 Rapid evolution of flowering time by an annual plant in
response to a climate fluctuation. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 1278–1282.

6.   Shaw RG, Etterson JR. 2012 Rapid climate change and the rate of adaptation: insight from
experimental quantitative genetics. *New Phytol.* **195**, 752–765.

7.   Peschel AR, Boehm EL, Shaw RG. 2021 Estimating the capacity of Chamaecrista
fasciculata for adaptation to change in precipitation. *Evolution* **75**, 73–85.

8.   Geber MA, Dawson TE. 1993 Evolutionary responses of plants to global change. In *Biotic*
interactions and global change (ed Kareiva, PM and Kingsolver, JG and Huey, RB), pp.
179–197. Sunderland, Mass.: Sinauer.

9.   Etterson JR, Franks SJ, Mazer SJ, Shaw RG, Gorden NLS, Schneider HE, Weber JJ,
Winkler KJ, Weis AE. 2016 Project Baseline: An unprecedented resource to study plant
evolution across space and time. *Am. J. Bot.* **103**, 164–173.

10.   Franks SJ, Hamann E, Weis AE. 2018 Using the resurrection approach to understand
contemporary evolution in changing environments. *Evol. Appl.* **11**, 17–28.

11.   Dickman EE, Pennington LK, Franks SJ, Sexton JP. 2019 Evidence for adaptive responses
to historic drought across a native plant species range. *Evol. Appl.* (doi:10.1111/eva.12803)

12.   Anstett DN, Branch HA, Angert AL. 2021 Regional differences in rapid evolution during
severe drought. *Evol. Lett.* (doi:10.1002/evl3.218)

13.   Wooliver R, Tittes SB, Sheth SN. 2020 A resurrection study reveals limited evolution of
thermal performance in response to recent climate change across the geographic range of
the scarlet monkeyflower. *Evolution* (doi:10.1111/evo.14041)

14.   Kooyers NJ *et al.* 2021 Population responses to a historic drought across the range of the
common monkeyflower (Mimulus guttatus). *Am. J. Bot.* **108**, 284–296.

15.   Vtipil EE, Sheth SN. 2020 A resurrection study reveals limited evolution of phenology in

622   response to recent climate change across the geographic range of the scarlet monkeyflower.
623   *Ecol. Evol.* **10**, 14165–14177.

624 16. Haldane JBS. 1957 The cost of natural selection. *J. Genet.* **55**, 511.

625 17. Gomulkiewicz R, Houle D. 2009 Demographic and genetic constraints on evolution. *Am.*
626   *Nat.* **174**, E218-29.

627 18. Bell G. 2017 Evolutionary Rescue. *Annu. Rev. Ecol. Evol. Syst.* **48**, 605–627.

628 19. Lynch M, Lande R. 1993 Evolution and extinction in response to environmental change. In
629   *Biotic interactions and global change* (ed Kareiva, PM and Kingsolver, JG and Huey, RB),
630   pp. 234–250. Sunderland, Mass.: Sinauer.

631 20. Hufbauer RA, Szűcs M, Kasyon E, Youngberg C, Koontz MJ, Richards C, Tuff T,
632   Melbourne BA. 2015 Three types of rescue can avert extinction in a changing environment.
633   *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10557–10562.

634 21. Lindsey HA, Gallie J, Taylor S, Kerr B. 2013 Evolutionary rescue from extinction is
635   contingent on a lower rate of environmental change. *Nature* **494**, 463–467.

636 22. Evans MEK, Dennehy JJ. 2005 Germ banking: bet-hedging and variable release from egg
637   and seed dormancy. *Q. Rev. Biol.* **80**, 431–451.

638 23. Kalisz S, McPeek MA. 1992 Demography of an age-structured annual: Resampled
639   projection matrices, elasticity analyses, and seed bank effects. *Ecology* **73**, 1082–1093.

640 24. Evans MEK, Ferrière R, Kane MJ, Venable DL. 2007 Bet hedging via seed banking in
641   desert evening primroses (Oenothera, Onagraceae): demographic evidence from natural
642   populations. *Am. Nat.* **169**, 184–194.

643 25. Gottlieb LD. 1974 Genetic Stability in a Peripheral Isolate of Stephanomeria exigua ssp.
644   coronaria That Fluctuates in Population Size. *Genetics* **76**, 551–556.

645 26. Epling C, Lewis H, Ball FM. 1960 The Breeding Group and Seed Storage: A Study in
646   Population Dynamics. *Evolution* **14**, 238–255.

647 27. Templeton AR, Levin DA. 1979 Evolutionary Consequences of Seed Pools. *Am. Nat.* **114**,
648   232–249.

649 28. Hairston NG Jr, De Stasio BT Jr. 1988 Rate of evolution slowed by a dormant propagule
650   pool. *Nature* **336**, 239–242.

651 29. Levin DA. 1990 The Seed Bank as a Source of Genetic Novelty in Plants. *Am. Nat.* **135**,
652   563–572.

653 30. Blows MW, Hoffmann AA. 2005 A reassessment of genetic limits to evolutionary change.
654   *Ecology* **86**, 1371–1384.

655   31.   Barton N, Partridge L. 2000 Limits to natural selection. *Bioessays* **22**, 1075–1084.

656   32.   Falconer DS. 1996 *Introduction to quantitative genetics*. 4th edn. Essex, England: Pearson
657   Education.

658   33.   Hamann E, Weis AE, Franks SJ. 2018 Two decades of evolutionary changes in Brassica
659   rapa in response to fluctuations in precipitation and severe drought. *Evolution*
660   (doi:10.1111/evo.13631)

661   34.   Cook BI, Ault TR, Smerdon JE. 2015 Unprecedented 21st century drought risk in the
662   American Southwest and Central Plains. *Sci Adv* **1**, e1400082.

663   35.   Williams AP, Cook BI, Smerdon JE. 2022 Rapid intensification of the emerging
664   southwestern North American megadrought in 2020–2021. *Nat. Clim. Chang.* **12**, 232–234.

665   36.   Williams AP, Cook ER, Smerdon JE, Cook BI, Abatzoglou JT, Bolles K, Baek SH, Badger
666   AM, Livneh B. 2020 Large contribution from anthropogenic warming to an emerging North
667   American megadrought. *Science* **368**, 314–318.

668   37.   Eckhart VM, Geber MA. 1999 Character variation and geographic range in Clarkia xantiana
669   (Onagraceae): breeding system and phenology distinguish two common subspecies.
670   *Madrono*

671   38.   Webb RW. 1936 Kern Canyon Fault, Southern Sierra Nevada. *J. Geol.* **44**, 631–638.

672   39.   Brossy CC *et al.* 2012 Map of the late Quaternary active Kern Canyon and Breckenridge
673   faults, southern Sierra Nevada, California. *Geosphere* **8**, 581–591.

674   40.   Lewis H. 1962 Catastrophic Selection as a Factor in Speciation. *Evolution* **16**, 257–271.

675   41.   Eckhart VM, Geber MA, Morris WF, Fabio ES, Tiffin P, Moeller DA. 2011 The geography
676   of demography: long-term demographic studies and species distribution models reveal a
677   species border limited by adaptation. *Am. Nat.* **178**, S26-43.

678   42.   Siegmund G-F, Moeller DA, Eckhart VM, Geber MA. 2022 Bet hedging is not sufficient to
679   explain intraspecific variation in germination patterns of a winter annual plant. *bioRxiv*. ,
680   2022.09.15.508102. (doi:10.1101/2022.09.15.508102)

681   43.   Briscoe Runquist RD, Geber MA, Pickett-Leonard M, Moeller DA. 2017 Mating System
682   Evolution under Strong Pollen Limitation: Evidence of Disruptive Selection through Male
683   and Female Fitness in Clarkia xantiana. *Am. Nat.* **189**, 549–563.

684   44.   Kooyers NJ. 2015 The evolution of drought escape and avoidance in natural herbaceous
685   populations. *Plant Sci.* **234**, 155–162.

686   45.   Burnette TE, Eckhart VM. 2021 Evolutionary divergence of potential drought adaptations
687   between two subspecies of an annual plant: Are trait combinations facilitated, independent,
688   or constrained? *Am. J. Bot.* (doi:10.1002/ajb2.1607)

46. Ogburn RM, Edwards EJ. 2012 Quantifying succulence: a rapid, physiologically
meaningful metric of plant water storage. *Plant Cell Environ.* **35**, 1533–1542.

47. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671–675.

48. Weis AE. 2018 Detecting the "invisible fraction" bias in resurrection experiments. *Evol.*
Appl. **11**, 88–95.

49. R Foundation For Statistical Computing. 2021 *R: A language and environmnet for*
statistical computing.

50. Wickham H, Girlich M. 2022 tidyr: Tidy Messy Data. See https://tidyr.tidyverse.org,
https://github.com/tidyverse/tidyr.

51. Wickham H, François R, Henry L, Müller K. 2022 dplyr: A Grammar of Data
Manipulation. See https://dplyr.tidyverse.org, https://github.com/tidyverse/dplyr.

52. Wickham H. 2009 *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-
Verlag.

53. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

54. Fox J, Weisberg S. 2011 *An R Companion to Applied Regression*. Thousand Oaks, CA:
Sage.

55. Lenth R. 2018 Emmeans: Estimated marginal means, aka least-squares means. *R Package*
Version **1**.

56. Hadfield JD. 2010 MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. *J. Stat. Softw.* **33**, 1–22.

57. Kruuk LEB. 2004 Estimating genetic parameters in natural populations using the 'animal
model.' *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **359**, 873–890.

58. Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, Kruuk LEB,
Nussey DH. 2010 An ecologist's guide to the animal model. *J. Anim. Ecol.* **79**, 13–26.

59. de Villemereuil P. 2018 Quantitative genetic methods depending on the nature of the
phenotypic trait. *Ann. N. Y. Acad. Sci.* **1422**, 29–47.

60. Haller BC, Messer PW. 2022 SLiM 4: Multispecies eco-evolutionary modeling. *Am. Nat.*
(doi:10.1086/723601)

61. Anderson JT, Eckhart VM, Geber MA. 2015 Experimental studies of adaptation in Clarkia
xantiana. III. Phenotypic selection across a subspecies border. *Evolution* **69**, 2249–2261.

62. Beaumont MA. 2010 Approximate Bayesian Computation in Evolution and Ecology. *Annu.*

Rev. Ecol. Evol. Syst. (doi:10.1146/annurev-ecolsys-102209-144621)

63. Csilléry K, François O, Blum MGB. 2012 abc: an R package for approximate Bayesian
computation (ABC). *Methods Ecol. Evol.* **3**, 475–479.

64. Kalisz S, McPeek MA. 1993 Extinction dynamics, population growth and seed banks : An
example using an age-structured annual. *Oecologia* **95**, 314–320.

65. Pake CE, Venable DL. 1996 Seed Banks in Desert Annuals: Implications for Persistence
and Coexistence in Variable Environments. *Ecology* **77**, 1427–1435.

66. Box GEP. 1976 Science and Statistics. *J. Am. Stat. Assoc.* **71**, 791–799.

67. Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LEB, Sheldon BC. 2008
Adaptive phenotypic plasticity in response to climate change in a wild bird population.
Science **320**, 800–803.

68. Gould B, Moeller DA, Eckhart VM, Tiffin P, Fabio E, Geber MA. 2014 Local adaptation
and range boundary formation in response to complex environmental gradients across the
geographical range of Clarkia xantiana ssp . xantiana. *J. Ecol.* **102**, 95–107.

69. Eckhart VM, Geber MA, McGuire CM. 2004 Experimental studies of adaptation in Clarkia
xantiana. I. Sources of trait variation across a subspecies border. *Evolution* **58**, 59–70.

70. Mazer SJ, Dudley LS, Hove AA, Emms SK, Verhoeven AS. 2010 Physiological
Performance in Clarkia Sister Taxa with Contrasting Mating Systems: Do Early-Flowering
Autogamous Taxa Avoid Water Stress Relative to Their Pollinator-Dependent
Counterparts? *Int. J. Plant Sci.* **171**, 1029–1047.

71. Fulgione A *et al.* 2022 Parallel reduction in flowering time from de novo mutations enable
evolutionary rescue in colonizing lineages. *Nat. Commun.* **13**, 1461.

72. Metz J, Lampei C, Bäumler L, Bocherens H, Dittberner H, Henneberg L, de Meaux J,
Tielbörger K. 2020 Rapid adaptive evolution to drought in a subset of plant traits in a large-
scale climate change experiment. *Ecol. Lett.* (doi:10.1111/ele.13596)

73. Lustenhouwer N, Wilschut RA, Williams JL, van der Putten WH, Levine JM. 2018 Rapid
evolution of phenology during range expansion with recent climate change. *Glob. Chang.*
Biol. **24**, e534–e544.

74. Anderson JT, Inouye DW, McKinney AM, Colautti RI, Mitchell-Olds T. 2012 Phenotypic
plasticity and adaptive evolution contribute to advancing flowering phenology in response
to climate change. *Proc. Biol. Sci.* **279**, 3843–3852.

75. Franks SJ, Weis AE. 2009 Climate change alters reproductive isolation and potential gene
flow in an annual plant. *Evol. Appl.* **2**, 481–488.

76. Fox GA. 2003 Assortative mating and plant phenology: evolutionary and practical

consequences. *Evol. Ecol. Res.* **5**, 1–18.

77.   Godineau C, Ronce O, Devaux C. 2021 Assortative mating can help adaptation of flowering
time to a changing climate: Insights from a polygenic model. *J. Evol. Biol.*
(doi:10.1111/jeb.13786)