Bacterial strain-dependent dissociation of cell recruitment and cell-to-cell spread in early

M. tuberculosis infection

*B. Shoshana Zha[a,d], *Ludovic Desvignes[b], Tawania J. Fergus[b], Amber Cornelius[b], Tan-

Yun Cheng[c], D. Branch Moody[c], and Joel D. Ernst[d]#

[a]Division of Pulmonary, Critical Care, Allergy and Sleep Medicine, Department of

Medicine, University of California, San Francisco, California, USA

[b]Department of Medicine, New York University School of Medicine, New York, New

York, USA

[c]Division of Rheumatology, Immunity and Inflammation, Department of Medicine,

Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

[d]Division of Experimental Medicine, Department of Medicine, University of California,

San Francisco, California, USA

*Co-first authors; the order was determined by the leadership role in writing the paper

#Corresponding Author: Joel Ernst, MD

1001 Potrero Ave, Room 601

UCSF Box 1234

joel.ernst@ucsf.edu

Running title: *M. tuberculosis* dissemination and T cell priming

Keywords: Mycobacterium tuberculosis, Innate Response, T cell priming, Beijing strain

Word Count, Abstract: 139

Word Count, Text: 4,242

Abbreviations

Early secretory antigenic target secretion system (ESX), Mediastinal lymph node

(MdLN), Mononuclear phagocyte (MNP), Monocyte-derived dendritic cells (moDC),

Monocyte-derived recruited macrophages (RM), *Mycobacterium tuberculosis* (*M.*

tuberculosis), neutrophils (neut), Phthiocerol dimycocerosates (PDIM), Phenolic

glycolipids (PGL)

ABSTRACT

In the initial stage of respiratory infection, *Mycobacterium tuberculosis* traverses from

alveolar macrophages to phenotypically diverse monocyte-derived phagocytes and

neutrophils in the lung parenchyma. Here, we compare the *in vivo* kinetics of early

bacterial growth and cell-to-cell spread of two strains of *M. tuberculosis*: a lineage 2

strain, 4334, and the widely studied lineage 4 strain H37Rv. Using flow cytometry, live

cell sorting of phenotypic subsets, and quantitation of bacteria in cells of the distinct

subsets, we found that 4334 induces less leukocyte influx into the lungs but

demonstrates earlier population expansion and cell-to-cell spread. The earlier spread of

4334 to recruited cells, including monocyte-derived dendritic cells, is accompanied by

earlier and greater magnitude of $CD4^+$ T cell activation. The results provide evidence

that strain-specific differences in interactions with lung leukocytes can shape adaptive

immune responses *in vivo*.

IMPORTANCE

Tuberculosis is a leading infectious disease killer world-wide and is caused by

Mycobacterium tuberculosis. After exposure to *M. tuberculosis,* outcomes range from

apparent elimination to active disease. Early innate immune responses may contribute

to differences in outcomes, yet it is not known how bacterial strains alter the early

dynamics of innate immune and T cell responses. We infected mice with distinct strains

of *M. tuberculosis* and discovered striking differences in innate cellular recruitment, cell-

to-cell spread of bacteria in the lungs, and kinetics of initiation of antigen-specific CD4 T

cell responses. We also found that *M. tuberculosis* can spread beyond alveolar

macrophages even before a large influx of inflammatory cells. These results provide

evidence that distinct strains of *M. tuberculosis* can exhibit differential kinetics in cell-to-

cell spread which is not directly linked to early recruitment of phagocytes but is

subsequently linked to adaptive immune responses.

INTRODUCTION

Mycobacterium tuberculosis is a facultative intracellular bacterium that resides in

tissue mononuclear phagocytes (MNP) and granulocytes (1, 2). Previous investigations

have shown that *M. tuberculosis* first enters alveolar macrophages (3, 4). After

replication in alveolar macrophages, the bacteria subsequently spread to multiple

subsets of phagocytes in the lung parenchyma, largely recruited from circulating

monocytes (5–7). Subsequently, monocyte-derived lung dendritic cells (moDC) acquire

the bacteria and transport them to draining lymph nodes (2), where they transfer

antigens to lymph node resident dendritic cells that prime antigen-specific T cells (8, 9).

After proliferating and differentiating, effector T cells traffic to the lungs and arrest

progression of the infection (10). *M. tuberculosis* antigen-specific T cell responses

require 14-17 days to develop after aerosol infection of mice with the commonly-used

H37Rv strain (2, 10), and an average of 6 weeks is required for development of

adaptive immune responses after infection in humans (11, 12). Considerable evidence

indicates that the rate-limiting step in initiating *M. tuberculosis* antigen-specific CD4 T

cell responses is acquisition of the bacteria by dendritic cells in the lung, and transport

of live bacteria to local draining lymph nodes (8, 10, 13–16).

The mechanisms of cell-to-cell spread of *M. tuberculosis* are incompletely

understood, although evidence suggests a role for necrosis of infected cells and release

of viable bacteria that are subsequently ingested by other cells (17). Virulent strains of

M. tuberculosis can inhibit apoptosis (18) through multiple mechanisms including

upregulation of the antiapoptotic protein Mcl-1 (19), inhibition of NOX2-induced reactive

oxygen species formation (20), upregulation of soluble TNF receptor 2 (21), and

inhibition of apoptotic envelope stabilization (22), although induction of apoptosis still

occurs and is important for control of *M. tuberculosis in vivo* (23). Conversely, virulent

M. tuberculosis can promote necrosis through mitochondrial transmembrane potential

disruption (24), depletion of host cellular NAD^+ (25), and inhibition of plasma membrane

repair with induction of lipoxin A4 (17, 26), among other mechanisms.

The early secretory antigenic target secretion system (ESX)-1, a type VII

secretion system encoded by the RD1 locus of *M. tuberculosis* (27, 28), is implicated in

multiple mechanisms for its role in *M. tuberculosis* virulence (29–32). These

mechanisms include recruitment of macrophages (33), activation of neutrophils (34),

and secretion of immunodominant T cell antigens (ESAT-6 and CFP-10) (35–37). In

addition, ESX-1 has been linked to induction of macrophage necrosis (38–40). ESX-1's

role in cellular recruitment and induction of cell death leads to the hypothesis that it

plays a key role in *M. tuberculosis* cell-to-cell transfer.

The majority of studies of early innate responses and bacterial cell-to-cell transfer

focus on *M. tuberculosis* lineage 4 strains, especially H37Rv or Erdman. However, there

are 6 other human-adapted lineages (41) which vary genetically, phenotypically, and

degree of induction of inflammatory responses *in vitro* (42). Lineage 2, which includes

the Beijing family, is thought to have enhanced pathogenicity (i.e. ability to cause

disease) and less protection by BCG vaccination (43, 44). The Beijing family is defined

by region of difference (RD) 105, and has 5 sublineages (45). Sublineage 207 has

demonstrated increased pathogenicity in guinea pigs, and the strain 4334 within this

sublineage was the source of more secondary cases of TB than any other in San

Francisco, California (45).

Increasing evidence indicates that distinct *M. tuberculosis* strains exhibit

differential interactions with innate immune responses, as reflected by differences in

cytokine release (46, 47) and monocyte activation (48). The goal of this work was to

characterize the kinetics of *M. tuberculosis* growth and cell-to-cell spread early in

infection, to compare the results between two distinct strains of *M. tuberculosis*, and to

assess the relationship of cell recruitment and bacterial spread to initiation of CD4 T cell

responses. We found that H37Rv and 4334 differ in their recruitment of leukocytes to

the lungs, spread from alveolar macrophages to other leukocytes, and the kinetics of

antigen-specific CD4 T cell activation. Our work demonstrates strain dependent

dissociation of inflammatory cell recruitment and bacterial cell-to-cell spread, and that

the timing of cell-to-cell spread impacts the dynamics of antigen-specific CD4 T cell

responses.

RESULTS

Differential early cell dynamics after aerosol infection with** M. tuberculosis **strains

H37Rv, 4334, and H37RvΔRD1.

To test the hypothesis that distinct *M. tuberculosis* strains vary in their initial

interactions with the host, we quantitated live bacteria in the lungs of mice infected with

H37Rv, 4334, or H37Rv lacking the virulence locus RD1 (H37RvΔRD1) which encodes

the ESX-1 type VII secretion system (27, 28) implicated in cell recruitment and

inflammation (29, 30, 33, 37). Mice were infected via aerosol with 200-300 colony

forming units (CFU) of each bacterial strain and analyzed at frequent time points 1-14

days post-infection, a short time frame that is known to be prior to development of

detectable T cell responses. H37Rv exhibited a lag of approximately 72h in which the

number of bacteria did not increase in the lungs, similar to that reported in other studies

(49, 50). In contrast, strain 4334 grew without a similar lag, and exhibited an increase in

the lung bacterial burden that was already detectable by day 3 (Fig. 1A). After day 6, the

populations of both strains expanded steadily during the first 2 weeks of infection and

were equivalent by day 10 post-infection. In contrast to the two ESX-1-replete strains,

H37RvΔRD1 experienced a prolonged lag in expansion (28). The three strains also

differed in their induction of increased lung cellularity: H37Rv induced the most marked

increases in cellularity; 4334 responses were intermediate; and H37RvΔRD1 induced

no detectable changes in the number of lung cells in the initial 14 days of infection (Fig.

1B).

We next compared and characterized the cell populations in the lungs during the

first two weeks of infection with the three *M. tuberculosis* strains. For this, we used

established cell phenotypic markers and flow cytometry to differentiate subsets in the

CD45$^+$ cell population: CD11b$^{neg/lo}$CD11c^{pos} alveolar macrophages; CD11b^{pos}CD11c^{pos}

which includes moDC; Gr-1^{hi}CD11c^{neg} neutrophils; Gr-1^{int}CD11c^{neg} monocytes; and Gr-

1^{neg}CD11c^{neg} monocyte-derived recruited macrophages (RM) (Fig. S1) (6). As shown in

Figure 2A, CD11b$^{neg/lo}$CD11c^{pos} alveolar macrophages and CD11b^{pos}CD11c^{pos} moDC

varied little during the first two weeks and did not differ in lungs of mice infected with

4334 or H37Rv. While Gr-1^{hi}CD11c^{neg} neutrophils did not increase in the lungs during

the first two weeks of infection with any strain, numbers were consistently lower in the

lungs of mice infected with 4334 and H37RvΔRD1 than with H37Rv. The most striking

difference between strains was in Gr-1^{int}CD11c^{neg} monocytes and Gr-1^{neg}CD11c^{neg} RM.

As previously reported (2), mice infected with H37Rv showed a progressive increase in

Gr-1^{int}CD11c^{neg} monocytes and Gr-1^{neg}CD11c^{neg} RM in the lungs over the first two

weeks, while neither population increased significantly in lungs of mice infected with

4334 or H37RvΔRD1. These findings indicate that the initial cellular inflammatory

response to infection with virulent *M. tuberculosis* varies considerably when distinct

bacterial strains infect genetically homogeneous hosts.

Mycobacterium tuberculosis *strain-dependent dynamics of spread to recruited*

lung cells.

To compare the spread of H37Rv and 4334 from alveolar macrophages to

recruited inflammatory cells during the initial phase of infection, we used BSL3-

contained flow cytometry sorting of live cells belonging to the previously-identified lung

leukocyte subsets (2, 5, 8). We sorted the individual cell populations from infected lungs

at frequent intervals and plated the resulting sorted cell fractions on solid media for

bacterial quantitation. Consistent with recently-published results generated by flow

cytometry detection of cells that harbored fluorescent protein-expressing *M. tuberculosis*

(3, 4), we found live bacteria exclusively in alveolar macrophages for the first days post-

infection (Fig. 2B, Fig. S2B). On day 6 post-infection, live *M. tuberculosis* were

detectable in Gr-1^{hi}CD11c^{neg} neutrophils in 2 of the 10 mice infected with strain 4334,

but in none of the 15 mice infected with H37Rv, and the average number of live bacteria

per cell remained higher in 4334 compared to H37Rv over time (Fig. S2A). Similarly, on

day 8 post-infection, 2 of the 10 mice in the group infected with 4334 and none of the 15

infected with H37Rv had detectable live *M. tuberculosis* in Gr-1^{int}CD11c^{neg} monocytes,

again with higher average number of live bacteria per cell until day 14. Within

CD11b^{pos}CD11c^{pos} moDC, a significantly higher number of live *M. tuberculosis* were

detected for strain 4334 than H37Rv at 8 days post-infection, but the average number of

live bacteria per cell in 4334 was higher than H37Rv only until day 10. In Gr-

1^{neg}CD11c^{neg} RM, the number of live *M. tuberculosis* per cell was not significantly

different between the two strains, although the number of live bacteria per cell was

slightly (but not statistically significantly) higher in 4334 than H37Rv at days 10 and 14

post infection.

H37RvΔRD1 expanded in alveolar macrophages at a rate similar to wild-type

H37Rv between days 1 and 10. H37RvΔRD1 also spread to CD11b^{pos}CD11c^{pos} moDC

and Gr-1^{hi}CD11c^{neg} neutrophils in quantities comparable to that of wild-type H37Rv by

day 10. However, H37RvΔRD1 did not expand further between days 10 and 14. Of

note, live H37RvΔRD1 was recoverable in only 2 out of 5 mice in Gr-1^{int}CD11c^{neg}

monocytes at day 14, and one mouse out of 5 at day 8 in Gr-1^{neg}CD11c^{neg} RM.

One known difference with functional importance between *M. tuberculosis*

isolates is the presence or absence of phenolic glycolipids (PGL), which are cell wall

lipids that modulate virulence (51). PGLs have been implicated in macrophage

recruitment (52), dendritic cell uptake, and reduction in inflammatory pathways (53).

PGLs are absent from H37Rv and many clinical isolates but are expressed in HN878, a

prototypical strain used to study the Beijing family. As 4334 is a member of the Beijing

family (44), we determined if 4334 has PGLs. Using electrospray ionization-quadrupole

time-of-flight-mass spectrometry (ESI-QTOF-MS), there was no detectable PGL in

either 4334 and H37Rv (Fig. 3A,B) under conditions in which PGL was detected in

HN878, and the structurally related long chain polyketide phthiocerol dimycocerosates

(PDIM) was ionized and detected. This result excludes PGL production as the cause of

differential cell-to-cell transfer in these two strains.

***Mycobacterium tuberculosis* 4334 is associated with less alveolar macrophage**

death than H37Rv.

Considering the absence of a lag and immediate expansion of 4334 in alveolar

macrophages *in vivo*, we examined the intracellular growth dynamics of 4334 and

H37Rv. We infected *ex vivo* murine alveolar macrophages with 4334 or H37Rv at an

MOI of 1, and quantitated CFU at 48 and 72h post-infection. In cultured alveolar

macrophages, H37Rv did not expand compared with the input by 48h but expanded

approximately 5-fold by 72h. In contrast, 4334 expanded 4-fold by 48h, and nearly 10-

fold by 72h (Fig. 4A). These growth kinetics mimic the initial phase of infection *in vivo*, in

which there was a delay in growth of H37Rv for the first 3 days while 4334 expanded

without a similar lag (Fig. 1B).

To determine if the apparent difference in growth of 4334 and H37Rv in alveolar

macrophages was due to differential cell death, we quantitated infection, early

apoptosis, and late cell death 48h post-infection. Using *M. tuberculosis* strains

expressing DsRed fluorescent protein and flow cytometry, we observed a higher

frequency of infection in cultured alveolar macrophages by 4334 than H37Rv (Fig.

4B,C), consistent with the results obtained by CFU plating. We used a viability marker

(Zombie Aqua, ZA) and a phosphatidylserine-binding marker (Annexin V, AV) to

quantitate the viability of alveolar macrophages infected *ex vivo* (Fig. 4B). In this

scheme, ZA^-/AV^- cells are viable, ZA^-/AV^+ cells are considered apoptotic, ZA^+/AV^- are

necrotic, and ZA^+/AV^+ represent late death, regardless of the mode of death (54). We

found significantly less cell death in alveolar macrophages infected by either strain

compared to uninfected controls, with quantitatively lower rates of cell death with 4334

compared to H37Rv. Moreover, infection with strain 4334 was associated with a lower

frequency of apoptotic cells than H37Rv (Fig. 4D). There was no significant difference in

apoptosis or late death between the two strains in bystander cells (that are defined as

cells in the same well but lacking DsRed-expressing bacteria).

Cell-to-cell spread of *M. tuberculosis* is thought to involve bacteria released by

dying cells. However, our results indicate that 4334 spreads more efficiently despite

being associated with lower rates of death of alveolar macrophages. We therefore

sought alternative explanations for the early spread of 4334. Since the ESX-1 type VII

secretion system has also been implicated in cell-to-cell spread of *M. tuberculosis*, we

used liquid chromatography-mass spectrometry (LC-MS/MS) to analyze culture filtrate

(i.e. secreted) proteins of 4334 and H37Rv. This revealed significantly higher levels of 3

ESX-1 substrates (EspA, EspB, EspF) and relative higher levels of 3 additional

substrates (EspC, ESAT-6, CFP-10) in culture filtrates of 4334 compared with H37Rv

(Fig. S3). In contrast, proteins secreted by non-ESX mechanisms (Ag85B and Ag85C)

did not differ between the two strains. The finding that cell-to-cell spread of *M.*

tuberculosis is associated with quantitative ESX-1 activity and not with the frequency of

cell death suggests an alternative mechanism of cell-cell spread and dissemination of

M. tuberculosis independent of cell death.

Mycobacterium tuberculosis *strain-dependent dynamics of antigen-specific CD4*

T cell priming.

M. tuberculosis induction of antigen-specific CD4 T cell responses requires

transport of live bacteria by migratory moDC from the lungs to the mediastinal draining

lymph node (8, 10, 13). Since we found more live bacteria in $CD11b^{pos}CD11c^{pos}$ moDC

from mice infected with 4334 than H37Rv as early as day 8 post-infection (Fig. 2B), we

hypothesized that this could result in earlier antigen-specific CD4 T cell priming in the

lung-draining lymph node. We first quantitated live bacteria in the mediastinal lymph

node (MdLN) on day 14 post-infection, corresponding to the onset of antigen-specific

CD4 T cell priming (10). This revealed approximately 3-fold more 4334 than H37Rv in

the lymph node (Fig 5A). We have previously established that a threshold number of

bacteria are required in the MdLN for priming of *M. tuberculosis* antigen-specific CD4 T

cells (10), and therefore hypothesized that 4334 might activate an earlier CD4 T cell

response than H37Rv. To test this, we adoptively transferred *M. tuberculosis* Ag85B-

specific TCR transgenic (P25TCR-Tg) naïve CD4 T cells (labeled with CellTrace Violet)

into mice 24h prior to aerosol infection. Examination of the adoptively transferred T cells

isolated from the MdLN revealed no proliferation at day 10 post-infection in mice

infected with either strain. In contrast, by day 14 post-infection, P25TCR-Tg CD4 T cells

had proliferated and expanded in mediastinal lymph nodes of mice infected with strain

4334 but not H37Rv (Fig. 5B-C). At day 17, P25TCR-Tg CD4 T cells had also

proliferated in mice infected with H37Rv, although at significantly lower frequency than

in mice infected with 4334.

IFNγ is essential for control of *M. tuberculosis* (55, 56). Therefore, we analyzed

both total IFNγ concentrations and CD4 T cell specific IFNγ production in the lungs of

infected mice of both strains. We found total IFNγ concentrations in lung homogenate

supernatants to be significantly higher in mice infected with 4334 than H37Rv at day 14

post-infection (Fig. S4A). We next quantitated IFNγ-producing T cells in the lungs (Fig.

S4B) and found that both endogenous and P25TCR-Tg IFNγ-producing T cells were

more abundant in mice infected with 4334 than H37Rv at day 14 post-infection, though

the difference was not statistically significant. By day 17 post-infection, P25TCR-Tg but

not endogenous, IFNγ-producing T cells were significantly more abundant in mice

infected with 4334 than H37Rv (Fig. 5D). Together, these results indicate that *M.*

tuberculosis strain 4334 spreads to moDC and initiates antigen-specific CD4 T cell

responses earlier than does strain H37Rv. This earlier spread to dendritic cells and T

cell priming happens despite strain 4334 inducing fewer inflammatory cells to the lung

early in infection, and is associated with accelerated priming of antigen-specific CD4 T

cells.

We investigated whether the more robust and earlier T cell response correlated

into superior long-term control of infection. Indeed, at 7 weeks post-infection, we found

a half-log reduction in CFU recovery from the lungs of mice infected with *M. tuberculosis*

strain 4334 versus H37Rv (Fig. S5A). Interestingly, while total cell numbers in the lungs

of mice infected with *M. tuberculosis* strain 4334 were significantly lower than H37Rv

infected mice (Fig. 2B), by 7 weeks this was reversed (Fig. S5B), and the total number

of lymphocytes appears to account for the difference in cellularity (Fig. S5D). The

increased cellularity also correlates with more granuloma-like lesions in the lungs of

mice infected with 4334 (Fig. S5C, S5E), resembling the pathologic findings reported in

guinea pigs (44).

DISCUSSION

The early host responses to pulmonary infection with *M. tuberculosis* are

beginning to be clarified (3, 4), although studies to date have not examined the potential

impact of bacterial strain diversity. We analyzed early cellular responses in mice

infected by aerosol with 4334, a lineage 2 strain (44), and compared this to responses

to H37Rv, from lineage 4. Using flow sorting of live lung cells, we compared cell

populations, bacterial growth, and bacterial spread from alveolar macrophages to

recruited cells in the first 2 weeks of infection. We found that 4334 induces less

inflammatory cell recruitment than H37Rv, resembling that of H37Rv lacking the RD1

locus. However, 4334 spreads beyond alveolar macrophages earlier and is present in

greater abundance in monocytes, neutrophils, and moDC. This results in earlier

trafficking of bacteria to the mediastinal lymph node, and in turn, is accompanied by

earlier and greater magnitude activation of Ag85B-specific CD4 T cells.

Recent studies have confirmed that alveolar macrophages are the first cells

infected in the lungs by *M. tuberculosis*, a fact that had long been speculated (6) but

only recently demonstrated (3, 4). Alveolar macrophages are initially permissive to *M.*

tuberculosis replication (4), likely through their inability to mount a large inflammatory

response while leading to the upregulation of self-preserving Nrf2 pathways (4) and lipid

metabolism (57). We did not find a difference in total $CD11b^{neg/lo}CD11c^{hi}$ alveolar

macrophages cell numbers after infection. However, there is an earlier expansion of

4334 with significantly higher CFU 8 days post-infection than in H37Rv infected mice,

due to a lag in initial H37Rv growth *in vivo* (49, 50). Since alveolar macrophages were

the only infected cells during this initial phase, we hypothesized that the differential

expansion of the two strains could be due to differences in intracellular replication in

alveolar macrophages. We confirmed that 4334 replicates at a higher rate than H37Rv

in cultured alveolar macrophages, and this is accompanied by a lower rate of cell death.

By 2 weeks after infection, alveolar macrophages have been found to translocate

from the alveolar space to the lung interstitium (3), providing the opportunity for spread

of *M. tuberculosis* to other cells. Extending our previous work as well as that of others

(3–5, 8), we observed spread of both H37Rv and 4334 to monocyte-derived dendritic

cells and macrophages. Here, we found differential spread to the $CD11b^{pos}CD11c^{pos}$

population, with 4334 appearing in this population at a higher rate than H37Rv 8 days

post-infection. Notably, this cell population includes moDC (5, 6), and although a small

percentage of alveolar macrophages also express CD11b (58, 59) with potential

upregulation during inflammation and infection (3, 60), those cells have not been found

to migrate to lymphoid tissues. Nonetheless, with the combined findings of an earlier

and higher rate of spread to neutrophils and monocyte-derived cells in mice infected

with 4334 versus H37Rv, our data implies greater spread of 4334 from alveolar

macrophages to recruited cells in spite of inducing recruitment of fewer inflammatory

cells than H37Rv.

The determinants and mechanisms of *M. tuberculosis* cell-to-cell spread are not

well understood, especially *in vivo*, but likely depend on multiple factors including host

cell recruitment, intracellular bacterial growth, cellular release, and survival in

extracellular spaces. In this work, we found that while 4334, a lineage 2 strain with

increased pathogenicity in guinea pigs (44) and the ability to induce higher levels of type

I interferons (61), recruits significantly fewer monocytes and cells that differentiated from

monocytes but demonstrates superior growth and/or survival rates in the cells it does

infect compared to H37Rv. This suggests that progression of infection is not solely

determined by cellular recruitment but also by the rates of intracellular replication and

transfer between host cells. Differential growth dynamics between *M. tuberculosis*

strains has been shown in multiple primary cell types, including human monocytes (62),

human monocyte-derived macrophages (63), and murine bone marrow-derived

macrophages (64). We hypothesize that the permissiveness of alveolar macrophages

for *M. tuberculosis* growth in the initial stage of infection (4, 57) maximizes the

intracellular growth variations between strains.

Multiple mechanisms can determine the intracellular survival and growth of *M.*

tuberculosis. We focused on differential host cell death since this is thought to play a

key role in cell-to-cell spread. Death-receptor induced apoptosis (23) and subsequent

efferocytosis (65) are key host mechanisms for control of *M. tuberculosis*. As opposed

to avirulent strains lacking the operon RD1 and thus functional ESX-1 machinery,

virulent strains of *M. tuberculosis* have been found to inhibit apoptosis (18) and promote

necrosis (66) through multiple mechanisms. Necrosis is hypothesized to be the major

mode of host cell death for bacterial cell-cell spread, as it allows release of bacilli into

the extracellular space for phagocytosis by permissive host cells, rather than being

contained in apoptotic vesicles and killed through efferocytosis (17, 33, 67, 68). In

contrast to that model, we found that 4334 sustains alveolar macrophage *ex vivo*

survival to a greater extent than does H37Rv, yet 4334 spreads more readily to other

cells in the lungs.

ESX-1 is implicated in the induction of cell death through the action of secreted

proteins including ESAT-6 (40), which is concentration dependent. We have shown that

4334 secretes higher levels of ESX-1 substrates compared to H37Rv, but this is not

correlated with activation of cell death of infected alveolar macrophages *ex vivo*. One

hypothesis of these discordant findings is the limitation of using *ex vivo* infection

modeling, however, the significant protection of alveolar macrophage viability in 4334-

infected cells over H37Rv-infected cells argues against differential induction of cell

death as a means of cell-to-cell spread. Rather, these results are compatible with a

model in which nonlytic release contributes to cell-to-cell spread of *M. tuberculosis* in

the lungs (70). Nonlytic release has been demonstrated in *Dictyostelium* amoebae,

where *M. marinum* and *M. tuberculosis* egress via ejection rather than lysis of the

amoebae (71), and *in vivo* in zebrafish where nonlytic cell-to-cell transfer of *M. marinum*

has been observed (72). Furthermore, *M. tuberculosis* can grow extracellularly (73, 74),

adding an additional possibility that 4334 survives extracellularly more effectively than

H37Rv, allowing uptake by diverse phagocytic cells. Teasing these differenitals is of

essential importance to understand the mechanism of tuberculosis cell spread.

We also found significantly more live 4334 *M. tuberculosis* than H37Rv in the

draining mediastinal lymph node 14 days post-infection. This correlates with the timing

and speed of 4334 spread to moDC, which transport *M. tuberculosis* to the draining

mediastinal lymph node (2, 9). Of note, we focused on CD11b^{pos}CD11c^{pos} DCs in this

study. During homeostasis, conventional DCs (cDCs) have been classified into two

populations in mice: cDC-1 (CD26$^+$CD11c$^+$CD103$^+$CD11b$^-$) and cDC-2

(CD26$^+$CD11b$^+$CD11c$^+$CD172a$^+$) (75, 76). cDC-1 have been characterized as inducing

CD8 T cell activation (77) while cDC-2 are implicated in CD4 T cell activation (78). In

previous work, we found that cells resembling cDC-2 carry live *M. tuberculosis* to the

draining mediastinal lymph node (2, 5), and cDC-1 do not appear to be significantly

infected by *M. tuberculosis* in the first two weeks of infection (4). We have previously

shown cDC-2-like cells are derived from monocytes (5), and therefore have termed this

population moDC. However, CD11b^{pos}CD11c^{pos} migratory DCs do not prime CD4 T

cells efficiently during *M. tuberculosis* infection (9), and we and others have previously

demonstrated that infected moDC transfer antigen to resident mediastinal lymph node

DC that in turn activate T cells (8, 9). Here, we observe a significant increase of T cell

activation at 17 days post-infection in mice infected with 4334 which correlates with a

higher bacterial burden in the draining lymph node at day 14 post-infection. These

findings are consistent with the finding of earlier spread of the bacteria to

CD11b^{pos}CD11c^{pos} moDC in the lungs, but do not indicate whether the greater

magnitude of Ag85B-specific CD4 T cell activation in the lymph node is the

consequence of more moDC presenting antigen, or more antigen presented per moDC.

Earlier T cell activation can lead to superior control of *M. tuberculosis* infection

(13). Here, we demonstrate that 4334 induces an earlier antigen-specific T cell

response which is associated with a reduction in CFU 7 weeks post-infection compared

to H37Rv infected mice. In addition, the quantitatively greater accumulation of

lymphocytes in the lungs of mice infected with 4334 is associated with an increased

number of inflammatory lesions in the lung at this time point.

A limitation of this work is the resolution of the flow sorting strategy. As the work

prioritized identification and characterization of the cell populations that harbor live *M.*

tuberculosis in the initial days after infection when the bacterial burdens are low, the

number of surface markers used for sorting and the resolution of certain cell subsets

was limited. Nevertheless, the resolution of the subsets was sufficient to allow tracking

of the spread of *M. tuberculosis* from alveolar macrophages to recruited inflammatory

cells over time. An additional limitation is that the genetic and molecular differences

between H37Rv and 4334 were not identified, although we do rule out the lineage 2

specific virulence lipid, PGL, as one potential difference. While both strains were

isolated from patients with pulmonary TB, H37Rv was initially isolated in 1905 (79), and

4334 was isolated in the mid-2000s (44). It is not known whether or how passage and

storage of the two strains has resulted in genotypic or phenotypic differences from the

original patient isolates. However, we do rule out the lineage 2 specific virulence lipid,

PGL, and comparative genetic analysis of these strains, as well as future work on the

recognized differences between lineage 2 and 4 strains can now be considered in light

of these detailed findings during early in vivo infection.

In conclusion, we found that *M. tuberculosis* strain-dependent differences in the

rate of spread from alveolar macrophages to recruited leukocytes in the lungs is

associated with lower rates of alveolar macrophage death and with a lower rate and

extent of initial leukocyte recruitment to the lungs. The finding that enhanced spread

from alveolar macrophages is associated with lower rates of alveolar macrophage cell

death suggests that nonlytic mechanisms are likely to contribute to cell-to-cell spread of

M. tuberculosis in vivo. The data also reveal that the strains examined differ in their

ESX-1- (type VII) but not SecA (type I)-mediated protein secretion activity, which

correlates with enhanced spread, suggesting that Esx-1 may contribute to nonlytic

spread of *M. tuberculosis* in the early stages of infection in the lungs. Understanding the

mechanisms underlying these findings will provide further insights into the virulence of

M. tuberculosis.

METHODS

Mice and Care

C57BL/6 mice were purchased from the Jackson Laboratory. P25 TCR Tg mice

whose CD4 T cells recognize peptide 25 (aa 240-254) of *M. tuberculosis* antigen 85B in

complex with mouse MHC II I-A^b (10, 80) were crossed in-house with congenic CD45.1$^+$

(B6.SJL-Ptprc^a Pepc^b/BoyJ) and with Rag1$^{-/-}$ mice. For infections with *M. tuberculosis*,

mice were housed under barrier conditions in the ABSL-3 facility at the NYU School of

Medicine. All mice were between 8 and 12 weeks of age at the beginning of the

experiment, and mice of both sexes were used. Mice were euthanized by CO_2

asphyxiation followed by cervical dislocation. All experiments were performed with the

prior approval of the NYU Institutional Animal Care and Use Committee (IACUC).

Cell isolation for *in vitro* and *in vivo* experiments

Alveolar macrophages (AM) were harvested by bronchoalveolar lavage (81), and purity confirmed through flow cytometry analysis with CD11c$^+$CD11b$^-$ cells (2) above 90% and autofluorescence (82). AM were cultured for up to 4 days in RPMI 1640 with 10% heat-inactivated FBS, 2mM L-glutamine, 1mM sodium pyruvate, 1x β-mercaptoethanol (Gibco), 10 mM HEPES, 10ng/mL of recombinant murine granulocyte-macrophage colony-stimulating factor (Peprotech), and 10 U/ml penicillin – 10 µg/ml streptomycin that was washed out prior to infection.

P25 TCR Tg CD4 T cells were isolated from the secondary lymphoid organs of P25TCR-Tg/CD45.1/Rag1$^{-/-}$ mice by magnetic cell sorting using anti-CD4 conjugated microbeads and an AutoMACS Classic sorter (Miltenyi Biotec), according to the manufacturer's recommendations.

ical *Mycobacterium tuberculosis* strains, growth, and infections

M. tuberculosis H37Rv was grown as previously described (83). 4334 was obtained and maintained as previously described (61). Mice were infected via the aerosol route, using an inhalation exposure system (Glas-Col) (2). The infectious dose was quantitated on day 1 by plating whole lung homogenates from 3–5 mice on Middlebrook 7H11 agar. To determine the bacterial load throughout the infection, lungs were harvested, homogenized, and serial dilutions were plated on Middlebrook 7H11 agar. Colony Forming Units (CFUs) were counted after incubation of plates at 37°C for 3 weeks.

For *in vitro* infections, mycobacteria were grown to mid-log phase, pelleted at 3750 g, resuspended in PBS + 0.5% Tween 80, re-pelleted and excess Tween 80

washed off by centrifugation in PBS. The final pellet was re-suspended in RPMI 1640

with 10% serum, the bacterial density was evaluated by measuring absorbance at 580

nm, and the multiplicity of infection was adjusted to the mammalian cell density.

Flow cytometry and cell sorting

Lungs were removed and processed into single-cell suspensions for flow

cytometry and cell sorting as previously described (2, 84). Antibodies conjugated to

various fluorophores and directed against surface markers were:

Antibody	Clone	Source
B220-PE	RA3-6B2	BioLegend
CD4-BUV395	RM4-5	BD Biosciences
CD4-APC/Cy7	RM4-5	BioLegend
CD8-APC	53-6.7	BioLegend
CD8-BV421	53-6.7	BioLegend
CD11c-PerCP	HL3	BioLegend
CD11b-APC/Cy7	M1/70	BD Biosciences
CD19-FITC	6D5	BioLegend
CD45-PE	30-F11	BD Biosciences
CD45-FITC	30-F11	BD Biosciences
CD45.1-AF700	A20	BioLegend
CD45.2-PerCp/Cy5.5	104	BD Biosciences
CD45.2-FITC	104	BD Biosciences
CD90.2-PerCP/Cy5.5	30-H12	BioLegend
Gr-1-APC	RB6-8C5	BD Biosciences
Ly6C-FITC	AL-21	BD Biosciences
Ly6G-AF647	1A8	BioLegend
Siglec-F-PE	E50-2440	BD Biosciences

For flow cytometry analysis, stained samples were fixed overnight in 1%

paraformaldehyde. A minimum of 200,000 events per sample, gated on single cells

using forward and side scatter parameters, were acquired using an LSRII and

FACSDiva software (BD Biosciences).

For live cell sorting, fluorescently labeled live cells were acquired using a BSL3-

contained iCyt/Sony Synergy cell sorter. Cells were first separated in non-hematopoietic

cells (CD45$^-$), AM (CD45$^+$CD11b$^-$CD11c$^+$), monocyte-derived dendritic cells (moDC,

CD45$^+$CD11b$^+$CD11c$^+$) and other myeloid cells (CD45$^+$CD11b$^+$CD11c$^{neg/low}$). Then, this

latter fraction was re-acquired and sorted into neutrophils (Neut, Gr1^{hi}CD11c$^-$),

monocytes (Gr1^{int}CD11c$^-$) and recruited macrophages (Gr-1$^-$CD11c^{low}). All cell fractions

were collected and plated for CFU recovery and quantitation on agar as described

above. For flow cytometry and cell sorting, acquisition data were analyzed using FlowJo

software (TreeStar).

***Mycobacterium tuberculosis*-antigen specific T cell adoptive transfer,**

proliferation, and IFNγ stimulation

P25TCR-Tg/CD45.1/Rag1$^{-/-}$ CD4$^+$ T cells were stained with CellTrace Violet

(ThermoFisher Scientific), according to the manufacturer's recommendations. Cells

were re-suspended at 2-3x10^6 cells/100 μl in sterile PBS for intravenous injection via

the retro-orbital route into anesthetized CD45.2$^+$ C57BL/6 recipient mice. Recipient mice

were infected with aerosolized *M. tuberculosis* 24h post-T cell transfer. T cell

proliferation was assessed at selected time points post-infection by quantitating

CellTrace Violet dilution by flow cytometry. Whole lung supernatants were analyzed for

IFNγ levels by ELISA.

Quantification of Necrotic and Apoptotic Cells

AM were infected with *M. tuberculosis* expressing DsRed for designated times,

after which cells were washed and then lifted with cold PBS containing 0.5 mM EDTA.

Cells were fixed, washed, and stained with Stain Zombie Aqua (Biolegend) to quantify

necrosis and AnnexinV-APC to quantify apoptosis (BD Biosciences) per manufacturer

protocols, washed with PBS, and analyzed by flow cytometry for DsRed (bacteria),

Annexin V, and Zombie Aqua.

Analysis of PDIM and PGL

Bacterial strains were grown shaking at 37°C for ~7 days to an OD580 of 0.6 in

Middlebrook 7H9 broth supplemented with 10% Oleic acid/dextrose/catalase and 0.05%

Tween 80. Cultures were centrifuged, supernatant removed, and pellets washed twice,

resuspended in 1mL of CH_3OH and contacted with 25mL $CHCl_3/CH_3OH$ (2:1) overnight.

Lipid was then extracted, dissolved in $CHCl_3:CH_3OH$ at 1mg/mL, and ran on an Agilent

Technologies 6520 Accurate-Mass Q-Tof and 1200 series HPLC as previously

described (85).

ESX-1 Secretion Assay

Bacterial strains were grown shaking at 37°C for ~7 days in Middlebrook 7H9

broth supplemented with 10% Oleic acid/dextrose/catalase and 0.05% Tween 80 to an

OD580 0.4 – 0.7. Cultures were centrifuged, washed thrice in Sauton's media (0.5 g

KH2PO4, 0.5 g MgSO4, 4.0 g L-asparagine, 60 ml glycerol, 0.05 g Ferric ammonium

citrate, 2.0 g citric acid, 0.1 ml 1% ZnSO4, dH20 to 900 ml, 2.5 ml 20% Tween-80), and

incubated for an additional 24 hours. Bacteria were centrifuged at 2500 g for 10 min,

and supernatants passed through 0.2 µm syringe filters to obtain filtrates. Samples were

reconstituted in 200uL of 2M urea, reduced with DTT at 57°C for 1h, alkylated with 0.5M

iodacetamide for 45 min at RT, followed by trypsin digestion and cleansed as previously

described (86). After SpeedVac concentration, samples reconstituted in 0.5% acetic

acid, and loaded onto an Acclaim PepMap trap column in line with an EASY Spray

50cm x 75µm ID PepMap C18 analytical HPLC column with 2µm bead size using the

auto sampler of an EASY-nLC 1000 HPLC (ThermoFisher) and solvent A (2%

acetonitrile, 0.5% acetic acid). The peptides were gradient eluted into a Q Exactive

(Thermo Scientific) mass spectrometer using a 2h linear gradient from 2% to 40%

solvent B (95% acetonitrile, 0.5% acetic acid), followed by 10 min from 40% to 100%

solvent B. Solvent B was held at 100% for another 10 min for column wash. Spectra

were acquired using the following parameters: resolution of 70,000, an automatic gain

control of 1e6, with a maximum ion time of 120 ms, and scan range of 300 to 1500 m/z.

Following each full MS scan twenty data-dependent high resolution HCD MS/MS

spectra were acquired.   All MS/MS spectra were collected using the following

instrument parameters: resolution of 17,500, AGC target of 2e5, maximum ion time of

250 ms, one microscan, 2 m/z isolation window, fixed first mass of 150 m/z, and

normalized collision energy of 27.

All acquired MS/MS spectra were searched against a combined database for

H37Rv and NITR203 on UniProt, using Andromeda search algorithm and MaxQuant for

quantitation (87). The data set was filtered to remove proteins with only one unique

peptide and that were not detected in all three replicates of at least one strain. LFQ

intensity values were $\log_2$ transformed. A two-sided t-test and correcting for multiple

testing by controlling for FDR at 5% using Benjamini-Hochberg's method. In addition, z-

scores were calculated and used to perform hierarchical clustering.

Histopathology

The left lung was excised, fixed in 10% buffered formalin for one week at room

temperature, then embedded in paraffin. 5 μm sections were cut and stained with

hematoxylin and eosin. Whole lung sections were scanned at 40X by the NYU School of

Medicine Experimental Pathology Research Laboratory, using a Leica SCN400 F

whole-slide scanner. Digital images were used for the quantitation of lung

histopathology using the open-source image processing software Fiji, with two

independent approaches. Briefly, the total surface area of the lung section was

calculated, then the contributions of airways and blood vessels to that area were

subtracted, leaving the contribution of the structural tissues. The proportion of

inflammatory infiltrates within those tissues was then quantitated, either by manual

contouring or using automated color density contouring. For each 40X section, the final

percentage of lung inflammatory infiltrate was calculated as the average of the values

obtained by each method.

Statistical analysis

Experiments were performed at least twice, with exception of experiments

focusing on H37RvΔRD1 and 7 weeks post-infection which were performed once.

Results are expressed as mean and standard deviation (SD). Unless otherwise stated,

parametric Student two-tailed t test and Holm multiple comparisons correction, with a

95% confidence interval was used to compare experimental groups, with $p < 0.05$

considered significant.

ACKNOWLEDGMENTS

We thank Diane Ordway and Midori Kato-Maeda for the *M. tuberculosis* strain

4334. Cynthia Portal-Celhay and Thais Klevorn aided in experiments. We acknowledge

the expertise and support of Beatrix Ueberheide and Jessica Chapman-Lim at the NYU

Proteomics Laboratory, supported in part by NYU Langone Health and the Laura and

Isaac Perlmutter Cancer Center Support Grant P30CA016087 from the National Cancer

Institute, for proteomics assays. Cell sorting and flow cytometry were performed by

Michael Gregory from the NYU Cytometry and Cell Sorting Laboratory, which is

supported in part by grant P30CA016087 from the National Institutes of Health/National

Cancer Institute.

This project was supported by NIH grants R01 AI051242 and R01 AI049313, as

well as an award from the Stony Wold-Herbert Fund.

L.D. and J.D.E. designed the experiments, B.S.Z, L.D., and J.D.E. analyzed the

results; T.J.F., L.D., and A.C. conducted experiments; T.Y and D.B.M. provided the

lipidomic analysis and reviewed the manuscript; B.S.Z, L.D., and J.D.E. wrote the

manuscript.

REFERENCES

1.   Sabin FR, Doan CA. 1927. The relation of monocytes and clasmatocytes to early

575   infection in rabbits with bovine tubercle bacilli. J Exp Med 46:627–644.

576   2.   Wolf AJ, Linas B, Trevejo-Nuñez GJ, Kincaid E, Tamura T, Takatsu K, Ernst JD.

577   2007. Mycobacterium tuberculosis infects dendritic cells with high frequency and

578   impairs their function in vivo. J Immunol 179:2509–2519.

579   3.   Cohen SB, Gern BH, Delahaye JL, Adams KN, Plumlee CR, Winkler JK, Sherman

580   DR, Gerner MY, Urdahl KB. 2018. Alveolar macrophages provide an early

581   Mycobacterium tuberculosis niche and initiate dissemination. Cell Host Microbe

582   24:439–446.

583   4.   Rothchild AC, Olson GS, Nemeth J, Amon LM, Mai D, Gold ES, Diercks AH,

584   Aderem A. 2019. Alveolar Macrophages Generate a Noncanonical NRF2-Driven

Transcriptional Response to Mycobacterium tuberculosis in vivo. Sci Immunol

4:eaaw6693.

5. Norris BA, Ernst JD. 2018. Mononuclear cell dynamics in M. tuberculosis infection

provide opportunities for therapeutic intervention. PLoS Pathog 14:e1007154.

6. Srivastava S, Ernst JD, Desvignes L. 2014. Beyond macrophages: The diversity

of mononuclear cells in tuberculosis. Immunol Rev 262:179–192.

7. Skold M, Behar SM, Sköld M, Behar SM. 2014. Tuberculosis triggers a tissue-

dependent program of differentiation and acquisition of effector functions by

circulating monocytes. J Immunol 181:6349–6360.

8. Srivastava S, Ernst JD. 2014. Cell-to-cell transfer of M. tuberculosis antigens

optimizes CD4 T cell priming. Cell Host Microbe 15:741–752.

9. Samstein M, Schreiber HA, Leiner IM, Sušac B, Glickman MS, Pamer EG. 2013.

Essential yet limited role for CCR2+ inflammatory monocytes during

Mycobacterium tuberculosis-specific T cell priming. Elife 2:e01086.

10. Wolf AJ, Desvignes L, Linas B, Banaiee N, Tamura T, Takatsu K, Ernst JD. 2008.

Initiation of the adaptive immune response to Mycobacterium tuberculosis

depends on antigen production in the local lymph node, not the lungs. J Exp Med

205:105–115.

11. Poulsen A. 1950. Some clinical features of Tuberculosis. Acta Tuberc Scand

24:311–346.

12. Wallgren A. 1948. A time-table of Tuberculosis. Tubercle 29:245–251.

13. Chackerian AA, Alt JM, Perera T V, Dascher CC, Behar SM. 2002. Dissemination

of Mycobacterium tuberculosis is influenced by host factors and precedes the

initiation of T-cell immunity. Infect Immun 70:4501–4509.

14. Blomgran R, Ernst JD. 2011. Lung neutrophils facilitate activation of naïve

antigen-specific CD4+ T cells during Mycobacterium tuberculosis infection. J

Immunol 186:7110–7119.

15. Blomgran R, Desvignes L, Briken V, Ernst JD. 2012. Mycobacterium tuberculosis

inhibits neutrophil apoptosis, leading to delayed activation of naive CD4 T cells.

Cell Host Microbe 11:81–90.

16. Grace PS, Ernst JD. 2015. Suboptimal antigen presentation contributes to

virulence of Mycobacterium tuberculosis in vivo. J Immunol 196:357–364.

17. Divangahi M, Chen M, Gan H, Desjardins D, Hickman TT, Lee DM, Fortune S,

Behar SM, Remold HG. 2009. Mycobacterium tuberculosis evades macrophage

defenses by inhibiting plasma membrane repair. Nat Immunol 10:899–906.

18. Keane J, Remold HG, Kornfeld H. 2000. Virulent Mycobacterium tuberculosis

Strains Evade Apoptosis of Infected Alveolar Macrophages. J Immunol 164:2016–

2020.

19. Sly LM, Hingley-Wilson SM, Reiner NE, McMaster WR. 2003. Survival of

Mycobacterium tuberculosis in host macrophages involves resistance to

apoptosis dependent upon induction of antiapoptotic Bcl-2 family member Mcl-1. J

Immunol 170:430–437.

20. Briken V, Miller JL. 2008. Living on the edge: Inhibition of host cell apoptosis by

Mycobacterium tuberculosis. Future Microbiol 3:415–422.

21. Remold M Katarzyna Balcewicz-Sablinska HG, Keane J. 1998. Pathogenic

Mycobacterium tuberculosis evades apoptosis of host macrophages by release of

TNF-R2, resulting in inactivation of TNF-a. J Immunol 161:2636–2641.

22.   Gan H, Lee J, Ren F, Chen M, Kornfeld H, Remold HG. 2008. Mycobacterium

tuberculosis blocks crosslinking of annexin-1 and apoptotic envelope formation on

infected macrophages to maintain virulence. Nat Immunol 9:1189–1197.

23.   Stutz MD, Allison CC, Ojaimi S, Preston SP, Doerflinger M, Arandjelovic P,

Whitehead L, Bader SM, Batey D, Asselin-Labat M-L, Herold MJ, Strasser A,

West NP, Pellegrini M. 2021. Macrophage and neutrophil death programs

differentially confer resistance to tuberculosis. Immunity 54:1758–1771.

24.   Chen M, Gan H, Remold HG. 2006. A mechanism of virulence: Virulent

Mycobacterium tuberculosis strain H37Rv, but not attenuated H37Ra, causes

significant mitochondrial inner membrane disruption in macrophages leading to

necrosis. J Immunol 176:3707–3716.

25.   Sun J, Siroy A, Lokareddy RK, Speer A, Doornbos KS, Cingolani G, Niederweis

M. 2015. The tuberculosis necrotizing toxin kills macrophages by hydrolyzing

NAD. Nat Struct Mol Biol 22:672–678.

26.   Chen M, Divangahi M, Gan H, Shin DSJ, Hong S, Lee DM, Serhan CN, Behar

SM, Remold HG. 2008. Lipid mediators in innate immunity against tuberculosis:

Opposing roles of PGE 2 and LXA 4 in the induction of macrophage death. J Exp

Med 205:2791–2801.

27.   Stanley SA, Raghavan S, Hwang WW, Cox JS. 2003. Acute infection and

macrophage subversion by Mycobacterium tuberculosis require a specialized

secretion system. PNAS 100:13001–13006.

28.   Pym AS, Brodin P, Brosch R, Huerre M, Cole ST. 2002. Loss of RD1 contributed

to the attenuation of the live tuberculosis vaccines Mycobacterium bovis BCG and

Mycobacterium microti. Mol Microbiol 46:709–717.

29. Koo IC, Wang C, Raghavan S, Morisaki JH, Cox JS, Brown EJ. 2008. ESX-1-

dependent cytolysis in lysosome secretion and inflammasome activation during

mycobacterial infection. Cell Microbiol 10:1866–1878.

30. Mishra BB, Moura-Alves P, Sonawane A, Hacohen N, Griffiths G, Moita LF, Anes

E, Moura-Alves P, Sonawane A, Hacohen N, Griffiths G, Moita LF, Anes E. 2010.

Mycobacterium tuberculosis protein ESAT-6 is a potent activator of the

NLRP3/ASC inflammasome. Cell Microbiol 12:1046–1063.

31. Wassermann R, Gulen MF, Sala C, Perin SG, Lou Y, Rybniker J, Schmid-Burgk

JL, Schmidt T, Hornung V, Cole ST, Ablasser A. 2015. Mycobacterium

tuberculosis differentially activates cGAS- and inflammasome-dependent

intracellular immune responses through ESX-1. Cell Host Microbe 17:799–810.

32. Conrad WH, Osman MM, Shanahan JK, Chu F, Takaki KK, Cameron J,

Hopkinson-Woolley D, Brosch R, Ramakrishnan L. 2017. Mycobacterial ESX-1

secretion system mediates host cell lysis through bacterium contact-dependent

gross membrane disruptions. PNAS 114:1371–1376.

33. Davis JM, Ramakrishnan L. 2009. The role of the granuloma in expansion and

dissemination of early Tuberculous infection. Cell 136:37–49.

34. Welin A, Björnsdottir H, Winther M, Christenson K, Oprea T, Karlsson A, Forsman

H, Dahlgren C, Bylund J. 2015. CFP-10 from Mycobacterium tuberculosis

selectively activates human neutrophils through a pertussis toxin-sensitive

chemotactic receptor. Infect Immun 83:205–213.

35.   Brandt L, Oettinger T, Holm A, Andersen AB, Anderson P, Andersen P. 1996. Key

678   epitopes on the ESAT-6 antigen recognized in mice during the recall of protective

679   immunity to Mycobacterium tuberculosis. J Immunol 157:3527–33.

680   36.   Elhay MJ, Oettinger T, Andersen P. 1998. Delayed-type hypersensitivity

681   responses to ESAT-6 and MPT64 from Mycobacterium tuberculosis in the guinea

682   pig. Infect Immun 66:3454–3456.

683   37.   Ravn P, Demissie A, Eguale T, Wondwosson H, Lein D, Amoudy HA, Mustafa

684   AS, Kok Jensen A, Holm A, Rosenkrands I, Oftung F, Olobo J, von Reyn F,

685   Andersen P. 1999. Human T cell responses to the ESAT-6 antigen from

686   Mycobacterium tuberculosis. J Infect Dis 179:637–645.

687   38.   Wong KW, Jacobs WR. 2011. Critical role for NLRP3 in necrotic death triggered

688   by Mycobacterium tuberculosis. Cell Microbiol 13:1371–1384.

689   39.   Dallenga T, Repnik U, Corleis B, Eich J, Reimer R, Griffiths GW, Schaible UE.

690   2017. M. tuberculosis-induced necrosis of infected neutrophils promotes bacterial

691   growth following phagocytosis by macrophages. Cell Host Microbe 22:519–530.

692   40.   Francis RJ, Butler RE, Stewart GR. 2014. Mycobacterium tuberculosis ESAT-6 is

693   a leukocidin causing Ca2+ influx, necrosis and neutrophil extracellular trap

694   formation. Cell Death Dis 5:e1474.

695   41.   Brites D, Gagneux S. 2017. The nature and evolution of genomic diversity in the

696   Mycobacterium tuberculosis vomplex. Adv Exp Med Biol 1019:1–26.

697   42.   Coscolla M, Gagneux S. 2014. Consequences of genomic diversity in

698   mycobacterium tuberculosis. Semin Immunol 26:431–444.

43.   Kremer K, Van Der Werf MJ, Au BKY, Anh DD, Kam KM, Van Doorn HR,

Borgdorff MW, Van Soolingen D. 2009. Vaccine-induced immunity by typical

Mycobacterium tuberculosis Beijing strains. Emerg Infect Dis 15:335–339.

44.   Kato-Maeda M, Shanley CA, Ackart D, Jarlsberg LG, Shang S, Obregon-Henao

A, Harton M, Basaraba RJ, Henao-Tamayo M, Barrozo JC, Rose J, Kawamura

LM, Coscolla M, Fofanov VY, Koshinsky H, Gagneux S, Hopewell PC, Ordway

DJ, Orme IM. 2012. Beijing sublineages of Mycobacterium tuberculosis differ in

pathogenicity in the guinea pig. Clin Vaccine Immunol 19:1222–1227.

45.   Kato-Maeda M, Kim EY, Flores L, Jarlsberg LG, Osmond D, Hopewell PC. 2010.

Differences among sublineages of the East-Asian lineage of Mycobacterium

tuberculosis in genotypic clustering. Int J Tuberc Lung Dis 14:538–544.

46.   Reiling N, Homolka S, Walter K, Brandenburg J, Niwinski L, Ernst M, Herzmann

C, Lange C, Diel R, Ehlers S, Niemann S. 2013. Clade-specific virulence patterns

of Mycobacterium tuberculosis complex strains in human primary macrophages

and aerogenically infected mice. MBio 4:e00250-13.

47.   Portevin D, Gagneux S, Comas I, Young D. 2011. Human macrophage responses

to clinical isolates from the Mycobacterium tuberculosis complex discriminate

between ancient and modern lineages. PLoS Pathog 7:e1001307.

48.   Manca C, Reed MB, Freeman S, Mathema B, Kreiswirth B, Barry CE, Kaplan G.

2004. Differential monocyte activation underlies strain-specific Mycobacterium

tuberculosis pathogenesis. Infect Immun 72:5511–5514.

49.   Dietrich J, Roy S, Rosenkrands I, Lindenstrøm T, Filskov J, Rasmussen EM,

Cassidy J, Andersen P. 2015. Differential influence of nutrient-starved

Mycobacterium tuberculosis on adaptive immunity results in progressive

tuberculosis disease and pathology. Infect Immun 83:4731–4739.

50. Valdés I, Montoro E, Mata-Espinoza D, Asín O, Barrios-Payan J, Francisco-Cruz

A, Valdivia JA, Hernández-Pando R. 2014. Immunogenicity and protection

conferred by Mycobacterium habana in a murine model of pulmonary

tuberculosis. Tuberculosis 94:65–72.

51. Ly A, Liu J. 2020. Mycobacterial virulence factors: Surface-exposed lipids and

secreted proteins. Int J Mol Sci 21:3985.

52. Cambier CJ, Takaki KK, Larson RP, Hernandez RE, Tobin DM, Urdahl KB,

Cosma CL, Ramakrishnan L. 2014. Mycobacteria manipulate macrophage

recruitment through coordinated use of membrane lipids. Nature 505:218–222.

53. Tabouret G, Astarie-Dequeker C, Demangel C, Malaga W, Constant P, Ray A,

Honoré N, Bello NF, Perez E, Daffé M, Guilhot C. 2010. Mycobacterium leprae

Phenolglycolipid-1 Expressed by Engineered M. bovis BCG modulates early

interaction with human phagocytes. PLoS Pathog 6:e1001159.

54. Telford WG. 2018. Multiparametric analysis of apoptosis by flow cytometry, p.

167–202. In Methods in Molecular Biology. Humana Press Inc.

55. Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA, Bloom BR. 1993. An

essential role for Interferon-γ in resistance to Mycobacterium tuberculosis

infection. J Exp Med 173:2249–54.

56. Cooper AM, Dalton DK, Stewart TA, Griffin JP, Russell DG, Orme IM. 1993.

Disseminated Tuberculosis in Interferon 7 gene-disrupted mice. J Exp Med

178:2243–2247.

57. Huang L, Nazarova E V, Tan S, Liu Y, Russell DG. 2018. Growth of

Mycobacterium tuberculosis in vivo segregates with host macrophage metabolism

and ontogeny. J Exp Med 215:1135–1152.

58.   Guth AM, Janssen WJ, Bosio CM, Crouch EC, Henson PM, Dow SW. 2009. Lung

environment determines unique phenotype of alveolar macrophages. Am J

Physiol Lung Cell Mol Physiol 296:936–946.

59.   Lafuse WP, Rajaram MVS, Wu Q, Moliva JI, Torrelles JB, Turner J, Schlesinger

LS. 2019. Identification of an Increased Alveolar Macrophage Subpopulation in

Old Mice That Displays Unique Inflammatory Characteristics and Is Permissive to

Mycobacterium tuberculosis Infection . J Immunol 203:2252–2264.

60.   Duan M, Steinfort DP, Smallwood D, Hew M, Chen W, Ernst M, Irving LB,

Anderson GP, Hibbs ML. 2016. CD11b immunophenotyping identifies

inflammatory profiles in the mouse and human lungs. Mucosal Immunol 9:550–

563.

61.   Wiens KE, Ernst JD. 2016. The mechanism for Type I Interferon induction by

Mycobacterium tuberculosis is bacterial strain-dependent. PLoS Pathog

12:e1005809.

62.   Li Q, Whalen CC, Albert JM, Larkin R, Zukowski L, Cave MD, Silver RF. 2002.

Differences in rate and variability of intracellular growth of a panel of

Mycobacterium tuberculosis clinical isolates within a human monocyte model.

Infect Immun 70:6489–6493.

63.   Wu S, Howard ST, Lakey DL, Kipnis A, Samten B, Safi H, Gruppo V, Wizel B,

Shams H, Basaraba RJ, Orme IM, Barnes PF. 2004. The principal sigma factor

sigA mediates enhanced growth of Mycobacterium tuberculosis in vivo. Mol

Microbiol 51:1551–1562.

64.   Park JS, Tamayo MH, Gonzalez-Juarrero M, Orme IM, Ordway DJ. 2006. Virulent

clinical isolates of Mycobacterium tuberculosis grow rapidly and induce cellular

necrosis but minimal apoptosis in murine macrophages. J Leukoc Biol 79:80–86.

65.   Martin CJ, Booty MG, Rosebrock TR, Nunes-Alves C, Desjardins DM, Keren I,

Fortune SM, Remold HG, Behar SM. 2012. Efferocytosis is an innate antibacterial

mechanism. Cell Host Microbe 12:289–300.

66.   Michelet X, Tuli A, Gan H, Geadas C, Sharma M, Remold HG, Brenner MB. 2018.

Lysosome-mediated plasma membrane repair is dependent on the small GTPase

Arl8b and determines cell death type in Mycobacterium tuberculosis infection. J

Immunol 200:3160–3169.

67.   Mohareer K, Asalla S, Banerjee S. 2018. Cell death at the cross roads of host-

pathogen interaction in Mycobacterium tuberculosis infection. Tuberculosis

113:99–121.

68.   Behar SM, Divangahi M, Remold HG. 2010. Evasion of innate immunity by

mycobacterium tuberculosis: Is death an exit strategy? Nat Rev Microbiol 8:668–

674.

69.   Keane J, Remold HG, Kornfeld H. 2000. Virulent Mycobacterium tuberculosis

Strains Evade Apoptosis of Infected Alveolar Macrophages. J Immunol 164:2016–

2020.

70.   Byrd TF, Green GM, Fowlston SE, Lyons CR. 1998. Differential growth

characteristics and streptomycin susceptibility of virulent and avirulent

Mycobacterium tuberculosis strains in a novel fibroblast-mycobacterium

microcolony assay. Infect Immun 66:5132–5139.

71.   Hagedorn M, Ronde KH, Russell DG, Soldati T. 2009. Infection by tubercular

mycobacteria is spread by nonlytic ejection from their amoeba hosts. Science

323:1729–1733.

72.   Cambier CJ, O'Leary SM, O'Sullivan MP, Keane J, Ramakrishnan L. 2017.

Phenolic glycolipid facilitates mycobacterial escape from microbicidal tissue-

resident macrophages. Immunity 47:552–565.

73.   Woodruff CE. 1934. A free growth period of tubercle bacilli in the guinea pig

omentum as related to the hypersensitive state. Am J Pathol 10:739–756.

74.   Rich A. 1944. The Pathogenesis of Tuberculosis, 1st ed. Charles C. Thomas,

Springfield, IL.

75.   Bosteels C, Scott CL. 2020. Transcriptional regulation of DC fate specification.

Mol Immunol 121:38–46.

76.   Guilliams M, Ginhoux F, Jakubzick C, Naik SH, Onai N, Schraml BU, Segura E,

Tussiwand R, Yona S. 2014. Dendritic cells, monocytes and macrophages: A

unified nomenclature based on ontogeny. Nat Rev Immunol 14:571–578.

77.   Ng SL, Teo YJ, Setiagani YA, Karjalainen K, Ruedl C. 2018. Type 1 conventional

CD103+ dendritic cells control effector CD8+ T cell migration, survival, and

memory responses during influenza infection. Front Immunol 9:3043.

78.   Dudziak D, Kamphorst AO, Heidkamp GF, Buchholz VR, Trumpfheller C,

Yamazaki S, Cheong C, Liu K, Lee H-W, Park CG, Steinman RM, Nussenzweig

MC. 2007. Differential antigen processing by dendritic cell subsets in vivo.

Science 315:107–111.

79.   Steenken W, Gardner LU. 1946. HISTORY OF H37 STRAIN OF TUBERCLE

BACILLUS1. Am Rev Tuberc 54:62–66.

80.   Tamura T, Ariga H, Kinashi T, Uehara S, Kikuchi T, Nakada M, Tokunaga T, Xu

W, Kariyone A, Saito T, Kitamura T, Maxwell G, Takaki S, Takatsu K. 2004. The

role of antigenic peptide in CD4+ T helper phenotype development in a T cell

receptor transgenic model. Int Immunol 16:1691–1699.

81.   Rothchild A, Mai D, Aderem A, Diercks A. 2020. Flow cytometry analysis and

fluorescence-activated cell sorting of myeloid cells from lung and bronchoalveolar

lavage samples from Mycobacterium tuberculosis-infected mice. Bio Protoc

10:e3630.

82.   Havenith CEG, Breedijk AJ, van Miert PPMC, Blijleven N, Calame W, Beelen

RHJ, Hoefsmit ECM. 1993. Separation of alveolar macrophages and dendritic

cells via autofluorescence: phenotypical and functional characterization. J Leukoc

Biol 53:504–510.

83.   Banaiee N, Jacobs WR, Ernst JD. 2006. Regulation of Mycobacterium

tuberculosis whiB3 in the mouse lung and macrophages. Infect Immun 74:6449–

6457.

84.   Srivastava S, Grace PS, Ernst JD. 2016. Antigen export reduces antigen

presentation and limits T cell control of M. tuberculosis. Cell Host Microbe 19:44–

54.

85.   Layre E, Sweet L, Hong S, Madigan CA, Desjardins D, Young DC, Cheng TY,

Annand JW, Kim K, Shamputa IC, McConnell MJ, Debono CA, Behar SM,

Minnaard AJ, Murray M, Barry CE, Matsunaga I, Moody DB. 2011. A comparative

lipidomics platform for chemotaxonomic analysis of mycobacterium tuberculosis.

Chem Biol 18:1537–1549.

86.   Safavi-Hemami H, Gajewiak J, Karanth S, Robinson SD, Ueberheide B, Douglass

AD, Schlegel A, Imperial JS, Watkins M, Bandyopadhyay PK, Yandell M, Li Q,

Purcell AW, Norton RS, Ellgaard L, Olivera BM. 2015. Specialized insulin is used

for chemical warfare by fish-hunting cone snails. PNAS USA 112:1743–1748.

87.   Rgen Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. 2014. Accurate

proteome-wide label-free quantification by delayed normalization and maximal

peptide ratio extraction, termed MaxLFQ. Mol Cell Proteomics 13:2513–2526.

Figure Legends

Figure 1. *M. tuberculosis* 4334 grows more rapidly but induces less cell

recruitment than H37Rv in early infection. C57BL/6 mice were infected by aerosol

with 200-300 Colony Forming Units (CFU) of *M. tuberculosis* H37Rv, H37RvΔRD1

(ΔRD1), or 4334. Lungs were harvested at day 1, 3, 6, 8, 10, and 14 post-infection, and

processed to single cell suspensions. (**A**) Total CFU isolated from the lungs of mice

infected with the individual strains during the first 14 days of infection. (**B**) Total number

of cells isolated from the lungs of mice infected with the individual strains. Results are

shown as mean ± SD in 1(ΔRD1), 2 (4334), and 6 (H37Rv) pooled experiments, with

n=5 mice per strain per day in each experiment. Statistical significance was assessed

by multiple unpaired T-test for each day and Holm-Sidak multiple comparisons

correction, with a 95% confidence interval and *$p<0.05$, **$p<0.01$, ***$p<0.001$,

****p<0.0001. For clarity, asterisks above H37Rv and 4334 are comparing only these 2

strains, and below are ΔRD1 compared to H37Rv.

Figure 2. Differential strain-dependent spread of _M. tuberculosis_ from

CD11b$^{neg/lo}$CD11c^{pos} alveolar macrophages to recruited lung myeloid cells. Mice

were infected and lungs harvested as in Figure 1. (**A**) Cells were stained and analyzed

by flow cytometry for quantitation of CD11b$^{neg/lo}$CD11c^{pos} alveolar macrophages,

CD11b^{pos}CD11c^{pos} monocyte-derived dendritic cells, Gr-1^{hi}CD11c^{neg} neutrophils, Gr-

1^{int}CD11c^{neg} monocytes, and Gr-1^{neg}CD11c^{neg} monocyte-derived recruited macrophages

in the lungs (see Supplementary Figure 1 and Material and Methods for gating

strategy). (**B**) Total CFU in each leukocyte subset after live flow cytometry sorting and

plating of sorted cell populations on 7H11 solid media. Results are shown as mean ±

SD in 1 (ΔRD1), 2 (4334), and 6 (H37Rv) and pooled experiments, with n=5 mice per

strain per day in each experiment. Statistically significant difference between strains

was assessed multiple unpaired T-test for each day and Holm-Sidak multiple

comparisons correction, with a 95% confidence interval and *p<0.05, **p<0.01,

***p<0.001, ****p<0.0001. For clarity, asterisks above H37Rv and 4334 are comparing

only these 2 strains, and below are ΔRD1 compared to H37Rv.

Figure 3. _M. tuberculosis_ strains 4334 and H37Rv do not produce PGL. PGL and

PDIM analysis of _M. tuberculosis_ H37Rv, HN878, and 4334 in 7H9 cultures, using

electrospray ionization-quadrupole time-of-flight-mass spectrometry (ESI-QTOF-MS).

Shown are representative images of (**A**) extracted ion chromatograms of PGL

[C114H212O18+NH4]+ at *m/z* 1887.601 and PDIM [C91H180O5+NH4]+ at *m/z*

1371.417, and (**B**) mass spectra. Strain HN878 was included in the analysis as a known

PGL producing positive control.

Figure 4. *M. tuberculosis* strain 4334 replicates to a greater extent than H37Rv in

alveolar macrophages. (**A**) Cultured alveolar macrophages were infected for 48 or 72h

with an MOI of 1 with *M. tuberculosis* H37Rv or 4334. Results are shown as mean CFU

fold-increase over initial inoculum. (**B**) Representative flow cytometry analysis of

cultured alveolar macrophages infected *in vitro* for 48h with *M. tuberculosis* H37Rv or

4334, each expressing DsRed fluorescent protein and stained with markers of necrosis

(Zombie Aqua, ZA) and apoptosis (Annexin V-APC). The upper panels show the

frequency of infected cells (DsRed$^+$) and lower panels staining with ZA and AV for cells

infected with the indicated strain 48h post-infection. (**C**) Percentage of DsRed$^+$ cultured

alveolar macrophages as quantitated from flow cytometry analysis using the gating

shown in panel B. (**D**) Frequency of cultured alveolar macrophages that are viable (ZA$^-$

AV$^-$), necrotic (ZA$^+$AV$^-$), early apoptotic (ZA$^-$AV$^+$) or late cell death (ZA$^+$AV$^+$) 48h after

infection with the indicated strain expressing DsRed fluorescent protein; bystanders are

defined as DsRed negative cells. Results are shown as mean $\pm$ SD for n=3 per time

point and per strain. Data between cells infected with respective strains were analyzed

by unpaired Student's t-test for each day and Holm multiple comparisons correction,

with a 95% confidence interval. Asterisks H37Rv vs 4334 (*p<0.05, **p<0.01,

****p<0.0001), hashtag uninfected vs H37Rv or 4334 ($^{\#}$p<0.05, $^{\#\#}$p<0.01, $^{\#\#\#}$p<0.001).

Figure 5. Earlier dissemination of *M. tuberculosis* strain 4334 to lymph nodes is

associated with a greater magnitude of Ag85B-specific CD4 T cell priming. (**A**)

Bacterial load in mediastinal lymph nodes (MdLN) of mice infected 14 days earlier with

M. tuberculosis H37Rv or 4334. Results are shown as mean CFU fold-increase over

initial inoculum. Flow cytometry assessment (**B**) and absolute quantitation (**C**) of the

proliferation of CellTrace Violet-labeled adoptively transferred naïve P25TCR-

Tg/CD45.1 CD4$^+$ T cells in the MDLN 10-, 14-, and 17-days post-infection with H37Rv

or 4334. Percentages shown in panel B are the frequencies of adoptively transferred

P25TCR-Tg CD4$^+$ T cells in which CTV was diluted as a result of cell division (CTV^{dilute}).

Results in panel C are shown as mean $\pm$ SD, for n=4 mice per day and per strain. (**D**)

Quantitation of IFNγ$^+$ endogenous and P25TCR-Tg (as defined by congenic markers

CD45.1/2) CD4$^+$ T cells in the lungs of mice infected for 14 or 17 days with H37Rv or

4334. Results are shown as mean $\pm$ SD, for n=4 mice per day and per strain. Flow

cytometry analysis for these results is shown in Supplementary Figure 4B. Differences

between *M. tuberculosis* strains were assessed by Student's t-test for each day and

Holm multiple comparisons correction, with a 95% confidence interval and *p<0.05, and

**p<0.01.

Supplemental Material Legends

Supplementary Figure 1. Flow cytometry gating strategy for identification of lung

myeloid cell populations. After processing lung tissue into single-cell suspensions and

staining, cells were flow sorted first as CD45^{neg} (non-hematopoietic) or CD45^{pos}. Further

cellular identification occurred using CD11b and CD11c to distinguish between **(1)**

CD11b$^{neg/lo}$CD11c^{pos} alveolar macrophages and **(2)** CD11b^{pos}CD11c^{pos} monocyte-

derived dendritic cells. Further differentiation of CD11b^{pos} cells was done using CD11c

and Gr1, as **(3)** Gr1^{hi}CD11c^{neg} neutrophils, **(4)** Gr1^{int}CD11c^{neg} monocytes, and **(5)** Gr-

1^{neg}CD11c^{neg} monocyte-derived recruited macrophages.

Supplementary Figure 2. Strain-dependent spread of *M. tuberculosis* in recruited

myeloid cells. (**A**) Cells were stained and analyzed by flow cytometry for quantitation of

subset and plated on 7H11 media (Figure 2). Depicted are the ratio of the average CFU

over the average cell number at each time point obtained post-infection in mice infected

with H37Rv, 4344, or ΔRD1. (**B**) Relative frequency of CFU recovered from myeloid cell

populations in the lungs of mice infected with *M. tuberculosis* H37Rv, ΔRD1, or 4334

after live flow sorting and plating of sorted cell fractions on 7H11 solid media. Results

are shown as mean $\pm$ SD in 1(ΔRD1), 2 (4334), and 6 (H37Rv) pooled experiments,

with n=5 mice per day and per strain in each experiment.

Supplementary Figure 3. *M. tuberculosis* strains exhibit differential ESX-1 activity.

H37Rv and 4334 were grown to logarithmic phase in 7H9, resuspended in Sauton's

media for 1 day, and filtrate proteins were reduced, alkylated, and trypsin digested prior

to run on LC-MS/MS. Spectra were searched against a combined database for H37Rv

and NITR203 on UniProt, using Andromeda search algorithm and MaxQuant for

quantitation. Statistical analysis done by Student t-test with Holm multiple comparisons

correction, 95% confidence interval, *$p<0.05$, and **$p<0.01$.

Supplementary Figure 4. *M. tuberculosis* strain 4334 induces higher levels of IFNγ

in the lungs of mice than H37Rv after two weeks of infection. (**A**) Total IFNγ levels

in the lung homogenate supernatants of mice infected with *M. tuberculosis* H37Rv or

4334. Results are shown as mean $\pm$ SD 4 experiments. Statistical significance was

assessed by unpaired Student's t-test for each day and Holm multiple comparisons

correction, with a 95% confidence interval and **p<0.01 and ***p<0.001. (**B**) Flow

cytometry gating strategy for the quantitation of IFNγ-producing CD4$^+$ T cells in the

lungs of mice infected with *M. tuberculosis* H37Rv or 4334 for 17 days.

Supplementary Figure 5. *M. tuberculosis* **strain 4334 induces a quantitatively**

greater adaptive immune response compared to H37Rv that is sustained 7 weeks

after initial infection and results in more lesions in the lung. Mice were aerosol

infected with 200-300 Colony Forming Units (CFU) of *M. tuberculosis* H37Rv or 4334

and harvested 7 weeks later. Lungs were processed to single cell suspensions and

plated on solid media for total CFU determination **(A)**, counted for total cell number **(B),**

and stained for flow cytometry analysis of cellular subsets **(D)** (see Supplementary

Figure 1 and Material and Methods for gating strategy). In addition, the left lung of

infected mice was fixed and stained with hematoxylin and eosin (H&E). **(C)** Using Fiji

open-source software, percentage of lung inflammatory infiltrate was calculated by

taking the total surface area of lung, subtracting contributions of airways and blood

vessels. Inflammatory infiltrates were quantitated by both manual contouring and

automated color density contouring, with the average of these values utilized for the

final percentage calculation. **(E)** Representative images of 5 µM sections at 40x (Leica

SCN400 F whole-slide scanner) are shown. Results are shown as mean ± SD of 4

mice. Statistical significance was assessed by unpaired Student's t-test and Holm

multiple comparisons correction with a 95% confidence interval and *p<0.05,

***p<0.001, and ****p<0.0001.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5