MACROSCALE THALAMIC FUNCTIONAL ORGANIZATION DISTURBANCES AND UNDERLYING CORE CYTOARCHITECTURE IN EARLY-ONSET SCHIZOPHRENIA

Running title: Expansion of thalamic functional hierarchies in schizophrenia

Yun-Shuang Fan [1,2], Yong Xu [8], Şeyma Bayrak [2], James M. Shine [3], Bin Wan [2,5,7], Haoru Li [1], Liang Li [1,6], Siqi Yang [1], Yao Meng [1], Sofie Louise Valk [2,5*#], Huafu Chen [1,4*#]

1. The Clinical Hospital of Chengdu Brain Science Institute, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, China; 2. Otto Hahn Group Cognitive Neurogenetics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany; 3. Brain and Mind Center, The University of Sydney, Sydney, Australia; 4. MOE Key Lab for Neuroinformation, High-Field Magnetic Resonance Brain Imaging Key Laboratory of Sichuan Province, University of Electronic Science and Technology of China, Chengdu, China; 5. Institute of Neuroscience and Medicine (INM-7: Brain and Behavior), Research Centre Jülich, Jülich, Germany; 6. Academy for Advanced Interdisciplinary Studies, Peking University; 7. International Max Planck Research School on Neuroscience of Communication: Function, Structure, and Plasticity (IMPRS NeuroCom), Leipzig, Germany; 8. Department of Psychiatry, First Hospital/First Clinical Medical College of Shanxi Medical University, Taiyuan, China.

* Both last co-authors contributed equally.

Corresponding authors:
Huafu Chen, E-mail: chenhf@uestc.edu.cn & Sofie Louise Valk, E-mail: valk@cbs.mpg.de.

Title character count: 111
Running title character count: 55
Word count of abstract: 191
Word count of text body: 4809
Number of references: 58
Number of tables: 1
Number of figures: 4 (All colored)
Supplement material: 1

Abstract

Schizophrenia is a polygenetic mental disorder with heterogeneous positive and negative symptom constellations, and is associated with abnormal cortical connectivity. The thalamus has a coordinative role in cortical function and is key to the development of the cerebral cortex. Conversely, altered functional organization of the thalamus might relate to overarching cortical disruptions in schizophrenia, anchored in development. Here, we contrasted resting-state fMRI in 99 antipsychotic-naive first-episode early-onset schizophrenia (EOS) patients and 100 typically developing controls to study whether macroscale thalamic organization is altered in EOS. Employing dimensional reduction techniques on thalamocortical functional connectome, we derived lateral-medial and anterior-posterior thalamic functional axes. We observed increased segregation of macroscale thalamic functional organization in EOS patients, which was related to altered thalamocortical interactions both in unimodal and transmodal networks. Using an *ex vivo* approximation of core-matrix cell distribution, we found that core cells particularly underlie the macroscale abnormalities in EOS patients. Moreover, the disruptions were associated with schizophrenia-related gene expression maps. Behavioral and disorder decoding analyses indicated that the macroscale hierarchy disturbances might perturb both perceptual and abstract cognitive functions and contribute to negative syndromes in schizophrenia, suggesting a unitary pathophysiological framework of schizophrenia.

Keywords: cytoarchitectural; early-onset schizophrenia; functional hierarchy; genetic; thalamus.

<u>Introduction</u>

Schizophrenia is a polygenetic psychiatric illness characterized by a combination of psychotic symptoms and motivational/ cognitive deficits, which usually emerge during early adulthood (1). Over the past two decades, a wealth of neuroimaging studies has indicated that schizophrenia can be associated with pathological interactions across widely distributed brain regions, instead of focal brain damages. Accordingly, the overarching dysconnection hypothesis posits that schizophrenia results from brain structural and functional connectivity abnormalities (2). The thalamus, which is well-placed to arbitrate the interactions between distributed brain organization (3), might play a pivotal role in the pathophysiological process of schizophrenia (4, 5).

The thalamus is a cytoarchitecturally heterogeneous diencephalic structure that contains an admixture of Parvalbumin (PVALB)-rich 'Core' cells and Calbindin (CALB1)-rich 'Matrix' cells (6). Whereas core cells preferentially target granular layers (Layers III and IV) of unimodal primary regions, such as primary visual, auditory and somatosensory cortices, matrix cells target supragranular layers (Layers I-III) over wide areas in a diffuse pattern (7). This means that distinct thalamic cells may interact with cortical areas organized into different topological zones (8). Cellular-scale information of the thalamus may be a critical factor to understand the thalamocortical interactions that supports cognition and behavior. Indeed, the thalamocortical system has been suggested to form the basis for binding multiple sensory experiences into a single framework of consciousness (9). By coordinating the

modular architecture of cortical networks, the thalamus has been reported to be engaged in

integrating information processing within the whole cerebral cortex (10).

In accordance with its function coordinating cortical network organization, the thalamus

plays the central role in the development of the cerebral cortex (11). During brain

development, the thalamus changes in concordance with the cerebral cortex and disturbances

of this coordinated process relate to cognitive dysfunctions (12), serving as a precursor of

schizophrenia. Thalamocortical dysconnectivity patterns, characterized by hypoconnectivity

with prefrontal regions and hyperconnectivity with sensorimotor areas, have been reported

in both pediatric (13) and early-stage (14) patients with schizophrenia. The dysconnectivity

pattern has been hypothesized to arise from disturbed brain maturation, particularly during

the   transition   from   youth   to   adulthood   (15).   Intriguingly,   thalamo-prefrontal

hypoconnectivity is correlated with thalamo-sensorimotor hyperconnectivity in patients,

potentially implying a shared pathophysiological mechanism (14). However, few studies

have investigated thalamocortical connectivity in the still developing brain of schizophrenia

from a comprehensive perspective.

Recently, the application of dimension reduction techniques has emerged as a promising

strategy for holistic representations of brain connectivity. These novel data-driven methods

decompose high dimensional connectome into a series of low dimensional axes capturing

spatial gradients of connectivity variations (16, 17). The gradient framework describes a

continuous coordinate system, in contrast to clustering-based methods resulting in discrete

communities (18). Using these methods in the context of cortex-wide functional connectome,

previous studies observed a cortical hierarchy that spans from unimodal primary regions to

transmodal regions (19), which has a close link with cortical microstructure like

cytoarchitecture or myeloarchitecture (20). Their coupling along the unimodal-transmodal

axis has been reported to be genetically- and phylogenetically-controlled, supporting flexible

cognitive functions (21). Perturbed macroscale cortical functional hierarchies have been

reported in various neurological (22, 23) and psychiatric disorders (18) including

schizophrenia (24). Also, thalamic hierarchies have been previously derived from

thalamocortical connectome, identifying a lateral-medial (L-M) principal gradient and an

anterior-posterior (A-P) secondary gradient (25). The L-M axis captures thalamic anatomical

nuclei differentiation, while the A-P axis characterizes unimodal-transmodal functional

hiearchy. Also the coupling between core-matrix cytoarchitecture and functional connectome

has been shown to describe the unimodal-transmodal cortical gradients, and argued to play a

major role in shaping functional dynamics within the cerebral cortex (8). Given the possible

implication of the thalamus in schizophrenia, thalamic hierarchies may be altered during

brain maturation and could provide new insights into the disrupted thalamocortical

organization in schizophrenia.

Here, we leveraged a cohort of individuals with early-onset schizophrenia (EOS), a disorder

that is neurobiologically continuous with its adult counterpart (26), to examine whether

macroscale thalamic functional organization shows disturbances in the still developing brain

of schizophrenia, mirroring neocortical reports. To this end, we first evaluated functional

hierarchies of the thalamus by employing dimension reduction techniques on

thalamocortical functional connectome (27). We then embedded thalamic functional

hierarchies in a neurobiological context by spatially correlating the macroscale patterns with

gene expression maps from the Allen Human Brain Atlas (AHBA) (28). Last, we tested

whether the functional hierarchies could estimate clinical symptoms of patients using a

machine learning regression strategy.

Materials and Methods

Participants

Ninety-nine antipsychotic-naive first-episode EOS patients and 100 typically developing

(TD) controls were recruited from the First Hospital of Shanxi Medical University and the

local community through advertisements, respectively. All pediatric participants were 7–17

years old. The diagnosis of schizophrenia was in accordance with the Structured Clinical

Interview for DSM-IV, and was confirmed by at least one senior psychiatrist (Y.X.) through

a structured clinical interview after at least 6-months of follow-up. The psychiatric

symptomatology of 71 patients was evaluated using the Positive and Negative Syndrome

Scale (PANSS). Exclusion criteria for all subjects included 1) >= 18 years old; 2)

neurological MRI anomalies; 3) any electronic or metal implants; or 4) substance abuse. In

addition, EOS patients were excluded if they suffered from the illness for > 1 year, and TD

controls were excluded if they and their first-degree relatives had any history of psychiatric

disorder. This retrospective study was approved by the Ethics Committee of the First

Hospital of Shanxi Medical University. Written consent was obtained from every participant

and their parents or legal guardians.

Four patients and two controls were excluded from the study because of incomplete

scanning data, nine patients and six controls due to excessive head motion [mean frame-wise

displacement, (FD) > 0.2 mm or outliers > 50%], and one control due to poor quality of

intrasubject brain registration. Ultimately, 86 EOS patients and 91 demographically-matched

TD controls were included in the analysis (See **Table 1** for detailed demographic data).

Table 1. Demographics and clinical data.

Characteristic	EOS	TD	Group comparisons	
			Statistic-Values	P-Values
Demographic sample	n=86	n=91		
Sex (male/female)	31/55	33/58	0.0009 [a]	0.98
Age (years)	14.56 ± 1.95	14.34 ± 2.02	3711 [b]	0.55
Handedness (right/left)	86/0	91/0	–	–
Mean FD (mm)	0.07 ± 0.03	0.07 ± 0.03	3843 [b]	0.84
Clinical sample	n=65	–		
PANSS total scores	65.4 ± 17.41	–		
PANSS general scores	31.68 ± 8.71	–		
PANSS positive scores	15.14 ± 5.11	–		
PANSS negative scores	14.00 ± 5.55	–		

119   **Note:** Mean ± SD; [a] The χ2 value for gender distribution was obtained by chi-square test; [b]

120   The U values were obtained by Mann–Whitney tests.

121   **Abbreviations:** EOS, early-onset schizophrenia; TD, typically developing; FD, frame-wise

122   displacement; PANSS, Positive and Negative Symptom Scale.

MRI acquisition

Multimodal MRI data were collected using a 3 Tesla MRI scanner (MAGNETOM Verio; Siemens, Germany) in the First Hospital of Shanxi Medical University. T1-weighted data were acquired using a three-dimensional fast spoiled gradient-echo sequence [repetition time (TR) = 2,300 ms; echo time (TE) = 2.95 ms; flip angle = 9°; matrix = 256 × 240; slice thickness = 1.2 mm (no gap); and voxel size = $0.9375 \times 0.9375 \times 1.2$ mm^3, with 160 axial slices]. Resting-state functional MRI (rs-fMRI) data were obtained using a two-dimensional echo-planar imaging sequence [TR = 2,500 ms; TE = 30 ms; flip angle = 90°; matrix = 64 × 64; number of volumes = 198; slice thickness = 3 mm (1 mm gap); and voxel size = $3.75 \times 3.75 \times 4$ mm^3, with 32 axial slices].

MRI processing

T1-weighted structural data were preprocessed with FreeSurfer (v7.1.0, http://surfer.nmr.mgh.harvard.edu/), which included cortical segmentation and surface reconstruction. Rs-fMRI functional data were preprocessed with the CBIG pipeline (https://github.com/ThomasYeoLab/CBIG) based on FSL [v5.0.9, (29)] and FreeSurfer (v7.1.0), which included removal of the first four volumes, slice-timing, motion correction, and boundary-based registration to structural images. See **Supplement 1** for further details. Preprocessed images were then registered to MNI152 template and resampled to the cortical surface using Ciftify package [v2.3.3, (30)]. The thalamus was localized using the Gordon 333 Atlas (31), including 2536 voxels across both hemispheres.

Macroscale thalamocortical gradient identification

Analogous to previous work (25), gradients of thalamocortical functional connectome were generated to describe thalamic functional organization using the diffusion embedding algorithm in BrainSpace Toolbox (32). Thalamocortical functional connectivities were first calculated based on Pearson correlations between the thalamic and cortical rs-fMRI time-series for each subject (16, 23), and then converted into cosine similarity matrices (16, 17). Subsequently, nonlinear dimensionality reduction techniques were employed on similarity matrices to resolve connectome gradient, i.e., spatial axis in connectome variations (27). See **Supplement 1** for detailed analyses. We selected the first two gradients to represent the macroscale thalamic connectome space, which explained 44% of the total eigenvariance in functional connectome (**Figure S2**). The relative positioning of thalamic voxels along each organizational axis describes similarity of their functional connectivity profiles. To quantify the dispersion of each thalamic voxel in the two-dimensional gradient space, we computed eccentricity, i.e., the square root of the Euclidian distance from each thalamic voxel to the center of mass in the two-dimensional gradient space (33). For each individual, global eccentricity was calculated by averaging eccentricity values across all thalamic voxels, indicating overall dispersion of the gradient space. We additionally explored cortical-thalamic gradients by generating cortical similarity matrices from cortical-thalamic connectivity profile (see **Supplement 2** for further details).

Thalamic functional community division

To characterize the functional relevance of macroscale thalamic gradient space, we created a

thalamic functional atlas using winner-take-all representation approach (25, 34). Partial

correlations were computed between rs-fMRI time-series of each thalamic voxel and six

cortical functional networks (35) including the visual network (VIS), sensorimotor network

(SMN), dorsal attention network (DAN), ventral attention network (VAN), frontoparietal

network (FPN), and default mode network (DMN). The limbic network was excluded for

low signal quality in corresponding cortical areas of our data. Each thalamic voxel was

labeled by the functional network showing the highest partial correlation coefficient.

Additionally, to explore cortical correspondences of thalamic gradients, we projected

thalamic gradients onto the cerebral cortex. For each cortical vertex, gradient projection was

calculated by correlating its cortical-thalamic connectivity profiles with thalamic gradients.

These cortical maps were down sampled into 400 cortical parcels and grouped into

functional networks (36) to further validate thalamic functional community division.

The core-matrix cytoarchitecture

To delineate the core-matrix cytoarchitecture in the thalamus, we used the spatial maps of

mRNA expression levels for two calcium-binding proteins (CALB1 and PVALB)

(https://github.com/macshine/corematrix) (8) generated from post-mortem Allen Human

Brain Atlas (28). Thalamic voxels with positive CP values (CALB1-PVALB values) related

to matrix projection cells, and voxels with negative values related to core populations.

Interregional correlations between CP map and differential eccentricity map were employed

to reveal an association between the thalamic cytoarchitecture and disturbed functional

organization in EOS patients. The variogram-matching model was used to correct for the

spatial autocorrelation of brain maps (37). Subsequently, we projected CP maps onto the

cerebral cortex to reveal couplings between the core-matrix cytoarchitecture and functional

connectome. Cortical parcels with positive coupling values indicated as preferential

associations with matrix thalamic populations, and negative values suggested core

populations. To evaluate cognitive terms associated with gene-connectome coupling maps,

we further conducted topic-based behavioral decoding using NeuroSynth meta-analytic

database (38). See **Supplement 1** for detailed analysis steps.

Clinical correspondences

To investigate clinical significance of thalamic functional organization disturbances, we

further associated the macroscale functional phenotype with schizophrenia-related genetic

expression. Based on previous work (39), we selected out 28 protein-coding genes

particularly implicated in schizophrenia etiology or treatment (See **Table S1** for detailed

information). Within an overlapping thalamic mask (1969 voxels), estimated mRNA levels

were extracted from Allen Human Brain Atlas and then normalized using a robust sigmoid

function (40). Gene expression maps were then spatially correlated with differential

eccentricity map between the EOS and TD groups, corrected by the variogram-matching

model (1000 surrogate maps) as abovementioned.

Second, we estimated the relationship between thalamic functional hierarchies and clinical

presentations in EOS patients. Following a machine learning pipeline, we used the elastic

net regression model to predict clinical symptoms in EOS (41). Eccentricity values of the

two-dimensional gradient space were defined as input features, and PANSS positive and

negative scores were used as predictors. The model performance was evaluated by

comparison of observed and predicted clinical scores. See **Supplement 1** for detailed

analyses.

Group comparisons between the EOS and TD groups

Between-group differences on all kinds of measurements were assessed by using two-sample

t-tests with covariates including age, gender, and mean FD. The multiple comparison

corrections were conducted using three methods for different spatial scales: voxel-wise,

parcel-wise, and global-wise. For voxel-wise thalamic gradient and eccentricity values,

multiple comparison corrections were employed using the permutation-based threshold-free

cluster enhancement (TFCE, 10,000 permutations, $p < 0.005$) method (FSL-PALM), which

could improve sensitivity and interpretable than cluster-based thresholding method (42). For

parcel-wise gene-connectome coupling values, false discovery rate (FDR) corrections ($p <$

0.005) were used to control the effect of false positives. For global-wise index like

network-level gradient values and global eccentricity values, Bonferroni corrections were

conducted with a significant level of $p < 0.05$.

Results

Macroscale thalamic gradients in TD and EOS

The principal gradient (G1, 24.6% explained) of the thalamus revealed a L-M axis, and the

second gradient (G2, 11.3% explained) described an A-P axis, in line with previous work

(25). In **Figure S2**, we also showed the third gradient pattern running in ventral-dorsal

direction (9% explained). In EOS patients, we observed expansions at both anchors of the

G1; the lateral portions including the ventral lateral and ventral posterior thalamic nuclei, the

medial dorsal areas compared to TD controls (**Figure 1A**, TFCE, $p < 0.005/2$). We also

observed expansions along the G2 axis including the pulvinar and anterior nuclear groups

(**Figure 1B**, TFCE, $p < 0.005/2$). Combing G1 and G2 axes, we computed an eccentricity

score using the square root of the Euclidian distance from each thalamic voxel to the center

of mass in the two-dimensional gradient space (33). Global eccentricity was assessed for

each participant by averaging eccentricity values across all thalamic voxels. We found

significantly increased global eccentricity for thalamic voxels in EOS compared to TD (t =

2.34, $p = 0.02$), indicating a segregation of macroscale thalamic functional organization in

patients (**Figure 1C**). Additionally, we explored the spatial pattern of cortical-thalamic

gradients, reflecting the organization of cortical-thalamic connectivity in the cortex (**Figure**

S3). The first cortical-thalamic gradient was similar with previous cortical functional

gradients, showing a unimodal-transmodal transition pattern. See **Supplement 2** for further

discussions.

Figure 1. Thalamic gradients in typically developing (TD) controls and early-onset schizophrenia (EOS) patients. (**A**) The group-level primary gradients (G1) for TD and EOS, and their between-group differences. The G1 depicts a transition from lateral to medial portions of the thalamus. Thalamic voxels showing significant G1 score differences were surrounded by black contours [t-test, EOS vs. TD; threshold-free cluster enhancement (TFCE), $p < 0.005$]. The density map represents the distribution of G1 loading for EOS (green) or TD (gray). (**B**) The group-level secondary gradients (G2) for TD and EOS, and their differences. G2 separates the anterior thalamic portions from the posterior portions. Thalamic voxels with significant G2 differences were surrounded by black contours. The

density map represents the G2 loading for EOS (green) or TD (gray). (**C**) Gradient spaces
built on the group-level G1 and G2, separately for TD and EOS. Each point represents a
thalamic voxel embedded in the gradient space. Voxels are color coded based on their mean
eccentricity scores across subjects. Eccentricity score was computed by the square root of
the Euclidian distance from each thalamic voxel to the center of mass in the
two-dimensional gradient space. Higher eccentricity indicates greater segregation, e.g.,
larger dissimilarity of thalamocortical connectivity, in the gradient space. The density plot
depicts the distribution of eccentricity scores in EOS and TD.

Functional relevance

Utilizing a whole-brain functional network parcellation (35), thalamocortical functional

connectome was distributed into six functional communities (**Figure 2A**). We then averaged

gradient loadings within each community for each subject and compared them between the

EOS and TD groups (**Figure 2B**, Bonferroni correction, $p < 0.05/6$). EOS patients had

significantly higher mean G1 scores in the SMN (t = 5.37, $p < 0.0001$) and DAN (t = 2.94, p

= 0.004), and lower G1 scores in the VAN (t = −2.85, p = 0.005), FPN (t = −3.69, p =

0.0003), and DMN (t = −3.03, p = 0.003) relative to TD controls. For the G2 axis, patients

showed decreased gradient scores in the VIS (t = −4.08, $p < 0.0001$), and increased in the

DMN (t = 5.45, $p < 0.0001$). In the two-dimensional gradient space showing functional

relevance (**Figure 2C**), EOS patients had apparent dissociation of SMN-related thalamic

voxels along the G1 axis, and a larger dissociation of VIS- and DMN-related regions along

the G2 axis.

Moreover, rather than dividing the thalamus into functional communities, we also projected

thalamic gradients onto the cerebral cortex and evaluated their correspondence with cortical

functional networks (**FigureS4**). Both cortical projections of G1 and G2 tended to follow the

unimodal-transmodal transition. However, the VAN was located in the unimodal part of G1

projection, while the transmodal portion of G2 projection. Nevertheless, the dissociation of

SMN-related thalamic voxels in EOS was supported by results of G1 projection, and the

VIS-associated dissociation of G2 axis was also replicated.

Figure 2. Thalamic gradients distributed into functional communities. (A) Network-level representations of the thalamocortical connectome. (**B**) G1 and G2 scores within functional communities. Density maps indicate gradient scores (top left: G1, bottom left: G2) within six networks for EOS and TD. Network level differences between EOS and TD were accessed by t-tests, and significant differences are depicted by * (Bonferroni correction, $p < 0.05$) and ** ($p < 0.0001$). Thalamic G1 (top right) and G2 (bottom right) representations across all participants. Voxels showing higher gradient values in EOS are colored red, whereas voxels with lower gradient values are colored blue. (**C**) Gradient space representation of the thalamus together with functional communities for TD (left) and EOS (right). Thalamic voxels are situated based on their G1 (x-axis) and G2 (y-axis) scores, and are colored according to their network assignment (purple-VIS; blue-SMN; red-DMN; grey-all other networks). ► stands for the G1 score differences between EOS and TD; ◄ for the G2 score differences; ♦ for both G1 and G2 score differences; ● for no difference. VIS, the visual network; SMN, the sensorimotor network; DAN, the dorsal attention network; VAN, the ventral attention network; FPN, the frontoparietal network; DMN, the default mode network.

The core-matrix cytoarchitectural basis

Having established macroscale thalamic connectome gradients, we further investigated

whether connectome differences between patients and controls were specific to thalamic

cytoarchitectural features, namely core vs. matrix cells, motivated by a previous work (8).

Thus, mRNA expression levels for CALB1 (core type cells) and PVALB (matrix type cells)

were separately assessed at the thalamic voxel level (**Figure S5A**). The differential

expression level between CALB1 and PVALB, i.e., CP scores, was used to delineate

thalamic core/matrix type cell distribution, where positive/negative CP value indicated with

matrix/core cell type, respectively. Eccentricity was used to evaluate the gradient

space-embedded position of each thalamic voxel (**Figure S5B**). EOS patients showed

increased eccentricity in medial dorsal, ventral lateral, and ventral anterior nucleus compare

to TD controls, indicating their increased dispersion from the center of the gradient space

(TFCE, $p < 0.005$).

We then mapped the core-matrix cytoarchitectural features onto the two-dimensional

gradient space (**Figure 3A**). Core cells showed mildly increased global eccentricity in EOS

patients compared with core cells of TD ($t = 2.38$, $p = 0.02$), but matrix cells did not ($t = 1.62$,

$p = 0.11$). No significant difference was found between global eccentricity of core and

matrix populations within EOS ($t = 1.07$, $p = 0.29$) or TD ($t = 0.36$, $p = 0.72$). Next,

Pearson's correlations were used to quantify group-level associations between the

cytoarchitecture and the connectome gradient disturbances, and the variogram methods that

control for the spatial autocorrelations were used to determine statistical significance levels

(**Figure 3B**). A negative correlation was found between CP values and t values of

eccentricity (r = −0.29, p_{vario} = 0.002). In detail, increased dispersion of macroscale

connectome gradient space in patients was particularly related to core cells (r = 0.31, p_{vario} <

0.0001), rather than matrix cells (r = 0.02, p_{vario} = 0.84).

Cognitive relevance

Given the association between CP map and thalamocortical connectome, we further assessed

whether gene-connectome coupling contributes to cognitive processing. Thus, we projected

the core-matrix cytoarchitecture to the cerebral cortex and then conducted a behavioral

decoding using the NeuroSynth database (38). Couplings between functional connectome

and gene expression were computed and further down sampled into 400 cortical parcels. As

shown in **Figure S6**, core cell populations were mainly associated with unimodal regions

that subserve primary sensory and multisensory functions. Matrix cell populations were

associated with transmodal cortices characterized by more abstract cognition.

Compared with TD controls, EOS patients showed increased gene-connectome couplings in

the DMN including the inferior parietal lobule, posterior cingulate cortex, inferior prefrontal

cortex, temporal areas, and the FPN including the precuneus and inferior parietal sulcus, and

temporoparietal junction network (**Figure 3C,** FDR, $p < 0.005$). These abnormal coupling

increases were associated with seven cognitive topics including declarative memory,

autobiographical memory, working memory, verbal semantics, social cognition, language,

and visuospatial (z−statistic > 3.1), indicating its implications in higher-level cognitive

processes. Additionally, reduced gene-connectome couplings were observed in the VIS

including the extrastriate and inferior extratriate, and the DAN including postcentral regions

and superior parietal lobule, as well as the SMN. These reductions were characterized by

low-level visual sensory and motor functions involving visual perception, visual attention,

motor, action, eye movements, visuospatial, multisensory processing, and reading (**Figure**

3D).

A. Thalamic Core-Matrix cytoarchitecture in two-dimensional gradient space

B. Associations between gene expression levels and eccentricity abnormalities

C. Gene-connectome coupling abnormalities

D. Cognitive topics-decoding

Figure 3. The core-matrix cytoarchitecture of the thalamus. **(A)** Core-matrix cytoarchitectural features of thalamic voxels projected onto the two-dimensional gradient space for TD and EOS. Rainclouds present group comparisons of global eccentricity in four fashions: global eccentricity values across core cells in EOS vs. global eccentricity values across matrix cells in EOS; core cells in TD vs. matrix cells in TD; core cells in EOS vs. core cells in TD; matrix cells in EOS vs. matrix cells in TD. **(B)** Associations between eccentricity differences (t values; EOS vs. TD) and differential gene expression levels (CP, left) as well as single gene expression levels (CALB1 and PVALB, right). Correlations were obtained across thalamic voxels (Pearson r values) and their significances were tested using variogram approach (p_{vario} values). **(C)** Parcel-wise couplings between functional connectome and gene expression for TD and EOS, and their differences (t-test, EOS vs. TD).

Cortical parcels with positive coupling values (blue) indicate as preferential associations
with matrix thalamic cells, and negative coupling values (green) suggested core cells.
Parcels with significantly different coupling patterns in patients are surrounded by black
contours [false discovery rate (FDR) corrections, $p < 0.005$]. **(D)** Topic-based behavioral
decoding of regions with abnormal couplings in EOS. Cognitive terms in warm color
correspond to brain regions showing hyper-couplings in patients relative to controls, and
cool color represent hypo-couplings. In the word cloud, the size of a cognitive term is
proportional to its loading strength for decoding an input brain mask.

Clinical relevance

To reveal clinical significance of the macroscale functional gradients, we first explored its

associations with schizophrenia-related gene expression maps, based on a previous work (39)

(**Figure 4A**). In particular, spatial correlations were conducted to quantify their relationships,

significance levels of which were evaluated by the variogram methods. The difference map

of eccentricity scores between patients and controls was significantly correlated with mRNA

expression levels for Glutamatergic neurotransmission-related genes including GRIN2A

(r=0.34, p_{vario} =0.001), GRIA1 (r=-0.23, p_{vario} =0.03), Calcium signaling-related genes

RIMS1 (r=0.26, = p_{vario} 0.005), synaptic function and plasticity-related genes including

CNTN4 (r=0.32, p_{vario} =0.001) and SNAP91 (r=0.24, p_{vario} =0.02), other neuronal ion

channels-related genes including HCN1 (r=0.25, p_{vario} =0.03) and CHRNA5 (r=0.33, p_{vario}

=0.001), neurodevelopment-related genes including BCL11B (r=-0.23, p_{vario} =0.02) and

FAM5B(r=0.27, p_{vario} =0.006). No significant correlation was observed in the other 19 genes,

including therapeutic target-related genes.

Second, we investigated whether thalamic gradients could predict clinical symptoms in EOS.

We used eccentricity scores as input features in a linear regression model to estimate PANSS

positive and negative scores of patients. The optimal model parameter (L1 ratio) was 0.3 for

positive symptoms prediction, and 0.1 for negative symptoms (**Figure S7**). Eccentricity

scores of thalamic voxels performed moderately when predicting the severity of negative

symptoms (r = 0.17 ± 0.25, MAE = 4.47 ± 0.83), while poor in the prediction of positive

symptoms (r = 0.06 ± 0.23, MAE = 4.21 ± 0.69). For positive and negative symptoms,

predictive models with median performance were separately reported in **Figure 4B**. In the

model predicting negative symptoms, weights were heavier in the thalamic voxels regarding

to transmodal networks such as the VAN and DMN.

Figure 4. Clinical relevance of thalamic gradients in EOS. (A) The relationship between
the macroscale functional phenotype and 28 schizophrenia-related gene expressions.
Interregional correlations (Pearson r coefficients) were employed between eccentricity
differences (t values; EOS vs. TD) and gene expression levels, and their significances were
evaluated by variogram approach (★ represents p_{vario} value < 0.05). In the left panel, the
size of a square is in proportion to absolute value of corresponding Pearson r coefficient, and
its color is coded by the sign of r value (red-positive; blue-negative). The highest positive
correlation with eccentricity differences was found at a glutamatergic neurotransmission
protein-coding gene, i.e., GRIN2A, whose expression pattern is shown in the top corner. The
highest negative correlation was found at BCL11B, a neurodevelopment-related gene. **(B)**
The relationship between thalamic functional organization and clinical presentations.
Eccentricity values in patients were used to predict PANSS positive and negative scores
based on a linear regression model. The model performance was evaluated by comparing
observed and predicted clinical scores. This procedure including model learning and testing
was repeated 101 times, generating the distributions of Pearson r coefficients and mean
absolute error (MAE). The model with median performance was reported here, and absolute
value of feature coefficient was used as weight for each thalamic voxel (a point in the

two-dimensional gradient space). The size of a point is coded by predictive weight, and is
colored according to its network assignment (purple-VIS; blue-SMN; green-DAN;
rose-VAN; orange-FPN; red-DMN).
420

<u>Discussion</u>

In the current study we investigated macroscale thalamic functional organization in EOS

through dimensionality reduction techniques on thalamocortical functional connectivity. We

found both expansions along L-M principal axis and A-P secondary axis of thalamic

hierarchies in EOS, indicating connectivity profiles between both anchors showed higher

dissimilarities in patients versus controls. Disordered functional hierarchies of the thalamus

were related to altered thalamocortical interactions both in unimodal and transmodal

networks. To evaluate the cytoarchitectural underpinnings of the macroscale functional

organization disturbances in EOS, we compared alterations in functional organization within

core- and matrix-cells that derived from an independent transcriptomic atlas. We found that

in particular, functional organization of thalamic core cells was altered in EOS. Patients'

abnormal coupling patterns between a-priori map of the core-matrix cytoarchitecture and

functional connectome characterized a spectrum from perceptual to abstract cognitive

functions. Moreover, transcriptomic-informed analyses suggested a close relationship

between macroscale functional organization and schizophrenia etiology-related gene

expressions in the thalamus. Employing a machine learning strategy, we found that thalamic

functional organization was able to predict negative symptoms in EOS. In sum, the current

findings provide mechanistic evidence for disrupted thalamocortical system in EOS, and

point to alterations in functional networks associated with both perceptual and cognitive

functions, suggesting a unitary pathophysiology of heterogeneous symptoms in

schizophrenia.

In line with previous work on thalamic hierarchies (25), the principal functional gradient

described continuous transition from the ventral lateral nucleus to the anterior and pulvinar

groups, and the second axis delineated gradual transition from the anterior nuclei to the

pulvinar. Whereas L-M thalamic axis has been reported to correspond to the distribution of

gray matter morphology, i.e., low-to-high intensity of neural mass, and the A-P gradient was

strongly related to the intrinsic geometry of the thalamus. These findings suggested an

association between functional thalamic hierarchies and its structure. Albeit not directly

shown, L-M and A-P axes may reflect functional relevance in different dimensions, i.e., two

kinds of transitions across functional networks. Indeed, we found that the L-M principal axis

functionally segregated the VIS and SMN, similar to the second gradient of cortical

connectome (16). Conversely, the A-P axis of thalamic hierarchies segregated unimodal and

transmodal networks, in accord with previous findings (25). Taken together, beyond

supporting macroscale thalamic hierarchical framework, the current findings further broaden

our knowledge of functional specialization of thalamic L-M and A-P axes.

Thalamic ventral lateral and ventral posterior nuclei, as one end of the L-M organizational

axis, exhibited evident dissociation in EOS. Both nuclei receive neuronal input from the

sensory periphery, and project to the motor and somatosensory cortices, respectively (43).

The etiology of schizophrenia has been suggested to damage refinement of

motor/somatosensory-thalamic connectivity patterns that occurs during brain maturation

(15). Indeed, in adolescent patients relative to adult patients, structural abnormalities in the

sensorimotor cortex are reported to be particularly salient (44), but may gradually fade out

within a longitudinal period of observation (45). Moreover, motor performance has been

reported worse in adolescent patients relative to adult patients when accounting for

developmental factors (46). In line with this observation, a meta-analysis suggested that

motor deficits may precede the onset of schizophrenia and may constitute robust antecedents

of this mental disorder (47). In the context, we postulate sensorimotor-related segregation

along thalamic L-M hierarchy might underlie premorbid disturbances in motor development,

a marker distinct to schizophrenia.

Compared with TD, EOS patients had increased segregation in two extremes of the A-P axis,

i.e., visual-related pulvinar nucleus and default mode-associated anterior nuclear group.

Weaker functional connectivity between the VIS and DMN has been previously reported in

EOS (48, 49). In fact, given the central role of the thalamus in the development of the

cerebral cortex (11), abnormalities of the cerebral cortex in schizophrenia might occur

secondary to thalamic pathology (4, 50). Thus, the unimodal-transmodal thalamic hierarchy

expansion might further result in disturbed cortical differentiation of unimodal and

transmodal regions in EOS. A compression of the unimodal-to-transmodal cortical hierarchy

was recently found in chronic adult-onset schizophrenia (51), contrasting with our

observation of cortical-thalamic hierarchy expansion. Given age-dependent shifts in the

macroscale cortical hierarchy (52), this inconsistence might due to their disparate stage of

the illness and age of onset, or their usage of antipsychotic drugs. Further longitudinal works

are needed to chart functional organization abnormalities of the thalamus and the cerebral

cortex during the course of schizophrenia. Nevertheless, the current findings embed

thalamus into a cortical functional organization linked to differentiation of sensory from

abstract cognitive functions, paving the way to comprehensively reveal cognitive defects,

another well documented precursor of schizophrenia excepts for motor deficits (47).

Leveraging our observations of alterations of thalamic functional organization against a

proxy map of core/matrix cells based on post-mortem transcriptomic data (8), we observed

thalamic core cells to underlie expansive functional hierarchies in EOS, rather than matrix

cells. Core and matrix cells are two primary types of thalamic relay neurons which

separately exhibit immunoreactivity to the calcium binding proteins Parvalbumin and

Calbindin (9). Thalamic nuclei differ in the ratio of core and matrix neurons (53).

Specifically, sensory and motor relay nuclei, as well as the pulvinar nuclei and the

intralaminar nuclei are chiefly composed of core cells (54). Compared with matrix cells, the

larger core cells innervate middle cortical layers in a more area-restricted and

topographically-organized fashion (9). Functionally, Parvalbumin-rich core cells have been

reported to act as drivers of feed-forward activity, while Calbindin-rich matrix cells fulfil a

more modulatory function (7). Together, this may suggest that thalamic hierarchy

disturbances in EOS may relate to the "feed-forward" pathway that transmits information

from the sensory periphery, not the "feed-back" pathway (55).

A further inspection of gene-connectome coupling abnormalities in EOS by evaluating the

selective connection of the core-matrix cytoarchitecture with the cortex could show that,

patients' core thalamus dys-connected with both unimodal and transmodal cortices. In

healthy adults, core regions have preferential connections with unimodal primary regions,

and matrix areas with transmodal cortices incorporating the DMN, FPN, VAN and the limbic

network (8). Conversely in the current pediatric sample, we observed core-related

connections with wide-spread cortical regions including both unimodal and posterior

transmodal cortices for TD, whereas EOS had lager similarity with the previously reported

adult pattern, i.e., stronger core-related couplings in the VIS, SMN and DAN and weaker

couplings in the DMN and FPN. Together, our results imply putatively excessive maturation

in the thalamocortical feedforward pathway of schizophrenia. However, future

molecular-level work is undoubtedly needed to elaborate on the feedforward pathway

alterations in the still developing brain of schizophrenia.

It has been suggested that schizophrenic brain may not form connections according to gene

encoded blueprints which have been phylogenetically determined to be the most efficient

(56). In line with this, we observed that impaired thalamic hierarchy in EOS was highly

associated with schizophrenia-related gene expressions, especially genes encoding

Glutamatergic neurotransmission and neurodevelopmental proteins (39). Our findings reveal

a gene-connectome correspondence in the thalamocortical system of EOS, adding new

evidence for genetics of schizophrenia. However, there is an obvious shortage in our

gene-related analyses. The gene expression levels were assessed from post-mortem brain

tissue of adults (28), which might be different from the pediatric human brain. Limited by

the lack of pediatric transcriptional atlas, our findings about the association between

macroscale connectome topology and gene architecture should be carefully considered.

Behavioral decoding of the gene-connectome coupling pattern derived a sensory-cognitive

architecture, describing functions associated with primary sensory and multisensory

processing to working memory, cognitive control and motivation. Along the continuous

behavioral spectrum, perception (especially visual sensory), motor, and higher cognition

such as memory are particularly affected by schizophrenia. Consistently, thalamic hierarchy

abnormalities were sensorimotor-related, as well as visual/ default mode-associated along a

second axis. As the "cognitive dysmetria" theory suggested, the multitude and diversity of

behavior deficits in schizophrenia might be tied to an impaired fundamental cognitive

process mediated by the thalamus (56, 57). This impairment, i.e., cognitive dysmetria

referred to a disruption of the fluid and coordinated sequences of both thought and action,

leading to a decreased coordination of perception, retention, retrieval, and response

functions (58). In particular, our study suggested that abnormal thalamic hierarchy was

closely related to negative symptoms of schizophrenia, i.e., a diminution of functions related

to motivation and interest. Compared to positive symptoms (such as delusions), negative

symptoms are more complex and likely to be the result of systematic disruption. Effective

treatment of negative symptoms has long been a clinical challenge for its poor outcomes.

The current study provides a thalamic hierarchy framework for heterogeneous behavior

deficits related to negative symptoms in schizophrenia, which might denote future therapy of

the resistant symptoms.

In sum, the current study describes thalamic functional organization abnormalities in EOS,

which could be related to "feed-forward" core thalamocortical pathway. The macroscale

disruptions were related to schizophrenia-related genetic factors. Crucially, it might perturb

behaviors involving both low-level perception and high-level cognition, resulting in diverse

negative syndromes in schizophrenia.

Data and code availability

The data that support our findings are available from the corresponding author upon reasonable request. The estimated spatial maps of mRNA expression levels were downloaded at: https://www.meduniwien.ac.at/neuroimaging/mRNA.html. The code for functional gradient analysis was adapted from the MICA lab (http:/mica-mni.github.io) and is available at https://github.com/Yun-Shuang/Thalamic-functional-gradient-SZ. The code for behavioral decoding was adapted from https://github.com/NeuroanatomyAndConnectivity/gradient_analysis/blob/master/05_metaanalysis_neurosynth.ipynb. Statistical analyses were carried out using PALM (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/PALM) and BrainSMASH (https://brainsmash.readthedocs.io/). Machine learning analyses were based on scikit-learn package (https://scikit-learn.org/stable/modules/generated/sklearn.linear_model.ElasticNetCV. html). Results were visualized using Connectome Workbench (https://www.humanconnectome.org/software/connectome-workbench), and Seaborn (https://seaborn.pydata.org/) in combination with ColorBrewer (https://github.com/scottclowe/cbrewer2).

Acknowledgements

We are grateful to all the participants and their guardians in this study. This work was supported by the National Natural Science Foundation of China (82121003, 62036003, 62073058, 62173070), Innovation Team and Talents Cultivation Program of National Administration of Traditional Chinese Medicine (ZYYCXTD-D-202003). S.L.V. was also funded in part by Helmholtz Association's Initiative and Networking Fund under the

Helmholtz International Lab grant agreement InterLabs-0015, and the Canada First Research

Excellence Fund (CFREF Competition 2, 2015-2016) awarded to the Healthy Brains,

Healthy Lives initiative at McGill University, through the Helmholtz International BigBrain

Analytics and Learning Laboratory (HIBALL).

<u>Disclosures</u>

The authors declare that they have no conflict of interest.

References

1. Kinney DK, Matthysse S (1978): Genetic transmission of schizophrenia. *Annual review of medicine*. 29:459-473.

2. Stephan KE, Friston KJ, Frith CD (2009): Dysconnection in schizophrenia: from abnormal synaptic plasticity to failures of self-monitoring. *Schizophrenia bulletin*. 35:509-527.

3. Shine JM (2021): The thalamus integrates the macrosystems of the brain to facilitate complex, adaptive brain network dynamics. *Progress in neurobiology*. 199:101951.

4. Jones EG (1997): Cortical development and thalamic pathology in schizophrenia. *Schizophrenia bulletin*. 23:483-501.

5. Pergola G, Selvaggi P, Trizio S, Bertolino A, Blasi G (2015): The role of the thalamus in schizophrenia from a neuroimaging perspective. *Neuroscience and biobehavioral reviews*. 54:57-75.

6. Jones EG (2009): Synchrony in the interconnected circuitry of the thalamus and cerebral cortex. *Annals of the New York Academy of Sciences*. 1157:10-23.

7. Jones EG (2001): The thalamic matrix and thalamocortical synchrony. *Trends in neurosciences*. 24:595-601.

8. Muller EJ, Munn B, Hearne LJ, Smith JB, Fulcher B, Arnatkeviciute A, et al. (2020): Core and matrix thalamic sub-populations relate to spatio-temporal cortical connectivity gradients. *NeuroImage*. 222:117224.

9. Jones EG (1998): Viewpoint: the core and matrix of thalamic organization. *Neuroscience*. 85:331-345.

10. Hwang K, Bertolero MA, Liu WB, D'Esposito M (2017): The Human Thalamus Is an

Integrative Hub for Functional Brain Networks. *The Journal of neuroscience : the official journal of the Society for Neuroscience.* 37:5594-5607.

11. Lopez-Bendito G, Molnar Z (2003): Thalamocortical development: how are we going to get there? *Nature reviews Neuroscience.* 4:276-289.

12. Nakagawa Y (2019): Development of the thalamus: From early patterning to regulation of cortical functions. *Wiley interdisciplinary reviews Developmental biology.* 8:e345.

13. Zhang M, Palaniyappan L, Deng M, Zhang W, Pan Y, Fan Z, et al. (2021): Abnormal Thalamocortical Circuit in Adolescents With Early-Onset Schizophrenia. *Journal of the American Academy of Child and Adolescent Psychiatry.* 60:479-489.

14. Woodward ND, Heckers S (2016): Mapping Thalamocortical Functional Connectivity in Chronic and Early Stages of Psychotic Disorders. *Biological psychiatry.* 79:1016-1025.

15. Woodward ND, Karbasforoushan H, Heckers S (2012): Thalamocortical dysconnectivity in schizophrenia. *The American journal of psychiatry.* 169:1092-1099.

16. Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G, et al. (2016): Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences of the United States of America.* 113:12574-12579.

17. Paquola C, Vos De Wael R, Wagstyl K, Bethlehem RAI, Hong SJ, Seidlitz J, et al. (2019): Microstructural and functional gradients are increasingly dissociated in transmodal cortices. *PLoS biology.* 17:e3000284.

18. Hong SJ, Vos de Wael R, Bethlehem RAI, Lariviere S, Paquola C, Valk SL, et al. (2019): Atypical functional connectome hierarchy in autism. *Nature communications.* 10:1022.

19. Vazquez-Rodriguez B, Suarez LE, Markello RD, Shafiei G, Paquola C, Hagmann P, et al. (2019): Gradients of structure-function tethering across neocortex. *Proceedings of the National Academy of Sciences of the United States of America*. 116:21219-21227.

20. Huntenburg JM, Bazin PL, Margulies DS (2018): Large-Scale Gradients in Human Cortical Organization. *Trends in cognitive sciences*. 22:21-31.

21. Valk SL, Xu T, Paquola C, Park B-y, Bethlehem RAI, Vos de Wael R, et al. (2021): Genetic and phylogenetic uncoupling of structure and function in human transmodal cortex. *bioRxiv*.2021.2006.2008.447522.

22. Bayrak S, Khalil AA, Villringer K, Fiebach JB, Villringer A, Margulies DS, et al. (2019): The impact of ischemic stroke on connectivity gradients. *NeuroImage Clinical*. 24:101947.

23. Meng Y, Yang S, Chen H, Li J, Xu Q, Zhang Q, et al. (2021): Systematically disrupted functional gradient of the cortical connectome in generalized epilepsy: Initial discovery and independent sample replication. *NeuroImage*. 230:117831.

24. Dong D, Luo C, Guell X, Wang Y, He H, Duan M, et al. (2020): Compression of Cerebellar Functional Gradients in Schizophrenia. *Schizophrenia bulletin*.

25. Yang S, Meng Y, Li J, Li B, Fan YS, Chen H, et al. (2020): The thalamic functional gradient and its relationship to structural basis and cognitive relevance. *NeuroImage*. 218:116960.

26. Jacobsen LK, Rapoport JL (1998): Research update: childhood-onset schizophrenia: implications of clinical and neurobiological research. *Journal of child psychology and psychiatry, and allied disciplines*. 39:101-113.

27. Coifman RR, Lafon S (2006): Diffusion maps. *Applied and Computational Harmonic*

Analysis. 21:5-30.

28. Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, Shen EH, Ng L, Miller JA, et al. (2012): An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature*. 489:391-399.

29. Woolrich MW, Jbabdi S, Patenaude B, Chappell M, Makni S, Behrens T, et al. (2009): Bayesian analysis of neuroimaging data in FSL. *NeuroImage*. 45:S173-186.

30. Dickie EW, Anticevic A, Smith DE, Coalson TS, Manogaran M, Calarco N, et al. (2019): Ciftify: A framework for surface-based analysis of legacy MR acquisitions. *NeuroImage*. 197:818-826.

31. Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE (2016): Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cerebral cortex*. 26:288-303.

32. Vos de Wael R, Benkarim O, Paquola C, Lariviere S, Royer J, Tavakol S, et al. (2020): BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Communications biology*. 3:103.

33. Park BY, Bethlehem RA, Paquola C, Lariviere S, Rodriguez-Cruces R, Vos de Wael R, et al. (2021): An expanding manifold in transmodal regions characterizes adolescent reconfiguration of structural connectome organization. *eLife*. 10.

34. Zhang D, Snyder AZ, Fox MD, Sansbury MW, Shimony JS, Raichle ME (2008): Intrinsic functional relations between human cerebral cortex and thalamus. *Journal of neurophysiology*. 100:1740-1748.

35. Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. (2011):

The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of neurophysiology.* 106:1125-1165.

36. Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo XN, Holmes AJ, et al. (2018): Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral cortex.* 28:3095-3114.

37. Burt JB, Helmer M, Shinn M, Anticevic A, Murray JD (2020): Generative modeling of brain maps with spatial autocorrelation. *Neurolmage.* 220:117038.

38. Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD (2011): Large-scale automated synthesis of human functional neuroimaging data. *Nature methods.* 8:665-670.

39. Schizophrenia Working Group of the Psychiatric Genomics C (2014): Biological insights from 108 schizophrenia-associated genetic loci. *Nature.* 511:421-427.

40. Arnatkeviciute A, Fulcher BD, Fornito A (2019): A practical guide to linking brain-wide gene expression and neuroimaging data. *Neurolmage.* 189:353-367.

41. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. (2011): Scikit-learn: Machine Learning in Python. *J Mach Learn Res.* 12:2825-2830.

42. Smith SM, Nichols TE (2009): Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neurolmage.* 44:83-98.

43. Ramcharan EJ, Gnadt JW, Sherman SM (2005): Higher-order thalamic relays burst more than first-order relays. *Proceedings of the National Academy of Sciences of the United States of America.* 102:12236-12241.

44. Douaud G, Smith S, Jenkinson M, Behrens T, Johansen-Berg H, Vickers J, et al. (2007):

Anatomically related grey and white matter abnormalities in adolescent-onset schizophrenia. *Brain : a journal of neurology.* 130:2375-2386.

45. Douaud G, Mackay C, Andersson J, James S, Quested D, Ray MK, et al. (2009): Schizophrenia delays and alters maturation of the brain in adolescence. *Brain : a journal of neurology.* 132:2437-2448.

46. White T, Ho BC, Ward J, O'Leary D, Andreasen NC (2006): Neuropsychological performance in first-episode adolescents with schizophrenia: a comparison with first-episode adults and adolescent control subjects. *Biological psychiatry.* 60:463-471.

47. Dickson H, Laurens KR, Cullen AE, Hodgins S (2012): Meta-analyses of cognitive and motor function in youth aged 16 years and younger who subsequently develop schizophrenia. *Psychological medicine.* 42:743-755.

48. Peng Y, Zhang S, Zhou Y, Song Y, Yang G, Hao K, et al. (2021): Abnormal functional connectivity based on nodes of the default mode network in first-episode drug-naive early-onset schizophrenia. *Psychiatry research.* 295:113578.

49. Hilland E, Johannessen C, Jonassen R, Alnaes D, Jorgensen KN, Barth C, et al. (2022): Aberrant default mode connectivity in adolescents with early-onset psychosis: A resting state fMRI study. *NeuroImage Clinical.* 33:102881.

50. Selemon LD, Wang L, Nebel MB, Csernansky JG, Goldman-Rakic PS, Rakic P (2005): Direct and indirect effects of fetal irradiation on cortical gray and white matter volume in the macaque. *Biological psychiatry.* 57:83-90.

51. Dong D, Yao D, Wang Y, Hong S-J, Genon S, Xin F, et al. (2020): Altered Sensorimotor-to-Transmodal Hierarchical Organization in Schizophrenia.

*bioRxiv.*2020.2003.2006.980607.

52. Dong HM, Margulies DS, Zuo XN, Holmes AJ (2021): Shifting gradients of macroscale cortical organization mark the transition from childhood to adolescence. *Proceedings of the National Academy of Sciences of the United States of America.* 118.

53. Jones EG (2002): Thalamic circuitry and thalamocortical synchrony. *Philosophical transactions of the Royal Society of London Series B, Biological sciences.* 357:1659-1673.

54. Halassa MM, Sherman SM (2019): Thalamocortical Circuit Motifs: A General Framework. *Neuron.* 103:762-770.

55. Garcia-Cabezas MA, Zikopoulos B, Barbas H (2019): The Structural Model: a theory linking connections, plasticity, pathology, development and evolution of the cerebral cortex. *Brain structure & function.* 224:985-1008.

56. Andreasen NC, O'Leary DS, Cizadlo T, Arndt S, Rezai K, Ponto LL, et al. (1996): Schizophrenia and cognitive dysmetria: a positron-emission tomography study of dysfunctional prefrontal-thalamic-cerebellar circuitry. *Proceedings of the National Academy of Sciences of the United States of America.* 93:9985-9990.

57. Andreasen NC, Paradiso S, O'Leary DS (1998): "Cognitive dysmetria" as an integrative theory of schizophrenia: a dysfunction in cortical-subcortical-cerebellar circuitry? *Schizophrenia bulletin.* 24:203-218.

58. Andreasen NC (1999): A unitary model of schizophrenia: Bleuler's "fragmented phrene" as schizencephaly. *Archives of general psychiatry.* 56:781-787.