Botrytis cinerea identifies host plants via the recognition of antifungal
capsidiol to induce expression of a specific detoxification gene

Teruhiko Kuroyanagi[1], Abriel Bulasag[1,2], Keita Fukushima[1], Takamasa Suzuki[3], Aiko Tanaka[1],
Maurizio Camagna[1], Ikuo Sato[1], Sotaro Chiba[1], Makoto Ojika[1] and Daigo Takemoto[1*]

[1] Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya, 464-8601,
Japan

[2] College of Arts and Sciences, University of the Philippines Los Baños, College, Laguna,
Philippines 4031

[3] College of Bioscience and Biotechnology, Chubu University, Kasugai, Aichi 478-8501, Japan

*Daigo Takemoto

Email: dtakemo@agr.nagoya-u.ac.jp

Author Contributions: T.K., A.B., K.F., I.S, S.C., M.O., and D.T., designed research; T.K., A.B.,
K.F., T.S., M.O., and D.T. performed research; T.K., K.F., T.S, A.T, M.C., and D.T. analyzed data;
T.S., M.C., M.O., and D.T. contributed new reagents/analytic tools; M.C and D.T. wrote the
paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: capsidiol, detoxification, phytoalexin, polyxenous pathogen, Solanaceae plants.

18
19

Abstract

The gray mold pathogen *Botrytis cinerea* has a broad host range, causing disease in over 400 plant species, but it is not known how this pathogen evolved this polyxenous nature. *B. cinerea* can metabolize a wide range of phytoalexins, including the stilbenoid, resveratrol, and the sesquiterpenoids capsidiol in tobacco, and rishitin in potato and tomato. In this study, we analyzed the metabolism of sesquiterpenoid phytoalexins by *B. cinerea*. Capsidiol was dehydrogenated to capsenone which was then further oxidized, while rishitin was directly oxidized to epoxy- or hydroxy-rishitins indicating that *B. cinerea* has separate mechanisms to detoxify structurally similar sesquiterpenoid phytoalexins. RNAseq analysis revealed that a distinct set of genes were induced in *B. cinerea* when treated with capsidiol or rishitin, suggesting that *B. cinerea* can distinguish structurally similar phytoalexins to activate appropriate detoxification mechanisms. The gene most highly upregulated by capsidiol treatment encoded a dehydrogenase, designated *Bccpdh*. Heterologous expression of *Bccpdh* in a capsidiol-sensitive plant symbiotic fungus, *Epichloë festucae*, resulted in an acquired tolerance of capsidiol and the ability to metabolize capsidiol to capsenone, while *B. cinerea Δbccpdh* mutants became relatively sensitive to capsidiol. The *Δbccpdh* mutant showed reduced virulence on the capsidiol producing *Nicotiana* and *Capsicum* species but remained fully pathogenic on potato and tomato. Homologs of *Bccpdh* are found in taxonomically distant Ascomycota fungi but not in other *Botrytis* species, suggesting that *B. cinerea* acquired the ancestral *Bccpdh* by horizontal gene transfer, thereby extending the pathogenic host range of this polyxenous pathogen to capsidiol-producing plant species.

Significance Statement

B. cinerea can metabolize a wide range of phytoalexins, however, the extent to which phytoalexin detoxification contributes to pathogenicity is largely unknown. In this study, we have shown that *B. cinerea* recognizes structurally resembling sesquiterpenoid phytoalexins, rishitin and capsidiol, to activate appropriate detoxification mechanisms. We identify *Bccpdh*, encoding a dehydrogenase for capsidiol detoxification, which is upregulated in *B. cinerea* exclusively during the infection of capsidiol producing plant species, and is required to exert full virulence. Analysis of the *Bccpdh* locus suggests that the gene was acquired via horizontal gene transfer. This work highlights that the polyxenous plant pathogen *B. cinerea* can distinguish its host plants by its anti-microbial compounds, to activate appropriate mechanisms for enhanced virulence.

Main Text

Introduction

  The antimicrobial secondary metabolites produced in plants during the induction of disease resistance are collectively termed phytoalexins (1, 2). Several hundred phytoalexins of diverse structures have been identified from a wide range of plant species, which include terpenoids, flavonoids and indoles (3, 4). Many of these phytoalexins are considered to exhibit their antimicrobial activity by targeting the cell wall or cell membrane of pathogens (5, 6), but their mechanisms of action remain largely unknown.
  In plants belonging to the Solanaceae family, the major phytoalexins are sesquiterpenoids, such as capsidiol in *Nicotiana* and *Capsicum* species and rishitin in *Solanum* species (2, 7, 8). In *Nicotiana* sp., production of capsidiol is strictly controlled by regulating the gene expression for capsidiol biosynthesis, such as *EAS* (5-*epi*-aristolochene synthase) and *EAH* (5-*epi*-aristolochene dihydroxylase), encoding the enzymes dedicated to the production of capsidiol (9, 10). In *N. benthamiana*, expression of *EAS* and *EAH* genes is hardly detected in healthy tissues, but their expression is rapidly induced during infection by pathogens or the treatment with elicitor protein (11, 12). Produced phytoalexins are secreted locally via ABC (ATP-binding cassette) transporters

at the site of pathogen attack (13, 14). For plants, such temporal and spatial control of phytoalexins
is probably critical as the toxicity of phytoalexins is often not specific to microorganisms but is also
harmful to plant cells (13, 15, 16). For example, rishitin affects the permeability of plant liposomal
membranes and disrupts chloroplasts (17). Therefore, timely production and efficient transport of
phytoalexins to the site of pathogen attack are important for plants to apply these double-edged
weapons effectively.
It has been reported that various phytopathogenic fungi can metabolize and detoxify phytoalexins
(18). Although many studies have shown an approximate correlation between the ability of strains
to detoxify phytoalexins and their virulence (19–21), the importance of phytoalexin detoxification
for pathogen virulence is largely unproven for most plant-pathogen interactions. The best-studied
phytoalexin metabolizing enzyme is pisatin demethylase (PDA) of *Nectria haematococca*. Deletion
of the *PDA* gene resulted in reduced virulence of *N. haematococca* on pea, directly proving the
importance of PDA for the virulence of this pathogen (22). It has also been reported that
sesquiterpenoid phytoalexins are metabolized by pathogenic fungi. Capsidiol is metabolized to less
toxic capsenone via dehydration by pathogens such as the gray mold pathogen *Botrytis cinerea*
and *Fusarium oxysporum* (23). *Gibberella pulicaris*, the dry rot pathogen of potato tubers, oxidizes
rishitin to 13-hydroxy rishitin and 11,12-epoxyrishitin (24). However, the enzymes involved in the
detoxification of sesquiterpenoid phytoalexins have not been isolated to date, and their importance
for pathogenicity has not been demonstrated.
In this study, we first investigated some pathogens isolated from Solanaceae and non-Solanaceae
plants on their tolerance of capsidiol and rishitin, as well as their ability to detoxify/metabolize these
phytoalexins. Among the pathogens that can metabolize both capsidiol and rishitin, we chose
Botrytis cinerea for further analysis to investigate the importance of its ability to detoxify capsidiol
for the pathogenicity on plant species that produce capsidiol.
94
95
96   **Results**
97
Metabolization of Sesquiterpenoid Phytoalexins Capsidiol and Rishitin by Fungal Plant
Pathogens.
Four oomycetes and 12 fungal species were evaluated for their ability to detoxify/metabolize
sesquiterpenoid phytoalexins capsidiol and rishitin, produced by Solanaceae plant species. Four
Phytophthora species isolated from Solanaceae host plants, including potato late blight pathogen
P. infestans, *P. nicotiana* isolated from tobacco, *P. capsici* isolated from green pepper and *P.*
cryptogea isolated from nipplefruit (*Solanum mammosum*), are all sensitive to capsidiol and rishitin.
The amount of capsidiol and rishitin after the incubation with these oomycete pathogens did not
decrease, indicating that they are neither capable of metabolizing nor tolerant to capsidiol and
rishitin (Fig. 1 and Supplemental Fig. S1). Among 12 fungal plant pathogens (eight isolated from
Solanaceae plants), seven and eight fungal strains can metabolize capsidiol and rishitin,
respectively, and showed increased resistance to phytoalexins. In some cases, pathogens can
metabolize phytoalexins that are not produced by their host, indicating that the ability to detoxify
phytoalexins does not always correlate with their host range (Supplemental Figs. S1-3 and
Supplementary Notes 1 and 2). *Botrytis cinerea* and *Sclerotinia sclerotiorum* are well-known as
pathogens with a wide host range and are capable of metabolizing both capsidiol and rishitin. In
this study, *B. cinerea* was selected as the pathogen to further investigate the role of detoxification
of phytoalexins on the pathogenicity of this polyxenous pathogen.
116
Metabolization of Capsidiol and Rishitin by *B. cinerea*
Metabolization of capsidiol and rishitin by *B. cinerea* were evaluated by LC/MS. Under the
experimental conditions of this study, capsidiol was metabolized to capsenone by dehydrogenation
within 12 h, and oxidized capsenone was detected at 24 h, whereas oxidized capsidiol was not
detected (Fig. 1C and Supplemental Figs. S4). In contrast, rishitin was directly oxidized within 6 h
and at least four different forms of oxidized rishitin were detected within 24 h (Fig. 1C and

Supplemental Figs. S5), indicating that despite the structural similarity of these phytoalexins, *B.*
cinerea detoxifies/metabolizes capsidiol and rishitin by different mechanisms (Fig. 1C).
125
Unique Sets of Genes Are Upregulated in *B. cinerea* Treated with Capsidiol, Rishitin or
Resveratrol.
To identify *B. cinerea* genes upregulated during the detoxification of sesquiterpenoid phytoalexins,
RNAseq analysis was performed for mycelia of *B. cinerea* cultured in minimal media supplemented
with capsidiol or rishitin. A stilbenoid phytoalexin, resveratrol, produced in grape was also used for
comparison. Mycelia of *B. cinerea* were grown in liquid CM media supplemented with either 500
µM rishitin, 500 µM resveratrol, or 100 µM capsidiol, as 500 µM capsidiol caused significant growth
defects in *B. cinerea* (Fig. 1A, Supplemental Fig. S6). The mycelial tissue was then used to perform
an RNAseq analysis. Among 11,707 predicted *B. cinerea* genes (25), 25, 27 or 23 genes were
significantly upregulated (Log2 fold change >2, p <0.05) by the treatment with capsidiol, rishitin or
resveratrol, respectively. Unexpectedly, distinctive sets of genes were upregulated even between
B. cinerea treated with capsidiol and rishitin which resemble each other structurally (Fig. 2A,
Supplemental Table S1-3), indicating that *B. cinerea* can either distinguish the structural difference
of capsidiol and rishitin or the damage caused by these sesquiterpenoid phytoalexins. For instance,
two genes Bcin08g00930 and Bcin12g01750 encoding hypothetical proteins containing a motif for
dehydrogenases were specifically induced by capsidiol, while the treatment of rishitin significantly
induced Bcin07g05430, encoding a cytochrome P450 gene (Fig. 2B). Expression of *BcatrB*
encoding an ABC transporter involved in the tolerance of *B. cinerea* against structurally unrelated
phytoalexins resveratrol, camalexin and fungicides fenpiclonil and fludioxonil (26, 27), is
upregulated by treatment with rishitin and resveratrol, but not by capsidiol. In contrast,
Bcin15g00040, encoding a major facilitator superfamily (MFS)-type transporter was upregulated
specifically by capsidiol (Fig. 2B). Interestingly, treatment of *B. cinerea* with phytoalexins also
activated genes predicted to be involved in pathogenicity to plants. For example, capsidiol
treatment induced expression of a hydrophobin gene Bcin06g00510.1 (Bhp3), while rishitin
treatment activated the expression of Bcin01g00080.1 (Bcboa8), encoding an enzyme for
biosynthesis of a phytotoxin, botcinic acid (Supplemental Tables S1 and S2). Given that capsidiol
is metabolized to capsenone by a dehydrogenation reaction in *B. cinerea* (Fig. 1C, Supplemental
Fig. S4), the function of Bcin08g00930 and Bcin12g01750, both encoding a predicted short-chain
dehydrogenase/reductase (SDR) were further analyzed in this study.
155
Bcin08g00930 Encodes a Short-Chain Dehydrogenase for the Detoxification of Capsidiol.
To investigate the function of SDR genes induced by capsidiol treatment, an endophytic fungus
Epichloë festucae was employed for the heterologous expression of these genes. Bcin08g00930
and Bcin12g01750 were expressed in *E. festucae* under the control of the constitutive *TEF*
promoter (Vanden Wymelenberg et al., 1997). The growth of wild type and control *E. festucae*
expressing DsRed was severely inhibited in the 100 µM capsidiol (Fig. 3A and B). *E. festucae*
transformants expressing Bcin12g01750 also hardly grew in 100 µM capsidiol, however, *E.*
festucae became tolerant to capsidiol by the expression of Bcin08g00930 (Fig. 3A and B). The
amount of capsidiol was not altered by wild type, DsRed or Bcin12g01750 expressing *E. festucae*,
while capsidiol was metabolized to capsenone by the *E. festucae* transformant expressing
Bcin08g00930 (Fig. 3C). These results indicated that Bcin08g00930 encodes a dehydrogenase
that can detoxify capsidiol, thus the SDR encoded by Bcin08g00930 was designated as BcCPDH
standing for *B. cinerea* capsidiol dehydrogenase.
169
Expression of *Bccpdh* Is Specifically Induced by Capsidiol.
To further investigate the expression pattern of *Bccpdh*, *B. cinerea* was transformed with a
reporter construct for GFP expression under the control of the 1 kb proximal *Bccpdh* promoter
(P_*Bccpdh:GFP*). P_*Bccpdh:GFP* transformants showed no obvious expression of GFP in water,
but a significant increase of GFP fluorescence was detected in 500 µM capsidiol (Fig. 4A). The
P_*Bccpdh:GFP* transformant was incubated with other anti-microbial terpenoids produced by
Solanaceae species, including rishitin, debneyol (29), sclareol (30) and capsidiol 3-acetate (31).

Among the terpenoids tested, treatment of capsidiol and capsidiol 3-acetate induced expression of
GFP in the P_*Bccpdh:GFP* transformant. This result further indicated that *B. cinerea* specifically
reacts to capsidiol and its derivative capsidiol 3-acetate. As the *Bccpdh* promoter was activated by
capsidiol 3-acetate, we investigated whether BcCPDH could metabolize capsidiol 3-acetate to
capsenone 3-acetate. However, capsidiol 3-acetate was not metabolized by *E. festucae* expressing
Bccpdh (Bcin08g00930) (Supplemental Fig. S7A). Consistently, capsidiol 3-acetate was directly
oxidized by *B. cinerea* and at least two different forms of oxidized capsidiol 3-acetate, but not
capsenone 3-acetate, were detected (Supplemental Fig. S7B), indicating that capsidiol and
capsidiol 3-acetate were metabolized in *B. cinerea* by distinctive pathways.
Analysis of different lengths of *Bccpdh* promoters indicated that 250 bp of promoter sequence are
sufficient for specific activation of the promoter by capsidiol treatment, and the cis-element required
for capsidiol-specific expression was shown to be within -250 and -200 bp upstream of the start
codon (Supplemental Figs. S8 and 9). By further analysis using *B. cinerea* transformant
P_*Bccpdh:Luc* using luciferase as a quantitative marker, it was shown that *Bccpdh* promoter was
activated within 2 h of capsidiol treatment, and the expression activity decreased as capsidiol was
metabolized (Supplemental Figs. S10, Supplementary Notes 3).
193
Expression of Bccpdh Is Specifically Induced During the Infection of Capsidiol Producing
Plant Species.
Leaves of *Nicotiana benthamiana*, which produce capsidiol as the major phytoalexin (13), were
inoculated with conidia of the *B. cinerea* transformant P_*Bccpdh:GFP*. No GFP signal was detected
in germinating *B. cinerea* conidia grown on the surface of *N. benthamiana* leaves 2 h after the
inoculation, but obvious GFP expression was detected in the appressoria at 8 h after the inoculation
(Fig. 4B). Growing hyphae in the leaf tissue of *N. benthamiana* showed GFP fluorescence, and the
intensity of GFP fluorescence in hyphae was stronger near the edge of the lesion compared with
that of hyphae growing in the areas of dead tissue within the lesions (Fig. 4B), probably because
capsidiol was eventually detoxified in these areas, which are heavily infected with *B. cinerea*.
To examine the possibility whether polyxenous *B. cinerea* activates *Bccpdh* for detoxification of
other phytoalexin or anti-microbial compounds produced in other plant species, the *B. cinerea*
P_*Bccpdh:GFP* transformant was inoculated on a wide variety of plants. Over 50 plant species
were used for the inoculation test including 9 Solanaceae, 6 Brassicaceae, 6 Rosaceae, 5
Fabaceae and 6 Asteraceae plants, and development of disease symptoms was observed in all
tested plant species. Among the tested plants, expression of GFP under the control of *Bccpdh*
promoter was only detected during the infection in three *Nicotiana* and two *Capsicum* species (Fig.
4C, Supplemental Fig. S11 and Table S4), all of which are reported to produce capsidiol (8, 32–
34). These results further indicated that *B. cinerea* specifically recognizes capsidiol for the induction
of *Bccpdh*.
214
BcCPDH Metabolizes and Detoxifies Capsidiol to Capsenone in B. cinerea.
B. cinerea Bccpdh KO mutant (*Δbccpdh*) was produced to investigate the function of BcCPDH
(Supplemental Fig. S12). Mycelia of *B. cinerea* wild type and *Δbccpdh* were incubated with capsidiol
and the metabolites were detected by LC/MS. While capsidiol was metabolized to capsenone in
the wild type strain, most of the capsidiol remained unmetabolized two days after incubation with
the *Δbccpdh* strain (Fig. 5A). Instead, oxidized capsidiol, which was not detectable in the wild type
strain, was detected in *Δbccpdh* incubations (Supplemental Fig. S13 and Supplementary Notes 4).
While the growth of *B. cinerea* hyphae was not affected by the disruption of the *Bccpdh* gene in
100 μM capsidiol, growth of *Δbccpdh* was diminished compared with that of wild type *B. cinerea* at
higher concentrations of capsidiol (Fig. 5B). These results confirmed that BcCPDH is the enzyme
responsible for the detoxification of capsidiol in *B. cinerea*.
226
Epoxidation of Capsidiol and Capsenone is Mediated via Bcin16g01490
The incubation of the *Δbccpdh* knockout strain in capsidiol revealed the presence of a potential
secondary capsidiol degradation pathway, which resulted in the accumulation of oxidized capsidiol.
Since we were unable to find any indication for such a pathway in our RNAseq results for 100 μM

capsidiol treated incubations, we extended our search to a preliminary RNAseq analysis which
used 500 µM capsidiol treatments. We identified the gene Bcin16g01490 encoding a cytochrome
P450, which was significantly upregulated under these elevated capsidiol concentrations
(Supplemental Table S1 and Fig. S14A). Heterologous expression of Bcin16g01490 in *E. festucae*,
resulted in the conversion of capsidiol to oxidized capsidiol as detected during *Δbccpdh* strain
incubations (Supplemental Figs. S13 and 14, Supplementary Notes 4). Moreover, sequential
incubation of capsidiol with the *Bccpdh* expressing *E. festucae* transformant followed by a
Bcin16g01490 expressing transformant reproduced the reactions detected in *B. cinerea*
(Supplemental Figs. S4 and S14B). Structural analysis of oxidized capsenone indicated that the
end product of these reactions is capsenone 11,12-epoxide (Fig. 1 and Supplemental Fig. S15-17,
Supplementary Notes 5).
242
BcCPDH Is Required for the Pathogenicity of *B. cinerea* on Plant Species Producing
Capsidiol.
Pathogenicity of *B. cinerea Δbccpdh* transformants was tested for several plant species. On plant
species that produce capsidiol, such as *N. benthamiana*, *N. tabacum* and *C. annuum*, the
development of disease symptoms by *Δbccpdh* was significantly reduced compared with those
caused by wild type *B. cinerea*. In contrast, both wild type and *Δbccpdh* strains caused comparable
symptoms in potato, tomato, grape and eggplant, which is consistent with the finding that
expression of *Bccpdh* is not induced during the infection in these plant species (Fig. 5C,
Supplemental Fig. S18). In the complementation strain, the ability to metabolize capsidiol to
capsenone was restored, as was virulence against capsidiol producing plants (Supplemental Fig.
S19). These results indicated that BcCPDH is a dedicated enzyme for the virulence of *B. cinerea*
on capsidiol-producing plant species.
255
Distribution of *Bccpdh* Homologues in the Fungal Kingdom
The distribution of *Bccpdh* homologs in taxonomically related fungal species (Ascomycota,
Leotiomycetes) was investigated. A search for *Bccpdh* homologs in the genome sequences of 6
host-specialized phytopathogenic *Botrytis* species, such as *B. tulipae* (pathogen of tulip), *B.*
hyacinthi (hyacinth) and *B. paeoniae* (peony), indicated that among *Botrytis* species, *Bccpdh* is a
unique gene only found in *B. cinerea* (Supplemental Fig. S20). Consistently, 4 *Botrytis* species, not
including *B. cinerea*, were incubated with capsidiol, but no CPDH activity was detected for the
tested strains (Supplemental Fig. S21). Search in 11 draft genomes of Leotiomycete fungi also
indicated the absence of *Bccpdh* homologs in these species. Comparison of the corresponding
genome regions between *Botrytis* species and *S. sclerotiorum* (another polyxenous pathogen
belonging to Leotiomycetes) revealed that the approx. 4.9 kb sequence surrounding *Bccpdh* is
unique to *B. cinerea* (Fig. 6). This unique sequence shows a lower GC content compared to the
surrounding region (Supplementa Fig. S22), suggesting that *Bccpdh* might have been obtained via
horizontal gene transfer (35). A blast search using BcCPDH as query sequence revealed that
probable orthologs can be found only in some Pezizomycotina fungi belonging to Ascomycota
(Supplemental Fig. S20). Orthologs were found from a taxonomically diverse range of fungal
species, including animal and insect pathogens, and there was no correlation between their
homology and taxonomic relationship, which might indicate multiple horizontal gene transfer events
of the *cpdh* gene in the diversification of Ascomycota fungi (Supplemental Figs. 20, 23-25,
Supplementary Notes 6). Among plant pathogenic fungi, *Bccpdh* homologs were found in *Fusarium*
species, all of which are pathogens that infect plants that do not produce capsidiol. This result is
consistent with a previous report showing that some *Fusarium* species can metabolize capsidiol to
capsenone (36).
279
CPDH Activity Is Conserved Among *Botrytis cinerea* Strains
Although *B. cinerea* is a polyxenous fungus, it has been reported that different strains of *B. cinerea*
exhibit various degrees of virulence on different host plants (37, 38). Hence, we investigated
whether BcCPDH is conserved among strains isolated from diverse plant species. Twenty-four *B.*
cinerea strains isolated from 14 different plant species were incubated with capsidiol and resultant

capsenone was detected by GC/MS. All tested *B. cinerea* strains can metabolize capsidiol to
capsenone, indicating that BcCPDH is highly conserved among *B. cinerea* strains even though
their (most recent) host was not a producer of capsidiol (Fig. 7A, Supplemental Table S5). Similarly,
17 strains of *F. oxysporum* isolated from 7 plant species were subjected to the analysis for CPDH
activity. Eight out of 17 *F. oxysporum* strains showed CPDH activity, and the activity was not related
to the natural host of the strains (Fig. 7B, Supplemental Table S6). This result is consistent with the
finding that *Bccpdh* orthologues can be found in 5 out of 14 available genomes of *F. oxysporum*
(Supplemental Fig. S25, data not shown). These results indicate that the cpdh gene is randomly
distributed in *F. oxysporum* strains, whereas it is highly conserved among *B. cinerea* strains.
294
295   **Discussion**
296
To survive the exposure to microorganisms in the environment, different plant species have
developed diverse resistance mechanisms over the course of evolution. The structural variety of
phytoalexins is a prime example of such diversity: different plant families produce phytoalexins with
relatively similar basic structures, but their side-chain structures often differ from species to species
(4). These differences in structure between species may have contributed to the determination of
host specificity between plants and pathogens, as in some cases a pathogen may have acquired
resistance to a particular phytoalexin, but an analogous substance is not overcome by the same
resistance mechanism (39). However, plant pathogens with a broad host range such as *B. cinerea*
must employ strategies to counter a multitude of diverse phytoalexins. The prompt killing of plant
cells, or the presence of efflux pumps of extremely low specificity may present effective strategies
for such pathogens. Indeed, *B. cinerea* produces host-nonspecific phytotoxins (such as botrydial
and botcinic acid, 40, 41) upon infection, and activates transporters capable of effluxing diverse
substances such as PDR-type ABC transporter BcatrB and MFS transporter Bcmfs1 (26, 27, 42).
More recently, it has been shown that *B. cinerea* suppresses the immune response of different
plant species via gene silencing by delivering various small RNAs into host cells (43, 44).
In addition to effective infection mechanisms which generally facilitate infection of a wide range of
plants, this study revealed that *B. cinerea* responds to chemical cues from the host plant to adapt
its infection strategy to overcome specific plant resistance mechanisms. *B. cinerea* strictly
distinguishes structurally similar phytoalexins, such as capsidiol and rishitin, and activates
appropriate detoxification responses. Interestingly, treatment of *B. cinerea* with phytoalexins
activated not only genes involved in detoxification and efflux of toxic compounds, but also genes
predicted to be involved in pathogenicity to plants. These results support the notion that *B. cinerea*
utilizes phytoalexins as a means to identify a given host and adjust the method of infection.
Expression of the *Bccpdh* gene is activated specifically during infection of capsidiol-producing
plants, and the pathogenicity of the *bccpdh* mutant strains is compromised on plants producing
capsidiol, but does not suffer any disadvantages on plants that do not produce capsidiol. These
results suggest that BcCPDH is a critical component that specifically enables *B. cinerea* to infect
capsidiol producing plants. Despite capsidiol producing plants representing only a small fraction of
the over 400 host plants of *B. cinerea*, CPDH activity was maintained in all investigated *B. cinerea*
strains isolated from plants that do not produce capsidiol. This may hint at the presence of a
selection pressure against the loss of CPDH, despite it only affecting a limited number of host
plants. This poses the question of whether *B. cinerea* is able to maintain acquired resistances for
a prolonged time, which may explain how it evolved and establish itself as the polyxenous pathogen
that it is.
331
How does *B. cinerea* Recognize Phytoalexins with Similar Structures?
Although *B. cinerea* is known to have the ability to metabolize a wide variety of phytoalexins (18),
only a small portion of these detoxification mechanisms are required for the infection of any
particular plant. Thus, *B. cinerea* needs to strictly control those various detoxification mechanisms,
since maintaining sufficient levels of all these detoxification mechanisms represents a considerable
waste of resources. Elucidating the mechanism by which *B. cinerea* recognizes phytoalexins and
activates a specific set of genes is one of the important subjects for further research. Two major

scenarios can be envisioned: first, *B. cinerea* may possess receptors that can identify the chemical
structures of phytoalexins. In plants, various terpenes with diverse structures are employed as
hormones recognized by specific receptors, such as gibberellins, cytokinins, and abscisic acid (39).
However, the possibility that *B. cinerea* maintains receptors to perceive all of the myriad of
phytoalexins may not be realistic. The second scenario would be that *B. cinerea* recognizes the
damage caused by phytoalexins. However, as for capsidiol and rishitin, both are presumed to have
cell membranes as their targets, and these phytoalexins are relatively unspecific toxicants that
cause damage even to plant cells (5, 13, 17, 45), so it is not certain whether there exists a specific
target by which these two compounds can be distinguished. Given that some genes, such as
BcatrB encoding a multidrug resistance pump, are commonly induced by structurally unrelated
phytoalexins and fungicides treatments (26, 27), *B. cinerea* may possess both induction
mechanisms. Using the reporter system developed in this study, mutant strains defective in
capsidiol response could be isolated to elucidate the mechanism by which *B. cinerea* distinctly
identifies phytoalexins.
353
Does *B. cinerea* Possess Other Capsidiol Resistance Mechanism besides Detoxification by
BcCPDH?
Although the *bccpdh* knockout mutants showed reduced virulence on plant species producing
capsidiol, the mutant can develop the disease symptom on these plants and showed tolerance to
100 µM capsidiol, same as the wild type. In contrast, *Phytophthora* spp., *Alternaria* spp. and *E.*
festucae are sensitive to the same concentration of capsidiol. Although we also isolated
Bcin16g01490, which we found to oxidize capsidiol, its activity is fairly limited, indicating that it is
unlikely to confer sufficient protection against capsidiol. We therefore presumed that *B. cinerea* has
another mechanism for capsidiol tolerance other than detoxification. RNAseq analysis of *B. cinerea*
genes upregulated by capsidiol treatment identified 2 genes (Bcin15g00040.1 and
Bcin14g02870.1/Bcmfs1) encoding MFS transporters and a gene (Bcin01g05890.1/Bcbmr1)
encoding an ABC transporter (Supplemental Table S1). Bcmfs1 has been reported to be involved
in the resistance of *B. cinerea* to structurally different natural toxicants (camptothecin produced by
the plant *Camptotheca acuminata* and cercosporin produced by the plant pathogenic fungus
Cercospora kikuchii) and fungicides (sterol demethylation inhibitors, DMIs) (42). The *bcbmr1*
mutants showed an increased sensitivity to fungicides, polyoxin (a chitin synthetase inhibitor) and
iprobenfos (a choline biosynthesis inhibitor) (46). The expression of one or more of these
transporters may be involved in capsidiol efflux from B. cinerea cells, and regulated by signaling
that is common to *Bccpdh*.
373
Why and How Can *B. cinerea* Strains Stably Maintain BcCPDH?
CPDH activity was detected in all tested *B. cinerea* strains isolated from plants that do not produce
capsidiol, and the *Bccpdh* gene is conserved in the genomes of published *B. cinerea* strains. In *F.*
oxysporum, in contrast, there were strains with and without CPDH activity regardless of the host
plant, and consistently, some publicly available *F. oxysporum* genomes contain *cpdh* homologs
and others do not. As shown in this study, metabolic capacity (and tolerance) to capsidiol and
rishitin can be detected in some fungal strains regardless of the natural host of these pathogens,
and perhaps unused detoxification activities are maintained in the population of phytopathogenic
filamentous fungi for future use.
It is theoretically implausible that all *B. cinerea* strains maintain an enzyme required only upon
infection of capsidiol-producing plants, even if said gene is completely repressed in its expression
under normal circumstances. Given that *B. cinerea* has an extremely wide host range, some strains
will go for long periods where capsidiol detoxification ability is not a selection pressure. One
possibility is that BcCPDH has functions other than capsidiol degradation. For example, BcCPDH
might be involved in the detoxification of antimicrobial substances produced by insects, because it
has been reported that spores of *B. cinerea* are transmitted between plants via insects such as
thrips (47). Since homologous genes for *Bccpdh* are found in several insect-infecting fungi, it is
expected that CPDH-like enzymes in these species can metabolize insect-derived substances. We,
therefore, performed preliminary experiments to determine whether *Bccpdh* expression can be

induced by inoculation of *B. cinerea* P_*Bccpdh:GFP* transformant with several insects, but to date, no induction of GFP expression has been detected (data not shown). Alternatively, *B. cinerea* may have acquired the trait of having many hosts because of its ability to maintain rarely used virulence factors. Future clarification of the phytoalexin recognition mechanisms and comparative analysis of the genome with that of closely related *Botrytis* species are anticipated to elucidate unknown features of *B. cinerea* that have led to its evolution as a polyxenous fungi.

Materials and Methods

Comprehensive descriptions of the materials and methods used in this study, including, biological materials, vector construction, transformation of *E. festucae* and *B. cinerea*, confocal microscopy, pathogenicity tests and detection and structural analysis of phytoalexins and their derivatives, are available in Supplemental data.

Data Availability

RNA-seq data reported in this work are available in GenBank under the accession number DRA013980.

Acknowledgments

We thank Prof. Barry Scott (Massey University, New Zealand) for providing *E. festucae* strain Fl1 and critical reading of the manuscript. We also thank Dr. David Jones (The Australian National University, Australia) for *N. benthamiana* seeds, Ms. Kayo Shirai (Hokkaido Central Agricultural Experiment Station, Japan), and Dr. Seishi Akino (Hokkaido University, Japan) for providing *P. infestans* isolate 08YD1, Prof. Takashi Tsuge (Chubu University, Japan) for providing *Fusarium* strains, Dr. Haruhisa Suga for providing *Fusarium graminearum* strain, Mr. Masashi Matsusaki (Aichi Prefectural Agricultural Research Center, Japan) for providing *Fusarium oxysporum* f. sp. *lycopersici* strain, and Mr. Taku Kawakami (Mie Prefecture Agricultural Research Institute, Japan) for providing *B. cinerea* strains. We are also grateful to Prof. Kazuhito Kawakita for valuable suggestions, and Dr. Kenji Asano and Mr. Seiji Tamiya (National Agricultural Research Center for Hokkaido Region, Japan) and Mr. Yasuki Tahara (Nagoya University, Japan) for providing tubers of potato cultivars.

References

1. K. O. Müller, H. Börger, Experimentelle Untersuchungen über die Phytophthora: Resistenz der Kartoffel. *Arb. Biol. Anst. Reichsanst.* **23,** 189–231 (1940).
2. J. Kuc, Phytoalexins, stress metabolism, and disease resistance in plants. *Annu. Rev. Phytopathol.* **33,** 275–297 (1995).
3. C. J. Brooks, D. G. Watson, Phytoalexins. *Nat. Prod. Rep.* **2,** 427–459 (1985).
4. R. Hammerschmidt, Phytoalexins: What have we learned after 60 years? *Annu. Rev. Phytopathol.* **37,** 285–306 (1999).
5. M. Turelli, C. Coulomb, P. J. Coulomb, J. P. Roggero, M. Bounias, Effects of capsidiol on the lipid and protein content of isolated membranes of *Phytophthora capsici*. *Physiol. Plant Pathol.* **24,** 211–221 (1984).
6. E. E. Rogers, J. Glazebrook, F. M. Ausubel, Mode of action of the *Arabidopsis thaliana* phytoalexin camalexin and its role in *Arabidopsis*-pathogen interactions. *Mol. Plant-Microbe Interact.* **9,** 748–757 (1996).
7. N. Katsui, *et al.,* The structure of rishitin, a new antifungal compound from diseased potato tubers. *Chem. Commun.* **1,** 43–44 (1968).

8.   J. A. Bailey, R. S. Burden, G. G. Vincent, Capsidiol: An antifungal compound produced in
Nicotiana tabacum and *Nicotiana clevelandii* following infection with tobacco necrosis virus.
Phytochemistry **14,** 597 (1975).
9.   U. Vögeli, J. Chappell, Induction of sesquiterpene cyclase and suppression of squalene
synthetase activities in plant cell cultures treated with fungal elicitor. *Plant Physiol.* **88,** 1291–1296
(1988).
10.   Y. Shibata, K. Kawakita, D. Takemoto, Age-related resistance of *Nicotiana benthamiana*
against hemibiotrophic pathogen *Phytophthora infestans* requires both ethylene- and salicylic acid-
mediated signaling pathways. *Mol. Plant-Microbe Interact.* **23,** 1130–1142 (2010).
11.   S. Rin, *et al.,* Expression profiles of genes for enzymes involved in capsidiol production in
Nicotiana benthamiana. *J. Gen. Plant Pathol.* **86,** 340–349 (2020).
12.   S. Imano, *et al.,* AP2/ERF transcription factor NbERF-IX-33 is involved in the regulation of
phytoalexin production for the resistance of *Nicotiana benthamiana* to *Phytophthora infestans*.
Front. Plant Sci. **12,** 821574 (2021).
13.   Y. Shibata, *et al.,* The full-size ABCG transporters Nb-ABCG1 and Nb-ABCG2 function in pre-
and postinvasion defense against *Phytophthora infestans* in *Nicotiana benthamiana*. *Plant Cell* **28,**
1163–1181 (2016).
14.   Y. He, et al., The Arabidopsis pleiotropic drug resistance transporters PEN3 and PDR12
mediate camalexin secretion for resistance to *Botrytis cinerea*. *Plant Cell* **31,** 2206–2222 (2019).
15.   T. Shiraishi, H. Oku, M. Isono, S. Ouchi, The injurious effect of pisatin on the plasma
membrane of pea. *Plant Cell Physiol.* **16,** 939–942 (1975).
16.   DA. Smith, Toxicity of phytoalexins. In Phytoalexins, J.A. Bailey and J.W. Mansfield, eds.
(Glasgow/London: Blackie.), 218–252 (1982).
17.   G. D. Lyon, Evidence that the toxic effect of rishitin may be due to membrane damage. *J. Exp.*
Bot. **31,** 957–966 (1980).
18.   M. S. C. Pedras, P. W. K. Ahiahonu, Metabolism and detoxification of phytoalexins and
analogs by phytopathogenic fungi. *Phytochemistry* **66,** 391–411 (2005).
19.   M. Sbaghi, P. Jeandet, R. Bessis, P. Leroux, Degradation of stilbene-type phytoalexins in
relation to the pathogenicity of *Botrytis cinerea* to grapevines. *Plant Pathol.* **45,** 139–144 (1996).
20.   P. Suleman, A. M. Tohamy, A. A. Saleh, M. A. Madkour, D. C. Straney, Variation in sensitivity
to tomatine and rishitin among isolates of *Fusarium oxysporum* f. sp. *lycopersici*, and strains not
pathogenic on tomato. *Physiol. Mol. Plant Pathol.* **48,** 131–144 (1996).
21.   K. M. Weltring, K. Loser, J. Weimer, Genetic instability of rishitin metabolism and tolerance
and virulence on potato tubers of a strain of *Gibberella pulicaris*. *J. Phytopathol.* **146,** 393–398
(1998).
22.   C. C. Wasmann, H. D. VanEtten, Transformation-mediated chromosome loss and disruption
of a gene for pisatin demethylase decrease the virulence of *Nectria haematococca* on pea. *Mol.*
Plant-Microbe Interact. **9,** 793–803 (1996).
23.   E. W. B. Ward, A. Stoessl, Postinfectional inhibitors from plants. III. Detoxification of capsidiol,
an antifungal compound from peppers. *Phytopathology* **62,** 1186 (1972).
24.   H. W. Gardner, A. E. Desjardins, S. P. McCormick, D. Weisleder, Detoxification of the potato
phytoalexin rishitin by *Gibberella pulicaris. Phytochemistry* **37,** 1001–1005 (1994).
25.   J. A. L. van Kan, et al., A gapless genome sequence of the fungus *Botrytis cinerea*. *Mol. Plant*
Pathol. **18,** 75–89 (2017).
26.   H. Schoonbeek, G. del Sorbo, M. A. de Waard, The ABC transporter BcatrB affects the
sensitivity of *Botrytis cinerea* to the phytoalexin resveratrol and the fungicide fenpiclonil. *Mol. Plant-*
Microbe Interact. **14,** 562–571 (2001).
27.   T. Vermeulen, H. Schoonbeek, M. A. de Waard, The ABC transporter BcatrB from *Botrytis*
cinerea is a determinant of the activity of the phenylpyrrole fungicide fludioxonil. *Pest Manag. Sci.*
57, 393–402 (2001).
28.   A. J. vanden Wymelenberg, D. Cullen, R. N. Spear, B. Schoenike, J. H. Andrews, Expression
of green fluorescent protein in *Aureobasidium pullulans* and quantification of the fungus on leaf
surfaces. *Biotechniques* **23,** 686–690 (1997).

29. R. S. Burden, et al., Debneyol, a fungicidal sesquiterpene from TNV infected *Nicotiana*
debneyi. *Phytochemistry* **24,** 2191–2194 (1985).
30. Z. Guo, G. J. Wagner, Biosynthesis of labdenediol and sclareol in cell-free extracts from
trichomes of *Nicotiana glutinosa*. *Planta* **197,** 627–632 (1995).
31. R. Uegaki, S. Kubo, T. Fujimori, Stress compounds in the leaves of *Nicotiana undulata* induced
by TMV inoculation. *Phytochemistry* **27,** 365–368 (1988).
32. P.-M. Molot, P. Mas, M. Conus, H. Ferrière, P. Ricci, Relations between capsidiol
concentration, speed of fungal invasion and level of induced resistance in cultivars of pepper
(*Capsicum annuum*) susceptible or resistant to *Phytophthora capsici*. *Physiol. Plant Pathol.* **18,**
379–389 (1981).
33. J. Bohlmann, et al., Gene expression of 5-*epi*-aristolochene synthase and formation of
capsidiol in roots of *Nicotiana attenuata* and *N. sylvestris*. *Phytochemistry* **60,** 109–116 (2002).
34. M. Matsukawa, et al., *Nicotiana benthamiana* calreticulin 3a is required for the ethylene-
mediated production of phytoalexins and disease resistance against oomycete pathogen
Phytophthora infestans. Mol. Plant-Microbe Interact. **26,** 880–892 (2013).
35. M. Ravenhall, N. Škunca, F. Lassalle, C. Dessimoz, Inferring horizontal gene transfer. *PLoS*
Comput. Biol. **11,** e1004095 (2015).
36. A. Stoessl, C. H. Unwin, E. W. B. Ward, Postinfectional fungus inhibitors from plants: Fungal
oxidation of capsidiol in pepper fruit. *Phytopathology* **63,** 1225–1231 (1973).
37. A. Mercier, et al., The polyphagous plant pathogenic fungus *Botrytis cinerea* encompasses
host-specialized and generalist populations. *Environ. Microbiol.* **21,** 4808–4821 (2019).
38. C. Plesken, et al., Genetic diversity of *Botrytis cinerea* revealed by multilocus sequencing, and
identification of *B. cinerea* populations showing genetic isolation and distinct host adaptation. *Front.*
Plant Sci. **12,** 663027 (2021).
39. E. Pichersky, R. A. Raguso, Why do plants produce so many terpenoid compounds? *New*
Phytol. **220,** 692–702 (2018).
40. N. Deighton, I. Muckenschnabel, A. J. Colmenares, I. G. Collado, B. Williamson, Botrydial is
produced in plant tissues infected by *Botrytis cinerea*. *Phytochemistry* **57,** 689–692 (2001).
41. J. L. Reino, R. Hernández-Galán, R. Durán-Patrón, I. G. Collado, Virulence–toxin production
relationship in isolates of the plant pathogenic fungus *Botrytis cinerea. J. Phytopathol.* **152,** 563–
566 (2004).
42. K. Hayashi, H. jan Schoonbeek, M. A. de Waard, Bcmfs1, a novel major facilitator superfamily
transporter from *Botrytis cinerea*, provides tolerance towards the natural toxic compounds
camptothecin and cercosporin and towards fungicides. *Appl. Environ. Microbiol.* **68,** 4996–5004
(2002).
43. D. Baulcombe, Plant science. Small RNA - The secret of noble rot. *Science* **342,** 45–46 (2013).
44. M. Wang, A. Weiberg, E. Dellota, D. Yamane, H. Jin, Botrytis small RNA *Bc-siR37* suppresses
plant defense genes by cross-kingdom RNAi. *RNA Biol.* **14,** 421 (2017).
45. M. Camagna, M. Ojika, D. Takemoto, Detoxification of the solanaceous phytoalexins rishitin,
lubimin, oxylubimin and solavetivone via a cytochrome P450 oxygenase. *Plant Signal. Behav.* **15,**
1707348 (2020).
46. M. Nakajima, J. Suzuki, T. Hosaka, T. Hibi, K. Akutsu, Functional analysis of an ATP-binding
cassette transporter gene in *Botrytis cinerea* by gene disruption. *J. Gen. Plant Pathol.* **67,** 212–214
(2001).
47. M. Fermaud, R. E. Gaunt, Thrips obscuratus as a potential vector of *Botrytis cinerea* in
kiwifruit. *Mycol. Res.* **99,** 267–273 (1995).
544

 Figures and Tables

546
Figure 1. Sensitivities and metabolic capacities of *Botrytis cinerea* and *Phytophthora infestans* to
sesquiterpenoid phytoalexins.
(A) Mycelial blocks (approx. 1 mm^3) of indicated pathogen were incubated in 50 µl water, 500 µM
capsidiol or 500 µM rishitin. Outgrowth of hyphae from the mycelial block (outlined by dotted red
lines) was measured after 24 h of incubation (n = 6). Bars = 100 µm. **(B)** Residual capsidiol and
rishitin was quantified by GC/MS 2 days after the incubation. **(C)** Predicted metabolism of capsidiol
and rishitin by *B. cinerea*. Note that the structure of oxidized capsenone was determined in this
study. Data marked with asterisks are significantly different from control as assessed by the two-
tailed Student's *t*-test: **$P < 0.01$, *$P < 0.05$.
556

558
Figure 3. Bcin08g00930 encodes a capsidiol detoxification enzyme, capsidiol dehydrogenase
BcCPDH.
(A) Mycelia of *Epichloë festucae* wild type (WT) or a transformant expressing Bcin08g00930 were
incubated in water or 100 μM capsidiol and outgrowth of mycelia was observed 7 days after
inoculation. Bars = 100 μm. **(B)** Hyphal outgrowth of *E. festucae* WT, transformants expressing
DsRed, Bcin08g00930 (BcCPDH) or Bcin12g01750 in water, 100 μM or 500 μM capsidiol was
measured after 24 h of incubation. Data are mean ± SE (n = 6). Asterisks indicate a significant
difference from WT as assessed by two-tailed Student's *t*-test, **P < 0.01. **(C)** Mycelia of *E. festucae*
WT or transformant expressing Bcin08g00930 (*Bccpdh*) were incubated in 100 μM capsidiol or for
48 h. Capsidiol and capsenone were detected by LC/MS.
569

A

B. cinerea - P_Bccpdh:GFP

	Control	Capsidiol	Rishitin	Capsidiol 3-acetate	Debneyol	Sclareol

B

Nicotiana benthamiana

2 hpi   8 hpi   2 dpi

C

N. tabacum (Tobacco)   **Capsicum annuum** (Bell pepper)   **Solanum lycopersicum** (Tomato)   **S. tuberosum** (Potato)

S. melongena (Eggplant)   **Vitis spp.** (Grape)   **Fragaria x ananassa** (Strawberry)   **Cucumis sativus L.** (Cucumber)

570
Figure 4. Specific activation of the *B. cinerea Bccpdh* promoter by capsidiol and its derivative.
(A) Mycelia of *B. cinerea* transformant containing *GFP* gene under the control of 1 kb *Bccpdh*
promoter (P_*Bccpdh:GFP*) was incubated in CM media containing 500 µM of anti-microbial
terpenoids. Bars = 50 µm. **(B)** (left and middle) Leaves of *N. benthamiana* were inoculated with

conidia of *B. cinerea* P_*Bccpdh:GFP* transformant. Expression of GFP in germinating conidia on
the leaf surface was observed by confocal laser microscopy 2 or 8 h after inoculation (hpi). CW,
stained with calcofluor white. Arrowheads indicate the appressoria of *B. cinerea.* Bars = 20 µm.
(right) Leaves of *N. benthamiana* were inoculated with mycelia of *B. cinerea* P_*Bccpdh:GFP*
transformant and the edge of the lesion was observed by confocal laser microscopy 2 d after the
inoculation (2 dpi). Bar = 100 µm. **(C)** Leaves of indicated plant were inoculated with mycelia of *B.*
cinerea P_*Bccpdh:GFP* transformant and the edge of the lesion was observed by confocal laser
microscopy 2 d after the inoculation. Bars = 100 µm.

Figure 5. *B. cinerea* BcCPDH is essential for the detoxification of capsidiol and virulence in the
plant species producing capsidiol.
(A) Mycelial block (approx. 1 mm^3) of *B. cinerea* wild type (WT) or *Bccpdh* KO mutant strain
(*Δbccpdh*-52) was incubated in 100 μM capsidiol for 4 d. Capsenone and capsidiol was detected
by LC/MS. **(B)** Mycelial blocks of *B. cinerea* WT or *Δbccpdh*-52 were incubated in capsidiol and
outgrowth of hyphae was measured after 36 h of incubation. Data are mean ± SE (n = 10). Asterisks
indicate a significant difference from WT as assessed by two-tailed Student's *t*-test, **P < 0.01. **(C)**
Leaves, tubers or fruits of indicated plant were inoculated with mycelial block (5 mm^3) of *B. cinerea*
WT or *Δbccpdh*-52 and lesion size was measured between 4 and 7 days after inoculation (dpi).
Asterisks indicate a significant difference from WT as assessed by two-tailed Student's *t*-test. **P
< 0.01. Lines and crosses (x) in the columns indicate the median and mean values, respectively.

Figure 6. Comparison of the *B. cinerea Bccpdh* gene locus with corresponding genome region of
other *Botrytis* species and *Sclerotinia sclerotiorum*. Edge of conserved region among *Botrytis*
species and specific region for *B. cinerea* (red line) were indicated by red arrowheads.
602

Figure 7. CPDH activity in *F. oxysporum* and *B. cinerea* strains isolated form a variety of plants. CPDH activity in *F. oxysporum* and *B. cinerea* strains isolated form a variety of plants. Mycelial blocks (approx. 1 mm^3) were incubated in 500 µM capsidiol for 2 days and capsidiol or capsenone were detected by GC/MS. Each elution profile describes the capsidiol and capsenone content in the solution after incubation of the strain with capsidiol. Strains with a substantial peak for capsidiol indicate the absence of CPDH activity.