Mutual repression between JNK/AP-1 and JAK/STAT stratifies cell behaviors during tissue regeneration

Janhvi Jaiswal [1,2], Raphael Engesser [3], Andrea Armengol Peyroton [1], Vanessa Weichselberger [1,2], Carlo Crucianelli [1], Isabelle Grass [1,4], Jens Timmer [3,4,5] and Anne-Kathrin Classen* [1,4,5]

* corresponding author

1 Hilde-Mangold-Haus, University of Freiburg, Germany

2 Spemann Graduate School of Biology and Medicine (SGBM), University of Freiburg, Germany

3 Institute of Physics and Freiburg Centre for Data Analysis and Modeling, University of Freiburg, Germany

4 CIBSS Centre for Integrative Biological Signalling Studies, University of Freiburg, Germany

5 BIOSS Centre for Biological Signalling Studies, University of Freiburg, Germany

20 **Abstract**
21

Epithelial repair relies on the activation of stress signaling pathways to coordinate cellular

repair behaviors. Their deregulation is implicated in chronic wound and cancer pathologies.

Despite such translational importance, an understanding of how spatial patterns of signaling

pathways and repair behaviors arise in damaged tissues remains elusive. Using TNF-α/Eiger-

mediated inflammatory damage to *Drosophila* imaginal discs, we uncover that JNK/AP-1

signaling cells act as paracrine organizers and initiate a mutual repression network that

spatially segregates JNK/AP-1 and JAK/STAT signaling cells into distinct populations. While

JNK/AP-1 signaling cells produce JAK/STAT-activating Upd ligands, these signal-sending

cells suppress activation of JAK/STAT via Ptp61F. Conversely, responding cells with activated

JAK/STAT suppress JNK activation via Zfh2. The resulting bistable segregation of signaling

domains is associated with distinct cellular tasks and regenerative potential. While JNK/AP-1

signaling cells at the wound center act as paracrine organizers, their cell cycle is senescently

arrested. Thus, compensatory proliferation occurs exclusively in JAK/STAT signaling cells at

the wound periphery. This spatial stratification is essential for proper tissue repair, as co-

activation of JNK/AP-1 and JAK/STAT in the same cells creates conflicting inputs on cell cycle

progression, leading to excess apoptosis of senescently arrested organizer cells. Finally, we

demonstrate that bistable spatial segregation of JNK/AP-1 and JAK/STAT drives senescent

and proliferative behaviors in transient as well as chronic tissue damage models, and

importantly, in Ras^{V12}, *scrib* tumors under the influence of JNK/AP-1 activity. Revealing this

previously uncharacterized regulatory network between JNK/AP-1, JAK/STAT and associated

cell behaviors have important implications for our conceptual understanding of tissue repair,

chronic wound pathologies and tumor microenvironments, where both pathways are strongly

implicated.

Introduction

Wound repair programs rely on the coordination of distinct responses to damage [1, 2]. Upon damage, wound-derived factors initiate inflammation and stress-signaling pathways which drive cellular responses like apoptosis, proliferation, survival and tissue remodeling [3-6]. While these responses are essential, it is critical that they remain spatio-temporally restricted to avoid the establishment of chronic wounds [7-9]. Chronic wounds are characterized by sustained inflammation, and deregulated proliferation and apoptosis [10]. The finding that these are also hallmarks of tumor microenvironments supported the idea that tumors resemble chronic, non-healing wounds [11-13]. Yet, the tissue-level coordination of signaling pathways and repair behaviors upon wounding, and the conditions under which they turn pathological remain elusive.

A myriad of wound models identified specific signaling pathways required for regeneration, of which several have also been shown to drive tumor growth. In particular, the JNK/AP-1 and the JAK/STAT pathways are consistently implicated in regeneration [14-17] as well as in tumors [18-23]. In the context of wound-healing, JNK/AP-1 activation is indispensable for regeneration [24-26]. JNK/AP-1 is one of the earliest signaling responses to damage, proposedly activated by wound-derived ROS signals [27, 28]. In this role, JNK/AP-1 regulates a variety of conflicting cell behaviors including apoptosis [29, 30], survival [31] and compensatory proliferation [32, 33]. These paradoxical behaviors have been extensively characterized individually [34-36], yet how they are organized on a tissue scale to ensure regeneration is not known.

We recently reported that high JNK/AP-1 signaling facilitates survival in wounds and tumors by mediating a cell cycle stall in G2 which is characterized by anti-apoptotic and senescent features [31]. Forced exit of these cells from G2 into mitosis and G1 caused a substantial increase in apoptosis, suggestive of the JNK-mediated G2 stall being a protective state during regeneration (Fig. S1A) [31]. This model is supported by recent evidence that the universal integrator of molecular damage, p53, is activated by the G2/M kinase Cdk1 to induce apoptosis competence upon exit from G2 [37]. The G2 protective state is critical as high JNK/AP-1 signaling cells are shown to produce mitogens necessary to induce compensatory proliferation and regeneration [28, 38-40]. Paradoxically, this reveals a cell population at the center of wounds which produces pro-proliferative signals but itself is prohibited from proliferation. These findings highlight the necessity of a mechanism, which ensures that high JNK/AP-1 signaling cells maintain a G2 stall in the presence of mitogenic signals, thereby outsourcing proliferation to other cells. Alternatively, it requires the existence of a mechanism that promotes cycling as well as survival of JNK-signaling cells. This illustrates that the underlying

84   mechanisms and molecular effectors that resolve these conflicting cell behaviors downstream
85   of JNK/AP-1 signaling in the same tissues remains elusive.
86
Prominent cytokines produced by JNK/AP-1 signaling cells belong to the Unpaired (Upd) family
[41-45] and activate the JAK/STAT pathway which, in turn, plays a significant role in mediating
proliferation [18, 20, 28, 39, 46-50] and cell survival during tissue stress [51, 52]. Interestingly,
we have previously also shown that JAK/STAT is required in the disc to restrict the expansion
of JNK/AP-1 signaling [51]. This is essential as JNK/AP-1 signaling within the tissue can self-
propagate by promoting an increase in the expression of its own paracrine activators - the
TNF-α/Eiger (Egr) cytokine [53, 54], as well as ROS [28, 55, 56].
94
These observations indicate an interdependence between the JNK/AP-1 and JAK/STAT
signaling pathways, whose coexistence has been consistently reported in wounds [14-17] and
tumor tissues [18-23]. Yet the regulatory network of the two pathways, beyond the regulation
of *upd's* by JNK, remains so far unexplored. As JNK/AP-1 and JAK/STAT drive contradicting
yet critical cell behaviors - namely G2 stalling and proliferation, linked by their regulation of
survival - characterizing their regulatory interactions is essential to understand how these
pathways cooperatively organize tissue stress responses.

Results
103

104   **Figure 1**
105   **JNK/AP-1 and JAK/STAT segregate into distinct spatial domains upon tissue damage**
106
To characterize the JNK/AP-1 and JAK/STAT signaling pathways at wound sites in detail, we
analyzed signaling dynamics and repair behaviors induced by targeted expression of the TNF-
α homologue *eiger* (*egr*) in the wing disc pouch (Fig. S1.1B,C). Ectopic *egr*-expression has
been extensively used as a model to genetically induce wounds and study regeneration [57].
egr-expression activates the stress-signaling pathway JNK/AP-1, assessed by the sensitive
JNK/AP-1 reporter - *TRE>RFP* (Fig. 1A,B) [58], which leads to loss of polarity, epithelial barrier
dysfunction (Fig. S1.1D-E') and inflammation [24, 25, 59]. Specifically, expression of *egr* also
induces *upd1-3* cytokine expression [51], MMP-1 production [60-62], ROS production [8, 63]
and protective responses including UPR [64] and, ultimately, NF-κB signaling upregulation [65]
(Fig.S1.1F-O, see also Fig.1E,F) in high JNK/AP-1 signaling cells – which are all known
hallmarks of inflammatory, and by extension, chronic wounds [6, 10]. In addition, high JNK/AP-
1 activity at the center of the *egr*-expressing domain (Fig. 1A,B) induces a senescent-like G2
cell cycle stall (Fig. 1C,D) which is essential for cell survival (Fig. S1.1A), as assessed using
the Fly-FUCCI toolkit [31, 66]. Indeed, we found that proliferation was excluded from the

inflammatory JNK-signaling domain at the center of the disc and was restricted to the wound
periphery (Fig. 1C,D,N). These observations highlight a pronounced spatial segregation of
regenerative cell behaviors linked to JNK-induced tissue damage responses.

Upon tissue damage, *upd's* - the pro-mitogenic cytokines for the JAK/STAT pathway - are
generally upregulated at wound sites where JNK/AP-1 activity is high (Fig. 1E,F,M, see also
S1.1F,G) [20, 28, 49, 51]. Consistent with their paracrine function [67, 68], JAK/STAT signaling
was broadly induced in the pouch periphery and hinge of *egr*-expressing discs (Fig. 1G,H),
coinciding with regions of proliferation (Fig. 1K,L,N). Yet, high JNK-signaling cells in *egr*-
expressing discs that upregulated the transcriptional reporters for *upd1* and *upd3* (Fig.
S1.1F,G; Fig. 1E,F,M) [69] did not undergo proliferation and, importantly, did not exhibit
JAK/STAT activity, as assessed by the *10xStat92E>dGFP reporter* (Fig. 1I,J) [70]. This was in
stark contrast to wing disc development where ligand expression and pathway activation
patterns largely coincide [70-73]. Our unexpected finding that JAK/STAT signaling is low in
upd-producing cells, strongly suggests that a distinct regulatory network controls JAK/STAT
activation during JNK-driven responses to tissue damage.

To further support this conclusion, we first wanted to rule out that the observed JAK/STAT
activation pattern in the periphery of *egr*-expressing discs perpetuated from early hinge
patterns of developing discs due to a Dilp8-induced developmental delay (Fig. S1.2 A) [74].
Hence, we monitored temporal changes in JAK/STAT activity via a time course of 0h, 7h and
14h of *egr*-expression. We consistently observed ectopic induction of the JAK/STAT reporter
in the hinge, above developmental levels and with an altered pattern (Fig. S1.2 B-M). This data
confirmed that *upd1-3*-driven JAK/STAT activity is induced *de novo* in the hinge and pouch
periphery upon tissue damage.

In summary, we find that JNK/AP-1 activation in the *egr*-expressing pouch coincides with the
expression of *upd* cytokines. We show that *upd* cytokines activate JAK/STAT only non-
autonomously in a cell population spatially segregated from high JNK-signaling and *upd*-
producing cells. In addition to the well-described non-autonomous activation of JAK/STAT by
JNK/AP-1 activation, this strongly indicates that cell-autonomous repression of JAK/STAT
signaling may be an important consequence of JNK/AP-1 activation (Fig. 1O). This is
consistent with recent observations of JNK/AP-1 and JAK/STAT activity segregation during
tissue stress in other tissue types, such as epidermal wounds [14]. We therefore demonstrate
a recurring theme of spatial segregation of JNK/AP-1 and JAK/STAT signaling upon tissue
damage. Importantly, we also show that such spatial patterns of segregating JNK/AP-1 and

JAK/STAT signals strongly correlate with spatial segregation of regenerative cell behaviors,

specifically G2 stalling and proliferation, into spatially distinct signaling domains.

Figure 2

JNK/AP-1 signaling represses JAK/STAT activity

We wanted to first understand whether JNK/AP-1 directly represses JAK/STAT activity and

thereby restricts it to the wound periphery. We therefore tested if clonal expression of JNK/AP-

1 directly represses JAK/STAT activity, by co-expressing a constitutively activated form of the

JNKK Hep along with p35 to prevent cell death [75, 76]. Indeed, *hep^act* clones cell-

autonomously repressed JAK/STAT activity (Fig. 2A-D, S2A), both when non-autonomously

activating it in surrounding cells in the peripheral pouch, hinge or peripodium (Fig. 2A-C, S2A),

as well as when within developmentally patterned activity in the hinge (Fig. 2D) [72]. These

experiments demonstrate that JNK/AP-1 represses JAK/STAT signaling cell-autonomously,

even when JNK/AP-1 has the ability to activate JAK/STAT non-autonomously.

In these mosaic experiments, hinge cells activated JAK/STAT signaling more robustly than

cells of the central pouch (Fig. 2A-B), likely as a result of developmental pre-patterns conferring

distinct competence [72]. Namely, the transcription factors Nubbin (Nub) and Rotund (Rn)

reduce developmental Stat92E activity in the pouch [77]. To first exclude the possibility that

JNK/AP-1 represses JAK/STAT by inducing pouch-specific transcription factors [77], we

assessed Nub and Rn levels within *hep^act* clones. *hep^act* clones did not ectopically induce Nub

or Rn, in fact, Nub was distinctly downregulated (Fig. S2B-D). Furthermore, we expressed *egr*

in the posterior compartment of the wing disc and examined whether JAK/STAT could be

activated in the adjacent anterior pouch (Fig. S1.1B). The cells highly positive for the JNK/AP-1

target gene - MMP-1 [78] did not activate JAK/STAT signaling. However, JAK/STAT signaling

could be distinctly activated in the surrounding tissue, including the anterior pouch (Fig. 2E,F).

These experiments confirm that JAK/STAT signaling can *in principle* be activated *de novo* in

the pouch by cell non-autonomous activity of JNK/AP-1 signaling cells. Altogether, these

experiments demonstrate that tissue damage associated with high levels of JNK/AP-1

signaling repress JAK/STAT activity in a cell autonomous manner, independent of

developmental competence.

Accordingly, we wanted to better understand if JAK/STAT repression by JNK/AP-1 may

depend on JNK/AP-1 signaling strength or time. We therefore analyzed the temporal evolution

of JNK/AP-1 and JAK/STAT activity in the JNK-signaling domain of *egr*-expressing discs. A

time course analysis revealed that JAK/STAT was mildly elevated during short-term JNK-

signaling in the pouch by 7 h of *egr*-expression, and increased even further with 14 h of *egr*-
expression (Fig. 2K,L,N,O, Fig. S2E,F,H,I,K,). Interestingly, cells displayed co-existence of
both pathways at moderate levels, but also often displayed exclusive activation of either
reporter (Fig. S2L,M). After 24 h of sustained *egr*-expression, high JNK/AP-1 signaling was
observed in the pouch, but notably, JAK/STAT activity had now largely disappeared from this
domain (Fig. 2M,P, Fig. S2G,J,K). These observations reveal that short-term or moderate
activity of JNK/AP-1 facilitates local, intermixed activation of JAK/STAT. However, sustained
and high activation of JNK/AP-1 efficiently excludes JAK/STAT activity cell autonomously. As
a consequence, JNK/AP-1 and JAK/STAT signaling are increasingly segregated into distinct
cell populations within the tissue.

Notably, these signaling dynamics are synchronously followed by cell behavior dynamics. We
observed that prolonged JNK/AP-1 signaling correlated with an increased exclusion of
proliferation from the pouch (Fig. 2G-J; Fig. 1K,L,N). Altogether, these data suggest that
chronic and high levels of JNK/AP-1 activation drive spatial wound site signaling and cell
behavior patterns from a local intermixed state to a larger, segregated and bistable field.

Figure 3
JNK/AP-1 acts as a wound organizer by employing a mutual-repression network

We wanted to understand the type of regulatory interactions between JNK/AP-1 and
JAK/STAT pathways that establish spatially segregated signaling domains on a tissue scale
[79-82]. Regulatory networks describe how components of different signaling pathways
interact [83-85] or are integrated to define a cell's signaling state and establish distinct cellular
outcomes [86, 87]. How regulatory networks can also organize cellular behaviors to influence
tissue-level outcomes has been extensively described using mathematical modeling [88-90].

To characterize the rules of the regulatory network between JNK/AP-1 and JAK/STAT that
ultimately drives emergence of spatial segregation to organize tissue stress responses, we
developed a mathematical model using partial differential equations. These described the
spatio-temporal evolution of JNK/AP-1 and JAK/STAT activation depending on relative
concentrations of components in the system (Fig. 3A-C) [86, 91]. We defined a set of
experimentally determined rules, namely: (1) JNK/AP-1 effectors propagate JNK/AP-1 via
production of paracrine factors, such as *egr* or diffusible ROS [28, 54, 92]; (2) JNK/AP-1
effectors activate JAK/STAT via production of paracrine factors, such as *upd*'s [20, 28, 49, 51];
(3) Both pathways use positive feedback loops to enhance and stabilize their own activation
[42, 55, 56, 93] and (4) JAK/STAT represses JNK/AP-1 cell-autonomously, via the ZEB-1

homologue Zfh2 (Fig. S3A-C) [51]. To simplify the model (Fig. 3B), we excluded negative

feedback loops on JNK/AP-1 and JAK/STAT [94-96], as they often have a more modulatory

function in noise buffering, temporal processing or saturation dynamics [97].

As we found experimentally that JNK/AP-1 repressed JAK/STAT cell-autonomously, we also

designed a mutual repression model that, together with the aforementioned rules, included a

rule that JNK/AP-1 represses JAK/STAT signaling (Fig. 3A,B') [98]. Using an unbiased

sampling approach of a large parameter space, we compared the ability of the two models to

produce simple bistable patterns (Fig. 3D). We found that the mutual repression model

produced 140-fold more positive outcomes than a uni-directional model (10588 vs. 72

solutions) (Fig. 3E). When we switched from scoring simple bistable patterns (Fig. 3D) to

observed bistable patterns (Fig. 3D') of JNK/AP-1 and JAK/STAT signaling in *egr*-expressing

wing discs, the difference in competencies between the two models became even more

prominent (Fig. 3E'). The uni-directional model was not sufficient to recapitulate observed

bistable patterns, which suggests that the rule of JNK/AP-1 cell-autonomously repressing

JAK/STAT is necessary for generating the segregated JNK/AP-1 and JAK/STAT signaling

patterns seen during tissue stress. Accordingly, we show that a mutual-repression network lies

at the core of JNK/AP-1 and JAK/STAT signaling interactions in response to tissue damage.

Supporting this result, mutual repression networks have been extensively described to guide

the formation of bistable states [99-105]. Furthermore, these networks can be further enhanced

by self-activation loops, such as those integrated in our model [98, 102, 106]. Indeed, if

JNK/AP-1 and JAK/STAT cross-talk were to completely lack mutual repression, JNK/AP-1 and

JAK/STAT signaling would be expected to expand from the wound site in an unrestrained

manner. Thus, the mutual repression regulatory network also ensures that these signaling

pathways remain restricted during regeneration.

The length scale-independent space utilized by the model raises the possibility that this mutual

repression network could act at length scales ranging from neighboring cells to multicellular

tissues. *In vivo*, length scales may depend on the level of network activation and thus the

concentration of non-autonomously acting cytokines (Egr, Upd's), as suggested by

experiments shown in (Fig. 2K-P and Fig. S1.2). Indeed, we observed that the number of

segregating solutions produced by the model increased with increasing rates of JNK/AP-1

activity (Fig. S3D). This observation parallels the progressive segregation of signaling domains

found experimentally upon chronic activation of JNK/AP-1 signaling (Fig. 2K-P).

Taken together we propose that JNK/AP-1 initiates a mutual repression network between

JNK/AP-1 and JAK/STAT signaling that, via the production of Upd's, facilitates the spatial

segregation of these two pathways. The spatial segregation of JNK/AP-1 and JAK/STAT via
mutual repression allows the tissue to maintain a JNK-dependent G2 stall in the presence of
pro-mitogenic signals, and restricts proliferation to a separate cell population. Thus, induction
of JNK/AP-1 signaling upon tissue damage and mutual repression with JAK/STAT is sufficient
to set up temporal and spatial signaling patterns, and functionally distinct cell populations.
273

274 **Figure 4**
275 **Ptp61F represses JAK/STAT activity in JNK/AP-1 signaling cells**
276

To understand how JNK/AP-1 may cell-autonomously repress JAK/STAT activation, we first
asked if JNK/AP-1 downregulates core components of the JAK/STAT pathway. Our previous
study demonstrated that the core components *dome*, *hop* and *stat92E* [107] were not
transcriptionally altered in *egr*-expressing discs [51]. Supporting this observation, we found
that the expression of GFP-tagged Hop [108] and Stat92E proteins from functionally validated
lines were unaltered in *egr*-expressing cells (Fig. S4.1A-I). This suggested that repression of
JAK/STAT signaling was likely due to altered function of regulators of JAK/STAT signaling. We
thus analyzed known negative regulators in *egr*-expressing discs. Levels of dPIAS/Su(var)2-
10 [109] and Apontic [110] were unchanged, but *ken* expression [111] was elevated in *egr*-
expressing discs (Fig. S4.2A-F). However, neither knock-down of *ken* nor *dPIAS/Su(var)2-10*
increased JAK/STAT activity in *egr*-expressing cells (Fig. S4.2G-L). Similarly, we could not
reproduce findings that knockdown of *apontic* reinstated JAK/STAT signaling in *egr*-expressing
discs (Fig. S4.2M-O) [112].
290

We then turned to the tyrosine phosphatase *Ptp61F,* a known repressor of Stat92E activation
[113, 114] which is transcriptionally induced in JNK-dependent tumors and other stress
conditions [23, 115]. Repression of JAK/STAT signaling by Ptp61F may occur at the level of
receptor complex signaling or Stat92E activity [113, 114, 116]. We found that knock-down of
ptp61F caused upregulation of the Stat92E reporter activity in high JNK-signaling cells of *egr*-
expressing discs (Fig. 4A-E, Fig. S4.3A-C). This suggests that Ptp61F is specifically required
in JNK-signaling cells to suppress Stat92E activity. Hence, to test if dephosphorylation of
Stat92E in JNK-signaling cells is rate-limiting for JAK/STAT signaling, we overexpressed an
HA-tagged Stat92E [117] in *egr*-expressing cells to saturate the dephosphorylation capacity of
Ptp61F. Indeed, *egr,stat92E-HA*-coexpressing cells in the central pouch domain activated the
JAK/STAT reporter at higher levels when compared to *egr*-expressing discs (Fig. 4F-J, Fig.
S4.3D,E). In contrast, co-expression of the JAK *hop* failed to induce JAK/STAT reporter activity
in *egr*-expressing discs (Fig. S4.3F-I). Importantly, this demonstrates that while signal
transduction from Upd ligands down to the effector Stat92E is in principle active in JNK-

signaling cells, it is the Ptp61F-mediated dephosphorylation of Stat92E that becomes rate-
limiting for JAK/STAT signaling.

Figure 5
JAK/STAT repression is required for the G2 stall and protects central wound cells from
apoptosis

As spatial segregation of JNK/AP-1 and JAK/STAT signaling domains was a robust feature of
egr-expressing discs, we wondered if the mutual repression network was necessary for
regeneration. We therefore forced expression of Stat92E in high JNK/AP-1 signaling cells, and
closely monitored cell and tissue level responses, such as cell survival and proliferation. The
loss of JNK/AP-1 and JAK/STAT segregation led to a pronounced increase in apoptosis (Fig.
5A-D). Co-expression of UAS-GFP confirmed that these apoptotic cells originated from cells
where JNK/AP-1 and JAK/STAT were forcibly co-activated (Fig. S5A-D).

We thus hypothesized that JAK/STAT interfered with the protective G2-stall of high JNK-
signaling cells, and thereby increased apoptosis [31, 37], While Stat92E expression altered
the proportion of S-phase cells in undamaged control discs, we observed no changes in the
proportion of cells in G1 or G2 (Fig. S5E-H). In contrast, a cell cycle analysis of *egr,Stat92E-*
HA co-expressing domains revealed a substantial increase in G1 and S-phase cells, indicating
that JNK-signaling cells failed to stall in G2 (Fig. 5E-J). Thus, forced Stat92E activation in high
JNK-signaling cells overrides the protective JNK-induced G2 stall and consequently increases
apoptosis. Supporting these results, we found that knock-down of Ptp61F in *egr*-expressing
discs also led to an increase in cycling cells in the tissue (Fig. 5O,P) correlating with increased
levels of apoptosis (Fig. 5K-N). This highlights the necessity of JNK/AP-1 and JAK/STAT
signaling segregation within the tissue, as it maintains the G2 stalled cell population. Thus, the
JNK-induced mutual repression network ensures the establishment of two distinct and
indispensable cell populations - one that stalls in G2 to prevent apoptosis and secrete
necessary pro-mitogenic factors like Upd and other paracrine effectors like Dilp8 and ImpL2
[20, 74], and a second that is able to respond to mitogenic signals and undergoes proliferation
necessary for regeneration. The strong similarities with paracrine signaling centers regulating
spatial patterning networks linked to specific cell behaviors and fate during development
encourages us to propose that the JNK-induced mutual repression network with JAK/STAT
functions as a central wound organizer network [118, 119].

Figure 6
The wound organizer network segregates JNK/AP-1 and JAK/STAT signaling to drive oncogenic growth

Our observation that tissue repair processes after chronic damage support the establishment of a signaling-sending and a responding cell population, which are respectively G2-stalled or proliferative, led us to speculate that such a wound organizer network could well-support a tumor microenvironment. Interestingly, the coexistence of JNK/AP-1 and JAK/STAT pathways in tumors has been extensively described [18-23] and the existence of signal-sending G2-stalled cells controlling non-autonomous proliferation has been demonstrated [31]. However, their spatial patterns have not been well defined.

To understand if JNK/AP-1 and JAK/STAT also engage in a mutual repression network under tumor conditions, we analyzed discs with a reduction in the well-characterized *scrib* tumor-suppressor gene [120] or ectopic expression of oncogenic *Ras^{V12}* [121, 122]. To generate a large genetically homogenous tissue, we used the pouch specific *rn-GAL4* driver to express *Ras^{V12}* or *scrib-RNAi* in the entire wing pouch. A wing pouch expressing oncogenic *Ras^{V12}* for 44 h did not activate JNK/AP-1 or JAK/STAT reporters, demonstrating that Ras^{V12} alone does not co-opt either pathway (Fig. 6A,B, Fig. S6A,B). RNAi-driven knock-down of *scrib* induced barrier dysfunction (Fig. S6C,D) and, consistent with previous reports [123, 124], moderately activated both JNK/AP-1 and JAK/STAT in the pouch (Fig. S6E-H). A closer examination, however, revealed that the *scrib-RNAi* expressing pouch predominantly segregated JNK/AP-1 and JAK/STAT signaling at the level of small cell clusters, while only few cell clusters displayed co-activation of both pathways (Fig. 6C). This pattern mirrored our observations of short-term *egr*-expression (Fig. S2L,M), thus suggesting that different contexts of mild JNK-activation cause segregation at short distances. These observations indicate that JNK/AP-1 activation upon barrier dysfunction via disruption of cell polarity also induces the wound organizer network, which subsequently sets up a bistable segregation of both pathways even at a short range between neighboring cells.

Curiously, if Ras^{V12} and *scrib* mutations occur in the same cell, they cooperate to cause dramatic overproliferation of imaginal discs, which is not observed in single *Ras^{V12}* or *scrib* mutant tissues. This cooperativity is thought to be driven by activation of JNK/AP-1, as a consequence of *scrib* disrupting cell polarity. JNK/AP-1 signaling then induces *upd's* and JAK/STAT activation, which can be utilized by a *Ras^{V12}* controlled network to drive overgrowth [18, 122]. We thus wondered if Ras^{V12} may disable the mutual repression motif between JNK/AP-1 and JAK/STAT. This would allow autocrine activation of JAK/STAT by JNK signaling in the same cell, as generally described in the literature [18, 122]. Cells driven to cycle despite

activation of JNK/AP-1 may then be protected by anti-apoptotic functions of Ras^{V12}. We wanted

to test this hypothesis and closely monitored JNK/AP-1 and JAK/STAT signaling as well as

cell proliferation in Ras^{V12} and *scrib RNAi*-coexpressing discs. Surprisingly, we found that

JNK/AP-1 and JAK/STAT activation were still clearly segregated into exclusive signaling

domains (Fig. 6D-F, Fig. S6I) with proliferation still predominantly associated with JAK/STAT

(Fig. 6G-I). This suggests that even in a Ras^{V12} and *scrib* cooperativity model, where all cells

are genetically identical, bistability driven by the mutual repression network organizes the

pouch into distinct JNK/AP-1 or JAK/STAT-signaling domains. This observation aligns with

previous observations that cooperativity between Ras and *scrib* can rely, in principle, on non-

autonomous interactions [18, 125]. Crucially however, our observations demonstrate that non-

autonomous cooperation between JNK/AP-1-activating and *Ras^{V12}* mutation arises as a self-

organizing principle from a wound organizer network creating segregating signaling domains

in a genetically homogenous tumorigenic tissue. Moreover, it suggests that activation of this

wound organizer network and segregation of signaling and proliferative tasks may provide an

advantage to tumors over autocrine integration of JNK/AP-1 and JAK/STAT signaling in the

same cell.

Discussion

Previous studies established a role for JNK/AP-1 in promoting regeneration upon injury [35,

59]. This role is, in part, mediated by the expression of *upd's* which are essential to promote

proliferation and survival via JAK/STAT signaling in a non-autonomous manner [28, 35, 48,

51, 126-129]. By identifying a mutual repression network between JNK/AP-1 and JAK/STAT,

we conceptually establish JNK as a core organizer of tissue repair with parallels to organizers

of cell fate patterning in developing tissues. The rules of the wound regulatory network we

describe have important implications. Previous reports suggest that JNK/AP-1 activates itself

non-autonomously, for example via activation of paracrine *egr* or ROS [28, 54, 92]. This implies

that JNK/AP-1 activation could result in unchecked spatial expansion of JNK/AP-1-signaling,

and accordingly of JAK/STAT signaling. Their unrestrained co-expansion would likely result in

chronic inflammation. However, the mutual repression motif restrains the expansion of both

JNK/AP-1 and JAK/STAT signaling domains, thus defining stable regions of inflammatory

(mitogenic signals) and regenerative (pro-proliferative) responses at the start of the tissue

repair process. We demonstrate that molecular mediators of mutual repression are Zfh2

downstream of JAK/STAT [51], and Ptp61F downstream of JNK.

At the level of cell behaviors, JNK/AP-1 and JAK/STAT generate mutually exclusive responses

(Fig. 7B). JNK/AP-1 signaling supports wound center behaviors like tissue sealing [24, 25, 78],

cell fusion [14], establishment of paracrine signaling and importantly, an apoptosis-resistant
state that depends on a G2 cell cycle stall [31]. In contrast, JAK/STAT signaling supports
wound-distal behaviors, such as compensatory proliferation, potentially via upregulation of G1-
S cyclins such as CycD or CycE [130, 131].

By reducing Stat92E activation upon damage, JNK/AP-1 creates a permissive state for the G2
stall by preventing cycling, thereby protecting cells from apoptosis in a cytotoxic, inflammatory
environment (Fig. 7C; Fig. S1A). When high levels of both pathways are co-activated in the
same cell, the anti-apoptotic state of the G2 stall is in conflict with proliferative outcomes and
therefore active cycling which causes an increase in apoptosis. Consequently, apoptosis of
these wound center signaling cells could jeopardize wound organizer function and ultimately
tissue repair behaviors (Fig.7D). Therefore, the JNK-activated G2-stall and JAK/STAT-
activated cell cycling need to be spatially segregated. Importantly, our observations indicate
that, with exceptions [132], even successful tumors avoid coexistence of JNK and JAK/STAT
signaling in the same cell as they do not profit from the conflicting cell-autonomous inputs on
proliferation and survival decisions. Instead, tumors may highjack a wound organizer network
to chronically activate paracrine mitogenic signals but outsource proliferation to other cells as
a consequence of the senescent properties of JNK-signaling, G2-stalled cells [31].

We find that spatial patterns generated by the JNK/AP-1 and JAK/STAT regulatory network
act at different length scales, depending on the duration and strength of JNK/AP-1 activation
(Fig. 7A). Transient or low JNK/AP-1 activity generates short range spatial segregation of
JNK/AP-1 and JAK/STAT. Specifically, we observed their segregation between neighboring
cells upon transient expression of *egr* or mild disruption of barrier function in *scrib*-RNAi
expressing tissue. Cell-by-cell segregation may allow tissue repair processes to respond
dynamically. Switching between (1) a stalled, apoptosis-resistant state supporting survival and
paracrine signaling in an inflammatory environment or (2) a regenerative, proliferative
phenotype could facilitate continuous integration of the repair state. As wound healing
progresses and the JNK/AP-1 signaling field shrinks, shifting spatial probabilities of stalled and
cycling cells would ultimately resolve the damage.

In contrast, prolonged or high JNK/AP-1 activity segregates JAK/STAT signaling more robustly
and consequently into larger bistable fields, prolonging wound resolution and increasing the
risk for chronic wound pathologies. Indeed, prolonged JNK/AP-1 activity in *Ras^{v12}*,*scrib*-RNAi
tumors or by chronic expression of *egr* drives robust bistable segregation of JNK/AP-1 and
JAK/STAT signaling into larger domains. Similarly, chronic wounds in human patients display
striking segregation of cellular responses. In a central inflammatory domain, cells acquire

senescent phenotypes, arrest and fail to proliferate. In the periphery, a hyperproliferative edge
is evidence of chronic mitogenic signals emanating from the wound [3, 9, 10, 133]. These
clinical patterns resemble patterning in *egr*-expressing discs, thereby suggesting that
Drosophila imaginal discs may represent a suitable model to study wound pathologies driven
by prolonged activation of inflammatory signaling.

Materials and Methods

Fly stocks

All fly stocks and experimental crosses were maintained on standard media and raised at 18 °C or room temperature (22 °C) unless otherwise specified. For detailed genotypes, please refer to Table S1 and S2.

Table S1 Key Resources Table

Designation - Genetic reagent (*D. melanogaster*)	Source or Reference	Identifiers	Additional information
'RFP': *Act5C>FRT.CD2>GAL4, UAS-RFP*	PMID: 29494583	BDSC: 30558	
'apt-RNAi': *UAS-apt RNAi (TRiP.JF02134)*	PMID: 26320097	BDSC: 26236	
'Dif-GFP': *Pbac(Dif-GFP.FPTB)VK00033*		BDSC: 42673	
dome[G0441]	PMID: 11751581	BDSC: 12030	
'dome-GFP': *FlyFos (dome::TGVBF)*	PMID: 26896675	VDRC: 318098	
'E2F1-FUCCI, CycB-FUCCI'; *Ubi-GFP.E2f1*[1-230], *Ubi-mRFP1.NLS.CycB*[1-266]	PMID: 24726363	BDSC: 55123	
'egr[ts]': *tub-GAL80ts, UAS-egr*			I.Hariharan, University of California, Berkeley
'en>RFP': *enGAL4, UAS-mRFP.NLS*		BDSC: 30557	
GFP-HP-1	PMID: 18316477	BDSC: 30561	
gstD-GFP	PMID: 18194654		D. Bohmann, University of Rochester Medical Center
'hep[act]': *UAS-hep*[act]	PMID: 10903185	BDSC: 9306	
'hop': *UAS-hop.H*	PMID: 7796812	BDSC: 79033	
hop[34]	PMID: 3095163		N. Perrimon
'hop-GFP': *FlyFos (hop::TGVBF)*	PMID: 26896675	VDRC: 318158	
hsflp[122]			D. Bilder, University of California, Berkeley
'GFP': *UAS-GFP S56T*		BDSC: 1521	
'ken-LacZ': *P{PZ}ken*[02970]	PMID: 10471706	BDSC: 11244	
'ken-RNAi': *UAS-ken-RNAi (TRiP.HMS01219)*	PMID: 26320097		
'ptp61f-RNAi': *UAS-ptp61f-RNAi* [i2-5]	PMID: 19398577	BDSC: 56510	
'p35': *UAS-p35.H*	PMID: 9144202	BDSC: 5072	
'rn(ts)>': *rn*[GAL4-DeltaS], *tubGAL80*[ts]		BDSC: 8142	recombinant
'rn(ts)>egr': *rn*[GAL4-5], *UAS-egr, tubP-GAL80*[ts]		BDSC: 8142; BDSC: 7018	recombinant; I.Hariharan, University of California, Berkeley
'Ras[v12], scrib-RNAi': *UAS-Ras*[v12],*scrib-RNAi*			M. Uhlirova, University of Cologne
'Ras[v12]': *UAS-Ras*[v12]			H. Richardson, University of Melbourne
'rnE/F-EGFP': *rn(E/F)-EGFP (ds Red+)*	PMID: 26497147		P. Volkan, Duke University School of Medicine
'scrib-RNAi': *UAS-scrib RNAi*			M. Uhlirova, University of Cologne
'Stat92E': *UAS-Stat92E.ORF.3xHA*	PMID: 23583758	Fly-ORF F000750	
'Stat92E[85c9]': *FRT82B stat92E*[85c9]			E. Bach, New York University School of Medicine
'Stat92E>dGFP': *10xStat92E>dGFP*	PMID: 17008134		

'Stat92E>GFP': 10xStat92E>GFP	PMID: 17008134		
'Stat92E-GFP': PBac{Stat92E-GFP.FLAG}^{VK00037}		BDSC: 38670	
'su(var)2-10-GFP': PBac{Su(var)2-10-GFP.FPTB}^{VK00037}	PMID: 29284660	BDSC: 64795	
'su(var)2-10-RNAi': UAS- Su(var)2-10-RNAi $^{TRiP.HMS00750}$	PMID: 26320097	BDSC: 32956	
'TRE>RFP': TRE-DsRed.T4	PMID: 22509270		
'Xbp1-GFP': UAS-Xbp1-GFP.HG	PMID: 23160805	BDSC: 60731	
upd3.1-3-LacZ			I.Hariharan, University of California, Berkeley
upd-LacZ: upd>LacZ (PD1)	PMID: 8582614		Y. H. Sun, Academia Sinica, Taipei
w^{118}			
'zfh2': UAS-Zfh2[EAB]	PMID: 24172131		F. Diaz-Benjumea, Universidad Autónoma de Madrid

Antibodies / Commercial Kits	Source or Reference	Identifiers	Dilution
Chicken anti-GFP	Abcam	Cat. #: ab13970	(1:1000)
Mouse anti-H3-pS10	Abcam	Cat. #: ab14955	(1:2000)
Mouse anti-β-Galactosidase	Promega	Cat. #: Z3782	(1:1000)
Mouse monoclonal anti-MMP-1	DSHB	Cat. #: 3A6B4	(1:30)
Mouse monoclonal anti-MMP-1	DSHB	Cat. #: 3B8D12	(1:30)
Mouse monoclonal anti-MMP-1	DSHB	Cat. #: 5H7B11	(1:30)
Mouse monoclonal anti-Nubbin	DSHB	Cat. #: 2D4	(1:100)
Mouse monoclonal anti-α-Tubulin	Sigma-Aldrich	Cat. #: T9026	(1:5000)
Rabbit anti-cleaved Dcp-1	Cell Signaling	Cat. #: 9578	(1:200)
Rabbit monoclonal anti-GFP	Invitrogen	Cat. #: G10362	(1:200)
Rabbit polyclonal anti-apt	R. Schuh, MPI-MS, Göttingen	NA	(1:200)
Rat monoclonal anti-GFP	Chromotek	Cat. #: 3H9-100	(1:1000)
Rat monoclonal anti-HA	Monoclonal Antibody Core Facillity at the Helmholtz Zentrum München	Clone #: 3F10	(1:20)
Rat monoclonal anti-RFP		Clone #: 5F8	(1:20)
Click-iT Plus EdU Alexa Fluor 647 Imaging Kit	Invitrogen	Cat. #: C10640	

PMID – PubMed Identifier; BDSC – Bloomington Drosophila Stock Center; DSHB – Developmental Studies Hybridoma Bank, VDRC – Vienna Drosophila Stock Collection, FlyORF – University of Zurich ORFeome Project

466 ## Table S2. Detailed Genotypes

Figure	Panel	Genotype	Temperature shift	Recovery
Figure 1	A	*TRE-RFP; 10xStat92E>dGFP*	24h 30°C, D7 AED	-
Figure 1	B	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 1	C	*Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 1	D	*Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 1	E	*TRE-RFP; Upd3.1-3-LacZ/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 1	F,M	*TRE-RFP; Upd3.1-3-LacZ/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 1	G,I,K	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 1	H,J,L,N	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.1	D	*rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-

Figure S1.1	E	*rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.1	F	*Upd-LacZ/X; TRE-RFP; TM6B, tub-GAL80*	24h 30°C, D7 AED	-
Figure S1.1	G	*Upd-LacZ/X; TRE-RFP; rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.1	H	*SuVar2-10 GFP/rn-GAL4, tub-GAL80ts*	24h 30°C, D7 AED	-
Figure S1.1	I	*SuVar2-10 GFP/rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.1	J	*gstD-GFP; rn-GAL4, tub-GAL80ts*	24h 30°C, D7 AED	-
Figure S1.1	K	*gstD-GFP; rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.1	L	*UAS-Xbp1-GFP.HG/ rn-GAL4, tub-GAL80ts*	24h 30°C, D7 AED	-
Figure S1.1	M	*UAS-Xbp1-GFP.HG/ rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.1	N	*TRE-RFP; Dif-GFP/rn-GAL4, tub-GAL80ts*	24h 30°C, D7 AED	-
Figure S1.1	O	*TRE-RFP; Dif-GFP/rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S1.2	B,E	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts*	0h 30°C, D7 AED	-
Figure S1.2	C,F	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts*	7h 30°C, D7 AED	-
Figure S1.2	D,G	*TRE-RFP; 10xStat92E>dGFP/TM6B, tub-GAL80ts*	14h 30°C, D7 AED	-
Figure S1.2	H,K	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	0h 30°C, D7 AED	-
Figure S1.2	I,L	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	7h 30°C, D7 AED	-
Figure S1.2	J,M	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	14h 30°C, D7 AED	-

Figure 2	A	*hsflp122; UAS-p35; 10xStat92E>dGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	10' HS 37°C, D7 AED	28h 22°C
Figure 2	B,C	*hsflp122; UAS-p35/UAS-hepact; 10xStat92E>dGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	10' HS 37°C, D7 AED	28h 22°C
Figure 2	D	*hsflp122; UAS-p35/UAS-hepact; 10xStat92E>dGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	7' HS 37°C, D5 AED	48h 22°C
Figure 2	E	*TRE-RFP; 10xStat92E>dGFP*	24h 30°C, D7 AED	-
Figure 2	F	*enGAL4-mRFP.NLS/TRE-RFP; 10xStat92E>dGFP/tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure 2	G	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts*	7h 30°C, D7 AED	-
Figure 2	H,K,N	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	7h 30°C, D7 AED	-
Figure 2	I	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts*	14h 30°C, D7 AED	-
Figure 2	J,L,O	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	14h 30°C, D7 AED	-
Figure 2	M,P	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-

Figure S2	A	*hsflp122; UAS-p35; 10xStat92E>dGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	10' HS 37°C, D7 AED	28h 22°C
Figure S2	B	*hsflp122; UAS-p35; 10xStat92E>dGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	7' HS 37°C, D7 AED	28h 22°C
Figure S2	C	*hsflp122; UAS-p35/UAS-hepact; 10xStat92E>dGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	7' HS 37°C, D7 AED	28h 22°C
Figure S2	D	*hsflp122; UAS-p35; rn(E/F)-EGFP/Act5C.GAL4 (FRT.CD2), UAS-RFP*	7' HS 37°C, D7 AED	28h 22°C
Figure S2	E,F,G	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts*	0h,7h,14h 30°C, D7 AED	-
Figure S2	H,I,J,K	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	0h,7h,14h 30°C, D7 AED	-
Figure S2	L	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts*	7h 30°C, D7 AED	-
Figure S2	M	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80ts, UAS-egr*	7h 30°C, D7 AED	-

Figure S3	A, C	*rn-GAL4, tub-GAL80ts, UAS-egr*	24h 30°C, D7 AED	-
Figure S3	B, C	*rn-GAL4, tub-GAL80ts, UAS-egr; UAS-zfh2EAB-mCherry*	24h 30°C, D7 AED	-

Figure	Panel	Genotype	Conditions	
Figure 4	A	*10xStat92E>dGFP/UAS-ptp61F RNAi^{i2-5}; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 4	B,D,E	*10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 4	C,D,E	*10xStat92E>dGFP/UAS-ptp61F RNAi^{i2-5}; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 4	F,J	*CycE-lacZ/10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 4	G,J	*CycE-lacZ/10xStat92E>dGFP; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 4	H,J	*CycE-lacZ/10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 4	I,J	*CycE-lacZ/10xStat92E>dGFP; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.1	A	*dome^{G0441};;	dome^{G0441};;dome-GFP*	24h 30°C, D7 AED	-
Figure S4.1	B	*hop^{34};;	hop^{34};;hop-GFP*	24h 30°C, D7 AED	-
Figure S4.1	C	*stat^{85c9}/TM6c	stat^{85c9}/stat^{85c9}	Stat92E-GFP; stat^{85c9}/TM6c	Stat92E-GFP; stat^{85c9}/stat^{85c9}*	24h 30°C, D7 AED	-
Figure S4.1	D	*Stat92E-GFP	GFP-HP1	w$^{[118]}$*	D7 AED	-
Figure S4.1	E	*w$^{[118]}$*	24h 30°C, D7 AED	-
Figure S4.1	F	*hop-GFP/TM6B, tub-GAL80*	24h 30°C, D7 AED	-
Figure S4.1	G	*hop-GFP/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.1	H,J,M	*Stat92E-GFP;*	24h 30°C, D7 AED	-
Figure S4.1	I	*Stat92E-GFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.1	K,M	*UAS-p35/Stat92E-GFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	A	*Su(var)2-10-GFP; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S4.2	B	*Su(var)2-10-GFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	C	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S4.2	D	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	E	*ken-LacZ^{02970};*	24h 30°C, D7 AED	-
Figure S4.2	F	*ken-LacZ^{02970}; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	G	*10xStat92E>dGFP/UAS-ken-RNAi; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S4.2	H,K	*10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	I	*10xStat92E>dGFP/UAS-ken-RNAi; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	J	*10xStat92E>dGFP; UAS-Su(var)2-10-RNAi/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S4.2	L	*10xStat92E>dGFP; UAS-Su(var)2-10-RNAi/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.2	M	*10xStat92E>dGFP; UAS-apt-RNAi/rn-GAL4, tub-GAL80^{ts}*	48h 30°C, D5 AED	-
Figure S4.2	N	*10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	48h 30°C, D5 AED	-
Figure S4.2	O	*10xStat92E>dGFP; UAS-apt-RNAi/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	48h 30°C, D5 AED	-
Figure S4.3	A,C	*CyO; 10xStat92E>dGFP*	D7 AED	-
Figure S4.3	B,C	*enGAL4, UAS-mRFP.NLS/UAS-ptp61F RNAi; 10xStat92E>dGFP*	D7 AED	-
Figure S4.3	D	*UAS-GFP; rn-GAL4, tub-GAL80^{ts}*	12h 30°C, D7 AED	-
Figure S4.3	E	*UAS-GFP; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}*	12h 30°C, D7 AED	-
Figure S4.3	F	*10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S4.3	G	*10xStat92E>dGFP/UAS-hop.H; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-

Figure	Panel	Genotype	Conditions	
Figure S4.3	H	*10xStat92E>dGFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S4.3	I	*10xStat92E>dGFP/UAS-hop.H; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	A,C,D	*TRE-RFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	B,C ,D	*TRE-RFP; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	E,G,H	*Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 5	F,G,H	*Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	I	*UAS-GFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	J	*UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	K,O,M,N	*UAS-GFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure 5	L,P,M ,N	*UAS-GFP/UAS-ptp61F RNAi $^{i2-5}$; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S5	A	*UAS-GFP; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S5	B	*UAS-GFP; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S5	C	*UAS-GFP; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S5	D	*UAS-GFP; UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S5	E	*Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S5	F	*Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$; rn-GAL4, tub-GAL80^{ts}, UAS-egr*	24h 30°C, D7 AED	-
Figure S5	G	*UAS-GFP; rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure S5	H	*UAS-Stat92E-3xHA/rn-GAL4, tub-GAL80^{ts}*	24h 30°C, D7 AED	-
Figure 6	A	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure 6	B	*UAS-Ras^{v12}/TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure 6	C,C',G	*UAS-scrib-RNAi/TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure 6	D,D',E,F,H,I	*UAS-Ras^{v12},scrib-RNAi/TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure S6.1	A,E	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure S6.1	B	*UAS-Ras^{v12}/TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure S6.1	C	*10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	48h 30°C, D6 AED	-
Figure S6.1	D	*UAS-scrib-RNAi; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	48h 30°C, D6 AED	-
Figure S6.1	F	*UAS-scrib-RNAi/TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure S6.1	G,I	*UAS-Ras^{v12},scrib-RNAi/TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}*	44h 30°C, D6 AED	-
Figure S6.1	H	*TRE-RFP; 10xStat92E>dGFP/rn-GAL4, tub-GAL80^{ts}, UAS-egr*	44h 30°C, D6 AED	-

Fly genetics

Fosmids carrying GFP-tagged *hop* (VDRC 318158) and *Stat92E* (BDSC 38670) alleles, expressed proteins at expected molecular weights and, at least partially complemented respective null alleles, suggesting that both GFP-fusion proteins are functional. A potential Dome-GFP fosmid (VDRC 318098) failed to give rise to a GFP protein by immunofluorescence and Western blot analysis and failed to complement a *dome* allele, thus Dome levels could not directly be analyzed. Genetic complementation assays were carried out by crossing the GFP-tagged fosmids into the background of LOF mutants (*dome^{G0441}, hop^{9p5}* and *stat^{85C9}*). For X-linked *dome^{G0441}* and *hop^{9p5}* alleles, we calculated the percentage of hemizygous viable adult males emerging from control and experimental crosses (Fig. S4.1A,B). For the autosomal *stat^{85C9}* allele, we determined pupariation success of homozygous mutant larvae by counting the number of pupariated larvae in control and experimental crosses (Fig. S4.1C).

We validated RNAi lines by demonstrating that they caused changes to the developmental pattern of JAK/STAT activity in the hinge if expressed throughout normal wing disc development. Crosses were set as previously described, using the *en-GAL4* driver to express the RNAi in the posterior compartment. Larvae were kept at 18°C till D7 AEL. On D7, early L3 staged larvae were dissected, showing characteristic JAK/STAT reporter activity pattern in the hinge. Following dissection and staining, sibling controls and RNAi expressing discs were mounted on the same slide. All discs were imaged at the same settings and the JAK/STAT reporter intensities were quantified in selected regions in the anterior and posterior hinge (Fig. S4.3C). Intensity values from the anterior compartment (not expressing the RNAi) was used as the internal control.

Flip-out clones

GAL4/UAS-driven 'flip-out' overexpression experiments utilized heat-shock-driven expression of a flippase to clonally express *a UAS* construct of choice. After 6h of egg collections, heat-shock was induced on developmental day 5 or 6 at 37°C for 7-10 min . Larvae were dissected at wandering 3rd instar stage or as indicated (28h or 48h after heat-shock, Fig. 2A-D, S2.A-D).

Genetic cell ablation using Gal4/UAS/Gal80ts

To induce expression of *egr*, experiments were carried out as described in [31, 38, 51] with few modifications. Briefly, larvae of genotype *rn-GAL4, tub-GAL80^{ts}* and carrying the desired *UAS*-transgenes were staged with a 6h egg collection and raised at 18°C at a density of 50 larvae/vial. Overexpression of transgenes was induced by shifting the temperature to 30°C for 24h at D7 after egg deposition (AED), and larvae dissected at recovery time point R0 h (Fig.

S1.1C) unless noted otherwise. For time course and tumor experiments, transgenes were

induced by shifting the temperature to 30°C for 0h, 7h, 14h or 24h at D7 (Fig. S1.2, 2 and S2),

or 44h at D6 (Fig. 6 and S6) respectively. Larvae were subsequently dissected for analysis or

allowed to recover at 22°C for the indicated time. All images represent R0 h unless noted

otherwise. Control genotypes were either *rn^{ts}>*, or sibling animals (*+/TM6B, tubGAL80 or*

+/TM6c) (Smith-Bolton et al. 2009). All experiments were performed with ≥ 2 biological

replicates.

Immunohistochemistry

Wing discs from third instar larvae were dissected and fixed for 15 min at room temperature in

4% paraformaldehyde in PBS. Washing steps were performed in PBS containing 0.1%

TritonX-100 (PBT). Discs were then incubated with primary antibodies (described in **Table S1)**

in PBT, gently mixing overnight at 4°C. Tissues were counterstained with DAPI (0.25 ng/µl,

Sigma, D9542), Phalloidin-Alexa Fluor 488/647 (1:100, Life Technologies) or Phalloidin-

conjugated TRITC (1:400, Sigma) during incubation with cross-absorbed secondary antibodies

coupled to Alexa Fluorophores (Invitrogen or Abcam) at room temperature for 2h. Tissues were

mounted using SlowFade Gold Antifade (Invitrogen, S36936). Whenever possible,

experimental and control discs were processed in the same vial and mounted on the same

slides to ensure comparability in staining between different genotypes. Images were acquired

using the Leica TCS SP8 Microscope, using the same confocal settings and processed using

tools in Fiji. Of note, all wing discs expressing the *10xStat92E>dGFP* reporter were boosted

with an anti-GFP antibody.

EdU Labelling

EdU incorporation was performed using the Click-iT Plus EdU Alexa Fluor 647 Imaging Kit,

(described in **Table S1)** prior to primary antibody incubation. Briefly, larval cuticles were

inverted in Schneider's medium and incubated with EdU (10µM final concentration) at RT for

15 minutes. Cuticles were then fixed in 4% PFA/PBS for 15 minutes, washed for 30 minutes

in PBT 0.5%. EdU-Click-iT labeling was performed according to manufacturer's guidelines.

Tissues were washed in PBT 0.1%, after which immunostainings, sample processing and

imaging were carried out as described above.

Western Blots

Cell lysates from third instar wing imaginal discs and brains were prepared in lysis buffer (50

mM Tris-HCl pH 7.5, 300mM NaCl, 0.1mM EDTA, 1% Triton X-100, 0.1% SDS, 5% Glycerol,

1mM PMSF, 1/10 tablet of Complete Mini Protease Inhibitor Cocktail) on ice. Protein samples

were loaded onto a 10% polyacrylamide gel, along with Chameleon Duo Pre-stained Protein

Ladder (LI-COR, P/N 928-60000) as a molecular weight ladder and run at 150V. After SDS-
PAGE, proteins were transferred onto a nitrocellulose membrane (Bio-Rad, 162-0115,
0.45µm) in transfer buffer (25 mM Tris, 192 mM glycine, 20% (v/v) methanol) using the wet-
tank method with a current density of 300mA for 1h. Prior to antibody incubation, membranes
were blocked with 5% milk in PBS. Membrane was incubated with primary antibodies in PBS-T
(1% Triton-X in 1xPBS) - rat anti-GFP (Chromotek, 3H9-100, 1:1000) and mouse anti-α-
tubulin (Sigma, T9026, 1:5000) on a rotative plate at 4°C overnight. Primary antibodies were
removed and washed in dH_2O twice for 5 mins each. Membrane was then incubated with
secondary antibodies diluted in PBS-T - donkey α-mouse secondary (LI-COR, 926-68072,
1:20000) labelled with a 700nm IRDye and goat α-rat (LI-COR, 926-32219, 1:20000) labelled
with an 800nm IRDye. Membrane was washed in dH_2O twice for 5 mins each before detection
using the Image Studio Software on the Odyssey SA system (LI-COR) (Fig. S4.1D).

Image Analysis and Quantification

General comments
Images were processed, analyzed and quantified using tools in Fiji (ImageJ v2.0.0) [134].
Extreme care was taken to apply consistent methods (i.e. number of projected sections,
thresholding methods, processing) for image analysis. Figure panels were assembled using
Affinity Designer v1.10. Statistical analyses were performed in Graphpad Prism or R v3.3.3
(www.R-project.org).

Total disc area quantification
Maximum intensity projections of z-stacks for each disc were obtained in Fiji. Masks were
generated after applying a fixed threshold (10-255) to the DAPI channel and noise was reduced
using the 'Despeckle' function. Area measurement from the resulting masks were obtained
using the 'Measure' function.

***10xStat92E>dGFP* quantification**
Maximum intensity projections of selected confocal sections were taken, excluding the
peripodium and carefully chosen to capture signaling activity within the disc proper. In the *egr-*
expressing or *egr,transgene*-coexpressing discs, a mask of the pouch domain was generated.
Region Of Interest (ROI) outlines were either manually drawn on the DAPI channel to include
the pouch and hinge domains till the medial hinge fold or by thresholding Nubbin staining.
Centroid co-ordinates were identified using the 'Measure' function and marked with a point
tool. To measure the fluorescence intensity of the *Stat92E>dGFP* reporter, a square ROI

(25x25μm) was placed centrally over the centroid and mean intensity within the ROI was
obtained using the 'Measure' function in Fiji. (Fig. 4)

Dcp-1 area quantification
Maximum intensity projections of stacks for each disc were obtained in Fiji. Masks were
obtained by thresholding for signal of interest. Signal to noise ratio was enhanced by 'Remove
outliers' (bright, radius = 1.5) or 'Despeckle' functions. The '3D object counter' function was
used to obtain surface area of the generated mask. To control for differences arising from the
total size of the discs, ratios between the Dcp-1 area and total WID area were obtained for
each disc and reported as percentage values. (Fig. 5C,D,M,N)

Tracing JNK, JAK/STAT, EdU and Upd reporters profiles
Partial max projections from confocal stacks of the disc proper were generated, carefully
excluding any peripodial signal. For hinge profiles of the JAK/STAT reporter, size-matched
discs were chosen and line traces were manually drawn along the hinge using the polygon
selection tool. Reporter intensity values from the 'Plot profile' function were averaged for 0h,
7h and 14h control and *egr*-expressing discs and plotted. (Fig. S1.2)
For reporter patterns along disc center to disc periphery, JNK/AP-1 masks were made using
thresholding tools and ROI outlines were generated. Based on ROIs, the centroid value was
determined and *n≥15* tracks were drawn from the centroid outwards towards the hinge for a
fixed distance - covering the JNK/AP-1 domain, the JNK/AP-1-JAK/STAT interface, and the
JAK/STAT signaling domain respectively. Using the 'Plot profile' function in Fiji, the
fluorescence intensity values of each reporter was obtained along these tracks and averaged.
Reporter patterns for each disc was graphed by scaling the averaged values from each track,
between 0 to 1 and plotted on the Y-axis. This was done independently for n≥3 discs, and a
representative image was selected for visualization of the spatial trends of the reporters. (Fig
1M,N)

Density plots for JNK/AP-1 and JAK/STAT reporters
For combined pouch and hinge regions (Fig. 1I,J), the ROI outlines were manually drawn on
the DAPI channel to include the pouch and hinge domains till the medial hinge fold. Partial
max projections of z-sections from the disc proper were generated, carefully excluding any
peripodial signal.

For time course experiments with *egr*-expressing discs (S2.H-K), a central z-section was
chosen and ROIs generated by thresholding the *TRE-RFP* channel to include only the JNK-
signaling pouch domain. After applying a 'Gaussian blur' filter (sigma=2), masks were

generated by applying a low fixed threshold (value = 10) based on early, 7h *egr*-expressing

discs. Noise removal was done using the 'Despeckle' function, followed by 'Fill Hole' function.

ROI outlines were generated using 'Create selection'. All ROIs were placed on the source

TRE>RFP and *Stat92E>dGFP* reporter channels. 'Save X-Y co-ordinates' function was used

to obtain pixel fluorescence intensity values for both reporters within the ROI, and the

distribution of fluorescence intensities across the samples were graphed by generating a

binned (bins = 16) 2D histogram plot using R. After all NA values were set to 0, data density

per bin for each disc was calculated as a percentage. Values were averaged across

corresponding bins for all discs within a timepoint to generate the average fluorescence

intensity distribution profile. To distinguish regions with low, medium and high signaling for

each pathway, the graphs were subdivided into sections s1-s9 by defining a threshold to

distinguish low, medium and high JNK/AP-1 and JAK/STAT signal intensities (Fig. S2K).

Percentage of datapoints within each section was determined from the averaged fluorescence

intensity distribution profile. Ordinary 2-way ANOVA with Dunnett's multiple comparisons test

was carried out for statistical analysis. Alpha was set at 0.05

Cell cycle profiles

Representative sections were selected per disc and a square ROI area ($75 \times 75 \mu m^2$) was placed

over the central pouch cells. Only viable cells stained with characteristic euchromatic and

heterochromatic DAPI staining were chosen. Using the 'Multi point' tool, cells showing a G1

profile (green), G2 profile (yellow+red) or neither (black) were counted. Percentage values for

the gap phases were calculated for n≥3 discs per genotype and visualized. T-tests were

performed on the calculated G1:G2 ratios per disc. (Fig 5G,H)

JAK/STAT reporter quantifications for testing RNAi lines

Crosses were setup as previously described and wing discs from larvae were dissected on D7

AED. Single sections at similar focal planes were carefully chosen during imaging. Using

drawing tools in Fiji, ROIs were generated to include JAK/STAT signaling cells in the hinge

within A and P compartments. The mean JAK/STAT reporter intensity within the ROI was

obtained from control and RNAi expressing discs using the 'Measure' function in Fiji. For each

dataset, paired t-tests were performed between A and P fluorescence intensity values for the

same genotype. (Fig. S4.3A-C)

Nuclear translocation of Stat92E-GFP

Images from the pouch (n=12) and hinge (n=9) domains of control discs and pouch (n=16) and

hinge (n=12) domains of *p35+egr* expressing discs were obtained to track the intra-cellular

localization of *Stat92E-GFP* within the nucleus and cytoplasm. Images were taken at high

magnification (63x). After thresholding ('Huang'), the selection tool was used to generate a

nuclear ROI using the DAPI channel. The inverse selection tool was used to generate a

corresponding cytoplasmic ROI for each image (Fig S4.1L). These nuclear and cytoplamic

ROIs were then placed on the *Stat92E-GFP* channel and mean fluorescence intensity values

were obtained for each subcellular fraction and unpaired t-tests were performed. (Fig. S4.1M)

MMP-1 intensity quantification

To measure the fluorescence intensity of MMP-1 in *egr*-expressing and *egr,zfh2-co*expressing

discs, maximum intensity projections of confocal sections were taken. Sections were carefully

chosen to capture JNK/AP-1 signaling activity in the disc proper, avoiding the peripodium and

cell debris. Square ROIs of fixed length (30 μm) were placed in 3 non-overlapping regions in

the pouch domain, and mean intensity was calculated using the 'Measure' function in Fiji. Mean

fluorescence intensities obtained from ROIs were averaged per disc and values from each disc

per genotype were used for statistical analysis. (Fig. S3A-C)

JNK$^+$ and JNK$^-$ Area and intensity quantifications (*Ras^{v12},scrib-RNAi* tumors)

A representative z-section was selected per disc. As our interest was in quantitating the spatial

segregation, only JNK/AP-1 or JAK/STAT signaling regions in the tumor were chosen. After

applying a 'Gaussian blur' filter (sigma=2), 'Moments' thresholding and 'Despeckle' function

for noise correction was applied to generate masks for regions showing *TRE>RFP* (JNK) and

Stat92E>dGFP (JAK/STAT) reporter activity. From the 'Image calculator' function, 'AND' and

'Subtract' operations on JNK/AP-1 and STAT masks were carried out to obtain the JNK$^+$STAT$^+$

mask, and the exclusive JNK$^+$ (JNK-STAT) or exclusive JNK$^-$(STAT-JNK) masks respectively.

Area of each mask was obtained using the 'Measure' function. Summed area of these masks

were considered as the total area and each area was then represented as a percentage of

total. One-way paired ANOVA with Holm-Šídák's multiple comparison test was performed to

test for statistical significance. Mean intensity for *TRE>RFP* or *10xStat92E>dGFP* reporters

were measured inside the JNK/AP-1 mask (JNK$^+$) and the exclusive JNK$^-$ masks. Paired t-

tests were performed to test for statistical significance. (Fig. 6E,F and S6I)

686
687
688

ph3 counts

Partial maximum intensity projection of three central z-sections were taken per disc. To
generate masks for regions showing *Stat92E>dGFP* reporter activity (STAT⁺), a 'Gaussian
blur' filter (sigma=1), 'Moments' threshold and 'Despeckle' function for noise correction was
applied. The DAPI channel was used to manually outline the pouch domain and generate
masks. Using the 'Image calculator' function, STAT⁻ mask was 'Subtract'-ed from the pouch
mask to generate the STAT⁻ mask. Areas of the STAT⁺ and STAT⁻ regions in the pouch were
obtained using the 'Measure' function. Then, the ph3 channel was duplicated and 'Li' threshold
was applied. 'Remove outlier' (radius=5 threshold=50) and 'Despeckle' function for noise
correction, followed by 'Watershed' function was applied to generate a ph3 mask. Using the
'AND' operator, the ph3 mask was overlayed with the STAT⁺ and STAT⁻ masks. '3D object
counter' function was then used to obtain ph3 'Object counts' within the STAT⁺ and STAT⁻
regions and divided by area of each ROI to get the ph3 counts per mm². Paired t-tests were
performed to test for statistical significance. (Fig. 6I)

Mathematical modelling

To test for the existence of a regulatory motif within our signaling network, a mathematical model was derived to describe the temporal dynamics of the concentration of specific molecules over a fixed 2D space. The system was modeled as a set of ordinary differential equations (ODEs) (see also Fig. 3C). Two sets of modeling equations were established, that reflected conditions which include or exclude repression of JAK/STAT by JNK. The term describing repression of JAK/STAT (referred to as 'JAK' in the modeling equations and hereafter) by JNK/AP-1 is highlighted in red.

Uni-directional repression model (see also Fig. 3B):

$$\frac{d\mathrm{EIG}(t,x)}{dt} = b_{EIG}(x) - k_{deg,EIG}\mathrm{EIG} + k_{act,EIG}\frac{\mathrm{JNK}^{n_1}}{K_{m,act,EIG}^{n_1} + \mathrm{JNK}^{n_1}} + D_{EIG}\nabla^2 EIG$$

$$\frac{d\mathrm{JNK}(t,x)}{dt} = b_{JNK} - k_{deg,JNK}\mathrm{JNK} + k_{act,JNK}\frac{\mathrm{EIG}^{n_2}}{(K_{m,act,JNK}^{n_2} + \mathrm{EIG}^{n_2})(1 + k_{inh,JNK,by,JAK}^{ni1}\mathrm{JAK}^{ni1})}$$

$$\frac{d\mathrm{JAK}(t,x)}{dt} = b_{JAK} - k_{deg,JAK}\mathrm{JAK} + k_{act,JAK}\mathrm{UPD}\frac{\mathrm{JAK}^{n_3}}{(K_{m,act,JAK}^{n_3} + \mathrm{JAK}^{n_3})}$$

$$\frac{d\mathrm{UPD}(t,x)}{dt} = b_{UPD} - k_{deg,UPD}\mathrm{UPD} + k_{act,UPD}\frac{\mathrm{JNK}^{n_4}}{K_{m,act,UPD}^{n_4} + \mathrm{JNK}^{n_4}} + D_{UPD}\nabla^2 UPD$$

Mutual repression model (see also Fig. 3B'):

$$\frac{d\mathrm{EIG}(t,x)}{dt} = b_{EIG}(x) - k_{deg,EIG}\mathrm{EIG} + k_{act,EIG}\frac{\mathrm{JNK}^{n_1}}{K_{m,act,EIG}^{n_1} + \mathrm{JNK}^{n_1}} + D_{EIG}\nabla^2 EIG$$

$$\frac{d\mathrm{JNK}(t,x)}{dt} = b_{JNK} - k_{deg,JNK}\mathrm{JNK} + k_{act,JNK}\frac{\mathrm{EIG}^{n_2}}{(K_{m,act,JNK}^{n_2} + \mathrm{EIG}^{n_2})(1 + k_{inh,JNK,by,JAK}^{ni1}\mathrm{JAK}^{ni1})}$$

$$\frac{d\mathrm{JAK}(t,x)}{dt} = b_{JAK} - k_{deg,JAK}\mathrm{JAK} + k_{act,JAK}\mathrm{UPD}\frac{\mathrm{JAK}^{n_3}}{(K_{m,act,JAK}^{n_3} + \mathrm{JAK}^{n_3})(1 + k_{inh,JAK,by,JNK}^{ni2}\mathrm{JNK}^{ni2})}$$

$$\frac{d\mathrm{UPD}(t,x)}{dt} = b_{UPD} - k_{deg,UPD}\mathrm{UPD} + k_{act,UPD}\frac{\mathrm{JNK}^{n_4}}{K_{m,act,UPD}^{n_4} + \mathrm{JNK}^{n_4}} + D_{UPD}\nabla^2 UPD$$

Characteristic units are represented by state (u), space (x) and time (t). The mathematical terms include, among others, basal rate of production (*b*), activation rate (k_{act}), linear degradation rate (k_{deg}), Michaelis–Menten constant (K_m), inhibition constant (k_{inh}), diffusion co-efficients (D_{EIG}, D_{UPD}), Hill co-efficients to model cooperativity ($n_1, n_2, n_3, n_4, ni_1, ni_2$) and Hill kinetics to describe self-amplification and mutual repression.

The first equation describes the dynamics of the concentration of Egr, produced at a basal rate dependent on the spatial co-ordinate x and degraded at a linear rate. Hill kinetics describe the production of Egr induced by JNK/AP-1 and its diffusion away from the source. The second equation describes the dynamics of the concentration of the JNK/AP-1 TF, produced at a basal rate and degraded linearly. Hill kinetics describe production of JNK/AP-1 induced by *egr* as well as inhibition of JNK/AP-1 by JAK [51]. The third equation describes the dynamics of the

concentration of the JAK TF, produced at a basal rate and degraded linearly. Hill kinetics

describe positive feedback dependent on Upd. Inhibition of JAK by JNK/AP-1 is described

and the model is tested with and without this mathematical term (highlighted in red) to check if

this interaction exists within the system. The fourth equation describes the dynamics of the

concentration of Upd, produced at a basal rate and degraded linearly. Hill kinetics describe the

production of Upd induced by JNK/AP-1 and its diffusion [28, 48, 51].

Technically, the modeling was done based on a non-dimensionalized version of the model,

rescaled to reduce the number of free parameters. The system was then defined as follows:

$$
\begin{aligned}
\frac{d\mathrm{EIG}(t,x)}{dt} &= \gamma\left(1(x) - \mathrm{EIG} + k_{act,EIG}\frac{\mathrm{JNK}^{n_1}}{K_{m,act,EIG}^{n_1} + \mathrm{JNK}^{n_1}}\right) + \nabla^2 \mathrm{EIG} \\[2mm]
\frac{d\mathrm{JNK}(t,x)}{dt} &= \gamma\left(1 - k_{deg,JNK}\mathrm{JNK} + k_{act,JNK}\frac{\mathrm{EIG}^{n_2}}{(K_{m,act,JNK}^{n_2} + \mathrm{EIG}^{n_2})(1 + k_{inh,JNK,by,JAK}^{ni1}\mathrm{JAK}^{ni1})}\right) \\[2mm]
\frac{d\mathrm{JAK}(t,x)}{dt} &= \gamma\left(1 - k_{deg,JAK}\mathrm{JAK} + k_{act,JAK}\mathrm{UPD}\frac{\mathrm{JAK}^{n_3}}{(K_{m,act,JAK}^{n_3} + \mathrm{JAK}^{n_3})(1 + k_{inh,JAK,by,JNK}^{ni2}\mathrm{JNK}^{ni2})}\right) \\[2mm]
\frac{d\mathrm{UPD}(t,x)}{dt} &= \gamma\left(1 - k_{deg,UPD}\mathrm{UPD} + k_{act,UPD}\frac{\mathrm{JNK}^{n_4}}{K_{m,act,UPD}^{n_4} + \mathrm{JNK}^{n_4}}\right) + d\nabla^2 \mathrm{UPD}
\end{aligned}
$$

743 For each dimensionless parameter, the parameter dependencies and the scanned parameter

744 range are represented in the table below:

Dimensionless parameter	Functional relation to systems parameter	Scanned parameter range
γ	$k_{deg,EIG}\, L^2 / D_{EIG}$	$0.01 \ldots 10000$
$\hat{k}_{deg,JNK}$	$k_{deg,JNK} / k_{deg,EIG}$	$0.01 \ldots 10000$
$\hat{k}_{deg,JAK}$	$k_{deg,JAK} / k_{deg,EIG}$	$0.01 \ldots 10000$
$\hat{k}_{deg,Upd}$	$k_{deg,Upd} / k_{deg,EIG}$	$0.01 \ldots 10000$
$\hat{k}_{act,EIG}$	$k_{act,EIG} / b_{EIG}$	$0.00001 \ldots 10$
$\hat{k}_{act,JNK}$	$k_{act,JNK} / b_{JNK}$	$0.01 \ldots 10000$
$\hat{k}_{act,JAK}$	$k_{act,JAK}\, D_{EIG} / L^2 / b_{UPD} / b_{JAK}$	$0.01 \ldots 10000$
$\hat{k}_{act,UPD}$	$k_{act,UPD} / b_{UPD}$	$0.01 \ldots 10000$
$\hat{K}_{m,act,EIG}$	$K_{m,act,EIG}\, D_{EIG} / L^2 / b_{EIG}$	$0.00001 \ldots 10$
$\hat{K}_{m,act,JNK}$	$K_{m,act,JNK}\, D_{EIG} / L^2 / b_{JNK}$	$0.01 \ldots 10000$
$\hat{K}_{m,act,JAK}$	$K_{m,act,JAK}\, D_{EIG} / L^2 / b_{JAK}$	$0.01 \ldots 10000$
$\hat{K}_{m,act,UPD}$	$K_{m,act,UPD}\, D_{EIG} / L^2 / b_{UPD}$	$0.01 \ldots 10000$
$\hat{k}_{inh,JAK,by,JNK}$	$k_{inh,JAK,by,JNK}\, b_{JAK}\, L^2 / D_{EIG}$	$0.01 \ldots 10000$
$\hat{k}_{inh,JNK,by,JAK}$	$k_{inh,JNK,by,JAK}\, b_{JNK}\, L^2 / D_{EIG}$	$0.01 \ldots 10000$
$\hat{n}_i$	n_i	$1 \ldots 4$
d	D_{UPD} / D_{EIG}	$1.5 \ldots 4$

We performed a global analysis by sampling approach introduced in *Rausenberger et. al* [135].
The models were simulated on a 1D spatial domain x = [0 1]. To represent the wound site, a
high basal production of Egr was fixed within a finite space at x = [0 0.05], while basal rate of
production of Egr for the rest of the region was set to zero. The rate of production of Upd was
defined as a function of JNK/AP-1 activity [18, 20, 39, 51]. For all states, initial condition was
set as u (t=0,x) = 0.1. Solutions for each model (with and without repression) were simulated
using 10^6 parameter sets, drawn uniformly from an interval from 10^{-3} to 10^{+2} on a logarithmic
scale, and tested under different conditions of diffusion – $D_{Upd} > D_{Egr}$, $D_{Upd} < D_{Egr}$ and $D_{Upd} =$
D_{Egr}. The resulting solutions were automatically scored for countergradient endpoint features
extracted from the experimentally observed gradient patterns.
758
759 Using this approach [135], positive solutions for a generic bistable pattern from the scanned
760 parameter range were automatically extracted based on the following features:
761
762 i. JAK at x = 0 lower than JAK at x = 1
763 ii. JNK/AP-1 at x = 0 higher than JNK/AP-1 at x = 1
764 iii. Relative difference of JAK higher than 10%
765 iv. Relative difference of JNK/AP-1 higher than 10%
766
767 Positive solutions for a specific bistable pattern which mimics biologically observed patterns
768 were automatically extracted based on the additional features:
769 v. JAK has a local maximum which is not at the boundary of the simulated domain
770 vi. Height of local JAK maximum at least 5%
771
772
773

Author Contributions

Conceptualization JJ, VW, AKC

Validation JJ, AA, IG, AKC

Investigation JJ, RE, AA, CC, AKC

Writing JJ, VW, RE, JT, AKC

Visualization JJ, RE, VW, AKC

Supervision JT, AKC

Funding Acquisition AKC

Acknowledgements

We thank the LIC facility (University of Freiburg) for technical help with imaging. We thank Erika Bach, David Bilder, Dirk Bohmann, Stephen Cohen, Fernando Diaz-Benjumea, Iswar Hariharan, Norbert Perrimon, Helena Richardson, Reinhardt Schuh, Y. Henry Sun, Mirka Uhlirova and Pelin Volkan for sharing reagents and the Bloomington Drosophila Stock Center (BDSC), the Vienna Drosophila Stock Collection (VDRC), the University of Zurich ORFeome Project (FlyORF), the Developmental Studies Hybridoma Bank (DSHB) and the Monoclonal Antibody Core Facility at the Helmholtz Zentrum Munich for providing fly stocks and antibodies. Thanks to Florian Steinberg (University of Freiburg) for support with protocol, reagents and imaging equipment for Western blot assays. We also thank the SGBM graduate school for support.

Funding

Funding for this work was provided by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy (CIBSS – EXC-2189 – Project ID 390939984), the Emmy Noether Programme (CL490/1-1), the Heisenberg Program (CL490/3-1) and by the Boehringer Ingelheim Foundation (Plus3 Programme).

References

1.   Shaw TJ, Martin P. Wound repair at a glance. J Cell Sci. 2009;122(Pt 18):3209-13. Epub 2009/09/04. doi: 10.1242/jcs.031187. PubMed PMID: 19726630; PubMed Central PMCID: PMCPMC2736861.

2.   Marchiando AM, Graham WV, Turner JR. Epithelial barriers in homeostasis and disease. Annu Rev Pathol. 2010;5:119-44. doi: 10.1146/annurev.pathol.4.110807.092135. PubMed PMID: 20078218.

3.   Eming SA, Martin P, Tomic-Canic M. Wound repair and regeneration: mechanisms, signaling, and translation. Sci Transl Med. 2014;6(265):265sr6. Epub 2014/12/05. doi: 10.1126/scitranslmed.3009337. PubMed PMID: 25473038; PubMed Central PMCID: PMCPMC4973620.

4.   Neves J, Demaria M, Campisi J, Jasper H. Of flies, mice, and men: evolutionarily conserved tissue damage responses and aging. Dev Cell. 2015;32(1):9-18. Epub 2015/01/15. doi: 10.1016/j.devcel.2014.11.028. PubMed PMID: 25584795; PubMed Central PMCID: PMCPMC4450349.

5.   Niethammer P. The early wound signals. Curr Opin Genet Dev. 2016;40:17-22. Epub 20160605. doi: 10.1016/j.gde.2016.05.001. PubMed PMID: 27266971; PubMed Central PMCID: PMCPMC5278878.

6.   Hotamisligil GS, Davis RJ. Cell Signaling and Stress Responses. Cold Spring Harb Perspect Biol. 2016;8(10). Epub 2016/10/05. doi: 10.1101/cshperspect.a006072. PubMed PMID: 27698029.

7.   Ayello EA, Dowsett C, Schultz GS, Sibbald RG, Falanga V, Harding K, et al. TIME heals all wounds. Nursing. 2004;34(4):36-41; quiz, -2. Epub 2004/07/13. doi: 10.1097/00152193-200404000-00040. PubMed PMID: 15247672.

8.   Schafer M, Werner S. Oxidative stress in normal and impaired wound repair. Pharmacol Res. 2008;58(2):165-71. Epub 20080619. doi: 10.1016/j.phrs.2008.06.004. PubMed PMID: 18617006.

9.   Wilkinson HN, Hardman MJ. Wound healing: cellular mechanisms and pathological outcomes. Open Biol. 2020;10(9):200223. Epub 20200930. doi: 10.1098/rsob.200223. PubMed PMID: 32993416; PubMed Central PMCID: PMCPMC7536089.

10.   Martin P, Nunan R. Cellular and molecular mechanisms of repair in acute and chronic wound healing. Br J Dermatol. 2015;173(2):370-8. Epub 20150714. doi: 10.1111/bjd.13954. PubMed PMID: 26175283; PubMed Central PMCID: PMCPMC4671308.

11.   Dvorak HF. Tumors: wounds that do not heal. Similarities between tumor stroma generation
and wound healing. N Engl J Med. 1986;315(26):1650-9. Epub 1986/12/25. doi:
10.1056/NEJM198612253152606. PubMed PMID: 3537791.

12.   Coussens LM, Werb Z. Inflammation and cancer. Nature. 2002;420(6917):860-7. Epub
2002/12/20. doi: 10.1038/nature01322. PubMed PMID: 12490959; PubMed Central PMCID:
PMCPMC2803035.

13.   Rybinski B, Franco-Barraza J, Cukierman E. The wound healing, chronic fibrosis, and cancer
progression triad. Physiol Genomics. 2014;46(7):223-44. Epub 2014/02/13. doi:
10.1152/physiolgenomics.00158.2013. PubMed PMID: 24520152; PubMed Central PMCID:
PMCPMC4035661.

14.   Lee JH, Lee CW, Park SH, Choe KM. Spatiotemporal regulation of cell fusion by JNK and
JAK/STAT signaling during Drosophila wound healing. J Cell Sci. 2017;130(11):1917-28. Epub
20170419. doi: 10.1242/jcs.187658. PubMed PMID: 28424232.

15.   Purice MD, Ray A, Munzel EJ, Pope BJ, Park DJ, Speese SD, et al. A novel Drosophila injury
model reveals severed axons are cleared through a Draper/MMP-1 signaling cascade. Elife. 2017;6.
Epub 2017/08/22. doi: 10.7554/eLife.23611. PubMed PMID: 28825401; PubMed Central PMCID:
PMCPMC5565368.

16.   Ahmed-de-Prado S, Diaz-Garcia S, Baonza A. JNK and JAK/STAT signalling are required for
inducing loss of cell fate specification during imaginal wing discs regeneration in Drosophila
melanogaster. Dev Biol. 2018. Epub 2018/06/06. doi: 10.1016/j.ydbio.2018.05.021. PubMed PMID:
29870691.

17.   Worley MI, Alexander LA, Hariharan IK. CtBP impedes JNK- and Upd/STAT-driven cell fate
misspecifications in regenerating Drosophila imaginal discs. Elife. 2018;7. Epub 2018/01/27. doi:
10.7554/eLife.30391. PubMed PMID: 29372681.

18.   Wu M, Pastor-Pareja JC, Xu T. Interaction between Ras(V12) and scribbled clones induces
tumour growth and invasion. Nature. 2010;463(7280):545-8. Epub 2010/01/15. doi:
10.1038/nature08702. PubMed PMID: 20072127; PubMed Central PMCID: PMCPMC2835536.

19.   Woodfield SE, Graves HK, Hernandez JA, Bergmann A. De-regulation of JNK and JAK/STAT
signaling in ESCRT-II mutant tissues cooperatively contributes to neoplastic tumorigenesis. PLoS
One. 2013;8(2):e56021. doi: 10.1371/journal.pone.0056021. PubMed PMID: 23418496; PubMed
Central PMCID: PMCPMC3572140.

20. Bunker BD, Nellimoottil TT, Boileau RM, Classen AK, Bilder D. The transcriptional response to
tumorigenic polarity loss in Drosophila. Elife. 2015;4. Epub 20150226. doi: 10.7554/eLife.03189.
PubMed PMID: 25719210; PubMed Central PMCID: PMCPMC4369581.

21. Atkins M, Potier D, Romanelli L, Jacobs J, Mach J, Hamaratoglu F, et al. An Ectopic Network
of Transcription Factors Regulated by Hippo Signaling Drives Growth and Invasion of a Malignant
Tumor Model. Curr Biol. 2016;26(16):2101-13. Epub 2016/08/02. doi: 10.1016/j.cub.2016.06.035.
PubMed PMID: 27476594.

22. Yang SA, Portilla JM, Mihailovic S, Huang YC, Deng WM. Oncogenic Notch Triggers
Neoplastic Tumorigenesis in a Transition-Zone-like Tissue Microenvironment. Dev Cell.
2019;49(3):461-72 e5. Epub 2019/04/16. doi: 10.1016/j.devcel.2019.03.015. PubMed PMID:
30982664; PubMed Central PMCID: PMCPMC6504601.

23. Hamaratoglu F, Atkins M. Rounding up the Usual Suspects: Assessing Yorkie, AP-1, and Stat
Coactivation in Tumorigenesis. Int J Mol Sci. 2020;21(13). Epub 20200627. doi:
10.3390/ijms21134580. PubMed PMID: 32605129; PubMed Central PMCID: PMCPMC7370090.

24. Ramet M, Lanot R, Zachary D, Manfruelli P. JNK signaling pathway is required for efficient
wound healing in Drosophila. Dev Biol. 2002;241(1):145-56. Epub 2002/01/11. doi:
10.1006/dbio.2001.0502. PubMed PMID: 11784101.

25. Bosch M, Serras F, Martin-Blanco E, Baguna J. JNK signaling pathway required for wound
healing in regenerating Drosophila wing imaginal discs. Dev Biol. 2005;280(1):73-86. Epub
2005/03/16. doi: 10.1016/j.ydbio.2005.01.002. PubMed PMID: 15766749.

26. Mattila J, Omelyanchuk L, Kyttala S, Turunen H, Nokkala S. Role of Jun N-terminal Kinase
(JNK) signaling in the wound healing and regeneration of a Drosophila melanogaster wing imaginal
disc. Int J Dev Biol. 2005;49(4):391-9. Epub 2005/06/22. doi: 10.1387/ijdb.052006jm. PubMed PMID:
15968584.

27. Finkel T. Signal transduction by reactive oxygen species. J Cell Biol. 2011;194(1):7-15. Epub
2011/07/13. doi: 10.1083/jcb.201102095. PubMed PMID: 21746850; PubMed Central PMCID:
PMCPMC3135394.

28. Santabarbara-Ruiz P, Lopez-Santillan M, Martinez-Rodriguez I, Binagui-Casas A, Perez L,
Milan M, et al. ROS-Induced JNK and p38 Signaling Is Required for Unpaired Cytokine Activation
during Drosophila Regeneration. PLoS Genet. 2015;11(10):e1005595. Epub 20151023. doi:
10.1371/journal.pgen.1005595. PubMed PMID: 26496642; PubMed Central PMCID:
PMCPMC4619769.

29.   Moreno E, Yan M, Basler K. Evolution of TNF signaling mechanisms: JNK-dependent
apoptosis triggered by Eiger, the Drosophila homolog of the TNF superfamily. Curr Biol.
2002;12(14):1263-8. Epub 2002/08/15. PubMed PMID: 12176339.

30.   Kanda H, Miura M. Regulatory roles of JNK in programmed cell death. J Biochem.
2004;136(1):1-6. Epub 2004/07/23. doi: 10.1093/jb/mvh098. PubMed PMID: 15269233.

31.   Cosolo A, Jaiswal J, Csordas G, Grass I, Uhlirova M, Classen AK. JNK-dependent cell cycle
stalling in G2 promotes survival and senescence-like phenotypes in tissue stress. Elife. 2019;8. Epub
20190208. doi: 10.7554/eLife.41036. PubMed PMID: 30735120; PubMed Central PMCID:
PMCPMC6389326.

32.   Ryoo HD, Gorenc T, Steller H. Apoptotic cells can induce compensatory cell proliferation
through the JNK and the Wingless signaling pathways. Dev Cell. 2004;7(4):491-501. Epub
2004/10/08. doi: 10.1016/j.devcel.2004.08.019. PubMed PMID: 15469838.

33.   Morata G, Shlevkov E, Perez-Garijo A. Mitogenic signaling from apoptotic cells in Drosophila.
Dev Growth Differ. 2011;53(2):168-76. Epub 2011/02/23. doi: 10.1111/j.1440-169X.2010.01225.x.
PubMed PMID: 21338343.

34.   Tournier C. The 2 Faces of JNK Signaling in Cancer. Genes Cancer. 2013;4(9-10):397-400.
Epub 2013/12/19. doi: 10.1177/1947601913486349. PubMed PMID: 24349637; PubMed Central
PMCID: PMCPMC3863340.

35.   Pinal N, Calleja M, Morata G. Pro-apoptotic and pro-proliferation functions of the JNK pathway
of Drosophila: roles in cell competition, tumorigenesis and regeneration. Open Biol. 2019;9(3):180256.
Epub 2019/03/07. doi: 10.1098/rsob.180256. PubMed PMID: 30836847; PubMed Central PMCID:
PMCPMC6451367.

36.   La Marca JE, Richardson HE. Two-Faced: Roles of JNK Signalling During Tumourigenesis in
the Drosophila Model. Front Cell Dev Biol. 2020;8:42. Epub 2020/03/03. doi:
10.3389/fcell.2020.00042. PubMed PMID: 32117973; PubMed Central PMCID: PMCPMC7012784.

37.   Ruiz-Losada M, Gonzalez R, Peropadre A, Gil-Galvez A, Tena JJ, Baonza A, et al.
Coordination between cell proliferation and apoptosis after DNA damage in Drosophila. Cell Death
Differ. 2021. Epub 20211125. doi: 10.1038/s41418-021-00898-6. PubMed PMID: 34824391.

38.   Smith-Bolton RK, Worley MI, Kanda H, Hariharan IK. Regenerative growth in Drosophila
imaginal discs is regulated by Wingless and Myc. Dev Cell. 2009;16(6):797-809. doi:
10.1016/j.devcel.2009.04.015. PubMed PMID: 19531351; PubMed Central PMCID:
PMCPMC2705171.

39.   Jiang H, Patel PH, Kohlmaier A, Grenley MO, McEwen DG, Edgar BA. Cytokine/Jak/Stat
signaling mediates regeneration and homeostasis in the Drosophila midgut. Cell. 2009;137(7):1343-
55. Epub 2009/07/01. doi: 10.1016/j.cell.2009.05.014. PubMed PMID: 19563763; PubMed Central
PMCID: PMCPMC2753793.

40.   Bergantinos C, Corominas M, Serras F. Cell death-induced regeneration in wing imaginal
discs requires JNK signalling. Development. 2010;137(7):1169-79. doi: 10.1242/dev.045559. PubMed
PMID: 20215351.

41.   Harrison DA, McCoon PE, Binari R, Gilman M, Perrimon N. Drosophila unpaired encodes a
secreted protein that activates the JAK signaling pathway. Genes Dev. 1998;12(20):3252-63. Epub
1998/10/24. doi: 10.1101/gad.12.20.3252. PubMed PMID: 9784499; PubMed Central PMCID:
PMCPMC317220.

42.   Hombria JC, Brown S, Hader S, Zeidler MP. Characterisation of Upd2, a Drosophila
JAK/STAT pathway ligand. Dev Biol. 2005;288(2):420-33. Epub 2005/11/10. doi:
10.1016/j.ydbio.2005.09.040. PubMed PMID: 16277982.

43.   Agaisse H, Petersen UM, Boutros M, Mathey-Prevot B, Perrimon N. Signaling role of
hemocytes in Drosophila JAK/STAT-dependent response to septic injury. Dev Cell. 2003;5(3):441-50.
Epub 2003/09/12. doi: 10.1016/s1534-5807(03)00244-2. PubMed PMID: 12967563.

44.   Gilbert MM, Weaver BK, Gergen JP, Reich NC. A novel functional activator of the Drosophila
JAK/STAT pathway, unpaired2, is revealed by an in vivo reporter of pathway activation. Mech Dev.
2005;122(7-8):939-48. Epub 2005/06/01. doi: 10.1016/j.mod.2005.03.004. PubMed PMID: 15925495.

45.   Toggweiler J, Willecke M, Basler K. The transcription factor Ets21C drives tumor growth by
cooperating with AP-1. Sci Rep. 2016;6:34725. Epub 2016/10/08. doi: 10.1038/srep34725. PubMed
PMID: 27713480; PubMed Central PMCID: PMCPMC5054425.

46.   Classen AK, Bunker BD, Harvey KF, Vaccari T, Bilder D. A tumor suppressor activity of
Drosophila Polycomb genes mediated by JAK-STAT signaling. Nat Genet. 2009;41(10):1150-5. Epub
20090913. doi: 10.1038/ng.445. PubMed PMID: 19749759; PubMed Central PMCID:
PMCPMC2782793.

47.   Davie K, Jacobs J, Atkins M, Potier D, Christiaens V, Halder G, et al. Discovery of
transcription factors and regulatory regions driving in vivo tumor development by ATAC-seq and
FAIRE-seq open chromatin profiling. PLoS Genet. 2015;11(2):e1004994. doi:
10.1371/journal.pgen.1004994. PubMed PMID: 25679813; PubMed Central PMCID:
PMCPMC4334524.

48.   Katsuyama T, Comoglio F, Seimiya M, Cabuy E, Paro R. During Drosophila disc regeneration,
JAK/STAT coordinates cell proliferation with Dilp8-mediated developmental delay. Proc Natl Acad Sci
U S A. 2015;112(18):E2327-36. Epub 20150420. doi: 10.1073/pnas.1423074112. PubMed PMID:
25902518; PubMed Central PMCID: PMCPMC4426433.

49.   Pastor-Pareja JC, Wu M, Xu T. An innate immune response of blood cells to tumors and
tissue damage in Drosophila. Dis Model Mech. 2008;1(2-3):144-54; discussion 53. Epub 20080918.
doi: 10.1242/dmm.000950. PubMed PMID: 19048077; PubMed Central PMCID: PMCPMC2562178.

50.   Lin G, Xu N, Xi R. Paracrine Unpaired Signaling through the JAK/STAT Pathway Controls
Self-renewal and Lineage Differentiation of Drosophila Intestinal Stem Cells. J Mol Cell Biol. 2009.
Epub 2009/10/03. doi: mjp028 [pii]

10.1093/jmcb/mjp028. PubMed PMID: 19797317.

51.   La Fortezza M, Schenk M, Cosolo A, Kolybaba A, Grass I, Classen AK. JAK/STAT signalling
mediates cell survival in response to tissue stress. Development. 2016;143(16):2907-19. Epub
20160706. doi: 10.1242/dev.132340. PubMed PMID: 27385008.

52.   Betz A, Ryoo HD, Steller H, Darnell JE, Jr. STAT92E is a positive regulator of Drosophila
inhibitor of apoptosis 1 (DIAP/1) and protects against radiation-induced apoptosis. Proc Natl Acad Sci
U S A. 2008;105(37):13805-10. Epub 20080908. doi: 10.1073/pnas.0806291105. PubMed PMID:
18779571; PubMed Central PMCID: PMCPMC2544535.

53.   Kauppila S, Maaty WS, Chen P, Tomar RS, Eby MT, Chapo J, et al. Eiger and its receptor,
Wengen, comprise a TNF-like system in Drosophila. Oncogene. 2003;22(31):4860-7. Epub
2003/08/02. doi: 10.1038/sj.onc.1206715. PubMed PMID: 12894227.

54.   Muzzopappa M, Murcia L, Milan M. Feedback amplification loop drives malignant growth in
epithelial tissues. Proc Natl Acad Sci U S A. 2017;114(35):E7291-E300. Epub 20170814. doi:
10.1073/pnas.1701791114. PubMed PMID: 28808034; PubMed Central PMCID: PMCPMC5584413.

55.   Khan SJ, Abidi SNF, Skinner A, Tian Y, Smith-Bolton RK. The Drosophila Duox maturation
factor is a key component of a positive feedback loop that sustains regeneration signaling. PLoS
Genet. 2017;13(7):e1006937. Epub 2017/07/29. doi: 10.1371/journal.pgen.1006937. PubMed PMID:
28753614; PubMed Central PMCID: PMCPMC5550008.

56.   Pinal N, Martin M, Medina I, Morata G. Short-term activation of the Jun N-terminal kinase
pathway in apoptosis-deficient cells of Drosophila induces tumorigenesis. Nat Commun.
2018;9(1):1541. Epub 2018/04/20. doi: 10.1038/s41467-018-04000-6. PubMed PMID: 29670104;
PubMed Central PMCID: PMCPMC5906466.

57.   Fox DT, Cohen E, Smith-Bolton R. Model systems for regeneration: Drosophila. Development.
2020;147(7). Epub 20200406. doi: 10.1242/dev.173781. PubMed PMID: 32253254; PubMed Central
PMCID: PMCPMC7157589.

58.   Chatterjee N, Bohmann D. A versatile PhiC31 based reporter system for measuring AP-1 and
Nrf2 signaling in Drosophila and in tissue culture. PLoS One. 2012;7(4):e34063. Epub 2012/04/18.
doi: 10.1371/journal.pone.0034063. PubMed PMID: 22509270; PubMed Central PMCID:
PMCPMC3324472.

59.   Hariharan IK, Serras F. Imaginal disc regeneration takes flight. Curr Opin Cell Biol.
2017;48:10-6. Epub 20170401. doi: 10.1016/j.ceb.2017.03.005. PubMed PMID: 28376317; PubMed
Central PMCID: PMCPMC5591769.

60.   Trengove NJ, Stacey MC, MacAuley S, Bennett N, Gibson J, Burslem F, et al. Analysis of the
acute and chronic wound environments: the role of proteases and their inhibitors. Wound Repair
Regen. 1999;7(6):442-52. Epub 2000/01/13. doi: 10.1046/j.1524-475x.1999.00442.x. PubMed PMID:
10633003.

61.   Yager DR, Nwomeh BC. The proteolytic environment of chronic wounds. Wound Repair
Regen. 1999;7(6):433-41. Epub 2000/01/13. doi: 10.1046/j.1524-475x.1999.00433.x. PubMed PMID:
10633002.

62.   Caley MP, Martins VL, O'Toole EA. Metalloproteinases and Wound Healing. Adv Wound Care
(New Rochelle). 2015;4(4):225-34. Epub 2015/05/07. doi: 10.1089/wound.2014.0581. PubMed PMID:
25945285; PubMed Central PMCID: PMCPMC4397992.

63.   Moseley R, Hilton JR, Waddington RJ, Harding KG, Stephens P, Thomas DW. Comparison of
oxidative stress biomarker profiles between acute and chronic wound environments. Wound Repair
Regen. 2004;12(4):419-29. Epub 2004/07/21. doi: 10.1111/j.1067-1927.2004.12406.x. PubMed PMID:
15260807.

64.   Chipurupalli S, Samavedam U, Robinson N. Crosstalk Between ER Stress, Autophagy and
Inflammation. Front Med (Lausanne). 2021;8:758311. Epub 20211105. doi:
10.3389/fmed.2021.758311. PubMed PMID: 34805224; PubMed Central PMCID: PMCPMC8602556.

65.   Wu C, Chen C, Dai J, Zhang F, Chen Y, Li W, et al. Toll pathway modulates TNF-induced
JNK-dependent cell death in Drosophila. Open Biol. 2015;5(7):140171. Epub 2015/07/24. doi:
10.1098/rsob.140171. PubMed PMID: 26202785; PubMed Central PMCID: PMCPMC4632500.

66.   Zielke N, Korzelius J, van Straaten M, Bender K, Schuhknecht GFP, Dutta D, et al. Fly-FUCCI:
A versatile tool for studying cell proliferation in complex tissues. Cell Rep. 2014;7(2):588-98. Epub
2014/04/15. doi: 10.1016/j.celrep.2014.03.020. PubMed PMID: 24726363.

67.   Wright VM, Vogt KL, Smythe E, Zeidler MP. Differential activities of the Drosophila JAK/STAT
pathway ligands Upd, Upd2 and Upd3. Cell Signal. 2011;23(5):920-7. Epub 20110122. doi:
10.1016/j.cellsig.2011.01.020. PubMed PMID: 21262354.

68.   Osman D, Buchon N, Chakrabarti S, Huang YT, Su WC, Poidevin M, et al. Autocrine and
paracrine unpaired signaling regulate intestinal stem cell maintenance and division. J Cell Sci.
2012;125(Pt 24):5944-9. Epub 20121004. doi: 10.1242/jcs.113100. PubMed PMID: 23038775.

69.   Jiang H, Grenley MO, Bravo MJ, Blumhagen RZ, Edgar BA. EGFR/Ras/MAPK signaling
mediates adult midgut epithelial homeostasis and regeneration in Drosophila. Cell Stem Cell.
2011;8(1):84-95. Epub 20101216. doi: 10.1016/j.stem.2010.11.026. PubMed PMID: 21167805;
PubMed Central PMCID: PMCPMC3021119.

70.   Bach EA, Ekas LA, Ayala-Camargo A, Flaherty MS, Lee H, Perrimon N, et al. GFP reporters
detect the activation of the Drosophila JAK/STAT pathway in vivo. Gene Expr Patterns. 2007;7(3):323-
31. Epub 20060822. doi: 10.1016/j.modgep.2006.08.003. PubMed PMID: 17008134.

71.   Rodrigues AB, Zoranovic T, Ayala-Camargo A, Grewal S, Reyes-Robles T, Krasny M, et al.
Activated STAT regulates growth and induces competitive interactions independently of Myc, Yorkie,
Wingless and ribosome biogenesis. Development. 2012;139(21):4051-61. Epub 20120919. doi:
10.1242/dev.076760. PubMed PMID: 22992954; PubMed Central PMCID: PMCPMC3472591.

72.   Hatini V, Kula-Eversole E, Nusinow D, Del Signore SJ. Essential roles for stat92E in
expanding and patterning the proximodistal axis of the Drosophila wing imaginal disc. Dev Biol.
2013;378(1):38-50. Epub 20130307. doi: 10.1016/j.ydbio.2013.02.016. PubMed PMID: 23499656;
PubMed Central PMCID: PMCPMC3776456.

73.   Recasens-Alvarez C, Ferreira A, Milan M. JAK/STAT controls organ size and fate specification
by regulating morphogen production and signalling. Nat Commun. 2017;8:13815. Epub 20170103. doi:
10.1038/ncomms13815. PubMed PMID: 28045022; PubMed Central PMCID: PMCPMC5216089.

74.   Colombani J, Andersen DS, Leopold P. Secreted peptide Dilp8 coordinates Drosophila tissue
growth with developmental timing. Science. 2012;336(6081):582-5. doi: 10.1126/science.1216689.
PubMed PMID: 22556251.

75.   Hay BA, Wolff T, Rubin GM. Expression of baculovirus P35 prevents cell death in Drosophila.
Development. 1994;120(8):2121-9. Epub 1994/08/01. PubMed PMID: 7925015.

76.   Weber U, Paricio N, Mlodzik M. Jun mediates Frizzled-induced R3/R4 cell fate distinction and
planar polarity determination in the Drosophila eye. Development. 2000;127(16):3619-29. Epub
2000/07/21. doi: 10.1242/dev.127.16.3619. PubMed PMID: 10903185.

77.   Ayala-Camargo A, Anderson AM, Amoyel M, Rodrigues AB, Flaherty MS, Bach EA.
JAK/STAT signaling is required for hinge growth and patterning in the Drosophila wing disc. Dev Biol.
2013;382(2):413-26. Epub 20130823. doi: 10.1016/j.ydbio.2013.08.016. PubMed PMID: 23978534;
PubMed Central PMCID: PMCPMC3795806.

78.   Stevens LJ, Page-McCaw A. A secreted MMP is required for reepithelialization during wound
healing. Mol Biol Cell. 2012;23(6):1068-79. Epub 20120119. doi: 10.1091/mbc.E11-09-0745. PubMed
PMID: 22262460; PubMed Central PMCID: PMCPMC3302734.

79.   Cheong R, Rhee A, Wang CJ, Nemenman I, Levchenko A. Information transduction capacity
of noisy biochemical signaling networks. Science. 2011;334(6054):354-8. Epub 20110915. doi:
10.1126/science.1204553. PubMed PMID: 21921160; PubMed Central PMCID: PMCPMC3895446.

80.   Balazsi G, van Oudenaarden A, Collins JJ. Cellular decision making and biological noise: from
microbes to mammals. Cell. 2011;144(6):910-25. Epub 2011/03/19. doi: 10.1016/j.cell.2011.01.030.
PubMed PMID: 21414483; PubMed Central PMCID: PMCPMC3068611.

81.   Khan I, Ponnusamy S, Rahman SU, Giubellinom A, Arany PR. Co-opting Developmental
Signaling Pathways to Promote Wound Healing. Melvin A. Shiffman, Low M, editors: Springer, Cham;
2018. pp 103-14 p.

82.   Ghilardi SJ, O'Reilly BM, Sgro AE. Intracellular signaling dynamics and their role in
coordinating tissue repair. Wiley Interdiscip Rev Syst Biol Med. 2020;12(3):e1479. Epub 20200208.
doi: 10.1002/wsbm.1479. PubMed PMID: 32035001; PubMed Central PMCID: PMCPMC7187325.

83.   Jordan JD, Landau EM, Iyengar R. Signaling networks: the origins of cellular multitasking.
Cell. 2000;103(2):193-200. Epub 2000/11/01. doi: 10.1016/s0092-8674(00)00112-4. PubMed PMID:
11057893; PubMed Central PMCID: PMCPMC3619409.

84.   Milo R, Shen-Orr S, Itzkovitz S, Kashtan N, Chklovskii D, Alon U. Network motifs: simple
building blocks of complex networks. Science. 2002;298(5594):824-7. Epub 2002/10/26. doi:
10.1126/science.298.5594.824. PubMed PMID: 12399590.

85.   Levine M, Davidson EH. Gene regulatory networks for development. Proc Natl Acad Sci U S
A. 2005;102(14):4936-42. Epub 20050323. doi: 10.1073/pnas.0408031102. PubMed PMID:
15788537; PubMed Central PMCID: PMCPMC555974.

86.   Azeloglu EU, Iyengar R. Signaling networks: information flow, computation, and decision
making. Cold Spring Harb Perspect Biol. 2015;7(4):a005934. Epub 2015/04/03. doi:
10.1101/cshperspect.a005934. PubMed PMID: 25833842; PubMed Central PMCID:
PMCPMC4382748.

87.   Tyson JJ, Chen KC, Novak B. Sniffers, buzzers, toggles and blinkers: dynamics of regulatory
and signaling pathways in the cell. Curr Opin Cell Biol. 2003;15(2):221-31. Epub 2003/03/22. doi:
10.1016/s0955-0674(03)00017-6. PubMed PMID: 12648679.

88.   Eungdamrong NJ, Iyengar R. Modeling cell signaling networks. Biol Cell. 2004;96(5):355-62.
Epub 2004/06/23. doi: 10.1016/j.biolcel.2004.03.004. PubMed PMID: 15207904; PubMed Central
PMCID: PMCPMC3620715.

89.   Umulis DM, Othmer HG. The role of mathematical models in understanding pattern formation
in developmental biology. Bull Math Biol. 2015;77(5):817-45. doi: 10.1007/s11538-014-0019-7.
PubMed PMID: 25280665; PubMed Central PMCID: PMCPMC4819020.

90.   Capek D, Muller P. Positional information and tissue scaling during development and
regeneration. Development. 2019;146(24). Epub 2019/12/22. doi: 10.1242/dev.177709. PubMed
PMID: 31862792.

91.   Kondo S, Miura T. Reaction-diffusion model as a framework for understanding biological
pattern formation. Science. 2010;329(5999):1616-20. Epub 2010/10/12. doi:
10.1126/science.1179047. PubMed PMID: 20929839.

92.   Khan SJ, Abidi SNF, Skinner A, Tian Y, Smith-Bolton RK. A positive feedback loop ensures
propagation of ROS production and JNK signalling throughout Drosophila tissue regeneration. 2017.
doi: 10.1101/152140.

93.   Shlevkov E, Morata G. A dp53/JNK-dependant feedback amplification loop is essential for the
apoptotic response to stress in Drosophila. Cell Death Differ. 2012;19(3):451-60. Epub 2011/09/03.
doi: 10.1038/cdd.2011.113. PubMed PMID: 21886179; PubMed Central PMCID: PMCPMC3278728.

94.   Martin-Blanco E, Gampel A, Ring J, Virdee K, Kirov N, Tolkovsky AM, et al. puckered encodes
a phosphatase that mediates a feedback loop regulating JNK activity during dorsal closure in
Drosophila. Genes Dev. 1998;12(4):557-70. doi: 10.1101/gad.12.4.557. PubMed PMID: 9472024;
PubMed Central PMCID: PMCPMC316530.

95.   Karsten P, Hader S, Zeidler MP. Cloning and expression of Drosophila SOCS36E and its
potential regulation by the JAK/STAT pathway. Mech Dev. 2002;117(1-2):343-6. doi: 10.1016/s0925-
4773(02)00216-2. PubMed PMID: 12204282.

96.   Callus BA, Mathey-Prevot B. SOCS36E, a novel Drosophila SOCS protein, suppresses
JAK/STAT and EGF-R signalling in the imaginal wing disc. Oncogene. 2002;21(31):4812-21. doi:
10.1038/sj.onc.1205618. PubMed PMID: 12101419.

97.   Madar D, Dekel E, Bren A, Alon U. Negative auto-regulation increases the input dynamic-
range of the arabinose system of Escherichia coli. BMC Syst Biol. 2011;5:111. Epub 20110712. doi:
10.1186/1752-0509-5-111. PubMed PMID: 21749723; PubMed Central PMCID: PMCPMC3163201.

98.   Li C, Zhang L, Nie Q. Landscape reveals critical network structures for sharpening gene
expression boundaries. BMC Syst Biol. 2018;12(1):67. Epub 20180613. doi: 10.1186/s12918-018-
0595-5. PubMed PMID: 29898720; PubMed Central PMCID: PMCPMC6001026.

99.   Jiang J, Struhl G. Complementary and mutually exclusive activities of decapentaplegic and
wingless organize axial patterning during Drosophila leg development. Cell. 1996;86(3):401-9. Epub
1996/08/09. doi: 10.1016/s0092-8674(00)80113-0. PubMed PMID: 8756722.

100.   Brook WJ, Cohen SM. Antagonistic interactions between wingless and decapentaplegic
responsible for dorsal-ventral pattern in the Drosophila Leg. Science. 1996;273(5280):1373-7. Epub
1996/09/06. doi: 10.1126/science.273.5280.1373. PubMed PMID: 8703069.

101.   Lee J, Lee J, Farquhar KS, Yun J, Frankenberger CA, Bevilacqua E, et al. Network of mutually
repressive metastasis regulators can promote cell heterogeneity and metastatic transitions. Proc Natl
Acad Sci U S A. 2014;111(3):E364-73. Epub 20140106. doi: 10.1073/pnas.1304840111. PubMed
PMID: 24395801; PubMed Central PMCID: PMCPMC3903259.

102.   Sokolowski TR, Erdmann T, ten Wolde PR. Mutual repression enhances the steepness and
precision of gene expression boundaries. PLoS Comput Biol. 2012;8(8):e1002654. Epub 20120830.
doi: 10.1371/journal.pcbi.1002654. PubMed PMID: 22956897; PubMed Central PMCID:
PMCPMC3431325.

103.   Theisen H, Haerry TE, O'Connor MB, Marsh JL. Developmental territories created by mutual
antagonism between Wingless and Decapentaplegic. Development. 1996;122(12):3939-48. Epub
1996/12/01. PubMed PMID: 9012514.

104.   Villa-Cuesta E, Modolell J. Mutual repression between msh and Iro-C is an essential
component of the boundary between body wall and wing in Drosophila. Development.
2005;132(18):4087-96. Epub 20050810. doi: 10.1242/dev.01977. PubMed PMID: 16093324.

105.   Zagorski M, Tabata Y, Brandenberg N, Lutolf MP, Tkacik G, Bollenbach T, et al. Decoding of
position in the developing neural tube from antiparallel morphogen gradients. Science.

2017;356(6345):1379-83. Epub 2017/07/01. doi: 10.1126/science.aam5887. PubMed PMID:

28663499; PubMed Central PMCID: PMCPMC5568706.

106.   Ferrell JE, Jr. Self-perpetuating states in signal transduction: positive feedback, double-

negative feedback and bistability. Curr Opin Cell Biol. 2002;14(2):140-8. Epub 2002/03/14. doi:

10.1016/s0955-0674(02)00314-9. PubMed PMID: 11891111.

107.   JAK/STAT Pathway Signalling in Drosophila Melanogaster [Internet]. Landes Bioscience.

108.   Sarov M, Barz C, Jambor H, Hein MY, Schmied C, Suchold D, et al. A genome-wide resource

for the analysis of protein localisation in Drosophila. Elife. 2016;5:e12068. Epub 2016/02/21. doi:

10.7554/eLife.12068. PubMed PMID: 26896675; PubMed Central PMCID: PMCPMC4805545.

109.   Niu GJ, Xu JD, Yuan WJ, Sun JJ, Yang MC, He ZH, et al. Protein Inhibitor of Activated STAT

(PIAS) Negatively Regulates the JAK/STAT Pathway by Inhibiting STAT Phosphorylation and

Translocation. Front Immunol. 2018;9:2392. Epub 20181026. doi: 10.3389/fimmu.2018.02392.

PubMed PMID: 30416501; PubMed Central PMCID: PMCPMC6212522.

110.   Saadin A, Starz-Gaiano M. Identification of Novel Regulators of the JAK/STAT Signaling

Pathway that Control Border Cell Migration in the Drosophila Ovary. G3 (Bethesda). 2016;6(7):1991-

2002. Epub 2016/05/14. doi: 10.1534/g3.116.028100. PubMed PMID: 27175018; PubMed Central

PMCID: PMCPMC4938652.

111.   Arbouzova NI, Bach EA, Zeidler MP. Ken & barbie selectively regulates the expression of a

subset of Jak/STAT pathway target genes. Curr Biol. 2006;16(1):80-8. doi: 10.1016/j.cub.2005.11.033.

PubMed PMID: 16401426.

112.   Harris RE, Stinchfield MJ, Nystrom SL, McKay DJ, Hariharan IK. Damage-responsive,

maturity-silenced enhancers regulate multiple genes that direct regeneration in Drosophila. Elife.

2020;9. Epub 20200603. doi: 10.7554/eLife.58305. PubMed PMID: 32490812; PubMed Central

PMCID: PMCPMC7299344.

113.   Muller P, Kuttenkeuler D, Gesellchen V, Zeidler MP, Boutros M. Identification of JAK/STAT

signalling components by genome-wide RNA interference. Nature. 2005;436(7052):871-5. Epub

2005/08/12. doi: 10.1038/nature03869. PubMed PMID: 16094372.

114.   Baeg GH, Zhou R, Perrimon N. Genome-wide RNAi analysis of JAK/STAT signaling

components in Drosophila. Genes Dev. 2005;19(16):1861-70. Epub 20050729. doi:

10.1101/gad.1320705. PubMed PMID: 16055650; PubMed Central PMCID: PMCPMC1186186.

115.   La Marca JE, Willoughby LF, Allan K, Portela M, Goh PK, Tiganis T, et al. PTP61F Mediates
Cell Competition and Mitigates Tumorigenesis. Int J Mol Sci. 2021;22(23). Epub 20211125. doi:
10.3390/ijms222312732. PubMed PMID: 34884538; PubMed Central PMCID: PMCPMC8657627.

116.   Willoughby LF, Manent J, Allan K, Lee H, Portela M, Wiede F, et al. Differential regulation of
protein tyrosine kinase signalling by Dock and the PTP61F variants. FEBS J. 2017;284(14):2231-50.
Epub 20170630. doi: 10.1111/febs.14118. PubMed PMID: 28544778.

117.   Schertel C, Huang D, Bjorklund M, Bischof J, Yin D, Li R, et al. Systematic screening of a
Drosophila ORF library in vivo uncovers Wnt/Wg pathway components. Dev Cell. 2013;25(2):207-19.
Epub 2013/04/16. doi: 10.1016/j.devcel.2013.02.019. PubMed PMID: 23583758.

118.   Meinhardt H. Cell determination boundaries as organizing regions for secondary embryonic
fields. Dev Biol. 1983;96(2):375-85. doi: 10.1016/0012-1606(83)90175-6. PubMed PMID: 6832478.

119.   Martinez Arias A, Steventon B. On the nature and function of organizers. Development.
2018;145(5). Epub 20180309. doi: 10.1242/dev.159525. PubMed PMID: 29523654; PubMed Central
PMCID: PMCPMC5868996.

120.   Bilder D, Li M, Perrimon N. Cooperative regulation of cell polarity and growth by Drosophila
tumor suppressors. Science. 2000;289(5476):113-6. Epub 2000/07/07. doi:
10.1126/science.289.5476.113. PubMed PMID: 10884224.

121.   Pagliarini RA, Xu T. A genetic screen in Drosophila for metastatic behavior. Science.
2003;302(5648):1227-31. Epub 2003/10/11. doi: 10.1126/science.1088474. PubMed PMID:
14551319.

122.   Dillard C, Reis JGT, Rusten TE. Ras(V12); scrib(-/-) Tumors: A Cooperative Oncogenesis
Model Fueled by Tumor/Host Interactions. Int J Mol Sci. 2021;22(16). Epub 20210818. doi:
10.3390/ijms22168873. PubMed PMID: 34445578; PubMed Central PMCID: PMCPMC8396170.

123.   Igaki T, Pastor-Pareja JC, Aonuma H, Miura M, Xu T. Intrinsic tumor suppression and
epithelial maintenance by endocytic activation of Eiger/TNF signaling in Drosophila. Dev Cell.
2009;16(3):458-65. Epub 2009/03/18. doi: 10.1016/j.devcel.2009.01.002. PubMed PMID: 19289090;
PubMed Central PMCID: PMCPMC2729686.

124.   Ohsawa S, Sugimura K, Takino K, Xu T, Miyawaki A, Igaki T. Elimination of oncogenic
neighbors by JNK-mediated engulfment in Drosophila. Dev Cell. 2011;20(3):315-28. Epub 2011/03/15.
doi: 10.1016/j.devcel.2011.02.007. PubMed PMID: 21397843.

125. Uhlirova M, Jasper H, Bohmann D. Non-cell-autonomous induction of tissue overgrowth by
JNK/Ras cooperation in a Drosophila tumor model. Proc Natl Acad Sci U S A. 2005;102(37):13123-8.
Epub 2005/09/10. doi: 10.1073/pnas.0504170102. PubMed PMID: 16150723; PubMed Central
PMCID: PMCPMC1201591.

126. Amoyel M, Anderson AM, Bach EA. JAK/STAT pathway dysregulation in tumors: a Drosophila
perspective. Semin Cell Dev Biol. 2014;28:96-103. Epub 20140328. doi:
10.1016/j.semcdb.2014.03.023. PubMed PMID: 24685611; PubMed Central PMCID:
PMCPMC4037387.

127. Dhanasekaran DN, Reddy EP. JNK-signaling: A multiplexing hub in programmed cell death.
Genes Cancer. 2017;8(9-10):682-94. Epub 2017/12/14. doi: 10.18632/genesandcancer.155. PubMed
PMID: 29234486; PubMed Central PMCID: PMCPMC5724802.

128. Trivedi S, Starz-Gaiano M. Drosophila Jak/STAT Signaling: Regulation and Relevance in
Human Cancer and Metastasis. Int J Mol Sci. 2018;19(12). Epub 20181214. doi:
10.3390/ijms19124056. PubMed PMID: 30558204; PubMed Central PMCID: PMCPMC6320922.

129. Herrera SC, Bach EA. JAK/STAT signaling in stem cells and regeneration: from Drosophila to
vertebrates. Development. 2019;146(2). Epub 20190129. doi: 10.1242/dev.167643. PubMed PMID:
30696713; PubMed Central PMCID: PMCPMC6361132.

130. Tsai YC, Sun YH. Long-range effect of upd, a ligand for Jak/STAT pathway, on cell cycle in
Drosophila eye development. Genesis. 2004;39(2):141-53. Epub 2004/06/02. doi:
10.1002/gene.20035. PubMed PMID: 15170700.

131. Du L, Wang J. JAK/STAT pathway promotes *Drosophila* neuroblast proliferation via the direct
CycE regulation. bioRxiv. 2020:2020.07.10.195875. doi: 10.1101/2020.07.10.195875.

132. Beira JV, Torres J, Paro R. Signalling crosstalk during early tumorigenesis in the absence of
Polycomb silencing. PLoS Genet. 2018;14(1):e1007187. Epub 2018/01/23. doi:
10.1371/journal.pgen.1007187. PubMed PMID: 29357360; PubMed Central PMCID:
PMCPMC5794193.

133. Wang Z, Shi C. Cellular senescence is a promising target for chronic wounds: a
comprehensive review. Burns Trauma. 2020;8:tkaa021. Epub 20200623. doi: 10.1093/burnst/tkaa021.
PubMed PMID: 32607375; PubMed Central PMCID: PMCPMC7309580.

134. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. doi:
10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMC3855844.

135.   Rausenberger J, Tscheuschler A, Nordmeier W, Wust F, Timmer J, Schafer E, et al.

Photoconversion and nuclear trafficking cycles determine phytochrome A's response profile to far-red

light. Cell. 2011;146(5):813-25. Epub 2011/09/03. doi: 10.1016/j.cell.2011.07.023. PubMed PMID:

21884939.

1244
1245

1246 **Figure 1**

1247 **JNK/AP-1 and JAK/STAT are activated in distinct spatial domains upon tissue damage**

1248

1249 **(A-B)** A control wing disc **(A)** and wing disc after 24h of *egr*-expression **(B)** in the pouch (R0) (see also **S1.1, B**)

1250 under the control of the inducible *rn-GAL4* (*rotund-GAL4*) driver (*rn^{ts}>*) in the pouch domain. Discs express the

1251 JNK/AP-1 reporter *TRE>RFP* and were stained for cleaved Dcp-1 to visualize apoptosis.

1252

1253 **(C-D)** A control wing disc **(C)** and wing disc after 24h of *egr*-expression **(D)** in the pouch (R0), under the control of

1254 the inducible *rn-GAL4* (*rotund-GAL4*) driver (*rn^{ts}>*). Discs also express the FUCCI cell cycle reporters *mRFP-NLS-*

1255 *CycB^{1-266}* (red) and *GFP-E2f1^{1-230}* (green) (see also **S1.1A**) and were assessed for DNA replication by EdU

1256 incorporation assay.

1257

1258 **(E-F)** A control **(E)** and *egr*-expressing disc **(F)** at R0, expressing the *upd3.1-3-LacZ* (gray) and the JNK/AP-1

1259 reporter *TRE>RFP* (magenta). Discs were stained for β-Galactosidase.

1260

1261 **(G-H)** A control **(G)** and *egr*-expressing disc **(F)** at R0, expressing the JNK/AP-1 reporter *TRE>RFP* (magenta) and

1262 the dynamic JAK/STAT reporter *Stat92E>dGFP* (green).

1263

(I-J) 2D density plots mapping pixel fluorescence intensity values for the *TRE>RFP* (JNK intensity, X-axis) and

Stat92E>dGFP (JAK/STAT intensity, Y-axis) reporters, measured in the combined wing pouch and hinge domains

(black dashed lines in **O**) in control, *rn-GAL4*-expressing discs **(I)** or *egr*-expressing discs **(J)** . Dashed lines

(cyan) represent visually defined thresholds distinguishing low or high reporter activity in the quadrants Q1,Q2,Q3

and Q4. Note that less than 1% of pixels show both high JNK and high JAK/STAT signaling activity in the

combined JNK and JAK/STAT signaling domains (i.e. pouch and hinge). Graphs represent pooled values from

n=4, control and *n=3, egr*-expressing discs.

1271

1272 **(K-L)** A control **(K)** and *egr*-expressing disc **(L)** assessed for S-phase activity using EdU incorporation assays. Discs

1273 also express the *TRE>RFP* (magenta) and *Stat92E>dGFP* (green) reporters.

1274

1275 **(M-N)** Fluorescence intensity profiles for *upd3.1-3LacZ* (gray) and *TRE>RFP* (magenta) reporters **(M)**, or *TRE>RFP*

1276 (magenta), *Stat92E>dGFP* (green) reporters and EdU (yellow) reporter intensities **(N)** traced from the pouch center

1277 (PC) to the disc periphery (DP) (yellow arrows in **F** and **O**). Graph represents mean ± SEM of reporter fluorescence

1278 intensity values from *n=15* tracks across a representative disc, scaled to the maximum measured value.

1279
1280

1281 **(O)** Schematic representation of undamaged control discs and spatially segregated patterns of JNK/AP-1 (magenta)

1282 and JAK/STAT (green) reporter activity in 24h *egr*-expressing discs (R0).

1283

1284 Maximum projections of multiple confocal sections are shown in **K-L**.

1285 Discs were stained with DAPI to visualize nuclei. Scale bars: 50 μm

1286
1287

Figure 2

JNK/AP-1 signaling represses JAK/STAT activity cell-autonomously

(A-C) RFP expression marks *p35*-expressing control clones in the pouch (magenta, **A**) or *p35,hep^act*-expressing JNK+ clones in the pouch (magenta, **B**) and peripodium (magenta, **C**) 28h after clone induction in discs expressing the stable *Stat92E>GFP* reporter (green). Yellow arrowheads indicate clones of interest. Note the cell-autonomous repression of *Stat92E>GFP* reporter activity in JNK+ clones in the pouch periphery and hinge **(B)**, as well as in the peripodium **(C)**, when compared to control clones in the pouch **(A)** and peripodium **(see S2A)**

(D) *p35,hep^act*-expressing JNK+ clones in a disc expressing the *Stat92E>GFP* reporter (green), stained with anti-MMP-1 (magenta) - a downstream target of the JNK/AP-1 signaling pathway. The yellow circle highlights a JNK+ clone of interest, which shows a strong cell-autonomous repression of the JAK/STAT reporter activity in the JAK/STAT-signaling competent hinge, 48h after clone induction.

(E-F) Control disc **(E)** and a disc expressing *egr* **(F)** in the posterior compartment using the *enGAL4* driver at R0. Dashed yellow line marks the A/P boundary. Both discs express the *Stat92E>dGFP* reporter (green) and are stained for the JNK/AP-1 target MMP-1 (magenta). White asterisk indicates JNK/AP-1 reporter activity in the posterior pouch region. Note the non-cell autonomous JAK/STAT reporter activity in the anterior pouch, indicated by yellow arrowheads in **F**.

(G-J) Control **(G,I)** and *egr*-expressing **(H,J)** discs at 7h or 14h after ablation expressing the *TRE>RFP* reporter (magenta) and assessed for S-phase activity using EdU incorporation assay (cyan). While the number of S-phase cells are uniformly distributed in the pouch domain of undamaged control discs **(G,I)**, note the successive decrease in the S-phase cells (cyan) from 7h **(H)** to 14h **(J)** of *egr*-expression, coinciding with increasing *TRE>RFP* reporter activity in the pouch. Insets show the same images, with settings adjusted to visualize the JNK/AP-1 activity in the pouch of 7h *egr*-expressing discs.

(K-M) A time-course analysis of *TRE>RFP* (magenta) and *Stat92E>dGFP* (green) reporter activity in *egr*-expressing discs after 7h **(K)**, 14h **(L)** and 24h **(M)** of ablation. Yellow squares show magnified regions. Yellow lines mark the interface between JNK/AP-1 and JAK/STAT signaling cells. Note the increasing dominance of JNK/AP-1 signaling and decreasing JAK/STAT activation inside the boundary of JNK/AP-1 signaling cells at the JNK/AP-1-JAK/STAT interface.

(N-P) Representative 2D density plots of pixel fluorescence intensity for the *TRE>RFP* (JNK intensity, X-axis) and *Stat92E>dGFP* (JAK/STAT intensity, Y-axis) reporters, measured exclusively within the JNK/AP-1 signaling pouch domain (magenta in Fig. **1O**) from 7h **(L)**, 14h **(M)** and 24h **(N)** *egr*-expressing discs. Black asterisks highlight the region of increase and subsequent decline in the density of co-localizing pixels with medium and high reporter activities from 7h to 14h **(N,O)**, and 14h to 24h **(O,P)** respectively.

Maximum projections of multiple confocal sections are shown in **E-J**.
Discs were stained with DAPI to visualize nuclei. Scale bars: 50 µm

1331 **Figure 3**

1332 **A mutual-repression loop promotes bistable segregation of JNK/AP-1 and JAK/STAT activation patterns**

1333

(A-E') Workflow to investigate the presence of a regulatory motif between the JNK/AP-1 and JAK/STAT signaling

pathways driving pattern formation in *egr*-expressing discs. Fluorescence intensity profiles for the *TRE>RFP*

(magenta) and *Stat92E>dGFP* (green) reporter patterns show bistable segregation in 24h egr-expressing discs.

Intensity profiles are traced from the pouch center (PC) to the disc periphery (DP) **(A)**. Schematic showing the

models tested – a uni-directional repression model acting only on JNK/AP-1 signaling **(B)** and a mutual repression

model acting on both JNK/AP-1 and JAK/STAT signaling **(B')**. Modeling equations were generated, and a global

analysis by sampling approach was used to analyze 21 selected parameters **(C)** (see also Fig.**S3**) and test for

outcomes that produce bistable, switch-like patterns **(D-D')**. Number of simple bistable outcomes **(D)** from the

mutual repression model (dark grey) are substantially greater than those from the uni-directional repression model

(light grey) **(E)**. Applying stringent selection criteria based on the biologically observed reporter activity pattern **(D')**,

positive outcomes reduce to 0 in the model without the mutual repression **(E')**.

1345

1346

1347 **Figure 4**

1348 **Ptp61F represses JAK/STAT activity in JNK/AP-1 signaling cells**

1349

1350 **(A-C)** A *ptp61F-RNAi*-expressing **(A)**, *egr*-expressing **(B)** and *egr,ptp61F-RNAi*-co-expressing disc **(C)** at R0. All

1351 discs also express the *Stat92E>dGFP* reporter (green). Yellow circles highlight the central domain of remaining *rn-*

1352 *GAL4* expressing cells. Note the derepression and ectopic JAK/STAT activity in the *egr,ptp61F-RNAi*-co-expressing

1353 disc.

1354

1355 **(D)** Quantification of total area of wing discs. Graphs display mean ± SEM for *n=8*, *egr*-expressing and *n=10*,

1356 *egr,ptp61F-RNAi*-co-expressing discs. T-tests were performed to test for statistical significance.

1357

1358 **(E)** Quantification of the *Stat92E>dGFP* reporter fluorescence intensity measured within the JNK/AP-1 signaling

1359 pouch domain. Black square in schematic shows measured region. Graphs display mean ± SEM for *n=8*, *egr-*

1360 expressing and *n=10*, *egr,ptp61F-RNAi*-co-expressing discs. U-tests were performed to test for statistical

1361 significance.

1362

1363 **(F-I)** A control **(F)**, *stat92E*-expressing **(G)**, *egr*-expressing **(H)** and *egr,stat92E*-co-expressing disc **(I)**. All discs also

1364 express the dynamic *Stat92E>dGFP* reporter (green). Staining for HA confirms expression and nuclear/cytoplasmic

1365 localization of the *UAS-stat92E-3xHA* construct. Yellow arrowheads point to the restricted, non-autonomously

1366 induced JAK/STAT activity in the peripheral pouch in control discs **(F,G)**. Yellow circles highlight the central domain

1367 of remaining *rn-GAL4* expressing cells **(H,I)**.

1368

1369 **(J)** Quantification of the *Stat92E>dGFP* reporter fluorescence intensity measured within the JNK/AP-1 signaling

1370 pouch domain. Black square in schematic **(E)** shows measured region. Graphs display mean ± SEM for *n=16*,

1371 control discs (*rn^{ts}>*), *n=9*, *stat92E*-expressing discs, *n=11*, *egr*-expressing and *n=12*, *egr,stat92E*-co-expressing

1372 discs. One-way ANOVA with multiple comparisons was performed to test for statistical significance.

1373

1374 Discs were stained with DAPI (magenta) to visualize nuclei. Scale bars: 50 μm

1375

1376

1377

1378

Figure 5

JAK/STAT repression by JNK/AP-1 is required for the G2 arrest and protects central wound cells from apoptosis.

(A-B) An *egr*-expressing **(A)** and *egr,stat92E*-coexpressing disc **(B)** at R0, stained for DAPI (cyan), and cleaved Dcp-1 (red) to visualize apoptosis.

(C-D) Quantification of total wing disc area **(C)** and normalized area occupied by cleaved Dcp-1 **(D)**. Graphs display mean ± SEM for *n=10, egr*-expressing and *n=9, egr,stat92E*-co-expressing discs. T-tests were performed to test for statistical significance.

(E-F) An *egr*-expressing **(E)** and *egr,stat92E*-coexpressing disc **(F)** at R0, also expressing the FUCCI reporter. Dashed white squares highlight the distinct shift from G2 (yellow cells) to G1 (green cells) in the pouch domain.

(G-H) Bar graph **(G)** representing the cumulative proportion of gap phase cells in G1 (green) and G2 (yellow) for each genotype. Graph **(H)** displays G1-phase:G2-phase ratios obtained from *n=5, egr*-expressing and *n=3, egr,stat92E*-co-expressing discs. Dashed white squares in **E-F** show measured regions. T-test with Welch's correction was performed to test for statistical significance.

(I-J) EdU incorporation assay to detect S-phase cells (gray) in *egr*-expressing **(I)**, or *egr,stat92E*-co-expressing **(J)** disc. Discs were stained with DAPI (red) to visualize nuclei.

(K-L) An *egr*-expressing **(K)** and *egr,ptp61F-RNAi*-coexpressing **(L)** disc at R0, stained for DAPI (cyan), and cleaved Dcp-1 (red) to visualize apoptosis.

(M-N) Quantification of total wing disc area **(M)** and normalized area occupied by cleaved Dcp-1 **(N)**. Graphs display mean ± SEM for *n=11, egr*-expressing and *n=15, egr,ptp61F-RNAi*-coexpressing discs. T-tests were performed to test for statistical significance.

(O-P) EdU incorporation assay to detect S-phase cells (gray) in an *egr*-expressing **(O)**, and *egr,ptp61F-RNAi*-coexpressing **(P)** disc. Discs were also stained with DAPI (red) to visualize nuclei.

Maximum projections of multiple confocal sections are shown in **A-B, I-L, O-P**. Scale bars: 50 µm

Figure 6

The wound organizer network segregates JNK/AP-1 and JAK/STAT signaling to drive oncogenic growth

____

(A-D) A control wing disc **(A)** and wing discs expressing *Ras^v12* **(B)**, *scrib-RNAi* **(C, C')**, or *Ras^v12,scrib-RNAi* **(D,D')**

for 44 h D6 AED. Discs also express the JNK/AP-1 reporter *TRE>RFP* (magenta) and the JAK/STAT reporter

Stat92E>dGFP (green). Yellow squares in mark area of magnification shown in **(C',D')**.

1422

(E) Schematic shows the *TRE>RFP*-positive (JNK$^+$, magenta), *Stat92E>dGFP*-positive (STAT$^+$, green) and

JNK/AP-1 and JAK/STAT double positive regions (JNK$^+$STAT$^+$, gray) selected for measurements. Graph represents

normalized area of measured regions based on the total signaling area. Note that only 15% of regions within the

tumor express both the JNK/AP-1 and JAK/STAT signaling reporters. Graph represents mean ± SEM for *n=6,*

Ras^v12,scrib-RNAi-expressing discs. One-way paired ANOVA with Holm-Šídák's multiple comparison test was

performed to test for statistical significance.

1429

1430 **(F)** Schematic shows *TRE>RFP*-positive (JNK$^+$, magenta) and *TRE>RFP*-negative (JNK$^-$, white) regions selected

1431 for measurements. Graph represents *Stat92E>dGFP* fluorescence intensity measured within selected JNK$^+$ and

1432 JNK$^-$ regions. Note the reduction in the reporter intensity from 71.91±4.32 to 36.66±1.74 within the JNK$^-$ vs. JNK$^+$

1433 region. Graph represents mean ± SEM for *n=6,* ras^{v12},scrib-RNAi-expressing discs. Paired t-tests was performed

1434 to test for statistical significance.

1435

1436 **(G-H)** A wing disc expressing *scrib-RNAi* **(G)** and *Ras^v12,scrib-RNAi* **(H)** at 44 h. Discs also express the

1437 *Stat92E>dGFP* reporter and were stained for phospho-Histone3 (ph3) to visualize cells in mitosis. Segmented

1438 masks of ph3 were generated for quantifications and an overlay of the *Stat92E>dGFP* reporter (green) and ph3

1439 (white) is represented. Orange dashed lines mark the STAT$^+$ regions within the discs.

1440

1441 **(I)** Schematic shows *Stat92E>dGFP*-positive (STAT$^+$, green) and *Stat92E>dGFP*-negative (STAT$^-$, white) regions

1442 selected for measurements. Graph represents ph3 counts normalized to area in the STAT$^+$ and STAT$^-$ regions for

1443 *n=5, Ras^v12,scrib-RNAi*-expressing discs.

1444

1445 Maximum projections of multiple confocal sections are shown in **G-H**

1446 Discs were stained with DAPI to visualize nuclei. Scale bars: 50 µm

1447

1448

1449

 Figure 7

Mutual repression between JNK/AP-1 and JAK/STAT sets up wound organizer activity

---

(A-B) Transient (7h *egr*-expression), and chronic (24h *egr*-expression or 44h *Ras^{v12},scrib-RNAi*-expression) activation of JNK/AP-1 signaling stratifies JNK-activating tissue into distinct JNK/AP-1 and JAK/STAT signaling domains **(A)** and cell behaviors **(B)**.

(C) *egr*-expressing discs spatially stratify tissue repair behaviors via a mutual repression motif. JNK/AP-1 represses JAK/STAT signaling activitiy via Ptp61F, while JAK/STAT represses JNK/AP-1 activation via Zfh2. JNK signaling ensures G2 arrest and cell survival, while JAK/STAT signaling ensures proliferation and supports survival in the absence of JNK.

(D) Disruption of the mutual repression motif by cell-autonomous activation of JNK/AP-1 and JAK/STAT signaling overrides the protective G2 stall and drives cells into mitotic cycling and G1, where p53 activation can lead to apoptotic outcomes.

Figure 1.

Figure 1

Figure 2

Figure 2

Figure 3

A

Biologically observed reporter activity patterns

Simple mathematical models

B Uni-directional repression model

B' Mutual repression model

C Modeling equations

$$\frac{d\text{EIG}(t,x)}{dt} = b_{EIG}(x) - k_{deg,EIG}\text{EIG} + k_{act,EIG}\frac{\text{JNK}^{n_1}}{K_{m,act,EIG}^{n_1} + \text{JNK}^{n_1}} + D_{EIG}\nabla^2 EIG$$

$$\frac{d\text{JNK}(t,x)}{dt} = b_{JNK} - k_{deg,JNK}\text{JNK} + k_{act,JNK}\frac{\text{EIG}^{n_2}}{(K_{m,act,JNK}^{n_2} + \text{EIG}^{n_2})(1 + k_{inh,JNK,by,JAK}^{ni1}\text{JAK}^{ni1})}$$

$$\frac{d\text{JAK}(t,x)}{dt} = b_{JAK} - k_{deg,JAK}\text{JAK} + k_{act,JAK}\text{UPD}\frac{\text{JAK}^{n_3}}{(K_{m,act,JAK}^{n_3} + \text{JAK}^{n_3})(1 + k_{inh,JAK,by,JNK}^{ni2}\text{JNK}^{ni2})}$$

$$\frac{d\text{UPD}(t,x)}{dt} = b_{UPD} - k_{deg,UPD}\text{UPD} + k_{act,UPD}\frac{\text{JNK}^{n_4}}{K_{m,act,UPD}^{n_4} + \text{JNK}^{n_4}} + D_{UPD}\nabla^2 UPD$$

Global analysis by sampling approach

Automated extraction of simulation outcomes based on selected inclusion criteria

D Simple bistable pattern

D' Observed bistable pattern

E

E'

Uni-directional repression model
Mutual repression model

Figure 3

Figure 4

Figure 4

Figure 5

Figure 5

Figure 7

A
transient JNK | high, chronic JNK | moderate, chronic JNK (tumor)

◆ JNK signaling
⬡ JAK/STAT signaling

B

⬡ JNK ⋙ G2 stall (Survival)
⬡ JAK/STAT ⋙ Proliferation (Survival)

C

JAK/STAT
Ptp61F   Zfh2
JNK

JNK   JAK/STAT
∨   ∨
G2 stall   Proliferation
Survival   Survival

D

JAK/STAT
Zfh2
JNK

JNK   JAK/STAT
∨   ∨
G1 cycling   Proliferation
Apoptosis   Survival