Activation of TnSmu1, an integrative and conjugative element, by an ImmR-like transcriptional regulator in *Streptococcus mutans*

Shawn King[1], Allison Quick[1], Kalee King[1], Alejandro R. Walker[2], and Robert C. Shields[1*]
[1] Department of Biological Sciences, Arkansas State University, Jonesboro, Arkansas, USA
[2] Department of Oral Biology, University of Florida, Gainesville, Florida, USA

*Corresponding Author: rshields@astate.edu

ORCiDs:
Shields: http://orcid.org/0000-0002-8823-9504
Walker: https://orcid.org/0000-0002-5839-6584

Keywords: *Streptococcus mutans*, integrative and conjugative element, TnSmu1, transcriptional regulator, ImmR

Funding information: National Institute of Dental and Craniofacial Research, Grant/Award Number: DE029882

Conflict of interest: the authors have no conflicts of interest to declare

Author contributions: S.K., A.Q., K.K., A.R.W., and R.C.S. were involved in the investigation, methodology, data curation, formal analysis, and revised the manuscript; S.K. and R.C.S. were involved in project administration; A.R.W. performed software development (implementation of code and supporting algorithms); R.C.S. was involved with funding acquisition, supervision, conceptualization of the study and drafted the manuscript.

Abstract

Integrative and conjugative elements (ICEs) are chromosomally encoded mobile genetic elements that can transfer DNA between bacterial strains. Recently, as part of efforts to determine hypothetical gene functions, we have discovered an important regulatory module encoded on an ICE known as TnSmu1 on the *Streptococcus mutans* chromosome. The regulatory module consists of a cl-like repressor with a helix-turn-helix DNA binding domain *immR*$_{Smu}$ (immunity repressor) and a metalloprotease *immA*$_{Smu}$ (anti-repressor). It is not possible to create an in-frame deletion mutant of *immR*$_{Smu}$ and repression of *immR*$_{Smu}$ with CRISPRi causes substantial cell defects. We used a bypass of essentiality (BoE) screen to discover genes that allow deletion of the regulatory module. This revealed that conjugation genes, located within TnSmu1, can restore the viability of an *immR*$_{Smu}$ mutant. Deletion of *immR*$_{Smu}$ also leads to production of a circular intermediate form of TnSmu1, that is also inducible by the genotoxic agent mitomycin C. To gain further insights into potential regulation of TnSmu1 by ImmR$_{Smu}$ and broader effects on *S. mutans* UA159 physiology we used CRISPRi and RNA-seq. Strongly induced genes included all the TnSmu1 mobile element, genes involved in amino acid metabolism, transport systems, and a Type I-C CRISPR-Cas system. Lastly, bioinformatic analysis shows that the TnSmu1 mobile element and its associated genes are well distributed across *S. mutans* isolates. Taken together, our results show that activation of TnSmu1 is controlled by the *immRA*$_{Smu}$ module, and that activation is deleterious to *S. mutans*, highlighting the complex interplay between mobile elements and their host.

INTRODUCTION

Streptococcus mutans is a Gram-positive bacterium that colonizes the human oral cavity [1]. Like many streptococci, *S. mutans* can cause disease when environmental conditions are favorable. Poor oral hygiene combined with frequent ingestion of simple sugars creates an environment in which *S. mutans* can cause dental caries (tooth decay) [2, 3]. If this disease is allowed to progress it causes the breakdown of teeth, with symptoms that include pain, difficulty eating and tooth loss [2]. Due to its widespread presence in the oral cavity and resulting disease burden it is necessary to identify targets for improved therapeutics and understand processes essential for the pathogen. With this goal in mind, we have recently applied transposon sequencing (Tn-seq) and CRISPR interference (CRISPRi) and have identified >200 *S. mutans* essential genes that are required for the viability of the organism [4, 5]. For *S. mutans*, most essential genes can be broadly sorted into three categories: processing of genetic information, energy production, and maintenance of the cell envelope [4].

It was during Tn-seq/CRISPRi experiments that we discovered an essential gene, SMU_218, which is annotated as a transcriptional regulator, and is designated *immR*$_{Smu}$ (immunity repressor) from here forwards. *immR*$_{Smu}$ resides in a two-gene operon with SMU_219/ *immA*$_{Smu}$ (anti-repressor). Probing *immR*$_{Smu}$ and *immA*$_{Smu}$ with SMART (Simple Modular Architecture Research Tool) [6], we found that the N-terminus of *immR*$_{Smu}$ contains a cl-like repressor of phage λ DNA-binding domain, often found in streptococcal phages. *immA*$_{Smu}$ contains a putative ImmA/IrrE family metallo-endopeptidase domain, based on possession of a conserved metalloprotease zinc-binding motif, HEXXH [7]. In addition, the toxin-antitoxin database RASTA [8] annotates this two-gene operon as a putative toxin-antitoxin module. We found this interesting as the antitoxin component (*e.g. immR*$_{Smu}$) of toxin-antitoxin modules are often essential, because they prevent the accumulation of the toxin protein. While we cannot exclude the hypothesis that this two-gene operon is a toxin-antitoxin module, exhaustive literature searches provide evidence for other functions of these genes. First, model λ cl repressors have an N-terminal domain that binds DNA and a 3' domain that functions in cl autoproteolysis, in conjunction with RecA [9]. In *S. mutans,* it is possible that these functions are separated into two genes, *immR*$_{Smu}$ and *immA*$_{Smu}$. Next, cl-like repressors regulate prophage induction in bacteria but they have also been shown to regulate integrative conjugative element (ICE) expression [10, 11]. Notably, *immR*$_{Smu}$ and *immA*$_{Smu}$ reside in the flanking region of the putative ICE designated as TnSmu1. This large region of DNA (23 kilobases) contains predicted conjugation genes, as well as many hypothetical proteins, and is found in several strains of *S. mutans* [12, 13]. Very little is known about the activity or the mobility of this element, but it shares similarities with ICE*St1*, ICE*St3*, Tn*916* and ICE*Bs1*. Although the regulatory regions of these different ICEs are substantially re-arranged compared to TnSmu1, ICE*St1* and ICE*Bs1* contain cl-like repressors that repress ICE expression [10, 11, 14]. Therefore, we have formulated two major hypotheses explaining the essential nature of *immR*$_{Smu}$: 1) *immR*$_{Smu}$ is an antitoxin in a toxin-antitoxin module or 2) *immR*$_{Smu}$ is involved in the activation of TnSmu1. For the first hypothesis CRISPR-mediated knockdown of *immR*$_{Smu}$ would lead to over-accumulation of *immA*$_{Smu}$, which would then act as a toxin. For the second hypothesis, loss of the repressor would lead to activation of TnSmu1 with loss of the element from the cell making *immR*$_{Smu}$ appear 'essential'.

Here we investigate both of our hypotheses for explaining the essentiality of *immR*$_{Smu}$. Using transposon mutagenesis, genome sequencing, RNA-seq, bioinformatics and standard molecular microbiology techniques we find that *immR*$_{Smu}$ is very likely to be a repressor of the ICE TnSmu1 and is related to the ICE*Bs1* immunity repressor *immR*. Removal of *immR*$_{Smu}$ repression causes upregulation of the TnSmu1 element leading to excision from the genome and formation of a circular intermediate. We found that TnSmu1 is activated by DNA damage

and we posit that this occurs via ImmA$_{Smu}$-mediated cleavage of the ImmR$_{Smu}$ repressor. We also provide evidence of broader effects on *S. mutans* physiology, including slowed growth and disrupted cell morphology when TnSmu1 is activated. In summary, we have discovered an important regulatory module controlling TnSmu1 activation, which extends prior findings made in ICE*Bs1*, and illustrates the complex relationship between mobile elements and their hosts.

RESULTS

Evidence that *immR*$_{Smu}$ is not an antitoxin.

Following our initial observations, we began by investigating the possibility that *immRA*$_{Smu}$ is a toxin-antitoxin module. We reasoned that we would be able to mutagenize both *immR*$_{Smu}$ and *immA*$_{Smu}$ at the same time, as this would not lead to the accumulation of the putative toxin. For other toxin-antitoxin modules (e.g. *mazEF* or *relBE*) double knockout of both genes is possible, whereas single knockout of the antitoxin module is not permitted. Under the conditions tested, we have not been able to obtain a double mutant of *immRA*$_{Smu}$ (Figure S1). However, deletion of the putative metalloprotease, *immA*$_{Smu}$, is permitted by *S. mutans* cells (Figure S1). A lack of viability for a double knockout provides some evidence that *immRA*$_{Smu}$ is not a toxin-antitoxin module. To confirm, we also created over-expression strains in *E. coli* to test the toxicity of each protein. The *immR*$_{Smu}$ and *immA*$_{Smu}$ genes were cloned into a pBAD protein expression vector, where the genes are induced by the addition of the monosaccharide arabinose. When the expression of *immA*$_{Smu}$ was induced, there was only a minor impact on the growth rate of *E. coli*, suggesting that *immA*$_{Smu}$ accumulation in *E. coli* is not toxic (Figure S2). Over-expression of the *immR*$_{Smu}$ protein did cause a moderate growth phenotype in *E. coli*. The growth phenotype was absent when *immR*$_{Smu}$ expression was not induced by arabinose. To summarize, the data do not support the hypothesis that ImmRA$_{Smu}$ constitute a toxin-antitoxin module.

Phenotypic defects caused by repression of *immR*$_{Smu}$

Next, we wanted to employ CRISPRi as a tool to characterize any phenotypic impacts caused by repression of *immR*$_{Smu}$. Phenotyping is performed by targeting the *immR*$_{Smu}$ gene with a short guide RNA (sgRNA) which acts together with dead Cas9 (dCas9) to block gene transcription; this system is also inducible with xylose [5]. Depletion of *immR*$_{Smu}$ causes a substantial growth defect compared with the control strain (targeting a non-essential gene *lacG*) (Figure 1A). Having shown that *immR*$_{Smu}$ repression leads to a strong growth defect, we next investigated if repression causes any cell morphology impacts. Using transmission electron microscopy (TEM) we were able to gather ultrastructural insights into cells experiencing *immR*$_{Smu}$ depletion. As shown in Figure 1B, when *immR*$_{Smu}$ was depleted, there was a sub-population of sgRNA-*immR*$_{Smu}$ cells with extreme cell morphology defects. Compared to the control strain (sgRNA-lacG) *immR*$_{Smu}$ depleted strains were bloated and appeared to be not dividing or not dividing correctly (at the cell septum) (Figure 1B). Overall, repression of *immR*$_{Smu}$ has a substantial impact on the normal physiology of *S. mutans*.

Repression of *immR*$_{Smu}$ with CRISPR inference strongly up-regulates TnSmu1.

To begin to understand the function of *immR*$_{Smu}$ we performed RNA-seq profiling of an *S. mutans* strain experiencing knockdown of the *immR*$_{Smu}$ gene using CRISPRi. As shown, when *immR*$_{Smu}$ is repressed by CRISPRi there is a strong growth defect (Figure 1A). RNA was collected from uninduced and induced (with 0.1% xylose) cultures grown to OD$_{600}$ ~0.4, and this RNA was then processed for RNA-seq. The RNA-seq analysis identified 134 genes whose log$_2$

fold change was ≥2 after repression of SMU_218 (FDR < 0.05; Table S1). Of these genes, 34 were repressed and 100 were induced (Figure 2A).

There are several notable trends identified by transcriptome analysis of the $immR_{Smu}$ depleted strain. Firstly, every gene within the TnSmu1 element is strongly induced (Figure 2A; Table S1). Most TnSmu1 genes were up-regulated by greater than 20-fold, with SMU_204c (hypothetical protein) induced by over 1000-fold (log_2 fold change = 10.03). Of the twenty most strongly induced genes, all belonged to the TnSmu1 region. Visual inspection of the TnSmu1 region, with reads shown and comparable to a non-induced strain, highlights the strong induction across the element (Figure 2B). This region is predicted to be an ICE and exists in the ICE database ICEberg 2.0 [15] but to our knowledge has not been experimentally verified. To be considered an ICE, these mobile elements must contain several important features: a recombinase (also known as an integrase), a specific site of attachment (typically a tRNA gene), conjugation machinery, a relaxase, and an *oriT* site [16]. Several of these features exist within the TnSmu1 region (Figure 2B). SMU_191c encodes for a putative tyrosine recombinase that may function to catalyze DNA breakage and rejoining. Immediately adjacent to the 5' end of SMU_191c is tRNA-leu, which is likely the integration site. The attachment (*att*) sites *attL* and *attR* have the sequence CTATACCGGCGGCCG. A putative MOB_T family relaxase is encoded by the gene SMU_207c and may function to convert a circular form of TnSmu1 into ssDNA prior to conjugation. Several genes within TnSmu1 are predicted to be members of a type IV secretion system (SMU_196c, SMU_197c, SMU_198c, SMU_199c, SMU_201c, and SMU_208c). A putative *oriT* site is located between SMU_207c and SMU_208c, with the sequence ACCCCCCTATTAGTATCGGGGGG. This shares similarities with the *oriT* site in the ICEs ICE*Bs1* (ACCCCCCCACGCTAACAGGGGGGT) and ICE*St3* (ACCCCCGATTTCTAATAGGGGGGT). The relaxase encoded by SMU_207c may recognize this *oriT* sequence, where it would nick the ICE DNA to form transfer DNA (T-DNA). Altogether, TnSmu1 contains many of the genes that are required for an ICE, and may allow excision, replication, transfer, and integration of the element. In addition, strong up-regulation of these genes upon $immR_{Smu}$ depletion is suggestive of $ImmR_{Smu}$ acting as a repressor of TnSmu1.

Worth highlighting is that two intergenic regions within TnSmu1 are strongly induced (Figure 2B). The first region is between SMU_217c and $immR_{Smu}$, and the second region is between $immA_{Smu}$ and SMU_220c. For the first region, we do not currently know what is being produced (*e.g.* small RNA, small peptide, antitermination system *etc*). The second region contains a chromosomally encoded type I toxin-antitoxin system [17]. Expression of a toxic peptide known as Fst-Sm, encoded by this system, is toxic when over-expressed in *S. mutans* (in the absence of the anti-toxin) [17]. The mechanism of action of the toxic peptide is unknown but plasmid encoded Fst peptides affect the integrity of the membrane, cause defects in cell division, and sensitize cells to nisin (by altering the cell membrane) [18, 19].

Several other genes are up or downregulated after $immR_{Smu}$ depletion. A broad overview of the effect on the *S. mutans* transcriptome is shown in Figure 2C with categorization by Cluster of Orthologous Groups (COGs). Several genes involved in amino acid transport or metabolism are up-regulated including a cysteine transport operon *tcyDEFGH*, *argD*, *cysK*, *opcD*, *pepQ*, *ilvE*, *gatA*, *hipO*, SMU_1216c (putative cystine transporter), SMU_1486c (putative histidinol-phosphatase), and SMU_1938c (putative methionine transporter). Other transport systems that are also up regulated include *treB* (trehalose PTS), *fruA* (fructose PTS), ammonium transporter (*nrgA*), bacteriocin/competence transporters (*comB*, SMU_1889c, SMU_1897, *comG*, *mutE2* and *mutG*), *tauC* and *msmK*. The expression of several purine metabolism genes is altered with *purDEK* up-regulated and *purC* down-regulated. All the genes in the Type I-C CRISPR-Cas system are up-regulated when $immR_{Smu}$ is repressed. Several genes are downregulated after

CRISPRi repression of *immR*_{Smu}, many of which are hypothetical proteins. Six translation genes are down-regulated, predominately 50S and 30S ribosomal proteins. Some, but not all, genes located in the genomic island TnSmu2 are downregulated (*mubD*, *mubC*, *mubH*, *mubG*, and *mubE*). This region encodes for a hybrid nonribosomal peptide (NRP) and polyketide (PK) system that produces mutanobactin [20]. A putative acyl carrier protein (SMU_27, *acpP*) is also down-regulated that may participate in PK synthesis [21].

A bypass of essentiality screen reveals genes that when inactivated allow the deletion of *immR*_{Smu}.

Under the conditions we have tested, we are not able to obtain mutants of *immR*_{Smu} or *immRA*_{Smu} but deletion of *immA*_{Smu} is permitted (Figure S1). We hypothesized that deletion of the *immRA*_{Smu} module might be possible if introduced into a transposon library previously generated by our laboratory [4]. This is because the Tn-seq library might contain mutants that interrupt a gene responsible for *immRA*_{Smu} essentiality and thus would tolerate *immRA*_{Smu} deletion. This is known as bypass of essentiality (BoE), with systematic studies of essential gene bypass having been conducted in yeast [22, 23]. BoE has been shown for several essential genes in bacteria including *zipA*, c-di-AMP null mutants, RNase E, *bamD*, and *ftsH* [24–28]. Utilizing a BoE screen, where we transformed an *immRA*_{Smu} deletion construct into the Tn-seq library, we were able to isolate eight double mutants capable of tolerating *immRA*_{Smu} deletion (Figure 3A). These mutants were then subjected to complete genome sequencing, to identify the location of the transposon insertion and identify suppressor mutations in the event these were also generated. Genome sequencing revealed the transposon insertions to be within the following regions: SMU_197c, SMU_201c (isolated twice), SMU_141c, *recN*, *perR*, the *murE* and SMU_1678 intergenic region, and the intergenic region between 16S rDNA and SMU_1750c (Table S2). The growth phenotypes of these strains were variable (Figure 2B). The intergenic transposon insertion into *murE* caused a considerable growth phenotype, as did an insertion in SMU_197c. Certain double mutant isolates grew almost the same as wild-type *S. mutans*, including *peR*::Tn, *recN*::Tn and SMU_141c::Tn.

We hypothesized that because of the strong selective pressure placed on cells during transformation, mutants could be created that are experiencing a duplication event; this is the loss of the essential module *immRA*_{Smu} but a copy of *immRA*_{Smu} is still present. To account for this, we used PCR to visualize the size of the *immRA*_{Smu} region. Gene duplication events occurred in the strains *perR*::Tn, *recN*::Tn, and the strain with a transposon insertion between 16S rDNA and SMU_1750c (Figure S3; Table S2). Double mutants with an *immRA*_{Smu} gene duplication grew better than strains that fully lacked *immRA*_{Smu} (Figure 2B). For the other five mutants, only one band for *immRA*_{Smu} was amplified and the size was as expected for a deletion mutant. As an additional control, PCR was also conducted to confirm that the Magellan6 transposon was inserted in the regions identified by whole genome sequencing. Transposon insertions were apparent for all strains except for SMU_141c::Tn where the results were inconclusive (Figure S4; Table S2).

Evidence of a circular intermediate form of TnSmu1.

Next, we noticed that in several double mutant backgrounds, read coverage (after genomic sequencing) was significantly elevated for a region between SMU_191c and SMU_220c (Figure 4; Figure S5). Read coverage depth in this region was ~300 for *S. mutans* UA159 but was elevated to as much as ~5000 reads in double mutant backgrounds. This increased read depth corresponds exactly to a region of the genome that is predicted to contain the putative ICE, TnSmu1. When excised from the host chromosome ICEs exist as circular DNA molecules [29].

For some ICEs these circular intermediate forms are capable of rolling-circle replication [30]. Increased read coverage depth suggests that loss of $immR_{Smu}$ is leading to excision and circularization of TnSmu1. In addition to the observation that sequencing reads increase when $immR_{Smu}$ is deleted, there was also evidence of unexpected genomic junctions in strains of *S. mutans* missing the $immR_{Smu}$ gene. There was evidence from the genome sequencing that *attL* and *attR* are combining, which is leading to a junction forming between SMU_191c and SMU_220c. This is additional evidence that TnSmu1 is excising from the host chromosome and forming a circular DNA molecule. To confirm these observations, we designed primers to test excision and circularization. For excision, these primers will only amplify a PCR product if TnSmu1 is no longer present on the host chromosome. For circularization, a PCR product will form if there is a junction between the ends of the ICE (*i.e.* SMU_191c and SMU_220c). Excision and circularization of TnSmu1 in wild-type *S. mutans*, grown to mid-log, occurs close to the limit of detection for the assay (Figure 5A and B). However, excision and circularization of TnSmu1 is apparent in $immR_{Smu}$ double mutant strains. For $\Delta immR_{Smu}$ strains carrying transposon insertions within putative TnSmu1 conjugation genes (SMU_197c and SMU_201c) there are multiple copies of TnSmu1 per cell. In these strains excision is occurring 1000-fold more compared to wild-type *S. mutans*. We also observed circularization/excision in two strains experiencing duplication of $immR_{Smu}$, *perR*::Tn and *recN*::Tn. We chose to examine these strains because they are much healthier than the strains with disrupted conjugation genes (Figure 3B). For these two strains, circularization of TnSmu1 was occurring but excision of the element was measured close to the limit of detection for the assay. Taken together, our observations of excision and circularization are consistent with TnSmu1 having this important life-cycle feature of ICEs.

Induction of TnSmu1 by the genotoxic agent mitomycin C.

The $immRA_{Smu}$ module shares similarities to cI-like phage repressors that are sensitive to DNA damage (via induction of the SOS response) [31]. Other ICEs contain similar regulatory modules, ICE*Bs1* being the most extensively characterized but also ICE*St3* and SXT [10, 32, 33]. To determine if DNA damage was an inducer of TnSmu1, possibly via cleavage of $ImmR_{Smu}$, we incubated *S. mutans* UA159 in the presence of genotoxic agent mitomycin C (MMC). When treated with MMC there was a clear dose-dependent response, with clear growth hindrance at 25 ng/mL (Figure 5C). Induction of TnSmu1 circularization was investigated at MMC concentrations of 3.125 ng/mL, 6.25 ng/mL and 12.5 ng/mL. With increasing concentrations of MMC, the relative abundance of circular TnSmu1 increased in a dose-dependent manner (Figure 5D). At a concentration of 12.5 ng/mL not all cells have a circular copy of TnSmu1 but circularization is 1000-fold above levels seen in untreated *S. mutans* cells.

Transcriptome analysis in *S. mutans* strains lacking $immR_{Smu}$

In order to provide further evidence that $immR_{Smu}$ represses activation of TnSmu1, we examined the transcriptomes of two strains generated during the BoE screens. RNA was extracted from SMU_197c::Tn $\Delta immRA_{Smu}$ and SMU_201c::Tn $\Delta immRA_{Smu}$ cultures grown to OD$_{600}$ ~0.4, and this RNA was then processed for RNA-seq. For the SMU_197c::Tn $\Delta immRA_{Smu}$ strain RNA-seq analysis identified 54 genes whose log$_2$ fold change was ≥ 2 (FDR < 0.05; Table S3). Of these genes, 7 were repressed and 47 were induced (Figure 6A). Similar results were obtained for the SMU_201c::Tn $\Delta immRA_{Smu}$ strain with RNA-seq analysis identifying 51 genes whose log$_2$ fold change was ≥ 2 (FDR < 0.05; Table S4); 7 genes were repressed and 44 were induced (Figure 6A). As with CRISPRi repression of $immR_{Smu}$, TnSmu1 expression is significantly induced in both $\Delta immR_{Smu}$ strains. In both strains SMU_205c (hypothetical protein residing in TnSmu1)

was the most strongly up-regulated gene by 968-fold (SMU_201c::Tn Δ*immR*$_{Smu}$) and 399-fold (SMU_197c::Tn Δ*immR*$_{Smu}$). Next, we generated a heat map of selected genes with the aim of comparing gene expression between Δ*immR*$_{Smu}$ strains and the sgRNA-218 strain (Figure 6B). Notably, up-regulation of TnSmu1, the Type I-C CRISPR-Cas system and the CslAB transporter system is consistent among all three strains. In addition, SMU_40 (hypothetical protein with a RelE toxin-antitoxin system domain) is moderately up-regulated (~5-fold) across the three strains. For the malolactic fermentation (*mle*) locus (SMU_137 to SMU_141)[34], down-regulation (~10-fold) was observed in the Δ*immR*$_{Smu}$ strains, and up-regulation (~3.5-fold) in the sgRNA-218 strain. The *mle* locus contains a malolactic enzyme (*mleS*; SMU_137), a malate permease (*mleP*; SMU_138), an oxalate decarboxylase (*oxdC*; SMU_139), a glutathione reductase (*gshR*; SMU_140) and a hypothetical protein (SMU_141); the role of this operon is in malolactic fermentation (conversion of L-malate to L-lactate). Taken together, this additional RNA-seq analysis shows that loss of Δ*immR*$_{Smu}$ is leading to activation of TnSmu1. In addition, there is a consistent trend showing activation of the Type I-C CRISPR-Cas system and the *cslAB* transporter system.

Distribution and conjugative transfer of TnSmu1 genes between *S. mutans* strains.

To begin to understand if TnSmu1 is capable of conjugative transfer, we wanted to determine if closely and distantly related strains of *S. mutans* carry this ICE. Notably, we have recently discovered that *S. mutans* strains carry CRISPR spacers against this element [35]. This finding suggests that *S. mutans* attempts to prevent transfer/acquisition of TnSmu1 via CRISPR-Cas mediated immunity, and it also provides evidence of potential transfer between strains. Figure 7 shows strains that carry TnSmu1 genes and is organized by strain relatedness. Distantly related strains, such as smu342 and UA159, carry full length TnSmu1 elements, which might be indicative of conjugal transfer. Although many *S. mutans* strains carry TnSmu1 genes, only nine strains carry the full element as organized on *S. mutans* UA159. ICEs have plasticity and lose/gain genes, often interacting with other mobile elements such as transposons, lysogenic phages and genomic islands. It is therefore not surprising that the structure of TnSmu1 changes as it transfers within the *S. mutans* species, particularly if its expression is lethal in other *S. mutans* hosts. Under normal conditions we anticipate that a very small sub-fraction of *S. mutans* cells would express TnSmu1 because of ImmR$_{Smu}$-mediated repression of the element. It is therefore likely that transfer of TnSmu1 from a donor strain to a recipient strain occurs at very low efficiency. Similar ICEs in other Firmicutes bacteria, like ICE*Bs1* and Tn*916*, exhibit considerable differences in conjugation rates. In the future we plan to explore conjugative transfer of this element between strains in greater detail.

DISCUSSION

In this study, we sought to determine the role of SMU_218/*immR*$_{Smu}$, a gene we previously identified as being 'essential' or non-mutable by Tn-seq. We hypothesized based on genomic context and protein domain similarities that *immR*$_{Smu}$ was either an antitoxin gene in a toxin-antitoxin system, or a transcriptional regulator involved in activation of the ICE TnSmu1. We provide evidence that *immR*$_{Smu}$ is not an antitoxin. Instead, we have concluded that *immR*$_{Smu}$ is a repressor that keeps TnSmu1 silent, except under certain environmental conditions. In exploring *immR*$_{Smu}$ we provide evidence that TnSmu1 can be induced, that it produces a circular intermediate, and the element, or parts of it, are distributed among *S. mutans* strains. We also uncovered that activation of TnSmu1 has a considerable impact on *S. mutans* physiology, leading to changes in gene expression, slowed growth and abnormal cell morphology.

Our conclusion that $immR_{Smu}$ is a repressor involved in the activation of TnSmu1 is based on several lines of evidence. Firstly, BLAST and other related bioinformatics searches show that $immR_{Smu}$ is a predicted helix-turn-helix DNA binding protein, and the downstream gene $immA_{Smu}$ is a predicted Zn^{2+} metalloprotease. BLAST searches also reveal that $immRA_{Smu}$ share similarities with other described transcriptional regulator and metalloprotease regulatory modules. Most importantly, the mobile element ICE*Bs1* contains a repressor, ImmR, that is cleaved by a protease ImmA, in a two gene organization like $immRA_{Smu}$ but in a different location within the ICE (Figure 8A) [11, 33]. Due to the similarities of the system described here, and ICE*Bs1 immRA*, we have named SMU_218 $immR_{Smu}$ and SMU_219 $immA_{Smu}$. Activation of ICE*Bs1* occurs when ImmR is cleaved by the protease ImmA, either through a cell-cell signaling pathway involving RapI and PhrI or via a RecA-dependent DNA damage response [11, 33]. TnSmu1 does not appear to contain a RapI/PhrI cell-cell signaling system but additional regulatory systems on top of $immRA_{Smu}$ cannot be ruled out, as TnSmu1 contains several hypothetical genes with unknown functions. ImmRA-like systems are encoded on other putative and studied mobile genetic elements. ICE*St1* and ICE*St3* contain a regulation module with a ImmR-like gene (*arp2*), a metalloprotease gene (*orfQ*) and a cI gene (*arp1*) [14]. In this module the metalloprotease gene, *orfQ*, is not downstream of the *immR*-like gene, *arp2*, but as with ICE*Bs1* these elements are activated by DNA damage, which is probably regulated via repressor proteolysis [14]. An ImmRA-like system is also required for the activation of a staphylococcal pathogenicity island SaPI3 [36]. For this regulation module, ImmR repression is partially alleviated when Sis (<u>S</u>aPI <u>i</u>nducer of <u>S</u>aPIs), a protein produced by other SaPIs, binds to ImmR but full activation only occurs with ImmA-mediated proteolytic cleavage of ImmR [36]. ImmRA modules are also encoded by lysogenic phages known to be activated by DNA damage [11]. These modules are therefore a common regulatory system governing activation of mobile genetic elements, and several have additional regulatory complexities in addition to ImmRA.

Aside from inducing excision of ICE elements, there is evidence that similar transcriptional regulator and metalloprotease modules are encoded in other systems. Recently, a system that parallels ImmRA was identified on CBASS (<u>c</u>yclic oligonucleotide <u>b</u>ased <u>a</u>nti-phage <u>s</u>ignaling <u>s</u>ystem) anti-phage systems [37]. Called CapH/CapP, this module regulates expression of a CBASS system in response to DNA damage. The transcriptional regulator CapH represses CBASS transcription until it is cleaved by the metallopeptidase CapP, which is stimulated by the presence of single-stranded DNA. Once the CBASS system is derepressed a cell killing pathway is induced which kills the host bacteria. Another ImmRA-like system, known as DdrO/IrrE, is encoded by *Deinococcus* spp. [38]. This system regulates a DNA damage response, with the repressor DdrO being cleaved by IrrE during radiation, leading to expression of DNA repair genes and an apoptotic-like cell death pathway [38]. There is a common theme among the ImmRA, CapHP and DdrO/IrrE modules in that they cause expression of genes upon sensing of DNA damage, with cleavage of a transcriptional repressor by an activated metalloprotease. In addition, both CapHP and DdrO/IrrE lead to host genome killing. That these systems have pathways that lead to cell-killing is notable because activation of TnSmu1 (via CRISPRi) caused growth defects and abnormal cell morphology. However, additional studies are warranted to determine if this is mediated directly or indirectly by TnSmu1.

Additional evidence that $immRA_{Smu}$ is a regulatory module was provided by RNA-seq which helped to identify genes activated in response to either depletion or deletion of $immR_{Smu}$. When $immR_{Smu}$ is repressed or deleted there is a clear and strong up-regulation of TnSmu1-associated genes by as much as 1000-fold. Although derepression of TnSmu1 genes by $ImmR_{Smu}$ would be most strongly confirmed by DNA-binding assays (*e.g.* electrophoretic mobility shift assay), activation of TnSmu1 genes upon loss of the ImmR repressor is consistent with similar systems described in ICE*Bs1*, CapH/CapP and DdrO/IrrE. In addition, deletion of

*immR*_{Smu} was found to activate excision/circularization of TnSmu1. Again, this would be consistent with a hypothesis that loss of the repressor leads to constitutive derepression of TnSmu1 followed by activation of the mobile element. Based on ICE*Bs1* ImmRA (and similar systems) we predict that ImmA$_{Smu}$ is a metalloprotease that is activated by DNA damage, leading to cleavage of ImmR$_{Smu}$ and derepession of TnSmu1. Although we do not provide direct evidence confirming this hypothesis, we do show that TnSmu1 circularization is induced in response to DNA damage by the genotoxic agent MMC. Taken together, activation by TnSmu1 as shown in RNA-seq studies and in response to MMC provides reliable evidence that ImmRA$_{Smu}$ functions as a regulatory module which has similarities to ICE*Bs1* ImmRA.

We were initially drawn to studying *immR*$_{Smu}$ because of Tn-seq data showing that the gene was 'essential'. With the evidence gathered here we are able to conclude that *immR*$_{Smu}$ is not a classical essential gene and does not participate in a core, required for survival, biological pathway. Instead, *immR*$_{Smu}$ appears 'essential' because deletion of *immR*$_{Smu}$ activates TnSmu1 leading to excision and eventual loss of the mobile element from the cell; thereby not making it possible to select a deletion strain. The observation that inactivation of nearby conjugation-related genes (SMU_197c and SMU_201c) allows *immR*$_{Smu}$ deletion provides evidence for this hypothesis. Here, disruption of the conjugation genes likely leads to TnSmu1/Δ*immR*$_{Smu}$ becoming 'trapped' in the cell, and selectable through antibiotic cassette replacement. However, we were able to describe significant impacts on cell growth, cell morphology and the transcriptome when *immR*$_{Smu}$ was depleted with CRISPRi. Despite in many cases ICEs carrying beneficial genes, such as virulence factors or antibiotic resistance genes, expression of these elements has been found to have a major impact on certain hosts. Induction of ICE*clc* in *Pseudomonas* spp. causes a sub-population of cells to differentiate into "transfer competent" cells that have arrested cell growth and cell lysis [39]. Activation of Tn*916* causes severe cell growth defects in *Enterococcus faecalis* and *Bacillus subtilis* [40]. Tn*916* only activates in a small percentage (0.1-3%) of cells and the lethality of Tn*916* was discovered by activating it in a much larger proportion of cells. The deleterious effects of Tn*916* were less impactful in cells lacking conjugation genes that reside in Tn*916*, and a gene, *yqaR* found within a defective phage-like element *skin* [40]. Intriguingly, Tn*916* can also cause cell death in *E. faecalis* and *B. subtilis*, species that do not encode for a *yqaR*-like gene. Future studies of the growth arrest/cell death mechanism in TnSmu1 would be helpful in understanding this generally conserved process for ICE transfer, and whether it is beneficial or costly for transfer efficiency.

We expect that under most conditions TnSmu1 will be quiescent because of ImmR$_{Smu}$-mediated repression of the mobile genetic element. This silence helps maintain vertical transmission of the element during non-stressful conditions. We anticipate that activation of TnSmu1 occurs when ImmA$_{Smu}$ becomes active and cleaves ImmR$_{Smu}$ leading to de-repression of TnSmu1 (Figure 8B). As with other ICEs, such as ICE*Bs1*, DNA damage causes activation of TnSmu1, as shown here with MMC treatment. For ICE*Bs1*, during DNA damage, ImmA is stimulated via an SOS (save our souls) response that causes RecA to activate ImmA cleavage of ImmR [11, 33]. Although our model of ImmRA$_{Smu}$-mediated regulation of TnSmu1 includes activation of ImmA by RecA, there may be substantial differences in the SOS response between *S. mutans* and *B. subtilis*. *Streptococcus* spp. have generally been thought to lack a classical SOS response because of work performed in *Streptococcus pneumoniae*, which lacks a LexA homologue, where genetic competence plays a central role in responding to DNA damage [41]. However, other streptococci such as *Streptococcus thermophilus* encode LexA-like repressors and competence development is antagonistic to the SOS response [42]. The *S. mutans* UA159 genome encodes two LexA-like repressors, SMU_1398 (*irvR*) and SMU_2027 (*hdiR*). IrvR does not participate in the SOS-response and instead is a stress-responsive biofilm regulator [43]. In comparison, *hdiR* is induced by SOS stress (MMC) in a *recA*-dependent manner [43]. Genetic

competence could play a role in an SOS-like response because *recA* is up-regulated as part of the late competence regulon [44]. Notably, other pathways may be involved in TnSmu1 regulation as RNA-seq studies show TnSmu1 up- or down-regulation in different conditions or mutant backgrounds [45–49]. Although our core model of TnSmu1 activation via $ImmR_{Smu}$ proteolysis borrows heavily from the ICE*Bs1* system we anticipate notable differences in how $ImmA_{Smu}$ becomes active.

To conclude, we show that *S. mutans* contains an ICE, TnSmu1, whose activation is likely controlled by an ImmRA-like regulatory module. Supporting this hypothesis is strong up-regulation of TnSmu1 genes and excision/circularization of the element upon loss or depletion of *immR_{Smu}*. The mobile element is also inducible via DNA damage, which is a conserved feature of ImmRA-like regulatory systems. Future studies will be directed to better understanding the steps involved in TnSmu1 activation, as well as identifying the mechanism that causes activation to be deleterious to host fitness and cell biology.

METHODS
Bacterial strains and culture conditions.
Streptococcus mutans strains were cultured from single colonies in Brain Heart Infusion (BHI, Difco) broth. Unless otherwise stated *S. mutans* was routinely cultured at 37°C in a 5% CO_2, microaerophilic atmosphere. *Escherichia coli* strains were routinely cultured in LB broth (Lennox formula; 10 g/L tyrptone, 5 g/L yeast extract and 5 g/L NaCl) at 37°C with aeration. Antibiotics were added to growth media at the following concentrations: kanamycin (1.0 mg/mL for *S. mutans*, 50 µg/mL for *E. coli*), spectinomycin (1.0 mg/mL for *S. mutans*), ampicillin (100 µg/mL for *E. coli*). A list of strains and plasmids (Table S5) and oligonucleotide primers (Table S6) can be found in the supplementary material.

Gene mutagenesis and plasmid cloning.
Standard DNA manipulation techniques were used to engineer deletion mutant strains [50]. A PCR ligation mutagenesis method was used to replace genes with non-polar kanamycin markers [51]. For each gene deletion, primers A and B were designed to amplify 500-600 bp upstream of the coding sequence (with ca. 50-bp overlapping the coding sequence of the gene). Primers C and D were designed to amplify 500 to 600 bp downstream of the coding sequence (with ca. 50-bp overlapping the coding sequence of the gene). Primers B and C contained *Bam*HI restriction enzyme sites for ligation of the AB and CD fragments to a non-polar kanamycin cassette digested from plasmid pALH124 [52]. Transformants were selected on BHI agar containing kanamycin. Double-crossover recombination, without introduction of nearby secondary mutations, was confirmed by PCR and Sanger sequencing using primers E and F, away from the site of recombination.

Plasmid cloning was conducted using the protein expression plasmid pBAD/His/A that contains an arabinose inducible promoter for tightly regulated protein production. PCR products for SMU_218, SMU_219 and SMU_218-219 were amplified and digested, before being ligated into digested pBAD/His/A plasmid. Correct in-frame insertion was verified with Sanger sequencing.

Transmission electron microscopy
Overnight cultures of CRISPRi strains were diluted 1:100 into 200 µL FMC-maltose without or with 0.025% xylose, and then incubated at 37°C in a 5% CO2 incubator for 16 h. Afterwards cells were rinsed with 0.1M sodium cacodylate buffer and then fixed in 3% glutaraldehyde overnight at 4˚C. The following day, cells were treated with 1.5% osmium tetroxide in the dark for 1 h at 4˚C. Afterwards, the cells were mixed in equal parts with 5% agarose in PBS, collected by centrifugation at 2000 x g, and cooled to 4°C. Small chunks of the bacterial pellet

plus agarose were then dehydrated in ethanol via the following steps: 30%, 50%, 70%, 80%, 90% each for 15 min, 99% 10 min, and then absolute ethanol 2 x 10 min. The dehydrated cells were embedded in an epoxy resin, sectioned and stained with uranyl acetate and lead citrate. Microscopy was conducted using a Hitachi H7600 transmission electron microscope.

Transposon interaction screen

To look for genetic interactions of $immR_{Smu}$ with other genes, a previously created Tn-seq library [4] was transformed with a $immRA_{Smu}$ deletion product, described above. Transformation of the library was completed in triplicate and each transformation reaction was screened on at least ten agar plates containing spectinomycin (transposon cassette) and kanamycin ($immRA_{Smu}$ deletion construct). Colonies that grew and were potentially viable $immRA_{Smu}$ deletions (plus a transposon cassette) were picked and plated onto fresh antibiotic-containing media. After this, colonies were cultured overnight and frozen in glycerol at -80°C. In addition, control experiments where conducted. Controls included attempting to delete $immRA_{Smu}$ with the deletion construct and selection of the Tn-seq library on kanamycin (to observe background spontaneous resistance).

Interaction screen genome sequencing analysis

Genomic DNA was isolated from strains using a MasterPure Gram Positive DNA (Epicentre) purification kit with modifications as previously described [53]. After DNA purification, total DNA concentration and purity was measured using a NanoDrop spectrophotometer (Thermo Fisher Scientific). Purified genomic DNA was sent to the Microbial Genome Sequencing Center, and samples were sequenced according to their protocol. Sequencing reads were compared to a *S. mutans* UA159 GenBank file (NC_004350.2) using Breseq. Following genomic sequencing, transposon insertion locations were verified with PCR and Sanger sequencing (primers available in Table S6). Correct deletion of $immRA_{Smu}$ was also verified using the primer pair 218_219E and 219F, followed by gel electrophoresis and comparison to expected product sizes.

qPCR measurements of TnSmu1 excision and circularization

Quantitative PCR (qPCR) was used to calculate the excision and circularization frequency of TnSmu1. Specific primers, which only create products during excision or circularization, were used in the qPCR assay (Table S6). Primers which amplified a chromosomal single copy gene, *sloR*, close to TnSmu1 were also used to standardize results. Primers were designed using the qPCR settings in the Primer3plus online application [54]. Cell samples were grown to mid-log in rich media and DNA was collected using a MasterPure Gram Positive DNA (Epicentre) purification kit with modifications as previously described [53]. Standard curves for each primer pair were generated using eight 10-fold dilutions of PCR products, starting with 10^8 copies/µl. Triplicates of standard curve DNAs, samples and cDNA controls were added to wells containing iQ SYBR Green Supermix (Bio-Rad) with primers (0.4 µM). Thermocycling was carried out using a CFX Touch Real-Time PCR Detection System (Bio-Rad) set to the following protocol: 40 cycles of 95°C for 10 s and 60°C for 45 s, with a starting cycle of 95°C for 30 s.

RNA-seq analysis

RNA was extracted from OD_{600} ~0.4-0.6 bacterial cultures using the RNeasy Mini Kit. RNA was extracted from biological triplicates of CRISPRi strains carrying a short guide RNA (sgRNA) targeting SMU_218 in the presence or absence of the dCas9 inducing molecule xylose. RNA was also extracted from wild-type UA159, SMU_197c::Tn $\Delta immRA_{Smu}$ and SMU_201c::Tn $\Delta immRA_{Smu}$. Next, RNA was sent to the Microbial Genome Sequencing Center who generated RNAseq libraries and sequenced samples according to their protocol. After sequencing, read counts were aligned to the *S. mutans* UA159 genome using bioinformatics tools hosted on the

Galaxy server maintained by the Research Computing Center at the University of Florida. Gene expression changes between samples were quantified with Degust (http://degust.erc.monash.edu/) using the edgeR methodology. The original RNA-seq data from this study was uploaded to the GEO database (https://www.ncbi.nlm.nih.gov/geo/) with the accession number GSE202804.

TnSmu1 distribution bioinformatics
Upon acknowledging the TnSmu1 operon presence in *S. mutans* UA159 we isolated the gene sequences of all the individual genes by constructing a BED type file, which was parsed to the getfasta command from the Bedtools suite [55]. The resulting individual FASTA files with each gene sequence were blasted against a collection of over 600 clinical isolate genomes in our possession. The blast output was then filtered with the objective of keeping only those strains showing both a percent identity greater than 75% and a ratio between the query length and the sequence length greater than 0.9. From this filtered blast output we isolated the PID column for each isolate across all genes and generated a matrix as input for the heatmap. In parallel we conducted a Multi-locus Sequence Typing (MLST) analysis with all resulting strains in the curated blast output. The MLST sequence was then parsed to PHYML [56] to construct a Maximum-likelihood phylogeny tree. Finally the phylogenies and the heatmap were plot together in R Statistical Language [57] with packages ggplot2 [58] and GGTREE [59].

Data availability statement: the data that supports the findings of this study are available from the corresponding author. Gene expression data was uploaded to the GEO database with the accession number GSE202804.

ACKNOWLEDGMENTS
We thank Alan Grossman at the Massachusetts Institute of Technology for helpful feedback related to this project. This work was supported by the National Institute of Dental and Craniofacial Research (NIDCR) grant DE029882 awarded to R.C.S.

References

1. **Loesche WJ**. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 1986;50:353–80.
2. **Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, *et al.*** Dental caries. *Nat Rev Dis Primer* 2017;3:17030.
3. **Sheiham A, James WPT**. Diet and Dental Caries: The Pivotal Role of Free Sugars Reemphasized. *J Dent Res* 2015;94:1341–1347.
4. **Shields RC, Zeng L, Culp DJ, Burne RA**. Genomewide Identification of Essential Genes and Fitness Determinants of *Streptococcus mutans* UA159. *mSphere* 2018;3:e00031-18.
5. **Shields RC, Walker AR, Maricic N, Chakraborty B, Underhill SAM, *et al.*** Repurposing the *Streptococcus mutans* CRISPR-Cas9 System to Understand Essential Gene Function. *PLoS Pathog* 2020;16:e1008344.
6. **Letunic I, Bork P**. 20 years of the SMART protein domain annotation resource. *Nucleic Acids Res* 2018;46:D493–D496.
7. **Jongeneel CV, Bouvier J, Bairoch A**. A unique signature identifies a family of zinc-dependent metallopeptidases. *FEBS Lett* 1989;242:211–214.
8. **Sevin EW, Barloy-Hubler F**. RASTA-Bacteria: a web-based tool for identifying toxin-antitoxin loci in prokaryotes. *Genome Biol* 2007;8:R155.
9. **Galkin VE, Yu X, Bielnicki J, Ndjonka D, Bell CE, *et al.*** Cleavage of bacteriophage lambda cI repressor involves the RecA C-terminal domain. *J Mol Biol* 2009;385:779–787.
10. **Bellanger X, Morel C, Decaris B, Guédon G**. Derepression of Excision of Integrative and Potentially Conjugative Elements from *Streptococcus thermophilus* by DNA Damage Response: Implication of a cI-Related Repressor. *J Bacteriol* 2007;189:1478–1481.
11. **Bose B, Auchtung JM, Lee CA, Grossman AD**. A conserved anti-repressor controls horizontal gene transfer by proteolysis. *Mol Microbiol* 2008;70:570–582.
12. **Maruyama F, Kobata M, Kurokawa K, Nishida K, Sakurai A, *et al.*** Comparative genomic analyses of Streptococcus mutans provide insights into chromosomal shuffling and species-specific content. *BMC Genomics* 2009;10:358.
13. **Waterhouse JC, Swan DC, Russell RRB**. Comparative genome hybridization of Streptococcus mutans strains. *Oral Microbiol Immunol* 2007;22:103–110.
14. **Carraro N, Libante V, Morel C, Decaris B, Charron-Bourgoin F, *et al.*** Differential regulation of two closely related integrative and conjugative elements from *Streptococcus thermophilus*. *BMC Microbiol* 2011;11:238.
15. **Liu M, Li X, Xie Y, Bi D, Sun J, *et al.*** ICEberg 2.0: an updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res* 2019;47:D660–D665.
16. **Wozniak RAF, Waldor MK**. Integrative and conjugative elements: mosaic mobile genetic elements enabling dynamic lateral gene flow. *Nat Rev Microbiol* 2010;8:552–563.
17. **Koyanagi S, Lévesque CM**. Characterization of a *Streptococcus mutans* intergenic region containing a small toxic peptide and its cis-encoded antisense small RNA antitoxin. *PloS One* 2013;8:e54291.
18. **Patel S, Weaver KE**. Addiction Toxin Fst Has Unique Effects on Chromosome Segregation and Cell Division in Enterococcus faecalis and Bacillussubtilis. *J Bacteriol* 2006;188:5374–5384.
19. **Weaver KE, Weaver DM, Wells CL, Waters CM, Gardner ME, *et al.*** Enterococcus faecalis Plasmid pAD1-Encoded Fst Toxin Affects Membrane Permeability and Alters Cellular Responses to Lantibiotics. *J Bacteriol* 2003;185:2169–2177.
20. **Wu C, Cichewicz R, Li Y, Liu J, Roe B, *et al.*** Genomic Island TnSmu2 of Streptococcus mutans Harbors a Nonribosomal Peptide Synthetase-Polyketide Synthase Gene Cluster Responsible for the Biosynthesis of Pigments Involved in Oxygen and H2O2 Tolerance. *Appl Environ Microbiol* 2010;76:5815–5826.

21. **Crosby J, Crump MP**. The structural role of the carrier protein – active controller or passive carrier. *Nat Prod Rep* 2012;29:1111–1137.
22. **Li J, Wang H-T, Wang W-T, Zhang X-R, Suo F, *et al.*** Systematic analysis reveals the prevalence and principles of bypassable gene essentiality. *Nat Commun* 2019;10:1002.
23. **van Leeuwen J, Pons C, Tan G, Wang JZ, Hou J, *et al.*** Systematic analysis of bypass suppression of essential genes. *Mol Syst Biol* 2020;16:e9828.
24. **Hart EM, Silhavy TJ**. Functions of the BamBCDE Lipoproteins Revealed by Bypass Mutations in BamA. *J Bacteriol*;202:e00401-20.
25. **Wang Y, Cao W, Merritt J, Xie Z, Liu H**. Characterization of FtsH Essentiality in Streptococcus mutans via Genetic Suppression. *Front Genet* 2021;12:659220.
26. **Pichoff S, Du S, Lutkenhaus J**. The bypass of ZipA by overexpression of FtsN requires a previously unknown conserved FtsN motif essential for FtsA–FtsN interaction supporting a model in which FtsA monomers recruit late cell division proteins to the Z ring. *Mol Microbiol* 2015;95:971–987.
27. **Anupama K, Leela JK, Gowrishankar J**. Two pathways for RNase E action in Escherichia coli in vivo and bypass of its essentiality in mutants defective for Rho-dependent transcription termination. *Mol Microbiol* 2011;82:1330–1348.
28. **Krüger L, Herzberg C, Rath H, Pedreira T, Ischebeck T, *et al.*** Essentiality of c-di-AMP in Bacillus subtilis: Bypassing mutations converge in potassium and glutamate homeostasis. *PLOS Genet* 2021;17:e1009092.
29. **Johnson CM, Grossman AD**. Integrative and Conjugative Elements (ICEs): What They Do and How They Work. *Annu Rev Genet* 2015;49:577–601.
30. **Wright LD, Johnson CM, Grossman AD**. Identification of a Single Strand Origin of Replication in the Integrative and Conjugative Element ICEBs1 of Bacillus subtilis. *PLOS Genet* 2015;11:e1005556.
31. **Casjens SR, Hendrix RW**. Bacteriophage lambda: Early pioneer and still relevant. *Virology* 2015;479–480:310–330.
32. **Beaber JW, Hochhut B, Waldor MK**. SOS response promotes horizontal dissemination of antibiotic resistance genes. *Nature* 2004;427:72–74.
33. **Auchtung JM, Lee CA, Monson RE, Lehman AP, Grossman AD**. Regulation of a Bacillus subtilis mobile genetic element by intercellular signaling and the global DNA damage response. *Proc Natl Acad Sci* 2005;102:12554–12559.
34. **Lemme A, Sztajer H, Wagner-Döbler I**. Characterization of mleR, a positive regulator of malolactic fermentation and part of the acid tolerance response in Streptococcus mutans. *BMC Microbiol* 2010;10:58.
35. **Walker AR, Shields RC**. Investigating CRISPR spacer targets and their impact on genomic diversification of Streptococcus mutans. 2022;2022.04.14.488371.
36. **Haag AF, Podkowik M, Ibarra-Chávez R, Gallego del Sol F, Ram G, *et al.*** A regulatory cascade controls Staphylococcus aureus pathogenicity island activation. *Nat Microbiol* 2021;6:1300–1308.
37. **Lau RK, Enustun E, Gu Y, Nguyen JV, Corbett KD**. *A conserved signaling pathway activates bacterial CBASS immune signaling in response to DNA damage*. Preprint; Biochemistry. Epub ahead of print 27 April 2022. DOI: 10.1101/2022.04.27.489752.
38. **Devigne A, Ithurbide S, Bouthier de la Tour C, Passot F, Mathieu M, *et al.*** DdrO is an essential protein that regulates the radiation desiccation response and the apoptotic-like cell death in the radioresistant Deinococcus radiodurans bacterium. *Mol Microbiol* 2015;96:1069–1084.
39. **Delavat F, Mitri S, Pelet S, van der Meer JR**. Highly variable individual donor cell fates characterize robust horizontal gene transfer of an integrative and conjugative element. *Proc Natl Acad Sci* 2016;113:E3375–E3383.

40. **Bean EL, McLellan LK, Grossman AD**. *Activation of the integrative and conjugative element Tn* 916 *causes growth arrest and death of host bacteria*. Preprint; Microbiology. Epub ahead of print 2 April 2022. DOI: 10.1101/2022.04.01.486793.

41. **Prudhomme M, Attaiech L, Sanchez G, Martin B, Claverys J-P**. Antibiotic Stress Induces Genetic Transformability in the Human Pathogen Streptococcus pneumoniae. *Science* 2006;313:89–92.

42. **Boutry C, Delplace B, Clippe A, Fontaine L, Hols P**. SOS response activation and competence development are antagonistic mechanisms in Streptococcus thermophilus. *J Bacteriol* 2013;195:696–707.

43. **Marx P, Sang Y, Qin H, Wang Q, Guo R, *et al.*** Environmental stress perception activates structural remodeling of extant Streptococcus mutans biofilms. *Npj Biofilms Microbiomes* 2020;6:1–11.

44. **Khan R, Rukke HV, Høvik H, Åmdal HA, Chen T, *et al.*** Comprehensive Transcriptome Profiles of Streptococcus mutans UA159 Map Core Streptococcal Competence Genes. *mSystems* 2016;1:e00038-15.

45. **Dmitriev A, Mohapatra SS, Chong P, Neely M, Biswas S, *et al.*** CovR-Controlled Global Regulation of Gene Expression in *Streptococcus mutans*. *PLoS ONE* 2011;6:e20127.

46. **Chattoraj P, Banerjee A, Biswas S, Biswas I**. ClpP of *Streptococcus mutans* Differentially Regulates Expression of Genomic Islands, Mutacin Production, and Antibiotic Tolerance. *J Bacteriol* 2010;192:1312–1323.

47. **Kaspar J, Shields RC, Burne RA**. Competence Inhibition by the XrpA Peptide Encoded Within the *comX* Gene of *Streptococcus mutans*. *Mol Microbiol* 2018;109:345–364.

48. **Kajfasz JK, Katrak C, Ganguly T, Vargas J, Wright L, *et al.*** Manganese Uptake, Mediated by SloABC and MntH, Is Essential for the Fitness of *Streptococcus mutans*. *mSphere*;5. Epub ahead of print 2020. DOI: 10.1128/mSphere.00764-19.

49. **Ahn S-J, Rice KC**. Understanding the *Streptococcus mutans* Cid/Lrg System through CidB Function. *Appl Environ Microbiol* 2016;82:6189–6203.

50. **Sambrook J, Russell D**. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 2001.

51. **Lau PCY, Sung CK, Lee JH, Morrison DA, Cvitkovitch DG**. PCR ligation mutagenesis in transformable streptococci: application and efficiency. *J Microbiol Methods* 2002;49:193–205.

52. **Ahn S-J, Burne RA**. The *atlA* operon of *Streptococcus mutans*: role in autolysin maturation and cell surface biogenesis. *J Bacteriol* 2006;188:6877–88.

53. **Shields RC, Mokhtar N, Ford M, Hall MJ, Burgess JG, *et al.*** Efficacy of a Marine Bacterial Nuclease against Biofilm Forming Microorganisms Isolated from Chronic Rhinosinusitis. *PloS ONE* 2013;8:e55339–e55339.

54. **Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, *et al.*** Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Res* 2007;35:W71-4.

55. **Quinlan AR, Hall IM**. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinforma Oxf Engl* 2010;26:841–842.

56. **Guindon S, Gascuel O**. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 2003;52:696–704.

57. R: a language and environment for statistical computing. https://www.gbif.org/tool/81287/r-a-language-and-environment-for-statistical-computing (accessed 12 April 2022).

58. **Valero-Mora PM**. ggplot2: Elegant Graphics for Data Analysis. *J Stat Softw* 2010;35:1–3.

59. **Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y**. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol Evol* 2017;8:28–36.

Figure 1. Growth profile and morphological changes in *immR*_{Smu} depleted strains. A CRISPRi strain with an sgRNA targeting $immR_{Smu}$ was generated and cultured in CDM containing xylose to repress $immR_{Smu}$ gene expression. In these experiments a strain with an sgRNA targeting *lacG* was constructed and serves as a control. (A) Cell densities were measured every 30 min for 16 h with the control strain depicted as light blue dots and the sgRNA-$immR_{Smu}$ strain depicted as dark blue dots. (2) Morphological changes were examined with transmission electron microscopy. Depletion of $immR_{Smu}$ led to a sub-population of cells to become noticeably larger (white arrows) with aberrant cell division.

Figure 2. Measuring the *S. mutans* transcriptome response to *immR*_{Smu} repression with CRISPRi-RNAseq. (A) Volcano plot of differential gene expression of *S. mutans* in a CRISPRi strain targeting *immR*$_{Smu}$ (with and without xylose *dcas9* induction). The expression difference was considered significant for a log2 fold change > 2 (vertical dotted line) and p value < 0.05 (horizontal dotted line). TnSmu1 genes are shown as light green dots. (B) RNAseq reads were visualized using the Integrated Genome Viewer, with the control sample shown in green, and the *immR*$_{Smu}$-targeting sample shown in blue. Read coverage is shown across the TnSmu1 mobile element. The *fstSm* and 217-218 intergenic region (IGR) small RNAs are highlighted. (C) The number of genes up- or downregulated with a Cluster of Orthologous Group (COG) assignment is indicated in a bar chart.

Figure 3. A transposon interaction screen identifies gene mutations that bypass the essentiality of *immR*Smu. (A) An *immRA*Smu::*aphA3* mutagenesis fragment was transformed into a *S. mutans* Tn-seq library with selection on antibiotic-containing agar. Deletion of *immRA*Smu is not possible in a wild-type strain of *S. mutans* (top panel). White *S. mutans* colonies were obtained when *immRA*Smu was transformed into the Tn-seq library (bottom panel). (B) After genome sequencing analysis, several strains were identified that allow *immRA*Smu deletion. Each strain, with the corresponding transposon insertion site, is shown in a single growth analysis chart (dark line, wild-type strain; light blue line, double mutant strain).

Figure 4. Mutations that bypass the essentiality of *immRA*_{Smu} cause an increase in genome sequencing reads from the ICE TnSmu1. Read coverage depth for both a wild-type (A) and a double mutant strain Δ*immRA*_{Smu}::*aphA3* SMU_197c::Tn (B) is shown. Genome sequencing read depth was substantially higher across the TnSmu1 region in the double mutant background. (C) Sequencing reads were visualized with the Integrated Genome Viewer for only the TnSmu1 region (and immediate up- and downstream genes). Read coverage is ca. 10-fold higher across the entire TnSmu1 region, except for the deleted *immRA*_{Smu} genes. The location of each transposon insertion, within putative conjugation genes, which allowed *immRA*_{Smu} deletion is shown as upside-down triangles. The TnSmu1 region is annotated with putative functions related to excision/recombination (integrase), conjugation, T-DNA formation (relaxase), strand nicking (*oriT*) and attachment sites (*attL* and *attR*).

Figure 5. TnSmu1 is capable of circularization and excision. Quantitative PCR was employed to measure the relative abundance of circular (A) and excised (B) copies of TnSmu1. The number of copies was compared to a nearby single-copy chromosomal gene, *sloR*. Induction of TnSmu1 circularization was also measured by inducing TnSmu1 with the genotoxic agent mitomycin C (MMC). The impact of different concentrations of MMC was quantified using a growth assay (C), and then different doses of MMC were used to induce TnSmu1 circularization (D).

Figure 6. RNA-seq analysis of strains lacking *immRA*Smu. RNA was extracted from
SMU_197c::Tn Δ*immRA*Smu and SMU_201c::Tn Δ*immRA*Smu cultures grown to OD$_{600}$ ~0.4, and
this RNA was then processed for RNA-seq. (A) Comparison of gene expression (log$_2$ fold
change) between SMU_197c::Tn Δ*immRA*Smu and SMU_201c::Tn Δ*immRA*Smu. Pearson's
correlation coefficients between transcriptomes are presented. Gene expression of TnSmu1,
Type I-C CRISPR-Cas and the *mle* locus are highlighted on the chart. (B) Heat map of selected
bacterial transcripts that are enriched or depleted in the chosen strains. Colors represent the
log$_2$ fold change compared to control conditions with a key shown to the right of the heat map.

Figure 7. TnSmu1 gene distribution across sequenced *S. mutans* genomes. *S. mutans* strains were searched for TnSmu1 genes, comparing to those carried on *S. mutans* UA159. Strains are organized by strain relatedness (*y*-axis) and gene ID is shown on the *x*-axis (SMU_191c to SMU_219). The percent sequence similarity for each gene is shown (compared to the gene on *S. mutans* UA159 TnSmu1). Red areas depict a lack of a TnSmu1 gene on a strain genome.

Figure 8. Comparison of ICE*Bs1* with TnSmu1 and proposed model for ImmRA$_{Smu}$ regulation of TnSmu1. (A) The genetic organization of ICE*Bs1* and TnSmu1 is shown. Genes are colored according to function with their names above or below. Genes with unknown functions are colored white. Both ICEs integrate at tRNA-leu. Note that the *immRA*$_{Smu}$ regulatory module is in a different location compared with ICE*Bs1*. (B) A diagram depicting the ImmRA$_{Smu}$ regulatory pathway for TnSmu1 activation is shown. Our model draws heavily from ICE*Bs1* ImmRA, and closely related systems, where DNA damage leads to activation of the protease ImmA. Activated ImmA cleaves the ImmR repressor leading to activation of TnSmu1. It is likely that stochasticity causes sub-population responses so that TnSmu1 activation can lead to transfer of the element, cell arrest/cell death and other phenotypes. The SOS pathway, leading to RecA-mediated activation of ImmA$_{Smu}$, is not well known in *S. mutans* and will need to be investigated further. Yet to be discovered signals may also lead to ImmA$_{Smu}$-mediated cleavage of ImmR$_{Smu}$.