Proliferation drives quorum sensing of microbial products in human macrophage populations

Nadia Rajab[1], Linden J. Gearing[2,3], Ruqian Lyu[4,5], Yair D.J. Prawer[1], Paul W. Angel[1], Sean M.

Grimmond[6], Andrew L. Laslett[7], Davis J. McCarthy[4,5], and Christine A. Wells[1#]

1. Centre for Stem Cell Systems, Department of Anatomy and Physiology, Faculty of Medicine,

Dentistry and Health Sciences, The University of Melbourne, VIC, 3010, Australia.

2. Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical Research,

Clayton, VIC, 3168, Australia.

3. Department of Molecular and Translational Sciences, Monash University, Clayton, VIC, 3168,

Australia.

4. Bioinformatics and Cellular Genomics, St Vincent's Institute of Medical Research, Fitzroy,

Australia

5. Melbourne Integrative Genomics, School of Biosciences – School of Mathematics & Statistics,

Faculty of Science, University of Melbourne, Melbourne, Australia

6. University of Melbourne Centre for Cancer Research, Faculty of Medicine, Dentistry and

Health Sciences, The University of Melbourne, VIC, 3010, Australia.

7. Australian Regenerative Medicine Institute, Monash University, Melbourne, VIC 3800,

Australia.

#Correspondence to CAW wells.c@unimelb.edu.au

20
Author Contributions

N.R. experimental design, data generation and analysis, writing

L.J.G. data analysis

R.L. SNP deconvolution and analysis

Y.D.J.P library preparation, data processing

P.W.A. data analysis

S.M.G experimental resources

A.L.L. experimental resources

D.M. experimental resources

C.A.W. experimental design, data analysis, writing, project funding

Conflict of Interests

33   NR and YDJP received training and subsidised reagents from 10X Genomics for the preparation

34   of samples described in this manuscript. The company did not seek to influence experimental

35   design, or interpretation of the data in this manuscript. No COI is declared for the other authors.

Funding

38   The project was run as part of the 10X Genomics Millenium Science Start Single Cell Fellowship

39   to NR and YDJP. C.A.W. is funded by NHMRC Synergy grant APP1186371.

Acknowledgements

The authors thank the Genomics Platform Group at The University of Melbourne Centre for

Cancer Research (UMCCR) sequencing core for sequencing of single-cell libraries, and Prof. Oliver

Hofmann for computational workflows and project discussions. The authors thank Dr Catherine

King (10x Genomics) and Dr Paul Gooding (Millenium Science) for their training and assistance in

preparation of 10x libraries. The authors thank Dr Jarny Choi and Dr Michael Clark, University of

Melbourne, for reading and commenting on the manuscript.

Abstract

Macrophages coordinate the initial host inflammatory response to tissue infection, as well as
mediating the reparative phase, by producing growth factors that promote tissue repair. One
model of this functional dichotomy is that peripherally recruited monocyte-derived macrophages
drive acute inflammatory responses to infection, whereas tissue-resident macrophages are
responsible for tissue repair. Alternatively, inflammation and repair may be inter-dependent
molecular programs, such that both recruited and resident cells have equivalent capacity to
contribute. Repeated exposure to pathogenic challenge results in innate tolerance, which may
also alter the contributions of discrete macrophage populations to inflammation or repair. In this
study a village model of tissue resident and recruited macrophages was created using induced
pluripotent stem cell-derived macrophages and peripheral blood monocyte-derived
macrophages, respectively. Population responses to repeated exposure to lipopolysaccharide
were assessed with single-cell RNA sequencing and donors demultiplexed with Vireo. A subset of
genes escaped classical tolerance programs in the iPSC, but not monocyte-derived macrophages,
and this was associated with differences in their proliferative capacity. This suggests that
targeting the proliferative resident macrophages would be most effective to limit inflammatory
signaling.

Key words:

Introduction

Multiple tissue-resident macrophage subsets have been described and defined by roles that are specialized to their tissue of residency, such as bone resorption by osteoclasts[1], neural pruning by microglia[2], lipid-associated macrophages[3], stem cell licensing in bone marrow[4] and intestine[5], or iron recycling in splenic macrophages[6]. Differences between macrophage subsets have also been described in the context of ontogeny (reviewed in [7]), anatomical location, or molecular phenotype[8]. Given the diverse roles of macrophages in tissue development, homeostasis, and in host defense, there is considerable interest in understanding mechanisms that might target macrophage subsets in tissue inflammation or repair.

Functional differences between tissue-resident macrophages (TRM) and monocyte-derived macrophages (MDM) recruited in response to inflammation are controversial. In some models, such as brain, the resident macrophages (microglia) play a neuroprotective role, while recruited MDM are thought to be the inflammatory mediator[9]. In contrast, others have described diverse monocyte and macrophage subsets associated with age-related inflammation, cancer or trauma[10], as well as inflammatory phenotypes associated with differences in IRF8 signaling across these subsets[11]. If macrophages have predefined roles in infection or inflammation, then this observed heterogeneity might well be explained by mixing of ontogenically and phenotypically distinct populations, exemplified by the progression of liver disease[12]. Alternatively, if both resident and recruited macrophage populations are intrinsically heterogeneous, then quorum-sensing will lead to roles that are randomly assigned across all populations, ensuring that the capacity for inflammation and repair is available regardless of the origin of the responding macrophage[13].

This idea of quorum sensing is supported by early studies of stochasticity of macrophage responsiveness showing heterogeneity in gene expression after exposure to lipopolysaccharide (LPS). Examples of this have been described in studies showing variability of LPS responses in clones of murine RAW264.7 cells[13,14]. In the first study, subclones demonstrated varying levels of gene expression after LPS stimulation, suggesting the coexistence of

macrophage subsets that could be propagated as subpopulations. In the second study,
heterogeneity in the timing of RelA recruitment to the TNF promoter in a RAW264.7 reporter line
was observed after LPS stimulation, but variability between cells could also be explained by
population density. These studies demonstrate that multiple mechanisms impact on the timing,
duration, and amplitude of macrophage activation, even in simple models receiving a highly
polarizing inflammatory signal, such as LPS.

It is obvious that, in a tissue setting, macrophages will be integrating signals from the local
environment and it is difficult to deconvolute the influence of these signals from ontogeny over
the course of an inflammatory event. Tissue culture models offer an opportunity to address cell-
intrinsic sources of macrophage diversity. Macrophages derived from human induced pluripotent
stem cells (PSCM) present new opportunities to model tissue residency and factors that shape
human macrophage development and activation. PSCMs are an excellent model of tissue
residency because they share molecular ontogeny with TRM [15,16] and are functionally equivalent
to TRM in engraftment models in brain and lung[17]. A recent comparison of macrophage
transcriptomes in our own Myeloid Atlas confirmed that PSCMs shared the transcriptional
phenotype of primary TRM including tissues such as brain, joint and liver[18].

Given PSCMs offer an *in vitro* representation of tissue-resident cells, they provide a relevant
experimental model to investigate sources of population heterogeneity in bacterial responses.
While others[19] have shown that PSCMs recognize and respond to LPS, it is not known whether
they can be tolerized on repeated exposure, which would more accurately mimic the regulatory
program seen after an infection than a single high dose pathogen exposure. We were also curious
to see how coordinated acute responses to LPS were across human macrophage populations, as
perhaps early versus late responders within a population could explain some of the previously
described heterogeneity in trained monocytes[20]. Here, we use single cell RNA-sequencing to
investigate population responses to high-dose LPS activation and then re-stimulation using a
coculture, or village, model of PSCMs and MDMs, as models of tissue residency versus recruited
macrophages respectively.

Results

<u>Establishing a 'village' of macrophages to assess macrophage heterogeneity at the single cell</u>

<u>level.</u>

The transcriptional heterogeneity of macrophage subsets was evident even in the unstimulated

populations (Figure 1). The use of a village model removed population density as a possible driver

of heterogeneity (Figure 1A), as all the cells were exposed to the same local environment,

maintained in CSF-1 (C, control), or exposed to LPS for 2 hours (A, acute) or 18 hours (R, rest),

Tolerised (T) macrophages were acutely stimulates as per (A and R), then washed after 18hours,

and rested for 2 hours then restimulated for 2 hours with LPS. PSCM and MDM formed distinct

clusters across the time course that were not explained by fractions of undifferentiated or

nonresponsive cells – MDM and PSCMs share similar phenotypes of CD14+ CD16low populations

after several days culture in CSF1 (Figure 1B). Cells collected at each time point were barcoded

for demultiplexing after single cell RNA-sequencing and donor origin of the cell was determined

by SNP deconvolution, to confirm that the heterogeneity was not driven by donor

(Supplementary Figure 1A). LPS exposure was a strongly polarizing signal in the acute time

periods, but population heterogeneity increased in the resting and restimulated conditions

(Figure 1C, D, E).

PSCM expressed hallmark genes of tissue resident macrophages (Figure 1F, Supplementary Table

1), including transcription factors associated with terminal macrophage differentiation, such as

MAF [21], and markers of perivascular and lymphatic-associated macrophages including *CD169*/

SIGLEC1, *LYVE1* and *FOLR2* [22,23]. PSCMs were transcriptionally similar to MDM and primary TRM

(Supplementary Figure 1B and C) using the benchmarking tool in the Stemformatics Myeloid

Atlas. MDM subsets were distinguished by expression of immunoregulatory glycoproteins

GPNMB, a negative regulator of TLR signaling [24], macrophage migratory factors such as *FUCA1*,

as well as chemokine receptors. However, the most striking difference between MDM and PSCM

clusters was the presence of a proliferative subset of PSCM in the control, acute and tolerized

conditions (Figure 1E).

The observations of ~40% of PSCM and <10% of MDM proliferating in the control condition is

consistent with the CSF1 responsiveness of tissue resident vs recruited macrophages. PSCM and

MDM were cultured in the presence of recombinant CSF1, which is a proliferative signal to tissue-

resident macrophages and essential for the maintenance of bone marrow and circulating

monocytes (reviewed in [25]). Monocytes are postmitotic cells in circulation, although a small

fraction have been identified proliferative in response to CSF1[26]. Just 10% of the PSCM were

proliferating by the resting or tolerized conditions (Figure 1E), consistent with a pro-survival but

antiproliferative consequence of TLR signaling. Both proliferating and nonproliferating PSCM

subsets expressed equivalent levels of components of the TLR4-LPS pathway, at similar levels to

the MDM control subsets (Figure 2A).

<u>Proliferating PSCM escape LPS tolerance</u>.

LPS is a synchronizing signal that results in high expression of core inflammatory mediators in all

cells receiving the signal. These included *TNF, IL1B, IL6, CXCL1, CXCL2, CXCL8, CXCL10, CCL3,*

CCL4L2, and members of the shared interleukin/TLR signaling pathway including *NFKBIA* and

NFKB1 (Figure 2, Supplementary Table 1)*,* and negative regulators of cytokine signaling such as

SOCS1 and *SOCS3.* These are predominantly targets of the TLR-MyD88-NF-κB pathway,

demonstrating that the PSCM expressed an active TLR4 signaling cascade. After 18 hours

activation, most inflammatory mediators have reached a baseline expression level in both

macrophage populations. Restimulation with LPS resulted in hypo-responsiveness in both MDM

and PSCM, particularly marked for *TNF* expression, and genes that are associated with TNF

signaling, including *NFKB1, NAMPT, TNFAIP1, 2* and *8, CXCL2, DUSP2,* and *TNFSF9*. These data

confirm that PSCM are LPS responsive and undergo NF-κB-associated tolerance to repeated LPS

exposure.

PSCM did not completely mimic the monocyte-derived tolerance program. PSCMs but not MDMs

expressed growth factors on LPS restimulation, including *FLT1, PDGFA*, *VEGFA*, *CSF1*, *CSF1R, CSF2*,

IL32, *IL36,* and *TGFB1* (Figure 2 and Supplementary Table 1). Less than 50% of PSCM were

recruited to the acute inflammatory response mediated by the MyD88-independent/IRF-

dependent pathway, including *IRF1, OAS1, IFIT2* compared to over 80% of MDM (Figure 2D).

Review of the pathways differentially expressed by MDM compared to PSCM confirmed that the

Reactome IFN signaling pathway scored significantly higher across the MDM time course (Figure

2E). Other targets of intracellular TLR signaling were absent entirely in PSCM, including *IFNB1*,

which was inducible in MDM but not PSCM populations. This may have implications for

proliferation differences between MDM and PSCM, as CSF1R is targeted by IFNB signaling in

monocytes[27], and further evidences negative feedback of CSF1 signaling in MDM after LPS

activation.

Genes that were completely tolerized in the MDM such as *IL1B*, and chemokines such as *CCL3*

and *CCL7* were expressed at high levels in acute or resting PSCM, and in the restimulated

conditions. These genes were expressed regardless of proliferating status and are predominantly

targets of post-transcriptional (AU-response element) regulation, which may be a mechanism

that supports acute activation in a macrophage that is also dividing (Figure 2, Supplementary

Table 2).

The current model of LPS tolerance is that of an altered chromatin landscape where NF-κB p50

homodimers are stabilized by BCL3, preventing recruitment and binding of NF-κB heterodimers

to LPS responsive cytokines[28]. BCL3 nuclear translocation and activation is regulated by

proliferation[29] and is differentially regulated in PCSM but not MDM across the acute phase of the

LPS time course, so it is plausible that proliferation of PSCM resets the BCL3-dependant program

of LPS tolerance, allowing for ongoing expression of inflammatory mediators in a subpopulation

of cells.

Discussion

In this study we tested two models of cell-intrinsic responses to LPS, identifying proliferation as a driver of macrophage heterogeneity in a minor subset of the MDM, whereas the proliferative subset represented about half of the PSCM population. Endotoxin tolerance in restimulated macrophages took two forms – either in the degree of induction of an inflammatory gene on re-stimulation with LPS or in the proportion of cells capable of expression of an inflammatory mediator. It was the 18 hour-stimulated and restimulated PSCM groups that continued to express cytokines, such as *IL1B*, and chemokines at a high level, possibly because these escaped tolerance in the cells proliferating on first exposure to LPS. TRM have the capacity to self-renew[30]. This homeostatic self-sufficiency of TRM enables the population to be maintained without requiring constant influx of monocytes (Figure 3).

Prior exposure to a TLR ligand causes hypo-responsiveness of macrophage populations to subsequent stimulations, and LPS/TLR4 drives the largest cohort of tolerized genes[31]. This so-called endotoxin tolerance is regulated by recruitment and stabilization of NF-κB p50 to inflammatory gene promoters, preventing engagement with the transcriptional machinery and silencing the locus[28]. Likewise, prior exposure to low dose pathogenic signals, cytokines, or apoptotic cells can induce 'trained' responses in innate immune cells, even days after the initial exposure (reviewed in [32]). While a key difference between the monocyte-derived MDM and laboratory-derived PSCM is the possibility of prior pathogen exposure in monocytes or precursors[20], we did not observe any evidence of so-called inflammatory macrophage signatures in the naïve MDM in this study. However, expression of IFN-related pathways in acutely activated MDM is consistent with 'training' of MYD88-independent signaling in the MDM rather than PSCM. Activation of IRF-related pathways appears to be part of the monocyte-macrophage transition[33], but was not recapitulated in the iPSC-derivation protocols.

The experimental design of our macrophage village tested whether LPS tolerance is a consequence of reduced numbers of cells responding to repeated LPS stimulation, or whether NF-κB target genes in the population are uniformly silenced. We observed maintenance of expression of a range of tolerizeable genes in PSCM 18 hour and re-stimulated groups.

Examples of this can be seen in PSCMs through several chemokine and inflammatory genes,

including *IL1B*, *CCL4L2, CCL3* and *IL6* expression, suggesting that in a subset of responsive cells,

LPS tolerance had been evaded. Genes previously shown to be stochastically expressed in

mouse RAW264.7 cells included *IL6* and *TNF*, with observations that *IL6* was expressed by all

RAW264 clones and *TNF* only expressed by 72% of them[13,14]. Similarly, we observed that *TNF*

was only expressed in about 75% of human PSCMs, but in 98% of MDMs after acute stimulation

with LPS. *IL6* was inducible at very low levels and in few cells in both *in vitro* human

macrophage models. We also observed proliferative differences between MDM as a model of

recruited and PSCM as a model of resident-tissue macrophages, consistent with the Muldoon

model of proliferative-dependent quorum licensing[13]. We hypothesize that the proliferation of

resident macrophages ensure that some subsets can evade tolerizing signals to remain

responsive to persistent infection and may therefore be the preferred target for persistent

inflammation.

Experimental Procedures

<u>Cell lines and ethics approvals</u>

Human macrophage derivation was carried out in accordance with The University of Melbourne

ethics committee HREC (approvals 1851831, 1646608). Monocytes were isolated from three

buffy coats, obtained from the Australian Red Cross Blood Service.

Human pluripotent stem cells (hPSCs) used were: PB001.1 obtained from the Stem Cell Core

Facility at the Murdoch Children's Research Institute (hPSCreg ID: MCRIi001- A [34]); HDF51

(CVCL_UF42) was kindly provided to Dr Andrew Laslett by Prof. Jeanne Loring (The Scripps

Research Institute, CA, USA); KOLF2 (Wellcome Trust Sanger Institute; hPSCreg ID: WTSli018-B)

cells were kindly provided by Prof. Paul Hertzog (Hudson Institute of Medical Research).

<u>Pluripotent Stem Cell Differentiation</u>

Human pluripotent stem cells were differentiated into macrophages based on protocol described

by [18]. Modifications were as follows: embryoid bodies were kept in rotational cultures without

transference to Matrigel plates for adherence, and progenitors collected from week 2 in RPMI-

1640 containing L-Glutamine (Life Technologies), 10% Fetal Bovine Serum and 100ng/mL CSF1

for macrophage differentiation (see macrophage differentiation).

<u>Peripheral Blood Monocyte Isolation</u>

Buffy Coat was diluted with PBS at a 1:3 dilution and underlayed with Histopaque®-1077 (Sigma-

Aldrich), centrifuged at 350g for 30 minutes at 24°C with no brake. Peripheral blood mononuclear

cells (PBMCs) were isolated from the interphase and washed twice by using MACS buffer with

0.5% heat inactivated Fetal Bovine Serum (FBS) and 2mM EDTA (Invitrogen™ UltraPure™ 0.5M

EDTA) and centrifuging at 400g for 5 minutes at 4°C, then resuspended in 40µl MACS buffer per

10^7 cells. Monocytes were positively selected using Human CD14 MicroBeads (Miltenyi Biotec)

and LS Columns, and then used for macrophage differentiation (see below). A cell count and

viability were determined using 0.4% Gibco™ Trypan Blue using a hemocytometer.

<u>Macrophage differentiation</u>

Monocytes or iPSC-derived myeloid progenitor cells were cultured in tissue-culture treated 6-

well plates. Cells were cultured in RPMI-1640 medium containing L-Glutamine (Life Technologies)

with 10% Fetal Bovine Serum and 100ng/ml recombinant Human M-CSF (R&D Systems) for 4

days.

<u>Flow Cytometry</u>

Cells were prepared for flow cytometry by first performing a blocking step with 5µl mouse serum

(Sigma-Aldrich) then cells were resuspended in FACS Buffer containing 0.5% Bovine Serum

Albumin (Sigma-Aldrich) in Hanks Balanced Salt Solution (no calcium, no magnesium, no phenol

red) (Thermo Fisher Scientific) and incubated with antibodies (see Table 1). Cell viability was

determined using 7-AAD (BioLegend) or PI staining solution (Invitrogen eBioscience). Gating was

performed based on Fluorescence Minus One. Isotypes were included as a control (Table 1).

Compensation was carried out by generating a matrix using compensation beads (BD™

CompBead Plus; Anti- 79 mouse Ig, κ). Cell analysis was carried out on a CytoFLEX S flow

cytometer (Beckman Coulter) using CytExpert acquisition software. Post-acquisition analysis was

performed with FCS Express 7 flow cytometry software.

Stimulation assay

100,000 cells of CD14+ HDF51-, PB001.1- and KOLF2-derived progenitors, or primary CD14+

monocytes from 3 different donors, were pooled then seeded into 6-well dishes in Gibco™

RPMI-1640 media (Thermo Fisher Scientific) with 10% heat inactivated FBS (Thermo Fisher

Scientific) and 100ng/ml M-CSF (R&D Systems). On day 4, media was changed, and cells left to

settle for 2 hours before stimulating with 10ng/mL LPS. After 18 hours, tolerance wells were

washed twice with Gibco™ PBS (Ca2+Mg2+ free; Thermo Fisher Scientific) before fresh media

was added and cells left to settle for 2 hours and re-stimulated with 10ng/mL LPS for 2 hours.

Cells were washed with Gibco™ PBS (Ca2+Mg2+ free; Thermo Fisher Scientific) before

incubating with macrophage cell release for 5 minutes at 37°C and collected in media,

centrifuged at 350g for 5 minutes and resuspended and put through a Flowmi™ strainer (before

cell counts were carried out for sample preparation, see 10x Illumina Single Cell Library

Preparation and RNA Sequencing).

10x Illumina Single Cell Library Preparation and RNA Sequencing

Single cell RNA-sequencing libraries were prepared using the Chromium Next GEM Single Cell 5'

v2 (Dual Index) protocol as per manufacturer's instructions. Cell suspensions were prepared for

a target recovery of 6000 cells per sample. Libraries were quality checked and quantified using

the Agilent Tapestation D5000 High Sensitivity Tape. Two sequencing runs were performed using

Novaseq6000 at University of Melbourne Centre for Cancer Research. Data processing and

merging of sequencing runs were carried out through CellRanger v6.0.2 pipeline to generate files

for analysis. Downstream analysis was carried out using Seurat package v4.0.5 [35].

Donor Deconvolution

The raw number of reads that were associated with each barcode were counted, and only cells

with more than 2000 reads associated were kept for further analysis. We applied cellSNP-lite

(with options: --minMAF 0.1, --minCOUNT 5)[36] for getting the allele counts for the list of

common SNPs in the population identified from the 1000 Genome Project[37]. We subsequently

applied Vireo[38] for demultiplexing the cell barcodes into 3 donors or cell lines used for each

sample. The same donors were then matched by their genotype similarity (Supplementary

Figure 1).

<u>Differential gene expression analysis</u>

A pseudo-bulk approach was used to identify differentially expressed genes[39]. The gene counts

in cells for each donor were aggregated to form bulk samples, and we applied differential

expression analysis using methods designed for bulk RNA-seq[40,41]. Briefly, 24 bulk samples were

created after summing gene counts for 3 donors in each time point, and lowly expressed genes

were filtered out. The gene counts for all samples were then normalized using TMM (trimmed

mean of M-values) [40]. Differential genes were identified by contrasting samples between

sample conditions, and the final list of DE genes in each contrasting group were obtained by the

"treat" method with log(2) fold change > 1[42].

<u>Gene Set and Pathway Enrichment Analysis and ARE Motif Analysis</u>

AU-rich element motifs were evaluated using the ARE motif database [43]. The search space was

restricted to 3'UTR motifs. Gene Set Enrichment Analysis (GSEA) was performed using the

MSigDB Hallmark gene sets [44]. Pathway enrichment used InnateDB [45] or Reactome [46] on

upregulated genes ≥1.5 Log2FC and p.value ≤0.05. Interferon module scores were calculated

with the Seurat function AddModuleScore using the Reactome interferon signaling pathway

gene set, obtained from the msigdbr package. Bar plots showing the average expression of

example genes were derived from the output of the Seurat DotPlot, taking the percentage of

cells per group expressing each gene of interest (bar height) and the unscaled average

expression per group (colour).

<u>Data and code availability</u>

Single cell sequencing data are available through (GSE210175). Code used for analysis is available

at:

https://github.com/wellslab/invitrosinglecell/blob/main/Analysiscode.R

References

1. Felix, R., Cecchini, M. G. & Fleisch, H. Macrophage colony stimulating factor restores in vivo

bone resorption in the op/op osteopetrotic mouse. *Endocrinology* **127**, 2592–2594 (1990).

2. Badimon, A. *et al.* Negative feedback control of neuronal activity by microglia. *Nature* **586**,

417–423 (2020).

3. Cox, N. *et al.* Diet-regulated production of PDGFcc by macrophages controls energy storage.

Science **373**, eabe9383 (2021).

4. Wattrus, S. J. *et al.* Quality assurance of hematopoietic stem cells by macrophages

determines stem cell clonality. *Science* **377**, 1413–1419 (2022).

5. Jardé, T. *et al.* Mesenchymal Niche-Derived Neuregulin-1 Drives Intestinal Stem Cell

Proliferation and Regeneration of Damaged Epithelium. *Cell Stem Cell* **27**, 646-662.e7

(2020).

6. Soe-Lin, S. *et al.* Nramp1 promotes efficient macrophage recycling of iron following

erythrophagocytosis in vivo. *Proc Natl Acad Sci U S A* **106**, 5960–5965 (2009).

7. Lee, C. Z. W. & Ginhoux, F. Biology of resident tissue macrophages. *Development* **149**,

dev200270 (2022).

8. Mulder, K. *et al.* Cross-tissue single-cell landscape of human monocytes and macrophages in

health and disease. *Immunity* **54**, 1883-1900.e5 (2021).

9.   Silvin, A. *et al.* Dual ontogeny of disease-associated microglia and disease inflammatory

macrophages in aging and neurodegeneration. *Immunity* **55**, 1448-1465.e6 (2022).

10. Young, A. M. H. *et al.* A map of transcriptional heterogeneity and regulatory variation in

human microglia. *Nat Genet* **53**, 861–868 (2021).

11. Zhou, Y. *et al.* Human and mouse single-nucleus transcriptomics reveal TREM2-dependent

and TREM2-independent cellular responses in Alzheimer's disease. *Nat Med* **26**, 131–142

(2020).

12. Daemen, S. *et al.* Dynamic Shifts in the Composition of Resident and Recruited

Macrophages Influence Tissue Remodeling in NASH. *Cell Reports* **34**, 108626 (2021).

13. Muldoon, J. J., Chuang, Y., Bagheri, N. & Leonard, J. N. Macrophages employ quorum

licensing to regulate collective activation. *Nat Commun* **11**, 878 (2020).

14. Ravasi, T. *et al.* Generation of Diversity in the Innate Immune System: Macrophage

Heterogeneity Arises from Gene-Autonomous Transcriptional Probability of Individual

Inducible Genes. *J Immunol* **168**, 44–50 (2002).

15. Buchrieser, J., James, W. & Moore, M. D. Human Induced Pluripotent Stem Cell-Derived

Macrophages Share Ontogeny with MYB-Independent Tissue-Resident Macrophages. *Stem

Cell Reports* **8**, 334–345 (2017).

16. Calvanese, V. *et al.* Mapping human haematopoietic stem cells from haemogenic

endothelium to birth. *Nature* **604**, 534–540 (2022).

17. Takata, K. *et al.* Induced-Pluripotent-Stem-Cell-Derived Primitive Macrophages Provide a

Platform for Modeling Tissue-Resident Macrophage Differentiation and Function. *Immunity*

52, 417–418 (2020).

18. Rajab, N. *et al.* An integrated analysis of human myeloid cells identifies gaps in in vitro

models of in vivo biology. *Stem Cell Reports* **16**, 1629–1643 (2021).

19. Alasoo, K. *et al.* Transcriptional profiling of macrophages derived from monocytes and iPS

cells identifies a conserved response to LPS and novel alternative transcription. *Sci Rep* **5**,

12524 (2015).

20. Zhang, B. *et al.* Single-cell RNA sequencing reveals induction of distinct trained-immunity

programs in human monocytes. *Journal of Clinical Investigation* **132**, e147719 (2022).

21. Lavin, Y. *et al.* Tissue-Resident Macrophage Enhancer Landscapes Are Shaped by the Local

Microenvironment. *Cell* **159**, 1312–1326 (2014).

22. Moura Silva, H. *et al.* c-MAF–dependent perivascular macrophages regulate diet-induced

metabolic syndrome. *Sci. Immunol.* **6**, eabg7506 (2021).

23. Dick, S. A. *et al.* Three tissue resident macrophage subsets coexist across organs with

conserved origins and life cycles. *Sci. Immunol.* **7**, eabf7777 (2022).

24. Ripoll, V. M., Irvine, K. M., Ravasi, T., Sweet, M. J. & Hume, D. A. *Gpnmb* Is Induced in

Macrophages by IFN-γ and Lipopolysaccharide and Acts as a Feedback Regulator of

Proinflammatory Responses. *J Immunol* **178**, 6557–6566 (2007).

25. Sehgal, A., Irvine, K. M. & Hume, D. A. Functions of macrophage colony-stimulating factor

(CSF1) in development, homeostasis, and tissue repair. *Seminars in Immunology* **54**, 101509

(2021).

26. Clanchy, F. I. L. Detection and properties of the human proliferative monocyte

subpopulation. *Journal of Leukocyte Biology* **79**, 757–766 (2006).

27. Tong, Y. *et al.* Concomitant type I IFN and M-CSF signaling reprograms monocyte

differentiation and drives pro-tumoral arginase production. *EBioMedicine* **39**, 132–144

(2019).

28. Carmody, R. J., Ruan, Q., Palmer, S., Hilliard, B. & Chen, Y. H. Negative Regulation of Toll-

Like Receptor Signaling by NF-κB p50 Ubiquitination Blockade. *Science* **317**, 675–678 (2007).

29. Zhang, M. Y., Harhaj, E. W., Bell, L., Sun, S. C. & Miller, B. A. Bcl-3 expression and nuclear

translocation are induced by granulocyte-macrophage colony-stimulating factor and

erythropoietin in proliferating human erythroid precursors. *Blood* **92**, 1225–1234 (1998).

30. Hashimoto, D. *et al.* Tissue-Resident Macrophages Self-Maintain Locally throughout Adult

Life with Minimal Contribution from Circulating Monocytes. *Immunity* **38**, 792–804 (2013).

31. Butcher, S. K., O'Carroll, C. E., Wells, C. A. & Carmody, R. J. Toll-Like Receptors Drive Specific

Patterns of Tolerance and Training on Restimulation of Macrophages. *Front. Immunol.* **9**,

933 (2018).

32. Divangahi, M. *et al.* Trained immunity, tolerance, priming and differentiation: distinct

immunological processes. *Nat Immunol* **22**, 2–6 (2021).

33. Song, R. *et al.* IRF1 governs the differential interferon-stimulated gene responses in human

monocytes and macrophages by regulating chromatin accessibility. *Cell Reports* **34**, 108891

(2021).

34. Vlahos, K. *et al.* Generation of iPSC lines from peripheral blood mononuclear cells from 5

healthy adults. *Stem Cell Research* **34**, 101380 (2019).

35. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

36. Huang, X. & Huang, Y. Cellsnp-lite: an efficient tool for genotyping single cells.

Bioinformatics **37**, 4569–4571 (2021).

37. The 1000 Genomes Project Consortium *et al.* A global reference for human genetic

variation. *Nature* **526**, 68–74 (2015).

38. Huang, Y., McCarthy, D. J. & Stegle, O. Vireo: Bayesian demultiplexing of pooled single-cell

RNA-seq data without genotype reference. *Genome Biol* **20**, 273 (2019).

39. Law, C. W. *et al.* RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. *F1000Res*

5, 1408 (2018).

40. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140

(2010).

41. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and

microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

42. McCarthy, D. J. & Smyth, G. K. Testing significance relative to a fold-change threshold is a

TREAT. *Bioinformatics* **25**, 765–771 (2009).

43. Bakheet, T., Hitti, E. & Khabar, K. S. A. ARED-Plus: an updated and expanded database of

AU-rich element-containing mRNAs and pre-mRNAs. *Nucleic Acids Research* **46**, D218–D220

(2018).

44. Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell*

Systems **1**, 417–425 (2015).

45. Breuer, K. *et al.* InnateDB: systems biology of innate immunity and beyond—recent updates

and continuing curation. *Nucleic Acids Research* **41**, D1228–D1233 (2013).

46. Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Research* **50**,

D687–D692 (2022).

Tables

Table 1 : Flow Cytometry Antibodies		
Marker	Antibody	Isotype control
CD14	Brilliant Violet 421™ anti-human CD14 Antibody (BioLegend; Cat. No. 325628)	Brilliant Violet 421™ Mouse IgG1, κ Isotype Ctrl Antibody (BioLegend: Cat. No. 400157)
CD16	Brilliant Violet 605™ anti-human CD16 Antibody (BioLegend: Cat. No. 302040)	Brilliant Violet 605™ Mouse IgG1, κ Isotype Ctrl Antibody (BioLegend: Cat. No. 400161)
CD45	APC anti-human CD45 Antibody (BioLegend: Cat. No. 368512)	APC Mouse IgG1, κ Isotype Ctrl (FC) Antibody (BioLegend: Cat. No. 400121)
CD11b	Brilliant Violet 711™ anti-human CD11b Antibody (BioLegend: Cat. No. 301344)	Brilliant Violet 711™ Mouse IgG1, κ Isotype Ctrl Antibody (BioLegend: Cat. No. 400167)
HLA-DR	Alexa Fluor® 488 anti-human HLA-DR Antibody (BioLegend: Cat. No. 307619)	Alexa Fluor® 488 Mouse IgG2a, κ Isotype Ctrl Antibody (BioLegend: Cat. No. 400233)

Figure 1 Proliferation drives macrophage heterogeneity in a village-in-a-dish model of repeated bacterial stimulation. (1A) Schematic of experimental design, including design of village coculture model for 3 MDM donors or 3 PSCM lines. Time course for LPS stimulation and multiplexed single cell library preparation. (1B) Representative surface expression of macrophage markers in MDM and PSCM measured by flow cytometry. Isotype control antibody staining (pink), signal (blue). (1C) Uniform Manifold Approximation and Projection (UMAP) of Seurat clusters (resolution 0.9) (1D) Seurat cluster group (columns) by sample barcode (rows). Groups are colour-coded according to (C) and labelled according to time point (C control, A acute, R 18 hours, T tolerance) and numbered from highest to lowest cell number per group. (1E) Genes that discriminated proliferating-cell clusters across LPS stimulation time course. Stacked bar plots indicate the proportion of proliferating-cell clusters (shaded bars) per barcode group. (1F) Key discriminating markers between UMAP clusters. Heat map of top genes (avg. Log2FC) in each subcluster. Legend denotes high (red) to low (blue) expression. See also Supplementary Figure 1, Supplementary Tables 1 and 2.

Figure 2 Inflammatory gene expression varied by the proportion of cells recruited to the
inflammatory phenotype, as well as average gene expression per cell. (2A) Schematic of the TLR
pathway and average expression of pathway members in unstimulated PSCM and MDM
populations. Bar plots coloured by cluster members (MDM Cluster C13, C17; PSCM clusters C8,
C11: see Figure 1C) to evaluate average expression in proliferating and nonproliferating control
clusters. (2B) Venn diagram of LPS-inducible genes after 2 hours (selected by Log2FC. > 1.5, Adj.
p value <0.05). P-values on graph indicate InnateDB pathway over-representation (Fishers exact
test). (2C) Percentage of cells per cluster expressing six representative targets of TLR4-LPS
stimulation. (2D) Percentage of cells per cluster expressing six representative targets of IRF
(MyD88-independent) stimulation. (2E) Seurat module scores of the Reactome Interferon
signaling pathway. Averaged cell score shown. (2F) Growth factor signaling persists in PSCM but
not MDM. Percentage of cells per cluster expressing four representative reparative factors. For
all bar plots, gene expression (average CPM / cell) indicated by colour where dark blue indicates
highest expression, light for lowest expression.

Figure 3 Cartoon of two macrophage populations describing escape from endotoxin tolerance

in subpopulations that are proliferating. Created in BioRender.

Supplementary Figure 1: Quality assessment of RNAseq libraries. A: donor contributions to PSCM
or MDM sequencing libraries. B: Correlation of sequencing tags labelled by experimental
condition (barcode) or donor ID. C: Screenshot of projection of PSCM and MDM libraries onto
the Stemformatics Myeloid atlas, and Cabyara analysis of most similar reference atlas samples.

Supplementary Material

Supplementary Table 1: Differential expression analysis of MDM and PSCM groups. Accompanies Figures 1 and 2.

Average and differential expression analysis between MDM and PSCM groups. Average expression, Log2FC, p value, and adjusted p value.

Supplementary Table 2: Average expression of TLR4 signaling components. Accompanies Figure 2A.

Average expression of components of TLR4 signaling pathway. Average expression shown for all identified seurat clusters.