Foxd1 dependent induction of temporal retinal character is required for visual
function.

María Hernández-Bejarano[1], Gaia Gestri[3^§], Clinton Monfries[4], Lisa Tucker[3^], Elena
I. Dragomir[3^], Isaac H. Bianco[3*], Paola Bovolenta[1,2], Stephen W. Wilson[3^§] and
Florencia Cavodeassi[1,4,§]

[1] Centro de Biología Molecular Severo Ochoa (CSIC-UAM) and [2] CIBER de
Enfermedades Raras (CIBERER), Nicolás Cabrera 1, 28049, Madrid, Spain
[3] Departments of Cell and Developmental Biology^ and Neuroscience, Physiology
and Pharmacology* University College London, Gower Street, WC1E 6BT, London,
UK
[4] St. George's, University of London, Cranmer Terrace, SW17 0RE, London, UK
14

16 § Correspondence: Florencia Cavodeassi (fcavodea@sgul.ac.uk)
17   Gaia Gestri (g.gestri@ucl.ac.uk)
18   Stephen W Wilson (s.wilson@ucl.ac.uk)
19

Running title: *foxd1* and visual function

Keywords: *foxd1*, *rx3*, retinal patterning, fovea, *area temporalis,* OKR, OMR,
zebrafish

Summary statement: This study provides a mechanistic link between eye patterning
and the establishment of functionally distinct retinal regions and reveals the temporal
retina preferentially controls specific aspects of visual function.

Abstract

Appropriate patterning of the retina during embryonic development is assumed to
underlie the establishment of spatially localised specialisations that mediate the
perception of specific visual features. For instance, in zebrafish, an area involved in
high acuity vision (HAA) is thought to be present in the ventro-temporal retina. Here
we show that the interplay of the transcription factor Rx3 with Fibroblast Growth
Factor and Hedgehog signals, initiates and restricts *foxd1* expression to the
prospective temporal retina, initiating naso-temporal regionalisation of the retina.
Abrogation of FoxD1 results in the loss of temporal and expansion of nasal retinal
character, and consequent absence of the HAA. These structural defects correlate with
severe visual defects as assessed in optokinetic and optomotor response assays. In
contrast, optokinetic responses are unaffected in the opposite condition in which nasal
retinal character is lost at the expense of expanded temporal character. Our study
indicates that the establishment of temporal retinal character during early retinal
development is required for the specification of the HAA, and suggests a prominent
role of the temporal retina in controlling specific visual functions.

Introduction

The position at which retinal ganglion neurons differentiate along the naso-temporal and dorso-ventral axes of the forming eye determines the location at which they innervate central targets, ensuring accurate retinotopic connectivity. Consequently, the mechanisms by which retinal neurons acquire their regional identity are critical for correct visual function. Regionally localised specialisations of the retina include the high acuity areas (HAAs) described in many diurnal organisms. In many primates and birds, this structure is called the fovea and is characterized by a morphological indentation, high density of RGCs and cone photoreceptors, absence of rod photoreceptors, specialised inner retina neuronal circuitry, and absence of vasculature (Bringmann et al., 2018; Bringmann, 2019; da Silva and Cepko, 2017; Kolb et al., 2020; Kozulin et al., 2009; Kozulin et al., 2010). In teleost fish such as the zebrafish, some features associated with the fovea are evident in the ventro-temporal region of the retina, suggesting this region is comparable to the fovea in birds and primates (Mangrum et al., 2002; Pita et al., 2015; Schmitt and Dowling, 1999; Yoshimatsu et al., 2020; Zhou et al., 2020; Zimmermann et al., 2018). During development, the human fovea shows precocious expression of neuronal differentiation genes (Hoshino et al., 2017). In the chick, DaSilva and Cepko (2017) identified low levels of retinoic acid (RA) activity and high levels of the Fibroblast Growth Factor ligand FGF8 in a highly circumscribed area of the retina, prefiguring the fovea, a pattern that is conserved in the human presumptive fovea (Cornish et al., 2004; Cornish et al., 2005; da Silva and Cepko, 2017).

HAAs are generally located in the temporal retina (Bringmann, 2019; Kolb et al., 2020; Pita et al., 2015), suggesting their specification may be dependent upon acquisition of temporal character. In the zebrafish, the subdivision of the retina in domains with nasal and temporal character is evident from early stages of optic vesicle evagination by the expression of *foxg1* in the future nasal half and *foxd1* in the future temporal half of the eye (Hernández-Bejarano et al., 2015; Picker et al., 2009). Initially, the naso-temporal subdivision is aligned with the dorso-ventral axis of the developing central nervous system. However, as eye morphogenesis progresses, the eye primordium rotates such that nasal and temporal retinae relocate to their final position, aligned with the anterior-posterior axis (Kwan et al., 2012; Picker et al., 2009).

Previous studies have shown that the establishment of nasal and temporal character requires the spatially localised activity of the Sonic-Hedgehog (Shh) and Fgf signalling pathways (Hernández-Bejarano et al., 2015; Picker and Brand, 2005; Picker et al., 2009). Shh, expressed in ventral midline structures promotes *foxd1* expression in the ventral (future temporal) half of the optic primordium. Fgfs, emanating from dorsal forebrain and adjacent tissues induce *foxg1* in the dorsal (future nasal) optic primordium and repress *foxd1* expression, contributing to its confinement ventrally. Cross-repression between FoxG1 and FoxD1 at the border between the dorsal and ventral halves of the eye primordium subsequently refines the NT subdivision (Hernández-Bejarano et al., 2015; Takahashi et al., 2009; Takahashi et al., 2003). When Shh signalling is absent, *foxd1* expression can be restored by suppressing Fgf signalling indicating that an additional factor(s) can promote *foxd1* expression independently of Shh and Fgf.

Here we elucidate the molecular mechanisms involved in establishing temporal fate
during retinal development and explore how alterations to naso-temporal retinal
patterning impact HAA formation and visual function. We identify the transcription
factor Rx3 (Loosli et al., 2003; Loosli et al., 2001) as being required for *foxd1*
expression in the forming eye. *rx3* is expressed in the eye field and is required for eye
formation in all studied vertebrates (Andreazzoli et al., 1999; Andreazzoli et al.,
2003; Bailey et al., 2004; Fish et al., 2014; Loosli et al., 2003; Loosli et al., 2001;
Mathers et al., 1997; Voronina et al., 2004). We show that the interplay of Rx3 with
Shh and Fgf activities initiates and restricts *foxd1* expression to the prospective
temporal half of the evaginating optic vesicles. We further show that the
establishment of a *foxd1* expression domain in the eye is linked to the formation of the
HAA. Larvae lacking FoxD1 function show normal retinal neuron differentiation and
lamination, but temporal retinal and HAA markers are absent, and nasal markers
expand throughout the retina. In contrast, fish in which nasal retinal character is
absent and temporal character is expanded, show a concomitant expansion of HAA
markers. In both conditions, retinotectal projections are severely perturbed. Despite
this, only fish lacking FoxD1 show significant visual impairment, which was revealed
by examining optokinetic and optomotor reflex behaviours and suggests impaired
perception of whole-field visual motion. Our study reveals a prominent role of the
temporal retina in controlling specific aspects of visual function and provides a
mechanistic link between early patterning of the eye primordium and efficient visual
performance.

Results

1- Rx3 provides competence for Shh to promote *foxd1* expression in the ventral
portion of the optic vesicle

Our previous studies (Hernández-Bejarano et al., 2015) showed that spatially
localised Shh and Fgf signalling establishes the discrete domains of *foxd1* and *foxg1*
expression in the presumptive retina. However, those studies also showed that
blocking the activity of both Shh and Fgf in otherwise wildtype embryos resulted in
expression of *foxd1* throughout the optic vesicle, suggesting that an additional factor
can promote *foxd1* expression in absence of normally critical signals.

The transcription factor Rx3 is specifically expressed in the eye field and is part of a
network of regulatory proteins that confers eye fate and promotes the evagination of
the optic vesicles. Lack of Rx function results in eye loss (Andreazzoli et al., 1999;
Andreazzoli et al., 2003; Loosli et al., 2003; Mathers et al., 1997; Voronina et al.,
2004). However, spatially localised expression of *rx3* (prospective retina marker)*,*
nkx2.1 (prospective hypothalamic marker) and *emx3* (prospective telencephalic
marker) is initially normal in the brain and eye-forming regions of *rx3*[-/-] embryos
(**Figure 1A-D**; Kennedy et al., 2004) indicating that in the mutants the eye-forming
territory is still present and molecularly distinct from surrounding anterior neural
structures. The presence of a prospective eye-forming domain has enabled us, in this
study, to assess the effect of loss of *rx3* on expression of genes that delineate NT
patterning.

foxg1, expressed in the telencephalon and nasal half of the eye primordium (**Figure**
1E-F; Hernández-Bejarano et al., 2015), was expanded throughout most of the

prospective eye domain in $rx3^{-/-}$ mutants (**Figure 1G-H**; Stigloher et al., 2006), while
foxd1 expression was much reduced (**Figure 1J-M**). We confirmed the changes in the
extent of *foxd1/foxg1* expression were localised to the prospective eye, by comparing
their expression domains to those of *rx3* itself (which labels the eye field) and *nkx2.1*
(**Figure 1A-B, I**). A previous transcriptomic analysis in $rx3^{-/-}$ mutants suggested that
foxd1 may be a downstream target for Rx3 (Yin et al., 2014). That study, together
with our observations, support a scenario in which Rx3 is required for *foxd1*
expression in the forming eye and provides cells with the competence to respond to
Shh in promoting the expression of *foxd1*. If this is the case, the widespread
expression of *foxd1* in the optic vesicle following loss of Shh/Fgf signalling should be
absent in $rx3^{-/-}$ embryos. To address this, we first assessed whether Shh and Fgf
signalling is overtly altered in $rx3^{-/-}$ mutants.

$rx3^{-/-}$ embryos showed normal levels of expression of genes encoding Fgf and Shh
ligands and target genes in the forebrain, suggesting that levels of activity of these
two pathways in $rx3^{-/-}$ mutants are similar to wild types. The ligand encoding genes
fgf8 and *shh* showed only slight changes at the most anterior portion of the forebrain
[compare the extent of *fgf8* (brackets, **Figure S1A-B**) and *shh* (arrows, **Figure S1G-**
H) expression between wildtype and $rx3^{-/-}$ embryos]. In addition, expression of *pea3*
and *erm* (direct targets of the Fgf pathway) and *gli1* and *patched 2* (direct targets of
the Shh pathway), was largely normal (**Figure S1C-F,I-L**). To abrogate Shh and Fgf
activity in $rx3^{-/-}$ embryos, we treated clutches of embryos obtained from $rx3^{+/-}$ in-
crosses simultaneously with two drugs: cyclopamine, which inhibits the Hh
transducer Smoothened; and SU5042, an inhibitor of Fgf receptors (as previously
described in Hernández-Bejarano et al., 2015).

In contrast to cyclopamine+SU5042-treated wildtype embryos in which *foxd1*
expression is expanded (**Figure 1R-S**; Hernández-Bejarano et al., 2015), the residual
expression of *foxd1* in $rx3^{-/-}$ embryos (**Figure 1L-M,T**) was completely lost when
embryos were treated with cyclopamine+SU5042 (**Figure 1T-V**; 30-50 embryos were
treated and processed together per experiment; $rx3^{-/-}$ embryos were recovered at the
expected mendelian proportions and the phenotypes observed were fully penetrant).
Remarkably, *foxg1*, which was lost in the optic vesicles of wildtype embryos treated
with cyclopamine+SU5402 (**Figure 1N-O**), was expressed throughout the eye field of
$rx3^{-/-}$ embryos regardless of treatment (**Figure 1P-Q,V**). This may be due to the lack
of the *foxg1* repressor FoxD1 in these conditions. Alternatively, it may be a
consequence of the progressive adoption of telencephalic character by the eye-
forming region of $rx3^{-/-}$ mutants (Stigloher et al., 2006). Consistent with the first
scenario, *foxg1* was not expressed in the prospective eye region of $rx3^{-/-}$ embryos
treated only with the Fgf inhibitor SU5402 (**Figure S2G-H**), in which there was
expanded *foxd1* expression (**Figure S2E-F**).

2- In absence of *foxd1*, retinae fail to develop temporal character

The results above indicate a critical role for Rx3 and Shh in the induction of *foxd1*
expression and the establishment of temporal character in the evaginating optic
vesicle. Studies in other model organisms have shown that FoxD1 has a prominent
role as determinant of retinal temporal character (Carreres et al., 2011; Hatini et al.,
1994; Herrera et al., 2004; Takahashi et al., 2009; Takahashi et al., 2003). To further
explore the role of *foxd1* in promoting temporal character, we generated a zebrafish

loss of function mutant using CRISPR-Cas9 technology (Prykhozhij et al., 2017; see
Materials and Methods). A 10 base-pair deletion just prior to the sequence encoding
the FoxD1 DNA binding domain led to a frame-shift and a stop codon in position 70
of the protein product (*foxd1^cbm16*; **Figure 2A-B**). This mutation likely produces a
non-functional form of the protein, truncated just prior to the DNA binding site.

foxd1^-/- embryos developed at least up to 9 dpf, showing no apparent morphological
malformations in the retina or elsewhere (data not shown). However, the temporal
retinal marker *epha4b*, and *foxd1* itself, were downregulated in the mutant eyes
(**Figure 2C-D,G-H**). This downregulation was accompanied by a mirror-image
duplication of expression of the nasal retinal markers *foxg1* and *ephA5a* in the
temporal half of the retina (**Figure 2E-F, I-J** and **Figure S3A-D**). Of note,
downregulation of *foxd1* expression in *foxd1^-/-* embryos was not due to non-sense
mediated decay, since expression in other regions of the embryo was not affected
(**Figure S4**). The retinal phenotype of *foxd1^-/-* mutants was phenocopied in F0
"crispants" generated by simultaneous injection of three guides targeting different
regions of the *foxd1* locus (**Figure S3G**; Kroll et al., 2021; Materials and Methods).
All genotyped crispant embryos showed gene editing (n=16; **Figure S3H**) and over
90% of the analysed embryos (65 out of 72) showed the expansion in *foxg1*
expression characteristic of *foxd1* mutants (**Figure S3E-F**).

Analysis of retinotectal projections revealed an aberrant pattern of tectal innervation
in *foxd1^-/-* mutants. In control conditions, temporally located RGCs, injected with DiI,
innervated the anterior portion of the contralateral tectum, while nasally located
RGCs, injected with DiO, innervated the posterior tectum (**Figure 2K**; inset shows
the corresponding eye, n=10 eyes). In *foxd1^-/-* mutants, RGC terminals (labelled as for
control conditions) from both nasal or temporal halves of the retina showed
overlapping arborisations across much of the tectum (**Figure 2L-M**; insets show the
corresponding eye, n=12 eyes). This result indicates that in the absence of RGCs with
temporal character, axons from RGCs with nasal character expand their arbours
throughout the tectum (Suetterlin et al., 2012). Overall, these results suggest that the
foxd1^-/- mutant retina has all-nasal character.

HAA specification is linked to naso-temporal patterning

In fish, the ventro-temporal region of the retina (also called the *area temporalis*;
Schmitt and Dowling, 1999) bears several structural specialisations that are thought to
provide high visual acuity and support prey detection: a high density of U-cones
(invested in detecting light in the ultraviolet spectrum) and low density of rods
(Yoshimatsu et al., 2020; Zimmermann et al., 2018), a specialised inner retina
neuronal circuit (Mangrum et al., 2002; Pita et al., 2015) and a characteristic number,
shape and position of synaptic terminals at the inner plexiform layer (Zhou et al.,
2020; Zimmermann et al., 2018). Immunostaining with anti-PKCα highlights the cell
anisotropies found in the inner nuclear layer and inner plexiform layer at the level of
the HAA (Zimmermann et al., 2018).
229
In contrast to wildtype zebrafish, where PKCα immunostaining was enriched in the
ventro-temporal region of the retina (**Figure 2N**), PKCα enrichment was lost in *foxd1^-*
^/- retinae (**Figure 2O**), suggesting that HAA-specific features are absent in *foxd1^-/-*

mutants. Despite this absence of HAA-specific features, *foxd1*[-/-] retinae showed an
otherwise normal PKCα expression and architecture (**Figure 2** and not shown).

To determine whether changes in HAA markers are disrupted in other conditions that
affect naso-temporal patterning, we analysed the opposite condition, in which
embryos present a retina with all-temporal character. To generate this condition, we
misexpressed Shh throughout the developing eye using the Gal4/UAS system (Distel
et al., 2010; Halpern et al., 2008; Paquet et al., 2009) as previously described
(Hernández-Bejarano et al., 2015). As shown in a previous report (Hernández-
Bejarano et al., 2015), *Tg{rx3:Gal4};UAS:Shh* embryos showed an expansion of
foxd1 expression throughout the retina at the expense of *foxg1*.

Tg{rx3:Gal4};UAS:Shh retinae showed a dorso-temporal expansion in the HAA-
associated enrichment of PKCα (**Figure 2P**, arrows; compare with wild type in
Figure 2N), suggesting that the HAA was not only present, but expanded in this
condition. Notably, an ectopic patch of PKCα enrichment also appeared to be present
in the ventro-nasal half of the retina (**Figure 2P**). Overall, these results suggest that
the establishment of nasal and temporal character is linked to the formation of retinal
specialisations such as the HAA.

Visual function is impaired in *foxd1* mutants

Perturbation of naso-temporal patterning and retinotectal projections in *foxd1*[-/-]
mutants is likely to affect visual function. We therefore examined two robust visually
guided stabilisation reflexes elicited by whole-field motion: the optokinetic and the
optomotor responses (OKR and OMR, respectively). The OKR is a visually driven
compensatory eye movement that minimises retinal image slip to stabilise gaze.
During OKR, animals perform slow phase eye rotations in the direction of visual
motion with intermittent fast phase saccades to reset eye position. In the OMR,
zebrafish larvae turn and swim in the direction of perceived whole field motion
(Neuhauss et al., 1999).

foxd1[-/-] mutants were substantially impaired in OKR assays. We presented 6 dpf
tethered larvae with rotating visual gratings and quantified the gain (ratio of eye
velocity to stimulus velocity) of the slow phase of the optokinetic nystagmus as a
function of stimulus spatial frequency, temporal frequency and contrast. Gain was
reduced by around 80% in all conditions in *foxd1*[-/-] mutants (**Figure 3B**, red) as
compared to wild types and heterozygote siblings (**Figure 3B**, black/blue). While this
could reflect deficits in visual perception, we also observed a reduction in oculomotor
range and in peak eye velocity (**Figure 3D**) suggesting that oculomotor defects might
contribute to the phenotype. We therefore tested a second stabilisation reflex evoked
by whole field visual motion, and measured OMR performance. Whereas wildtype
larvae robustly turned in the direction of whole-field motion, *foxd1* crispant larvae
were more prone to make mistakes and only performed about 67% turns in the correct
direction (**Figure 3F**). *foxd1* crispant larvae did not show evidence of locomotor
dysfunction as they performed a similar total number of swim bouts compared to wild
types and displayed a normal range of turn angles (**Figure 3G and Figure S5**). Taken
together, these data indicate that loss of FoxD1 function leads to impaired visual
motion perception.

To determine whether impairment in visual motion perception is characteristic of other conditions that affect naso-temporal patterning and retinotectal projections, we assessed visual performance in *Tg{rx3:Gal4};UAS:Shh* fish with double temporal retinae. RGCs positioned in the nasal region of *Tg{rx3:Gal4};UAS:Shh* embryos projected more anteriorly in the tectum, overlapping with the projections from temporally located RGCs (**Figure 3A**; Hernández-Bejarano et al., 2015). Despite altered naso-temporal character and retinotectal projections, we did not observe any deficits in OKR performance in *Tg{rx3:Gal4};UAS:Shh* larvae (**Figure 3C**, red; compare with uninjected *Tg{rx3:Gal4}* larvae in black). These results suggest that the difference in OKR performance in *foxd1⁻/⁻* mutants and *Tg{rx3:Gal4};UAS:Shh* embryos is associated with the absence or presence, respectively, of temporal retinal character and, potentially, associated to connectivity alterations.

Discussion

In this study we identify the eye field specification gene *rx3* as a regulator of *foxd1* expression, thereby further resolving the transcriptional and signalling pathways that lead to naso-temporal retinal pattern (**Figure 4**). We propose that *rx3* expression in the nascent eye field confers to this whole domain competence to express *foxd1*. *foxd1* is repressed in the dorsal half of the eye primordium by Fgf activity and promoted in the ventral half by Hh activity (this study and Hernández-Bejarano et al., 2015). Fgf activity also promotes the expression of the nasal determinant *foxg1* and FoxG1 and FoxD1 subsequently engage in a negative cross-regulatory relationship that refines and maintains the border between nasal and temporal retinal domains (Hernández-Bejarano et al., 2015).

The requirement of Rx3 to promote *foxd1* expression is most clearly revealed when the activity of other regulators is removed. In the absence of Fgf and Shh, *foxd1* expands throughout the optic vesicles. Removing Rx3 activity in this condition leads to loss of *foxd1* expression with a concomitant expansion of *foxg1* expression, confirming the requirement for Rx3 in the induction of *foxd1* expression.

An *area temporalis* in the zebrafish retina characterized by tightly packed cones was identified more than 20 years ago (Schmitt and Dowling, 1999). This area corresponds to the HAA, and more recent studies have shown that it bears a distinctive density of particular neuronal types and a specialised neuronal circuitry (Mangrum et al., 2002; Pita et al., 2015; Yoshimatsu et al., 2020; Zhou et al., 2020; Zimmermann et al., 2018). Our analysis suggests that the establishment of temporal retinal character is required for HAA formation. *foxd1⁻/⁻* mutants show an expansion of nasal retinal markers at the expense of temporal markers. Accumulation of PKCα in the ventro-temporal retina, which highlights the structural features of the HAA inner retina and inner plexiform layer (Zimmermann et al., 2018), is lost in *foxd1⁻/⁻* mutants, suggesting that the mutant retina is devoid of at least some HAA features.

Not only the HAA, but also other structural anisotropies in the temporal retina may be affected in *foxd1⁻/⁻* mutants. Recent studies suggest that the OKR is driven mainly by stimuli covering the central visual field (Dehmelt et al., 2021) and predict that the HAA specialisations in the temporal retina are dispensable for OKR. However, OKR and OMR tests in *foxd1⁻/⁻* mutants reveal defective whole-field motion perception in these larvae. This suggests that either the HAA or other regions of temporal retina

required for this visual response, are affected in *foxd1*$^{-/-}$ mutants. Conversely, larvae
with temporal character expanded into the nasal domain (*Tg{rx3:Gal4};UAS:Shh*)
show an overtly normal OKR. These data suggest that perturbation of nasal retinal
character does not interfere with this aspect of visual function, and supports the idea
that at least some aspects of the OKR are predominantly driven by retina with
temporal character.

Ablation and lesion experiments in both fish and other vertebrate models suggest that
the execution of the OKR is driven mainly by the pretectum and can occur in the
absence of an intact tectum (Flandrin and Jeannerod, 1981; Roeser and Baier, 2003).
Here we show that defective whole-field motion perception in *foxd1*$^{-/-}$ mutants, which
show extensive disruption of retinotectal projections, is perturbed, but our analysis
does not allow us to determine whether these two phenotypes are causally related as
innervation of other retinorecipient areas may also be perturbed in *foxd1*$^{-/-}$ mutants.
Indeed, some pretectal arborisation fields are preferentially innervated by the
temporal retina (Robles et al., 2014), and our analysis did not determine whether these
are affected in *foxd1*$^{-/-}$ mutants.

In summary, our study uncovers a mechanistic link between early naso-temporal
patterning events in the eye primordium and the establishment of functionally distinct
regions in the retina. Despite the importance of naso-temporal patterning for accurate
RGC connectivity to specific retino-recipient areas, analysis of the visual function in
foxd1$^{-/-}$ mutants and *Tg{rx3:Gal4};UAS:Shh* fish reveals a surprising robustness in the
ability of the larvae to perceive stimuli in the absence of nasal or temporal-specific
retinal character. These lines of fish provide an exciting new avenue to further our
understanding of the role of nasal and temporal retinal character in mediating specific
visual functions.

Materials and Methods

Fish lines and husbandry

AB and *tupl* wildtype zebrafish strains, the transgenic line *Tg{rx3::Gal4-VP16}*^{vu271Tg}
(Weiss et al., 2012), and mutant lines *chk*^{ne2611} (Loosli et al., 2003) and *foxd1*^{cbm16}
were maintained and bred according to standard procedures (Aleström et al., 2020;
Westerfield, 1993). All experiments conform to the guidelines from the European
Community Directive and the British (Animal Scientific Procedures Act 1986) and
Spanish (Real Decreto 53/2013) legislation for the experimental use of animals.

Generation of the foxd1 mutant

The sequence to target in the *foxd1* open reading frame (ENSDARG00000029179)
was selected using the UCSC Genome Browser (http://genome.uscs.edu). A DNA
oligonucleotide bearing the target sequence (GCTTGTAGGGGTCCCGTGC;
positions 435-417 on the reverse strand of the transcript), the T7 RNA polymerase
binding site and the sequence complementary to the oligoB universal primer was
commercially obtained, expanded with Expand High Fidelity DNA polymerase
(Roche) and transcribed with the Maxi script T7 kit (NEB), following manufacturers'
instructions. Cas9 mRNA was generated by linearisation and transcription from the
PCS2-nCas9n (Addgene, #47929) clone, purified (PCR cleanup kit, Roche) and

transcribed (SP6 mMessage mMachine Kit Ambion), following manufacturers'
instructions. F0 founders were generated by co-injection of guide RNA (25ng/µl) and
Cas9 mRNA (300ng/µl) in *AB/tupl* embryos at one cell stage. Cleavage efficiency
was assessed in pools of injected embryos by CRISPR-STAT analysis (Carrington et
al., 2015). Genotyping of F0 founders and their progeny was performed by CRISPR-
STAT or HRM analysis from genomic DNA samples obtained from tail fin biopsies.
The primers used are detailed in **Table S1**.

Generation of multi-guide foxd1 crispants

The phenotype of the *foxd1^{cbm16}* mutants was phenocopied by multi-guide injection,
following the protocol previously published by (Kroll et al., 2021). Three synthetic
RNA guides were designed (1AA, 1AB and 1AC guides, Supplementary Table 1) and
ordered to Integrated DNA Technologies (IDT). Guides were annealed to the
tracrRNA oligonucleotide (IDT#1072532), assembled with Cas9 protein
(IDT#1081058) and injected into one-cell stage wildtype embryos, as previously
described (Kroll et al., 2021). A subset of the injected embryos was genotyped by
HRMA using the HRMA primers described in **Table S1**, to confirm the presence of
gene editing.

Microinjection and drug treatments

Treatments with SU5402 and cyclopamine were performed as previously described
(Hernández-Bejarano et al., 2015). SU5402 (Calbiochem) and cyclopamine
(Calbiochem) were applied at a concentration of 10µM and 100µM, respectively, to
pools of embryos derived from the mating of *rx3$^{+/-}$* parental fish, at the corresponding
stage of embryonic development. 30-50 embryos were treated and processed together
per experiment/marker. *rx3$^{-/-}$* embryos were recovered at the expected mendelian
proportions and the phenotypes observed were fully penetrant.

Overexpression of *shh* in the optic vesicle under the control of the Gal4/UAS system
(Halpern et al., 2008) was performed by injecting 20-30ng of bidirectional
GFP:UAS:*shh* plasmid DNA into the cell of one-cell stage *Tg{rx3::Gal4-VP16}^{vu271Tg}*
embryos. Only embryos with homogeneous GFP expression in the optic vesicles were
selected and processed for analysis (as described in Hernández-Bejarano et al., 2015).

mRNA detection and immunolabelling

mRNA detection (preparation of RNA antisense probes and whole mount *in situ*
hybridisation) was performed as previously described (Hernández-Bejarano et al.,
2015). Immunolabelling of 8 dpf wildtype, *foxd1* and *Tg{rx3::Gal4};*UAS:*Shh* retinae
with anti-PKCα (Sigma, P4334, 1:100), Alexa-coupled secondary antibodies (Jackson
ImmunoResearch, 1:500) and the nuclear dye DAPI was performed on RPE-dissected
retinae as described in (Zimmermann et al., 2018).

Tracing of retinotectal projections

Nasal and temporal RGCs were labelled with DiO and DiI, respectively in 6 dpf *wild*
type, *foxd1$^{-/-}$* and *Tg{rx3:Gal4};*UAS:*Shh* retinae previously fixed in 4%
paraformaldehyde. Labelled larvae were incubated at room temperature for at least 24

hours before imaging. Each tectum and its corresponding eye were sequentially
imaged (Hernández-Bejarano et al., 2015).

Optokinetic response test (OKR)

6 dpf larvae were anesthetised with tricaine (MS-222, Sigma), embedded in 1% low-
melting point agarose (Sigma) and immersed in embryo medium. A small portion of
agarose was removed from around the eyes to allow movement. Once larvae had
recovered from anaesthesia, OKR performance was tested by presenting rotating
visual gratings that varied in size, contrast and speed. We presented a set of 12 unique
stimuli that was repeated twice for each animal. Horizontal eye movement was
tracked under infrared illumination (850 nm) with an AVT Pike camera (100 frames
per second). Stimulus presentation and machine vision were controlled using
LabVIEW (National Instruments). Slow phase gain, saccadic eye velocity and
oculomotor range were computed using custom MATLAB software
(https://IsaacBianco@bitbucket.org/biancolab/okrsuite.git). *foxd1*$^{-/-}$ mutants (n=8)
were compared to wildtype (n=5) and *foxd1*$^{+/-}$ heterozygotes (n=27) and
*Tg{rx3:Gal4};*UAS:*Shh* (n=8) were compared to *Tg{rx3:Gal4}* (n=10) uninjected
larvae. Significance was calculated with a two-way ANOVA test, followed by a
Bonferroni test (GraphPad Prism5). Graphical representation shows eye gain (0-1
range, *y* axis) at variable speed (cycles per screen), frequency (degrees per second) or
contrast (0-1 range; *x* axis).

Optomotor response assay (OMR)

6 dpf zebrafish larvae were individually placed in a 6 cm Petri dish and presented
from below via a projector with moving grating stimuli in a closed loop manner. The
customized stimulus protocol and tracking of the freely moving fish was implemented
using the Stytra software package (Štih et al., 2019). The time course of the assay
consisted of 5 repetitions of rightwards and leftwards moving gratings (20 seconds
each), separated by 10 seconds of static gratings (pause), for a total time of 5 minutes
per fish. 21 wildtype and 21 *foxd1*-crispant larvae were subjected to the OMR assay.
Custom behavioral analysis was implemented using Python and the bouter package
(Stih et al., 2022).

Imaging and data processing

DiI/DiO-traced embryos were embedded in low melting point agarose (Sigma) at 1-
1.5% in PBS for confocal imaging using a 40X (0.8NA) long-working distance water
immersion lens. A Zeiss LSM710 confocal microscopy system was used for image
acquisition. Immunolabelled 8dpf retinae were imaged using a Leica SP8 microscope
with at 25x water immersion lens.

In situ hybridised embryos and dissected eyes were mounted flat in a drop of glycerol
and dorsal images were acquired with a 20X (0.70NA) dry lens using a Leica CTR
5000 microscope connected to a digital camera (Leica DFC 500), and operated by
Leica software. Some embryos were embedded in gelatine/BSA for vibratome
sectioning as previously described (Hernández-Bejarano et al., 2015; Sánchez-
Arrones et al., 2013). Alternatively, embryos were cryoprotected in sucrose 30% and
embedded in OCT (Sakura Fintek) for cryo-sectioning as previously described

(Cavodeassi et al., 2013). Sections (16 µm) were obtained using a Leica VT1000S
vibratome or a Leica cryostat, mounted in glycerol, and imaged with a 40X (1.3 NA)
oil-immersion lens. Images in Supplementary Figure 3 were acquired using a 20X dry
lens using a Nikon Eclipse microscope connected to a digital camera (DS-Fi3) and
operated by Nikon software (NIS-Elements).

Raw confocal images were processed with ImageJ. Processed images were exported
as TIFF files and all figures were composed using Photoshop.

Acknowledgements

We would like to thank Alexander Picker, discussions with whom inspired and
prompted this study; and Kenzo Ivanovitch, Paride Antinucci and Paris Ataliotis for
reading and providing insightful feedback during the writing of this manuscript. The
animal facility at the CBMSO, the zebrafish facility at UCL and the Biological
Research Facility at St. Georges, University of London (SGUL) are acknowledged for
their help with animal care. The imaging facilities at the CBMSO, UCL and SGUL
are also acknowledged for their help with image acquisition.

Funding

This work has been funded by grants from the Spanish Government [BFU2014-
55918-P to FC; BFU2016-75412-R and BFU2016-81887-REDT to PB], the
Wellcome Trust [Seed Award in Sciences 213928/Z/18/Z to FC; Investigator Award
104682/Z/14/Z to SWW; Senior Research Fellowship 220273/Z/20/Z to IHB], the
Medical Research Council [MR/L003775/1 and MR/T020164/1to SWW and GG] and
the Fight for Sight Foundation [5153/5154 to FC]. An Institutional Grant from the
Ramón Areces Foundation to the CBMSO is also acknowledged.

Competing interests

The authors declare no competing or financial interests.

References
475
Aleström, P., D'Angelo, L., Midtlyng, P. J., Schorderet, D. F., Schulte-Merker,
S., Sohm, F. and Warner, S. (2020). Zebrafish: Housing and husbandry
recommendations. *Lab Anim* **54**, 213-224.
Andreazzoli, M., Gestri, G., Angeloni, D., Menna, E. and Barsacchi, G. (1999).
Role of Xrx1 in Xenopus eye and anterior brain development. *Development*
126, 2451-2460.
Andreazzoli, M., Gestri, G., Cremisi, F., Casarosa, S., Dawid, I. B. and
Barsacchi, G. (2003). Xrx1 controls proliferation and neurogenesis in
Xenopus anterior neural plate. *Development* **130**, 5143-5154.
Bailey, T. J., El-Hodiri, H., Zhang, L., Shah, R., Mathers, P. H. and Jamrich, M.
(2004). Regulation of vertebrate eye development by Rx genes. *Int J Dev Biol*
48, 761-770.
Bringmann, A. (2019). Structure and function of the bird fovea. *Anat Histol Embryol*
48, 177-200.

Bringmann, A., Syrbe, S., Görner, K., Kacza, J., Francke, M., Wiedemann, P.
and Reichenbach, A. (2018). The primate fovea: Structure, function and
development. *Prog Retin Eye Res* **66**, 49-84.
Carreres, M. I., Escalante, A., Murillo, B., Chauvin, G., Gaspar, P., Vegar, C.
and Herrera, E. (2011). Transcription factor Foxd1 is required for the
specification of the temporal retina in mammals. *J Neurosci* **31**, 5673-5681.
Carrington, B., Varshney, G. K., Burgess, S. M. and Sood, R. (2015). CRISPR-
STAT: an easy and reliable PCR-based method to evaluate target-specific
sgRNA activity. *Nucleic Acids Res* **43**, e157.
Cavodeassi, F., Ivanovitch, K. and Wilson, S. W. (2013). Eph/Ephrin signalling
maintains eye field segregation from adjacent neural plate territories during
forebrain morphogenesis. *Development* **140**, 4193-4202.
Cornish, E. E., Madigan, M. C., Natoli, R., Hales, A., Hendrickson, A. E. and
Provis, J. M. (2005). Gradients of cone differentiation and FGF expression
during development of the foveal depression in macaque retina. *Vis Neurosci*
22, 447-459.
Cornish, E. E., Natoli, R. C., Hendrickson, A. and Provis, J. M. (2004).
Differential distribution of fibroblast growth factor receptors (FGFRs) on
foveal cones: FGFR-4 is an early marker of cone photoreceptors. *Mol Vis* **10**,
1-14.
da Silva, S. and Cepko, C. L. (2017). Fgf8 Expression and Degradation of Retinoic
Acid Are Required for Patterning a High-Acuity Area in the Retina. *Dev Cell*
42, 68-81.e66.
Dehmelt, F. A., Meier, R., Hinz, J., Yoshimatsu, T., Simacek, C. A., Huang, R.,
Wang, K., Baden, T. and Arrenberg, A. B. (2021). Spherical arena reveals
optokinetic response tuning to stimulus location, size, and frequency across
entire visual field of larval zebrafish. *Elife* **10**.
Distel, M., Hocking, J. C., Volkmann, K. and Köster, R. W. (2010). The
centrosome neither persistently leads migration nor determines the site of
axonogenesis in migrating neurons in vivo. *J Cell Biol* **191**, 875-890.
Fish, M. B., Nakayama, T., Fisher, M., Hirsch, N., Cox, A., Reeder, R.,
Carruthers, S., Hall, A., Stemple, D. L. and Grainger, R. M. (2014).
Xenopus mutant reveals necessity of rax for specifying the eye field which
otherwise forms tissue with telencephalic and diencephalic character. *Dev Biol*
395, 317-330.
Flandrin, J. M. and Jeannerod, M. (1981). Effects of unilateral superior colliculus
ablation on oculomotor and vestibulo-ocular responses in the cat. *Exp Brain*
Res **42**, 73-80.
Halpern, M. E., Rhee, J., Goll, M. G., Akitake, C. M., Parsons, M. and Leach, S.
D. (2008). Gal4/UAS transgenic tools and their application to zebrafish.
Zebrafish **5**, 97-110.
Hatini, V., Tao, W. and Lai, E. (1994). Expression of winged helix genes, BF-1 and
BF-2, define adjacent domains within the developing forebrain and retina. *J*
Neurobiol **25**, 1293-1309.
Hernández-Bejarano, M., Gestri, G., Spawls, L., Nieto-López, F., Picker, A.,
Tada, M., Brand, M., Bovolenta, P., Wilson, S. W. and Cavodeassi, F.
(2015). Opposing Shh and Fgf signals initiate nasotemporal patterning of the
zebrafish retina. *Development* **142**, 3933-3942.

Herrera, E., Marcus, R., Li, S., Williams, S. E., Erskine, L., Lai, E. and Mason,
C. (2004). Foxd1 is required for proper formation of the optic chiasm.
Development **131**, 5727-5739.
Hoshino, A., Ratnapriya, R., Brooks, M. J., Chaitankar, V., Wilken, M. S.,
Zhang, C., Starostik, M. R., Gieser, L., La Torre, A., Nishio, M., et al.
(2017). Molecular Anatomy of the Developing Human Retina. *Dev Cell* **43**,
763-779.e764.
Kennedy, B. N., Stearns, G. W., Smyth, V. A., Ramamurthy, V., van Eeden, F.,
Ankoudinova, I., Raible, D., Hurley, J. B. and Brockerhoff, S. E. (2004).
Zebrafish rx3 and mab21l2 are required during eye morphogenesis. *Dev Biol*
270, 336-349.
Kolb, H., Nelson, R., Ahnelt, P., Ortuno-Lizaran, I. and Cuenca, N. (2020). The
Architecture of the Human Fovea.
Kozulin, P., Natoli, R., Bumsted O'Brien, K. M., Madigan, M. C. and Provis, J.
M. (2010). The cellular expression of antiangiogenic factors in fetal primate
macula. *Invest Ophthalmol Vis Sci* **51**, 4298-4306.
Kozulin, P., Natoli, R., O'Brien, K. M., Madigan, M. C. and Provis, J. M. (2009).
Differential expression of anti-angiogenic factors and guidance genes in the
developing macula. *Mol Vis* **15**, 45-59.
Kroll, F., Powell, G. T., Ghosh, M., Gestri, G., Antinucci, P., Hearn, T. J.,
Tunbak, H., Lim, S., Dennis, H. W., Fernandez, J. M., et al. (2021). A
simple and effective F0 knockout method for rapid screening of behaviour and
other complex phenotypes. *Elife* **10**.
Kwan, K. M., Otsuna, H., Kidokoro, H., Carney, K. R., Saijoh, Y. and Chien, C.
B. (2012). A complex choreography of cell movements shapes the vertebrate
eye. *Development* **139**, 359-372.
Loosli, F., Staub, W., Finger-Baier, K. C., Ober, E. A., Verkade, H., Wittbrodt,
J. and Baier, H. (2003). Loss of eyes in zebrafish caused by mutation of
chokh/rx3. *EMBO Rep* **4**, 894-899.
Loosli, F., Winkler, S., Burgtorf, C., Wurmbach, E., Ansorge, W., Henrich, T.,
Grabher, C., Arendt, D., Carl, M., Krone, A., et al. (2001). Medaka eyeless
is the key factor linking retinal determination and eye growth. *Development*
128, 4035-4044.
Mangrum, W. I., Dowling, J. E. and Cohen, E. D. (2002). A morphological
classification of ganglion cells in the zebrafish retina. *Vis Neurosci* **19**, 767-
779.
Mathers, P. H., Grinberg, A., Mahon, K. A. and Jamrich, M. (1997). The Rx
homeobox gene is essential for vertebrate eye development. *Nature* **387**, 603-
607.
Neuhauss, S. C., Biehlmaier, O., Seeliger, M. W., Das, T., Kohler, K., Harris, W.
A. and Baier, H. (1999). Genetic disorders of vision revealed by a behavioral
screen of 400 essential loci in zebrafish. *J Neurosci* **19**, 8603-8615.
Paquet, D., Bhat, R., Sydow, A., Mandelkow, E. M., Berg, S., Hellberg, S.,
Fälting, J., Distel, M., Köster, R. W., Schmid, B., et al. (2009). A zebrafish
model of tauopathy allows in vivo imaging of neuronal cell death and drug
evaluation. *J Clin Invest* **119**, 1382-1395.
Picker, A. and Brand, M. (2005). Fgf signals from a novel signaling center
determine axial patterning of the prospective neural retina. *Development* **132**,
4951-4962.

Picker, A., Cavodeassi, F., Machate, A., Bernauer, S., Hans, S., Abe, G.,
Kawakami, K., Wilson, S. W. and Brand, M. (2009). Dynamic coupling of
pattern formation and morphogenesis in the developing vertebrate retina. *PLoS*
Biol **7**, e1000214.
Pita, D., Moore, B. A., Tyrrell, L. P. and Fernández-Juricic, E. (2015). Vision in
two cyprinid fish: implications for collective behavior. *PeerJ* **3**, e1113.
Prykhozhij, S. V., Steele, S. L., Razaghi, B. and Berman, J. N. (2017). A rapid and
effective method for screening, sequencing and reporter verification of
engineered frameshift mutations in zebrafish. *Dis Model Mech* **10**, 811-822.
Robles, E., Laurell, E. and Baier, H. (2014). The retinal projectome reveals brain-
area-specific visual representations generated by ganglion cell diversity. *Curr*
Biol **24**, 2085-2096.
Roeser, T. and Baier, H. (2003). Visuomotor behaviors in larval zebrafish after
GFP-guided laser ablation of the optic tectum. *J Neurosci* **23**, 3726-3734.
Schmitt, E. A. and Dowling, J. E. (1999). Early retinal development in the zebrafish,
Danio rerio: light and electron microscopic analyses. *J Comp Neurol* **404**, 515-
536.
Stigloher, C., Ninkovic, J., Laplante, M., Geling, A., Tannhäuser, B., Topp, S.,
Kikuta, H., Becker, T. S., Houart, C. and Bally-Cuif, L. (2006).
Segregation of telencephalic and eye-field identities inside the zebrafish
forebrain territory is controlled by Rx3. *Development* **133**, 2925-2935.
Stih, V., Petrucco, L., Lavian, H. and Portugues, R. (2022). Bouter (v0.2.0).
Zenodo.
Suetterlin, P., Marler, K. M. and Drescher, U. (2012). Axonal ephrinA/EphA
interactions, and the emergence of order in topographic projections. *Semin*
Cell Dev Biol **23**, 1-6.
Sánchez-Arrones, L., Nieto-Lopez, F., Sánchez-Camacho, C., Carreres, M. I.,
Herrera, E., Okada, A. and Bovolenta, P. (2013). Shh/Boc signaling is
required for sustained generation of ipsilateral projecting ganglion cells in the
mouse retina. *J Neurosci* **33**, 8596-8607.
Takahashi, H., Sakuta, H., Shintani, T. and Noda, M. (2009). Functional mode of
FoxD1/CBF2 for the establishment of temporal retinal specificity in the
developing chick retina. *Dev Biol* **331**, 300-310.
Takahashi, H., Shintani, T., Sakuta, H. and Noda, M. (2003). CBF1 controls the
retinotectal topographical map along the anteroposterior axis through multiple
mechanisms. *Development* **130**, 5203-5215.
Voronina, V. A., Kozhemyakina, E. A., O'Kernick, C. M., Kahn, N. D., Wenger,
S. L., Linberg, J. V., Schneider, A. S. and Mathers, P. H. (2004). Mutations
in the human RAX homeobox gene in a patient with anophthalmia and
sclerocornea. *Hum Mol Genet* **13**, 315-322.
Weiss, O., Kaufman, R., Michaeli, N. and Inbal, A. (2012). Abnormal vasculature
interferes with optic fissure closure in lmo2 mutant zebrafish embryos. *Dev*
Biol **369**, 191-198.
Westerfield, M. (1993). *The Zebrafish Book: A Guide for the Laboratory Use of the*
Zebrafish (Brachydanio rerio). Eugene: University of Oregon Press.
Yin, J., Morrissey, M. E., Shine, L., Kennedy, C., Higgins, D. G. and Kennedy, B.
N. (2014). Genes and signaling networks regulated during zebrafish optic
vesicle morphogenesis. *BMC Genomics* **15**, 825.

Yoshimatsu, T., Schröder, C., Nevala, N. E., Berens, P. and Baden, T. (2020).
636   Fovea-like Photoreceptor Specializations Underlie Single UV Cone Driven
637   Prey-Capture Behavior in Zebrafish. *Neuron* **107**, 320-337.e326.

638   **Young, R. M., Hawkins, T. A., Cavodeassi, F., Stickney, H. L., Schwarz, Q.,**
639   **Lawrence, L. M., Wierzbicki, C., Cheng, B. Y., Luo, J., Ambrosio, E. M.,**
640   **et al.** (2019). Compensatory growth renders Tcf7l1a dispensable for eye
641   formation despite its requirement in eye field specification. *Elife* **8**.

642   **Zhou, M., Bear, J., Roberts, P. A., Janiak, F. K., Semmelhack, J., Yoshimatsu, T.**
643   **and Baden, T.** (2020). Zebrafish Retinal Ganglion Cells Asymmetrically
644   Encode Spectral and Temporal Information across Visual Space. *Curr Biol* **30**,
645   2927-2942.e2927.

646   **Zimmermann, M. J. Y., Nevala, N. E., Yoshimatsu, T., Osorio, D., Nilsson, D. E.,**
647   **Berens, P. and Baden, T.** (2018). Zebrafish Differentially Process Color
648   across Visual Space to Match Natural Scenes. *Curr Biol* **28**, 2018-2032.e2015.

649   **Štih, V., Petrucco, L., Kist, A. M. and Portugues, R.** (2019). Stytra: An open-
650   source, integrated system for stimulation, tracking and closed-loop behavioral
651   experiments. *PLoS Comput Biol* **15**, e1006699.
652
653

Figure legends

Figure 1: foxd1 expression is reduced in *rx3*[-/-] mutant optic vesicles and lost in *rx3*[-/-] upon combined abrogation of Shh and Fgf.

(A-U) Frontal sections at the level of the forming telencephalon, optic vesicles and hypothalamus, with dorsal up, genotype/treatment top right and genes analysed bottom left.
(A-D) Expression of *rx3* (A) and *nkx2.1* (B) in 4ss wildtype/*rx3*[-/-] embryos (indistinguishable at this stage with a representative embryo shown) and *emx3* in 10ss wildtype (C) and *rx3*[-/-] (D) embryos highlighting the prospective telencephalic (t), hypothalamic (hy) and eye-forming (ov) domains.
(E-M) Expression of *foxg1* (E-H) and *foxd1* (J-M) in 4ss (E,G,J,L) and 10ss (F,H,K,M) embryos (genotype top right). Panel (I) shows double *in situ* hybridisation of *foxd1* or *foxg1* and *nkx2.1* (red) to show relationship between optic vesicles and hypothalamus.
(N-U) 10ss stage embryos showing expression of *foxg1* (N-Q) or *foxd1* (R-U) in embryos treated with DMSO (N,P,R,T) or cyclopamine+SU5402 (O,Q,S,U).
(V) schematic representation of the conditions in (R-U).
Scale bar in this and all figures unless specified: 100μm. Dashed lines in this and all figures highlight the contour of the optic vesicles.
Abbreviations: ov, optic vesicles; hy, hypothalamus; dhy, dorsal hypothalamus; t, telencephalon.

Figure 2: **Loss of** *foxd1* **results in retinae with expanded nasal, and reduced temporal character.**

(A) *foxd1* sequence comparison between nucleotides 163 and a 190 of the open reading frame, highlighting the 10 base-pair deletion (Δ10, bottom row) in the *foxd1*[cbm16] mutant.
(B) Diagrammatic representation of the Foxd1 protein, depicting in red the deleted region in the *cbm16* allele. The asterisk in (A-B) marks the position at which a newly generated stop codon is present.
(C-J) Lateral views with nasal to the left of 48hpf (C-F) and 5dpf (G-J) retinae showing expression of nasal or temporal markers in *foxd1*[-/-] mutants and siblings. The markers assessed and the genotypes of the retinae are detailed bottom left and top right respectively. Asterisk in (D,F,H,J) highlights altered expression in *foxd1* mutants.
(K-M) Dorsal view of the tectum of *foxd1*[-/-] mutant (L-M) and sibling embryos (K), with nasal RGC arbors labelled with DiO (green) and temporal RGC arbors with DiI (red). Insets in (K-M) show dorsal views of the corresponding eye. Tectum and eye orientation in panels (K-M) is indicated by the double arrow in (K).
(N-P) PKCα immunostaining (red) in lateral views with nasal to the left of 8dpf wildtype (N), *foxd1*[-/-] mutant (O) and *tg{rx3:Gal4};UAS:Shh* (P) eyes. Nuclei are counterstained with dapi (blue). Insets in (N-P) are schematics of the eyes in the corresponding panels, highlighting the extent of temporal character (purple), as assessed by gene expression, and the areas of enriched PKCα (red). Arrows indicate the position of PKCα enrichment.

Figure 3: **OKR assays reveal severe visual deficiencies in** *foxd1$^{-/-}$* **but not** *tg{rx3:Gal4};UAS:Shh* **embryos.**

(A) Diagrammatic representation of the changes in naso-temporal pattern and retinotectal projections observed in the four categories of fish tested by OKR. Genotype is detailed at the top of each diagram.

(B-C) Graphs detailing eye gain (expressed in a 0-1 range) in response to changes in spatial frequency (expressed in cycles per screen; Bi,Ci), grating velocity (expressed in degrees per second; Bii,Cii) and contrast (expressed in a 0-1 range; Biii,Ciii) in wild type (black, n=5 in B), *foxd1$^{+/-}$* (blue, n=27 in B), *foxd1$^{-/-}$* (red, n=8; in B), *tg{rx3:Gal4}* (black [rx3:gal4], n=10 in C) and *tg{rx3:Gal4};UAS:Shh* (red, [+UAS:shh], n=8 in C). Graphical representation was generated with GraphPad Prism5 (ANOVA double test, followed by a Bonferroni test) and statistical significance calculated with a t-student test (***p<0.001).

(D-E) Range (expressed in degrees) of slow phase eye movement (left graph) and peak velocity (expressed in degrees per second) during fast phase eye movement (right graph) in *foxd1$^{-/-}$* versus wildtype and *foxd1$^{+/-}$* fish (D), and *tg{rx3:Gal4}* versus *tg{rx3:Gal4};UAS:Shh* fish (E).

(F) Fraction of correct turns for wildtype larvae (n=21) and *foxd1* crispants (*foxd1 F0*, n=21) when subjected to left- and right- oriented whole field motion stimuli. Only directional bouts (i.e. left- and rightward swims, without forward swims) were considered for quantification.

(G) Number of swim bouts during pause intervals and whole field motion stimuli for wildtype larvae (left, n=21) and *foxd1* crispants (right, n=21). Bar heights represent the means across all fish in each group; error bars represent the standard error of the mean (SEM).

*Figure 4***: Summary of roles for Rx3, Fgfs and Shh in establishment of nasal and temporal character in the developing eye primordium.**

(A) Schematic representation of wildtype and *rx3$^{-/-}$* embryos depicting the changes in *foxg1* (blue) and *foxd1* (magenta) expression upon abrogation of Shh and Fgf activities, singly or in combination.

(B) Diagram detailing the proposed regulatory network controlling NT patterning. Shh promotes *foxd1* expression in the ventral (future temporal) half of the optic primordium. Fgfs induce *foxg1* in the dorsal (future nasal) optic primordium and repress *foxd1* expression, contributing to its confinement ventrally. Cross-repression between FoxG1 and FoxD1 at the border between the dorsal and ventral halves of the eye primordium subsequently refines the naso-temporal subdivision. Although induction of *shh* and *fgf8* expression occurs independently, *fgf8* expression is lower in absence of Shh signalling (Hernández-Bejarano et al., 2015) suggesting Hh signalling initially promotes Fgf signalling. Rx3 induces *foxd1* expression and provides competence to optic vesicle cells to enhance *foxd1* expression in prospective temporal retina in response to Shh signalling.

(C) Schematic representation of a differentiated optic cup with nasal (blue) and temporal (magenta) character, and the HAA (red) highlighted. Nasal character is expanded and HAA and temporal character are absent in *foxd1$^{-/-}$* mutants (right) as compared to the wild type (left).

A

wild type   *rx3-/-*

DMSO

Shh-

Fgf-

Shh-
Fgf-

B

Fgf

foxg1

foxd1 ← **Rx3**

Shh

C

HAA

wild type   *foxd1-/-*