1   Complexities of recapitulating polygenic effects in natural populations: replication of genetic
2   effects on wing shape in artificially selected and wild caught populations of *Drosophila*
3   *melanogaster.*
4
5

6   Katie Pelletier*[1], William R. Pitchers[2,8]*, Anna Mammel[2,7], Emmalee Northrop-Albrecht[2,5],
7   Eladio J. Márquez[3,6], Rosa A. Moscarella[3,4], David Houle[3], Ian Dworkin[1,2]
8   *Co-first authors
9   Corresponding authors: dworkin@mcmaster.ca, dhoule@bio.fsu.edu

10   Author affiliations: 1 – Department of Biology, McMaster University

11   2 – Department of Integrative Biology, Michigan State University

12   3 – Department of Biological Science, Florida State University

13   4 – Current address: Department of Biology, University of Massachusetts

14   5 – Current address Division of Gastroenterology and Hepatology, Mayo
15   Clinic, Rochester MN
16   6 – Current address: Branch Biosciences, Waltham Cambridge MA
17
18   7 – Current address: Neurocode USA. 3548 Meridian St, Bellingham, WA
19   98225
20   8 – Current address: BiomeBank, Adelaide 5031, South Australia
21

Abstract

Identifying the genetic architecture of complex traits is important to many geneticists,

including those interested in human disease, plant and animal breeding, and evolutionary

genetics. Advances in sequencing technology and statistical methods for genome-wide

association studies (GWAS) have allowed for the identification of more variants with smaller

effect sizes, however, many of these identified polymorphisms fail to be replicated in

subsequent studies. In addition to sampling variation, this failure to replicate reflects the

complexities introduced by factors including environmental variation, genetic background, and

differences in allele frequencies among populations. Using *Drosophila melanogaster* wing

shape, we ask if we can replicate allelic effects of polymorphisms first identified in a GWAS

(Pitchers et al. 2019) in three genes: *dachsous (ds)*, *extra-macrochaete (emc)* and *neuralized*

(neur), using artificial selection in the lab, and bulk segregant mapping in natural populations.

We demonstrate that multivariate wing shape changes associated with these genes are aligned

with major axes of phenotypic and genetic variation in natural populations. Following seven

generations of artificial selection along the *ds* shape change vector, we observe genetic

differentiation of variants in *ds* and genomic regions containing other genes in the hippo

signaling pathway. This suggests a shared direction of effects within a developmental network.

We also performed artificial selection with the *emc* shape change vector, which is not a part of

the hippo signaling network, but showed a largely shared direction of effects. The response to

selection along the *emc* vector was similar to that of *ds*, suggesting that the available genetic

diversity of a population, summarized by the genetic (co)variance matrix (**G**), influenced alleles

captured by selection. Despite the success with artificial selection, bulk segregant analysis using

natural populations did not detect these same variants, likely due to the contribution of

environmental variation and low minor allele frequencies, coupled with small effect sizes of the

contributing variants.

Introduction

Dissecting the genetic architecture underlying complex traits remains challenging,

because of the joint contributions of many alleles of small effect, genotype-by-environment

interactions, and other factors. Progress in sequencing technology in conjunction with

development of GWAS statistical methodologies has enabled identification of loci contributing

to numerous complex traits and diseases. However, such mapping approaches identify only a

subset of loci contributing to trait variation (Visscher et al., 2017). In part, this reflects the low

power to detect rare alleles, and those with small effects (Tam et al., 2019). For alleles that are

relatively common in a population, replication rates between GWAS studies are high, even

when effect sizes are small (Marigorta et al., 2018). However, GWAS studies have failed to

replicate the effects observed in many candidate gene studies, in part due to the fact that many

alleles identified in these studies are rare in populations, and require very large cohorts to

detect (Fritsche et al., 2016; Ioannidis et al., 2011).

In cases where an association is replicated between studies, the magnitude of the effect

can vary substantially between different cohorts or populations (CONVERGE consortium, 2015;

Marigorta et al., 2018). Differences can arise because of genetic background due to epistatic

gene by gene (GxG) interactions, or due to gene-by-environment (GxE) interactions. The initial

estimates of effect size will be biased upwards if statistical testing in the initial cohort is used to

determine which SNPs are chosen for replication studies. It is important to understand which of

these causes of differences in effect size are of practical significance when we want to

generalize results to different populations or environments.

In this study, we focus on the issue of replication in a multivariate context, where the

joint inheritance of multiple features are simultaneously investigated. We will refer to the suite

of measured features as a 'multivariate trait' for convenience. In this case, what we want to

estimate is the vector of effects of each SNP on all measured features. Each SNP may have a

unique combination of effects. Univariate effects vary only in magnitude, as we can only infer

effects on a single feature. For a multivariate trait, estimated genetic effects vary in magnitude,

the sum of effects on all traits, and also in direction, how the total effect is allocated among

different features (Melo et al., 2019). The ability to study the direction along with the

magnitude of genetic effects provides an additional and important way of assessing

repeatability. For a univariate trait, there is a 50% chance that the replicate estimate will be in

the same direction as the original estimate, even with no true effect. By contrast, the

probability of a "replicated" genetic effect sharing a similar direction by chance alone decreases

as the number of measured features increases (Marquez and Houle, 2015; Stephens, 2013).

Studying genetic effects in a multivariate context is beneficial in other ways. First, it has

been demonstrated both empirically and via simulations, that genetic mapping for multivariate

traits generally increases statistical power over trait by trait analyses (Fatumo et al., 2019;

Pitchers et al., 2019; Porter and O'Reilly, 2017; Shriner, 2012). Second, some multivariate traits

cannot be sensibly reduced to a single measurement. The wing shape we study is a great

example of such a multivariate trait. We have good reason to believe that wing shape is

important for flight (Ray et al., 2016), but we cannot yet say that any feature, such as wing

length or width, is more or less important than any other. Natural selection on wing shape may

affect any or all combinations of measurements.

Perhaps most importantly, traits are not inherited in isolation, but are the joint outcome

of an integrated developmental process that results in extensive genetic correlations that can

have important effects on evolution. The main source of such correlations are the patterns of

pleiotropic effects generated by mutational effects. Multivariate studies of inheritance allow

pleiotropic effects to be estimated in a rigorous and justifiable manner (Melo et al., 2019). The

multivariate breeder's equation, $\Delta z = G\beta$, enables short term prediction of evolutionary

responses. Key to understanding how populations respond to selection in the short term

requires an understanding of properties of the genetic (co)variance matrix ($\mathbf{G}$), and in particular

the axis of greatest genetic variation, $\mathbf{g}_{max}$. Studies demonstrate that the direction of $\mathbf{g}_{max}$

influences evolutionary trajectories (Blows and McGuigan, 2015; McGuigan, 2006; Schluter,

1996). The degree to which genetic effects associated with particular variants align to major

axes of genetic (co)variance, expressed through $\mathbf{G}$, may provide insights into which alleles are

most likely to be "captured" by selection (Pitchers et al., 2019).

Due to the polygenic nature of complex traits, including multivariate ones, it is

important to consider not only the direction of effect for alleles in a single gene but also

correlated effects between genes contributing to the phenotype. Interestingly, initial

comparisons of directions of genetic effects among induced mutations in two *Drosophila*

melanogaster wing development pathways showed only partially correlated effects on wing

shape within and between pathways (Dworkin and Gibson, 2006). However, recent work has

demonstrated that despite large differences in magnitude, the direction of genetic effects of

variants segregating in populations are sometimes similar to those from validation experiments

using RNAi knockdown of those same genes (Pitchers et al., 2019). Additionally, Pitchers et al.

(2019) demonstrated this shared direction of effect could also be shared between a SNP and

RNAi knockdown of other genes in the same signaling pathway, such as those involved with

hippo signaling, a key pathway involved with wing growth and morphogenesis (Pan et al.,

2018).

Pitchers et al. (2019) identified over 500 polymorphisms contributing to wing shape

variation in the *Drosophila* genetic resource panel (DGRP). Among these, the hippo pathway

was over-represented in SNPs associated with wing shape (Pitchers et al., 2019). The degree to

which identified hippo pathway variants reflect allele specific effects, differences in magnitude

of genetic effects, and even the large statistical uncertainty associated with genetic effects of

small magnitude are unclear. Given common dominance patterns, and the likely non-linear

genotype-phenotype relationships of most genetic effects, small to moderate changes in gene

function may result in modest phenotypic effects (Green et al., 2017; Melo et al., 2019; Wright,

1934). Large effect mutants and many RNAi knockdown studies have moderate to large

phenotypic effects that are not reflective of the magnitude of genetic effects of SNPs

contributing to phenotypic variance in natural populations.

The expression of genetic effects also depends on genetic and environmental context,

with gene-by-gene (GxG) and gene-by-environment (GxE) interactions contributing to

phenotypic variation. The context-dependence of genetic effects for a multivariate trait has

been demonstrated for *Drosophila* wing shape. Variants in *Epidermal growth factor receptor*

(*Egfr*), influencing *Drosophila* wing shape are replicable in both lab reared, and wild-caught

cohorts (Dworkin et al., 2005; Palsson et al., 2005; Palsson and Gibson, 2004). However, in

replication studies, effect sizes of alleles were diminished in both outbred populations and wild

cohorts. In the latter case the same variant explained 1/10 of the phenotypic variance explained

in the initial study (Dworkin et al., 2005). Interestingly, in a series of experimental crosses

among strains, the effects of the SNP were replicable for direction and magnitude in multiple

experimental assays and crossing schemes. Despite this, the genetic effect on wing shape from

this SNP largely disappeared in one natural population (Palsson et al., 2005). A number of

reasons have been proposed for the failure to replicate genetic effects including environmental

effects, differences between controlled lab and natural environments (Dworkin et al., 2005),

and genetic background (Greene et al., 2009), among others. Because both environment and

genetic background likely affect the genotype-phenotype map in a non-linear fashion (Wright,

1934), it is important to test observed associations in other experimental contexts.

A promising approach to confirm the estimated effects of candidate genetic variants is

to test whether they respond to artificial selection in the direction of the inferred effect. This

approach is particularly relevant to evolutionary questions, but has rarely been used. In this

study, we use artificial selection and bulk segregant analysis (BSA), to replicate and validate

SNPs associated with three genes, previously identified in a GWAS of *Drosophila* wing shape

(Pitchers et al., 2019); *dachsous (ds), an atypical cadherin involved with hippo signaling;* the

transcriptional co-repressor *extra-macrochetae (emc),* and the E3 ubiquitin ligase *neuralized*

(neur), involved with Notch signaling. Using the vectors of shape change based on RNAi

knockdowns of each gene, we demonstrate that the direction of shape change for these genetic

effects is aligned with major axes of natural phenotypic and genetic variation. Using artificial

selection based on the direction of shape change defined by RNAi knockdown, we were able to

replicate the effects observed for *ds,* but not *emc*, likely due to the available genetic diversity in

the population. We then asked if these effects could be replicated in a natural population using

a bulk segregant approach, observing little evidence for replication in these samples. We

discuss our results in the context of the replicability of genetic effects and the shared direction

of genetic effects due to shared developmental processes.

Methods

Source Populations and phenotypic analysis

Drosophila strains

Phenotype data for the *Drosophila* genetic resource panel (DGRP) was collected for 184

strains as part of a GWAS study as described in Pitchers *et al* (2019). Genotype data for these

strains was obtained from freeze 2 of the DGRP (Huang et al., 2014). For replication using

artificial selection, 30 DGRP strains were used: DGRP-149, 324, 383, 486, 563, 714, 761, 787,

796, 801, 819, 821, 822, 832, 843, 849, 850, 853, 859, 861, 879, 887, 897, 900, 907, 911, 913.

These strains were selected to increase genetic variation at the *ds* locus (Supplemental Figure 1,

Table 1). Reciprocal pairwise crosses between the 30 selected DGRP strains were used to create

heterozygotes and these 30 heterozygous genotypes were successively pooled for 4

subsequent generations, allowing for recombination. After pooling, the synthetic outbred

population was maintained for approximately 47 subsequent generations (allowing for

recombination) before the start of artificial selection experiments.

For the replication in wild-caught populations using BSA, individuals were collected via

sweep-netting from orchards and vineyards in Michigan and after species identification, stored

in 70% ethanol. In 2013 and 2014, cohorts were collected from Fenn Valley Winery (FVW13 and

FVW14 respectively, GPS coordinates: 42.578919, -86.144936). Additionally in 2014, cohorts

were collected from Country Mill Orchard (CMO, GPS coordinates: 42.635270, -84.796706), and

Phillip's Hill Orchard (PHO, GPS coordinates: 43.117981, -84.624235). For all collected cohorts,

except for the FVW14 collection, only males were used in this study given difficulties

distinguishing *Drosophila melanogaster* and *D. simulans* females morphologically. For the

genomic analysis of the FVW14 wild caught population (below) we utilized both males and

females as the number of individuals was insufficient otherwise. For the collection where

females were included in the study, there is no evidence of contamination with *D. simulans* as

all dissected wings were classified as *D. melanogaster* using linear discriminant analysis (LDA).

LDA was trained using male wings from the collected *D. melanogaster* data set and males from

D. simulans. There was 100% agreement between the classification of females within each

species with our phenotypic classification, indicating that it is unlikely that *D. simulans* females

were included in our samples (Supplemental Figure 2).

<u>Morphometric Data</u>

Landmark and semi-landmark data were captured from black and white TIFF images

using the pipeline described in Houle et al. (2003). First, two landmark locations, the humeral

break and alula notch, were digitized using tpsDig2 (version 2.16). Wings (Van der Linde 2004–

2014 , v3.72) software was used to fit nine cubic B-splines, and manually correct errors. All

shape data was subjected to Procrustes superimposition (registration), removing the effects of

location, isometric scaling, and and minimizing effects of rotation, , via an iterative least

squares approach (Rohlf and Slice, 1990). Generalized Procrustes superimposition (registration)

and extraction of 14 landmarks and 34 semi landmarks was done using CPR v1.11 (Márquez

2012–2014, Figure 1). Sliding of semi-landmarks utilized minimization of Procrustes Distance as

the objective function. Superimposition results in the loss of 4 possible dimensions of variation

while semi-landmarks are constrained to vary along one "axis", restraining these points to

approximately a single dimension of variation each. This results in a total of ~58 available

dimensions of shape variation, that can be summarized using the first 58 Principal components

(PCs). Allometry was adjusted for in the analysis by fitting a model for landmark coordinates

onto centroid size, and using the residuals from this model (Klingenberg, 2022). By accounting

for the allometric component of shape, shape variation associated with size variation can be

accounted for (Supplemental Figure 3). For most analyses, 'allometry corrected' shape data

were used, with the exception of shape models fit using the Geomorph package in R, where

Procrustes landmarks were used and centroid size was included as a predictor in the model.

<u>Generation of shape vectors for artificial selection and bulk segregant analysis</u>

A panel of shape change vectors was estimated using the progesterone-inducible

Geneswitch GAL4, under the regulation of an ubiquitous *tubulin* driver, to drive the expression

of RNAi for genes of interest (*ds, emc, neur*), as previously described in Pitchers *et al.,* 2019.

GAL4 expression was induced throughout larval development by adding mifepristone, an

analog of progesterone, to the larval food. Knockdown was varied by assaying phenotypes at

mifepristone concentrations of 0.3, 0.9, and 2.7 μM, plus a control without mifepristone. Wing

shape change associated with knockdown of the gene of interest was estimated using

multivariate regression of shape on concentration of mifepristone. Shape change vectors

estimated from the RNAi experiments for *ds,emc* and *neur,* were used in this experiment

(Figure 1b, Supplemental Figure 4). The magnitude ('length') of the vector measures how much

shape change occurs per unit change in mifepristone. In general, vectors of greater magnitudes

enable better estimate of direction of effect for shape change. As reported in Pitchers et al.,

(2019), the magnitude (l^2-norm) of vectors for RNAi knockdown of these genes are 5.5 for *ds*,

2.8 for *neur*, and 0.44 for *emc*.

Shape data collected as part of a previous study (Pitchers et al., 2019) was used to

assess the relationships between shape change vectors from the RNAi titration and $\mathbf{g_{max}}$, the

first eigenvector of the **G** matrix estimated from DGRP line means. The effects of sex, centroid

size and their interaction were removed using a linear model and these residuals were used to

calculate shape score by projecting the data (see Supplemental Figure 5) onto the shape change

vector estimated in each knockdown experiment. To assess major axes of genetic variation

among DGRP strains, principal component analysis was performed on allometry adjusted model

residuals (Supplemental Figure 5B). PCA was done in a similar manner for individuals from the

wild caught cohorts. Correlations between the first three eigenvectors ("genetic PCs" including

$\mathbf{g_{max}}$), the first three PCs from the wild caught cohorts and the shape scores for *ds*, *emc* and

neur were calculated (Figure 1a, Supplemental Figure 5). From this, *ds, emc* and *neur* shape

change vectors were selected for further experiments given high correlation with directions of

natural genetic variation (Figure 1, Supplemental Figure 5). Note, as described below, while *ds*

and *emc* were used for artificial selection, due to the similar response between them, we used

ds and substituted *neur (*for *emc)* for the BSA.

<u>Artificial selection of synthetic outbred population</u>

The synthetic outbred population resulting from pooling DGRP lines was used as the

parent population for artificial selection. Both the *ds* and *emc* artificial selection experiment

were carried out with three independent replicates of each "up" and "down" selection regimes,

along with unselected control lineages. Each generation, wings of live flies were imaged using

the 'wingmachine' system and shape data collected (Houle et al., 2003, Van der Linde 2004–

2014 ,v3.72). Shape scores were calculated by projecting the data onto the *ds* or *emc* shape

change vector as described above, and the 40 individuals each with highest or lowest shape

scores, were selected to found the next generation (Supplemental Figure 5A). For the control

lineages, 40 individuals were randomly selected for the next generation within each replicate

lineage. Following seven generations of selection, 75 individuals from each lineage were

selected for pooled sequencing, described below. The response to selection was evaluated both

by computing Procrustes distance (PD) between average shape of wings between generations

one and seven, and using shape scores (projections) with a linear mixed effect model allowing

for the fixed effect factors of treatment and sex, continuous predictors of centroid size and

generation, with third order interactions among these effects. The effect of generation was

allowed to vary by replicate lineages (lmer(ds ~ (CS + Sex + line + gen0)^3 + (1 +

gen0|line:rep)). Realized heritabilities were estimated separately for up and down selection

lineages, from the slope of the regression of cumulative selection differentials on cumulative

selection response, averaging over sex and with a random effect of replicate lineage.

<u>Wild populations</u>

For the BSA, wings for wild caught individuals were dissected and mounted in 70%

glycerol in PBS. Images of wings were captured using an Olympus DP30B camera mounted on

an Olympus BX51 microscope (Olympus software V.3,1,1208) at 20X magnification. When

possible, both left and right wings were dissected, imaged and averaged to calculate an

individual's mean shape. For some individuals a wing was damaged so only one wing could be

used. Shape was captured as described above. The total number of individuals phenotyped

from each cohort can be found in Supplemental table 1.

To remove allometric effects in the data, shape was regressed onto centroid size and

the model residuals were used for all subsequent morphometric analysis. Only data from males

was used to compare shape in wild populations, although, including females from the FVW14

population and regressing shape onto centroid size and sex gave equivalent results

(Supplemental Figure 6). To test for shape differences between collection cohorts, the effect of

centroid size and collection cohort on shape were modeled (procD.lm(shape ~ CS + pop_year))

using the procD.lm function in Geomorph v 3.1.3. (Adams and Otárola-Castillo, 2013) and

distances between populations were calculated using the pairwise function. To select

individuals for sequencing, a 'shape score' was calculated using the method described above.

Shape data was projected onto the vector of shape change defined by the *ds* or *neur*

knockdowns. The *emc* projection vector was not used for BSA due to the high similarity with *ds*

shape change (Figure 1), and the similarity of the selection response. Its inclusion would result

in selection of largely the same cohorts of individuals for sequencing for both *ds* and *emc*. As an

alternative, we utilized the *neur* shape vector as it was largely uncorrelated with that of *emc*

and *ds*, but strongly correlated with natural variation in shape. The 75 most extreme individuals

on the shape score distribution, within each wild-caught cohort, were selected for pooled

sequencing. Allele frequencies within each population was estimated by sequencing 75 random

individuals within each cohort. The difference vector between mean shapes of selected pools

(within each population) was used to calculate Procrustes distance (PD) between pools and the

correlation of this shape change vector with the selection vector used. An estimate of genetic

distances between populations was calculated using allele frequencies (mapping pipeline

described below) in the pools of the 75 randomly selected individuals using Bray's distance with

the vegdist() function from the vegan package (v2.6-2) in R.

<u>Sequencing and Genomic Analysis</u>

296   DNA extractions from pools of selected individuals was performed using a Qiagen

DNeasy DNA extraction kit. Library preparation and Illumina sequencing was performed at the

research technology support facility at Michigan State University. All library samples were

prepared using the Rubicon ThruPLEX DNA Library Preparation kit, without a procedure for

automatic size selection of samples. Paired end libraries (150bp) were sequenced using Illumina

HiSeq 2500, with each sample (either one pool of 75 individuals in the BSA or one pooled

replicate lineage in the artificial selection) being run on two lanes.

303   Reads were trimmed with Trimmomatic (v0.36) to remove adapter contamination and

checked for quality using FastQC prior to alignment (Bolger et al., 2014). Trimmed reads were

aligned to the *Drosophila melanogaster* genome (v6.23) using BWA-MEM (v0.7.8)(Li and

Durbin, 2010). Sequencing replicates of the same biological samples were merged using

SAMtools (v1.11). PCR duplicates were removed using Picard with the MarkDuplicates tool (v

2.10.3) and reads with a mapping quality score less than 20 were removed using SAMtools (Li et

al., 2009). A local realignment around indels was performed using GATK using the

IndelRealigner tool (v3.4.46). For artificial selection experiments, reads were merged for all up,

down and control selection lines as replicates lineages were independent. For wild cohorts,

pools were not merged between populations. mpileup files were created using SAMtools and

used for subsequent genomic analysis. Highly repetitive regions of the *Drosophila* genome were

identified and subsequently masked in mpileup files using RepeatMasker (v4.1.1) with default

settings. INDELs and regions within 5bp of an indel were identified and masked using

popoolation2 scripts. Population genetic statistics were calculated using PoPoolation (v1.2.2)

and PoPoolation2 (v1.201) (Kofler et al., 2011b, 2011a).

For the BSA in the wild-caught cohorts, a modified Cochran-Mantel-Haenszel (CMH) test

was used to measure significantly differentiated sites between pools of individuals. Sampling

effects were accounted for using the ACER package (v.1.0) in R, assuming $N_e = 10^6$ with 0

generations of differentiation between selected pools (Spitzer et al., 2020). To adjust for

multiple testing, the p-value was corrected using a Benjamini-Hochberg correction (Benjamini

and Hochberg, 1995) with an adjusted alpha of 0.05. For each significant site from the CMH

test, using an adjusted p-value cut-off of 0.05, we identified the nearest gene using BEDtools

(v2.19.1) (Quinlan and Hall, 2010). In addition, to account for sampling variation, we sampled

genomic coverage to 75x for all samples, dropping sites that did not meet this threshold and

repeating the CMH test. We confirmed that there was no association between genetic and

shape differentiation between populations, and that the populations do not show strong

phenotypic differentiation based on either overall shape variation, or shape scores used to

identify selected individuals for BSA (Supplemental Figures 3 and 7). There was some variation

among populations in overall wing size (Supplemental Figure 8), however we (assuming

common allometry) adjusted for allometric effects on shape.

For artificial selection experiments, F_{ST} was calculated in 5000bp windows. We chose

this window size as it is expected that blocks of LD in the synthetic outbred population will be

much larger in comparison to that of the wild caught samples (King et al., 2012a; King et al.,

2012b; Marriage et al., 2014). This statistic was used to compare the "up" selected pools to the

"down" selected pools to help identify regions of differentiation between selected populations.

For the artificial selection comparisons, genes in regions of high F_{ST} were identified by

finding overlaps between outlier windows and annotated *Drosophila* genes using

GenomicRanges (v1.46.1) in Bioconductor. High F_{ST} was defined as F_{ST} values greater than three

standard deviations above the mean. GO terms associated with identified genes were

annotated using TopGO package (v2.34.0) (Alexa et al., 2006) in Bioconductor. GO enrichment

was then performed to identify those terms overrepresented in the identified list using TopGO

and a Fisher's exact test. Over representation of 2 GO terms in outlier windows (hippo

signaling, GO:0035329; negative regulation of hippo signaling GO:0035331) were tested using a

permutation test that randomly sampled genomic windows from the total windows for which

F_{ST} was calculated and the permutation was run 1000 times. The distribution of the ratio of

observed to expected genes annotated with the term of interest within randomly sampled

regions was compared to the number observed in the data.

<u>Verification of *ds* indel in DGRP</u>

Sanger sequencing was performed on individuals from a cross between DGRP lines predicted to

have the polymorphism (DGRP 195, 28, 96, 48, 59, 801) and those without (DGRP 129, 301, 69,

385, 75, 83, 491, 34, 774) crossed to a line carrying a deletion in the region of interest (BDSC

24960) to account for potential residual heterozygosity in otherwise inbred strains. DNA was

prepared by incubating flies in DNA extraction buffer (1mM EDTA, 25mM NaCl, 10 mM TrisHCl

pH 7.5) for 10 minutes, followed by storage at -20 C. PCR application of the region of interest

(Forward primer: ggagtacaagttgctcgaac: Reverse Primer: cagatcgtgttccctttagc) using Taq DNA

polymerase (Gene DirectX) (PCR mix: 1uL DNA, 1uL forward primer, 1uL reverse primer, 0.5uL

10mM dNTPs, 2uL 10x PCR Buffer, 0.1 uL taq, H20 to 20uL). PCR conditions were as follows: 5

minutes 95°, (30 seconds 95°, 30 seconds 55°, 30 seconds 72°)x30. Reactions were checked on a

gel and cleaned with the GenepHlow^{TM} Gel/PCR Kit (Geneaid). Sanger sequencing reactions

were performed by the Mobix Lab at McMaster University. All alignments were created using

ClustalOmega (Madeira et al., 2022).

Data Availability:

All code and processed data needed to complete the analysis is available on GitHub at:

https://github.com/DworkinLab/WingShapeBSA/. A static version of the repository is available

on figshare (https://doi.org/10.6084/m9.figshare.22141154.v1).

All raw sequence data available as part of the NCBI Short Read Archive, BioProject

PRJNA936488 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA936488/), with individual

sequence accessions SAMN33354503 - SAMN33354634.

Results

dachsous (ds) shape change is aligned with major axes of genetic and phenotypic variation in

natural populations

To assess the relationship between shape change vectors and axes of natural variation

described in the the DGRP, mean shape vectors were calculated for each DGRP strain, then

used in a PCA to summarize axes of variation among strains. Mean shape vectors for each strain

of DGRP were projected onto shape change vectors for _ds emc_, and _neur_, defined from the

RNAi knockdowns (see Supplementary Figure 5, which visually explains the procedure),

generating gene specific "shape scores". Correlations between shape scores for individual DGRP

projected onto the shape change vectors (Figure 1, Supplemental Figure 5), and with PC1

generated from the DGRP (PC1$_{DGRP}$) strains was estimated (PC1$_{DGRP}$-_ds_: r = -0.56; PC1$_{DGRP}$-_emc_: r

= -0.45; Figure 1). The correlation of the DGRP data, projected onto each of the _ds_ and _emc_

shape change vectors was also correlated (Figure 1, _ds-emc_: r = 0.69). This is likely due to the

correlation between gene specific shape change vectors themselves (r = 0.65), based on RNAi

titration experiments. Projections of the DGRP data onto the vector defining the _neur_ shape

change is aligned with PC1 (PC$_{DGRP}$-_neur_: r = -0.69) and PC3 (PC$_{DGRP}$-_neur_: r = -0.64), indicating

this as an important axis of shape variation in this population (Figure 1), that is moderately

similar to projections onto _ds_ (_ds-neur_: r = 0.56) and very similar to _emc_ (_neur-emc_: r = 0.83)

shape change vectors. Interestingly, the strength of the correlation for the DGRP strains

projected onto these vectors, differs from the magnitude of correlations for the RNAi titration

vector of _neur_ with that of _ds_ (r = 0.034) or _emc_ (0.3). Because of these observed correlations,

and previous associations observed (Pitchers et al., 2019) _ds, emc_ and _neur_ were selected as

focal genes for subsequent studies.

We also examined the relationship between direction of phenotypic effects with the

wild caught cohorts. For these samples, phenotypic variance for shape is due to the joint

contribution of genetic and environmental effects. To illustrate the difference in shape variance

in wild populations and the DGRP, we calculated correlations between the first three

eigenvectors for shape in the DGRP, the combined wild cohorts as well as the CMO cohort

alone. We observed low correlations between the DGRP eigenvectors and those estimated

from wild populations (Supplemental Table 2). As observed with the DGRP, there is a

substantial correlation between projections of shapes of individuals onto the *ds* shape change

vector and PC1 (defined by phenotypic variation among wild caught files, PC_{wild}) in most of the

sampled cohorts (*ds*-$PC1_{wild}$ PHO: r = 0.78; CMO: r = 0.87; FVW13: r = -0.22; FVW14: r = 0.95,

Figure 2, Supplemental Figure 9). In cohorts where the *ds* shape change vector was not

correlated with PC1, specifically the FVW13 collection, this vector is correlated with PC2 (*ds*-

$PC2_{wild}$ PHO: r = 0.12; CMO r = -0.44; FVW13: r = -0.63; FVW14: r = 0.19; Figure 2;

Supplemental Figure 5). The pattern for the wing shape from wild-caught individuals projected

onto the *emc* shape change vector was generally similar to that observed for *ds* (Figure 2). We

also observe a correlation between *neur* shape change and PC1 in most cohorts (*neur*-$PC1_{wild}$

PHO: r = 0.51; CMO: r = -0.051; FVW12: r = -0.95 FVW13; FVW14: r = -0.084; Figure 2;

Supplemental Figure 7). As with the *ds* shape change vector, in some cohorts such as the CMO

the stronger correlation is between the *neur* shape change vector and PC2 (PHO: r = 0.22; CMO:

r = -0.57; FVW13: r = -0.0.059; FVW14: r = 0.85; Figure 2; Supplemental Figure 7). Interestingly,

in the CMO cohort, the correlations between the projection of shape data onto the *ds* and *neur*

shape change vectors is low (*ds-neur*: r = 0.11, Figure 2).

<u>Multiple loci linked to hippo signaling - including *ds*- respond to artificial selection for *ds* and</u>

<u>*emc* shape changes.</u>

To examine if variants in *ds* are contributing to shape variation, and independently

replicate the findings of the earlier GWAS (Pitchers et al. 2019), we performed artificial

selection experiment for wing shape along the *ds* shape change vector, and examined the

genomic response to selection. By the final generation of selection, we observed a substantial

shape change in both the "up" (females: Procrustes Distance (PD) = 0.039, males: 0.044) and

"down" directions (females: PD = 0.022, males: PD = 0.022), compared to the base population

at the start of the experiment. In comparison, the shape change among unselected control

lineages was much smaller (females: PD = 0.005, males: PD = 0.005) (Figure 3, Supplemental

Figure 10). The direction of phenotypic shape change after seven generations of selection was

in a similar direction to the *ds* shape change vector (defined by RNAi knockdown) for both the

up (females: r = 0.90, males: r = 0.90) and down (females: r = -0.82, males: r = -0.77) selection

lineages. Realized heritabilities, averaged over sex and replicate were moderate (Supplemental

Figure 11, up = 0.38, 95% CI: 0.25 − 0.50; down = 0.28, 95% CI: 0.24 − 0.50). Hippo signaling,

including the effects of *ds*, is often associated with changes in size (Pan, 2007). However, we do

not observe a significant change of wing size in our selection lineages in either sex

(Supplemental Figure 12). It is possible that with more generations of selection we would have

observed a clear change in size, as there is a trend indicating such divergence (Supplemental

Figure 12).

Genome-wide patterns of F_{ST} were examined between up and down *ds* selection

lineages. We observed strong genetic differentiation linked with the *ds* locus (Figure 3,

Supplemental Figure 13), along with several other regions in the genome. One of the SNPs in

the intron of *ds* (2L:702560), identified in Pitchers et al. (2019) through GWAS, showed the

expected pattern of response to selection, with opposing sign in up and down selection

lineages, with the SNP going to high frequency in all three up selection lineages (Table 1). It

should be noted that this SNP is near a complex polymorphism including an insertion of 18bp

that may result in inaccurate genotyping at this locus (Supplemental Figure 14). Gene ontology

analysis for genes in regions of the genome with an F_{ST} greater than 0. 345 (three standard

deviations from mean F_{ST}), show enrichment for hippo signaling loci (Supplemental Table 3).

The top 20 enriched terms are all related to cell signaling and development. Of note is the

inclusion of the terms for 'negative regulation of hippo signaling' (GO:0035331), and 'hippo

signaling' (GO:0035329) in this list (Supplemental Table 3, Supplemental Figure 13). Using a

permutation test we confirmed these results, selecting random sets of genomic intervals equal

in size to the number of observed outlier windows, and measured the ratio of genes annotated

to the expected number of genes in these regions. The observed value for the terms for hippo

signaling (ratio = 4.76) and negative regulation of hippo signaling (ratio = 9.23) were in the

upper 99.5% percentile in comparison to the distributions under permutation (Supplemental

Figure 15).

For the artificial selection experiment based on the *emc* shape change vector we

observed phenotypic differentiation under artificial selection in both up (females: PD = 0.043,

males: PD = 0.040), and down directions (females: PD = 0.021, males: PD = 0.020), with little

change in control lineages (females: PD = 0.009, males: PD = 0.008) (Figure 4). The direction of

phenotypic change is correlated with the *emc* (RNAi knockdown) shape change vector in both

up (females: r = 0.75, males: r = 0.69) and down (females: r = -0.69, males: r = -0.75) directions.

Realized heritabilities, averaged over sex and replicate were calculated for both up and down

lineages (Supplemental Figure 16, up = 0.38, 95% CI: 0.29 – 0.47; down = 0.28, 95% CI: 0.21 –

0.35). Genetic differentiation linked to the *emc locus* was modest following selection, but we

again observed striking genetic differentiation linked to *ds* (Figure 4, Supplemental Figure 13).

Notably, as seen in Supplemental Figure 1, the site frequency spectrum (SFS) suggests modest

allelic variation at the *emc* locus in the synthetic outbred population. Using a three standard

deviation cut-off for F_{ST}, we did observe enrichment for various developmental GO terms, but

not of hippo signaling terms (Supplemental Table 4, Supplemental Figure 13).

<u>Bulk segregant analysis in wild caught cohorts does not recapitulate effects of the GWAS or</u>

<u>artificial selection</u>

Having demonstrated that variants in (or linked to) *ds* respond to artificial selection for

wing shape along the *ds* shape change vector, we next wanted to determine whether we could

recapitulate these findings with wild caught individuals. In addition to determining whether we

can replicate effects in wild cohorts, it provides the opportunity to identify causal SNPs because

of low LD generally observed in wild caught *Drosophila*. Wild caught populations introduce

considerably more environmental variation for shape along with a different site frequency

spectrum for variants contributing to shape variation (and *ds* like shape changes specifically). In

particular, it is known that several of the variants that the original GWAS detected in *ds* have

low minor allele frequency (MAF) (Pitchers et al., 2019) (Table 2). The SNP at 2L:702560 does

appear to be at intermediate frequency but it occurs both directly before and after an indel,

making alignment and variant calling in this region challenging (Supplementary Figure 14). We

have included the frequencies (Table 2), but these results should be interpreted with caution

due to the technical complexities of mapping and variant calling close to indels.

As we sampled multiple cohorts of wild-caught flies in different locations and years in

Michigan (USA), we wanted to confirm that any phenotypic differentiation among these

samples was modest and would not impact genomic analysis for the BSA. We observe modest,

statistically significant wing shape differences among cohorts from a Procrustes ANOVA,

utilizing permutations of the residuals for the relevant "null" model (Supplemental Table 5; R^2=

0.16, F = 351, Z_{RRPP} = 18.3, p = 0.001) (Collyer and Adams, 2018). This appears to be due to

differences in wing shape between the PHO population and other populations based on

pairwise Procrustes Distances (Supplemental Table 5, Supplemental Figure 3). In a joint PCA

including all populations, there is very modest separation between populations using allometry

adjusted shape (Supplemental Figure 3). Most relevant to the BSA approach we used, when we

project all wild caught individuals onto the *ds* and *neur* vectors, there is no clear separation

among sampling locales (Supplemental Figure 7). There is some variation in wing size between

populations (Supplemental Figure 8), but this is unlikely to influence downstream analysis as we

use size adjusted estimates. There is little evidence of genetic differentiation between

populations with the two collections from Fenn Valley Winery separating more on a Principal

Co-ordinate Analysis (PCoA)(Supplemental Figure 17) than other sampling locales. There is also

no relationship between genetic and phenotypic distances between samples (Supplemental

Figure 18). These results suggests that the multiple sampling locales should not influence

downstream genomic analysis as individuals used for generating pools were compared within

each population, and we observe little evidence for substantial differences among populations.

Because there is a single bout of phenotypic selection distinguishing pools for the BSA,

changes in shape and allele frequencies are expected to be modest. We observe shape

differences between the two pools within each population (PD = CMO: 0.033; PHO: 0.036;

FVW13: 0.040; FVW14: 0.041; Supplemental Figure 19). Correlations of the shape difference

vectors of the pools (i.e. difference between the two pools created from the extremes along

the *ds* shape change axis), and the direction of the *ds* shape change vector used for selection, is

high (CMO: 0.94, PHO: 0.79, FVW13: 0.92, FVW14: 0.90).

BSA genome scans show little evidence of genetic differentiation linked to the *ds gene*

(Figure 5). Across the genome, 15 sites were detected as significantly differentiated between

"up" and "down" selected pools based on a CMH test with FDR cut-off of 5% (Figure 5, Table 3).

The genes nearest to these sites are not associated with hippo signaling pathways or implicated

in the development of the *Drosophila* wing (Table 3). Because PHO had somewhat distinct

shape variation from the other populations and had a lower correlation of the difference vector

between selected pools and *ds* shape change vector, we repeated the CMH test with this

population left out. We observe significant differentiation at 174 sites between "up" and

"down" pools (Supplemental Table 6, Supplemental Figure 20). We identified the nearest genes

to these sites and GO analysis indicated enrichment for wing development terms, in particular

related to Wnt signaling, but not hippo signaling terms (Supplemental Table 6). Importantly, we

do not observe differentiation linked to *ds* or any other hippo loci. To ensure that the results we

obtained were not due to uneven coverage between samples, we down-sampled genomic

coverage to 75x for each sample, dropping sites that did not meet this threshold. Significant

differences were detected at 19 sites (Supplemental Figure 21, Supplemental Table 7), but none

of these overlapped with those identified using all the genomic data. Two of the significant sites

are located in the *dumpy* gene, a gene known to have a role in wing morphogenesis during

pupation (Etournay et al., 2015). F_{ST} between selected and random pools within each cohort are

generally low (Supplemental Figure 22).

In addition to the BSA selection based upon the *ds* shape change, we also selected pools

of individuals based on the *neur* shape change vector. We did not use *emc* shape change in this

experiment due to the high similarity between the *ds* and *emc* shape change vectors (r = 0.65),

and the similar response to selection reported above. We selected the *neur* shape change

vector as it is not aligned with *ds,* but does align with directions of natural variation, in wild

populations (Figure 1, 2, Supplemental Figure 9). Additionally, there is little relationship

between the *ds* and *neur* shape change axis (r = 0.12, Supplemental Figures 7 and 8) in the wild

caught cohorts. We observe shape changes between pools of individuals (PD = CMO: 0.027;

PHO: 0.028; FVW14: 0.041; FVW13: 0.038, Supplemental Figure 23). There is little evidence of

genetic differentiation between *neur* selected pools (Supplemental Figure 24). Only 4 sites were

identified as being significantly differentiated between pools and none of these sites are

associated with wing development (Supplemental Table 9). When population differentiation

between pools within populations is measured using F_{ST}, genetic differentiation remains low

across the genome (Supplemental Figure 25)

Discussion

The primary goal of this study was to determine whether we could recapitulate genetic

effects initially observed through a traditional GWAS using an "inverted" approach: artificially

selecting on phenotypes and observing changes in allele frequencies. We observed that shape

changes associated with the *ds*, *emc* and *neur* genes were associated with major axes of genetic

variation among a panel of wild type strains (DGRP) reared in the lab, and axes of phenotypic

variation among wild caught individuals (Figure 1, 2). After observing a strong response to

artificial selection along two shape change vectors (*ds* and *emc*), we examined patterns of

genomic differentiation and observed substantial changes in allele frequency for markers linked

with *ds* itself (figure 3), and markers linked to numerous genes associated with hippo signaling

(Supplemental Figures 13 and 15).

In contrast, our BSA experiments, using pools of wild caught individuals chosen to be

phenotypically divergent on the same shape vectors, did not detect differences in the loci

identified in the artificial selection experiments (Figure 5, Supplemental Figure 24). As we

discuss in detail below, these seemingly contradictory results are in fact not that surprising.

Following artificial selection based on *ds* shape change we observe allele frequency changes

not only at *ds* but also linked to a number of other hippo signalling loci (Figure 3, Supplemental

Table 3). The previous GWAS study identified a number of loci associated with wing shape

variation in the DGRP, however, this approach cannot predict which alleles are causative

(Pitchers et al., 2019). In our synthetic outbred population, we maximized variation among

haplotype blocks containing many of the candidate SNPs in *ds*, increasing our ability to detect

frequency changes at and near the implicated variants. Although LD blocks in the outcrossing

population from this study remain large, *ds* variants exist on multiple distinct haplotypes,

allowing for an examination of allele frequency changes for each. Of particular interest is SNP

2L:702560, previously identified though GWAS (Pitchers et al., 2019) as influencing wing shape

variation. It was driven to near fixation in each of the artificial selection lineages (Table 1).

Although this polymorphism is annotated as a SNP, this region may contain a complex

polymorphism (Supplemental Figure 14), making it difficult to accurately assess genotypic calls.

Because of this, the predicted allele frequency in the founding population and allele

frequencies in this region may be inaccurate. Previous studies demonstrate the importance of

alleles at intermediate frequency in founding populations to those contributing to responses to

selection over short timescales (Kelly and Hughes, 2019). If this polymorphism is at a more

intermediate frequency in the founding population, it would be more likely to be captured by

selection during these experiments. Additionally, haplotype blocks in the initial population are

large, and may contain many potential functional variants. However, based on the results of

both the current and previous studies, these *ds* variants associated with 2L:702560 are good

candidates for functional validation in future work.

When selecting on the *emc* shape change vector, which is similar to that of the *ds* shape

change, we observe only a modest allele frequency change at *emc*, and a more robust response

at *ds* (Figure 4). In hindsight, this is not particularly surprising and there are multiple

contributing factors. Given the increased genetic diversity at *ds* compared to *emc* in the

founding population, alleles in *ds* may have provided a more accessible genetic target, as

selection can only act upon the diversity available in the population. Additionally, if our

estimated direction of effects and selection for *emc* (based on RNAi knockdown) was not well

aligned with the actual direction of *emc* SNP effects, this could result in weaker selection on

variants at the *emc* locus. It is worthwhile pointing out the small magnitude of the *emc* shape

change vector (0.44) relative to *ds* (5.5). However, previous work has indicated that there is a

relationship between this estimated *emc* shape change vector (from RNAi) and the effect of

SNPs in *emc* on shape change (Pitchers et al., 2019).

In addition to a response on allele frequency associated with *ds*, our results suggest a

response on segregating variation at other hippo signaling loci in the *ds* artificial selection

experiment. Earlier work has suggested that the direction of effects within signaling pathways

are inconsistent for alleles of small effect (Dworkin and Gibson, 2006). However, allelic effect

sizes in the 2006 study were heterogeneous and may result in direction and magnitude being

confounded. In contrast, in both the current and the Pitchers et al. (2019) studies, we estimated

the direction of genetic effects by titrating gene knockdown. The strength of this approach is

highlighted in the result that segregating variation at multiple hippo loci was selected on

(Supplemental Figures 13 and 15). Our finding is consistent with models for the architecture of

complex traits that predict that many alleles of small effect will contribute to trait variation with

many genes within developmental pathways (Boyle et al., 2017; Wray et al., 2018). This

pathway response has also been demonstrated in human adaptation to pathogen resistance

(Daub et al., 2013) and high altitude (Gouy et al., 2017). These results are consistent with the

expectation that polymorphisms in the same developmental pathway would show correlated

phenotypic effects and therefore correlated genomic responses to selection. However, this may

not be reflective of all wild caught populations. In this study, we generated a population that

had high diversity at *ds*, while these variants are at much lower frequency in natural

populations (Table 2). The amount of selectable variation a variant provides, depends on both

effect size, *a*, and variant frequencies, *p*, as $V_A = 2p(1-p)a^2$. When allele frequencies are near 0

or 1, even variants with large effects will have only a small contribution to short term selection

response.   Therefore, the outcrossed population we created here is an ideal situation to

validate the existence of the measured effects. It is unlikely to be typical of natural populations

where functional variants may be rare.

Given the clear and robust response observed in the artificial selection experiment, it may

seem surprising that we do not observe allele frequency changes in the BSA using the wild

cohorts. Indeed, previous work has demonstrated that variants in *Egfr*, could be replicated in

wild caught samples (Dworkin et al., 2005; Palsson et al., 2005) and were also found in genome

wide associations (Pitchers et al., 2019). However, there are many explanations for why we may

not have been able to detect these allele frequency changes in our experiment. First, the

addition of environmental variation to the system introduces additional complications. In the

aforementioned example with *Egfr*, the genetic effect of the SNP in wild-caught cohorts was

~10% of the magnitude estimated in lab-reared flies. As discussed previously, the *ds* variants

implicated in the previous GWAS study are at low frequency in the natural cohorts (Table 2).

Given that natural populations of *Drosophila* are generally large and wing shape is likely under

weak selection (Gilchrist and Partridge, 2001), mutation-drift-selection balance may maintain

most variation, resulting in low minor allele frequencies at these sites. Because allelic

contribution to wing shape are expected to be both rare in wild populations and of small

phenotypic effect, we do not expect large allele frequency changes given only one "generation"

of selection. Using the approach of ACER (Spitzer et al., 2020) to account for sampling effects,

we observe few differentiated sites, and none in the *ds* gene, indicating that BSA may not be

well-suited to identify modest allele frequency changes, thus, not particularly effective for

polygenic traits. Although our approach was tailored to look for variants that had consistent

direction of frequency changes across the four collection cohorts, it is possible that different

loci were contributing variation within each cohort. We attempted to address this question by

examining allele frequency changes between selected pools within each cohort (Supplemental

Figures 22 and 24) but could not identify specific loci contributing to differences within any one

population. Previous successful BSA studies identified smaller numbers of contributing loci with

few polymorphisms contributing to the trait of interest. For example, in *Drosophila,* a number

of melanin synthesis genes contributing to variance in pigmentation between populations were

identified using a BSA (Bastide et al., 2013). Pigmentation may represent a relatively 'simpler'

genetic architecture (fewer variants of individually larger genetic effect, smaller impact of

environmental variation, smaller mutational target size) and if so, this may have enabled the

success of the BSA approach with such systems. In the case of wing shape, we know that many

alleles of small effect contribute to variation in the trait (Pitchers et al., 2019).

Our approach for the BSA was to perform the same phenotypic selection within each of

four distinct "populations". It is important to recognize that there was heterogeneity among

our populations, not only in allele frequencies, but in environmental variance and potentially

GxE, even though all were caught in locales in lower Michigan. We detected small degrees of

phenotypic and genetic differences between cohorts, however these effects are neither

correlated with one another, nor related to the *ds* and *neur* shape scores used for selecting

individuals (Supplemental Figures 3, 7, 18). The population from the Phillips Orchard (PHO) was

phenotypically distinct from the other populations. When we performed the BSA without this

population, we observed a larger set of variants associated with shape (Supplemental Figure

20), albeit still not showing any effects at *ds* or *neur* genes themselves. One possibility is that

the increased number of sites when the PHO sample is removed from analysis represents an

unknown statistical artefact we have not identified. However, a more likely explanation is that

there are some large unknown environmental influences (*E*), or that the genetic effects show a

degree of *GxE* (with a specific environment in PHO) that contributed to shape variation along

the *ds* direction in this population. Such obfuscating effects have been observed before with

the previously discussed *Egfr* example, where the SNP effect identified and validated in multiple

contexts (Dworkin et al., 2005; Palsson et al., 2005; Palsson and Gibson, 2004) could not be

detected in one natural population, despite being at intermediate frequencies in each sample

(Palsson et al., 2005). Importantly, we did detect differentiation at sites associated with

developmental processes in the wild cohorts, suggesting that the failure to detect variation

linked to *ds* or other hippo signaling loci (Table 3, Supplemental Table 6,7) is not due simply to a

lack of power.

The response to selection at *ds* and other hippo signaling loci in the artificial selection

experiment based on *ds* shape change indicates that this is an important axis of variation for

wing shape. Coupled with the alignment of phenotypic effects of perturbations in genes in this

pathway with directions of **G** and **P**, this finding may seem to suggest a developmental bias in

available variation. However, we caution against such interpretations based solely on the

findings in this study. The structure of the **G** matrix strongly influenced our findings as we

artificially created a population to maximize genetic diversity at *ds*. When another effect is

aligned with *ds* shape change, as in the case of *emc* shape change, we observed the same

response at the hippo signaling loci and not at *emc.* Only the genetic diversity in the starting

population was available to be selected on so this influenced selection towards the "spiked in"

ds variants, even if the inferred phenotypic effects of *emc* variants are very similar.

Alternatively, the inferred *emc* direction of effects (via RNAi knockdown) may be sufficiently

"distant" from true effects of *emc* variants. If this was the case, we were ineffectively selecting

for *emc* shape changes. In other cases where single genes are implicated in divergence between

multiple populations, such as *mc1r* in mice (Steiner et al., 2007) or *pitx1* in stickleback (Chan et

al., 2010), other factors such as low pleiotropy, developmental and mutational constraints and

history of selection in the population are used to explain why these genes are so often

implicated in evolutionary change (Gompel and Prud'homme, 2009; Martin and Orgogozo,

2013; Stern and Orgogozo, 2008). In our case, it is not *ds* itself that is special but rather the

orientation of the **G** matrix to align $\mathbf{g_{max}}$ with the direction of effect for *ds* that shapes our

results. Selection acts on variants aligned with the vector of selection (Reddiex and Chenoweth,

2021). By varying the orientation of $\mathbf{g_{max}}$ in the parental population, we would be able to

address questions about the repeatability of hippo overrepresentation and if this can be

explained by more than just the orientation of **G.**

Despite the need for skepticism about the potential for developmental bias influencing

directions of variation, the correlated response of sites linked to multiple other hippo signaling

genes is intriguing. Coupling of more traditional mapping approaches like GWAS with short

term artificial selection provides an additional route to validation and replication of genetic

effects. It also suggests that using multivariate data to address the distribution of genetic

effects will pay long-term dividends to our understanding of both inheritance and the evolution

of multivariate traits.

Acknowledgements:

We would like to thank the Hosken lab for *D. simulans* wing shape data. Yun Bo Xi aided

with the sequencing of *ds* haplotype. We thank Dr. Brian Golding for computational resources.

We thank Dr. Catherine Peichel and two anonymous reviewers for helpful feedback on the

manuscripts.

References:
Adams, D.C., Otárola-Castillo, E., 2013. geomorph: an R package for the collection and analysis
of geometric morphometric shape data. Methods Ecol. Evol. 4, 393–399.
https://doi.org/10.1111/2041-210X.12035

Alexa, A., Rahnenfuhrer, J., Lengauer, T., 2006. Improved scoring of functional groups from
gene expression data by decorrelating GO graph structure. Bioinformatics 22, 1600–
1607. https://doi.org/10.1093/bioinformatics/btl140
Bastide, H., Betancourt, A., Nolte, V., Tobler, R., Stöbe, P., Futschik, A., Schlötterer, C., 2013. A
Genome-Wide, Fine-Scale Map of Natural Pigmentation Variation in *Drosophila*
melanogaster. PLoS Genet. 9, e1003534. https://doi.org/10.1371/journal.pgen.1003534
Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. J. R. Stat. Soc. Ser. B Methodol. 57, 289–300.
https://doi.org/10.1111/j.2517-6161.1995.tb02031.x
Blows, M.W., McGuigan, K., 2015. The distribution of genetic variance across phenotypic space
and the response to selection. Mol. Ecol. 24, 2056–2072.
https://doi.org/10.1111/mec.13023
Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170
Boyle, E.A., Li, Y.I., Pritchard, J.K., 2017. An Expanded View of Complex Traits: From Polygenic to
Omnigenic. Cell 169, 1177–1186. https://doi.org/10.1016/j.cell.2017.05.038
Chan, Y.F., Marks, M.E., Jones, F.C., Villarreal, G., Shapiro, M.D., Brady, S.D., Southwick, A.M.,
Absher, D.M., Grimwood, J., Schmutz, J., Myers, R.M., Petrov, D., Jónsson, B., Schluter,
D., Bell, M.A., Kingsley, D.M., 2010. Adaptive Evolution of Pelvic Reduction in
Sticklebacks by Recurrent Deletion of a Pitx1 Enhancer. Science 327, 302–305.
https://doi.org/10.1126/science.1182213
Collyer, M.L., Adams, D.C., 2018. RRPP: An R package for fitting linear models to
high-dimensional data using residual randomization. Methods Ecol. Evol. 9, 1772–1779.
https://doi.org/10.1111/2041-210X.13029
CONVERGE consortium, 2015. Sparse whole-genome sequencing identifies two loci for major
depressive disorder. Nature 523, 588–591. https://doi.org/10.1038/nature14659
Daub, J.T., Hofer, T., Cutivet, E., Dupanloup, I., Quintana-Murci, L., Robinson-Rechavi, M.,
Excoffier, L., 2013. Evidence for Polygenic Adaptation to Pathogens in the Human
Genome. Mol. Biol. Evol. 30, 1544–1558. https://doi.org/10.1093/molbev/mst080
Dworkin, I., Gibson, G., 2006. Epidermal Growth Factor Receptor and Transforming Growth
Factor-β Signaling Contributes to Variation for Wing Shape in *Drosophila melanogaster*.
Genetics 173, 1417–1431. https://doi.org/10.1534/genetics.105.053868
Dworkin, I., Palsson, A., Gibson, G., 2005. Replication of an *Egfr*-Wing Shape Association in a
Wild-Caught Cohort of *Drosophila melanogaster*. Genetics 169, 2115–2125.
https://doi.org/10.1534/genetics.104.035766
Etournay, R., Popović, M., Merkel, M., Nandi, A., Blasse, C., Aigouy, B., Brandl, H., Myers, G.,
Salbreux, G., Jülicher, F., Eaton, S., 2015. Interplay of cell dynamics and epithelial tension
during morphogenesis of the *Drosophila* pupal wing. eLife 4, e07090.
https://doi.org/10.7554/eLife.07090
Fatumo, S., Carstensen, T., Nashiru, O., Gurdasani, D., Sandhu, M., Kaleebu, P., 2019.
Complimentary Methods for Multivariate Genome-Wide Association Study Identify New
Susceptibility Genes for Blood Cell Traits. Front. Genet. 10, 334.
https://doi.org/10.3389/fgene.2019.00334

Fritsche, L.G., Igl, W., Bailey, J.N.C., Grassmann, F., Sengupta, S., Bragg-Gresham, J.L., Burdon,
K.P., Hebbring, S.J., Wen, C., Gorski, M., Kim, I.K., Cho, D., Zack, D., Souied, E., Scholl,
H.P.N., Bala, E., Lee, K.E., Hunter, D.J., Sardell, R.J., Mitchell, P., Merriam, J.E., Cipriani,
V., Hoffman, J.D., Schick, T., Lechanteur, Y.T.E., Guymer, R.H., Johnson, M.P., Jiang, Y.,
Stanton, C.M., Buitendijk, G.H.S., Zhan, X., Kwong, A.M., Boleda, A., Brooks, M., Gieser,
L., Ratnapriya, R., Branham, K.E., Foerster, J.R., Heckenlively, J.R., Othman, M.I., Vote,
B.J., Liang, H.H., Souzeau, E., McAllister, I.L., Isaacs, T., Hall, J., Lake, S., Mackey, D.A.,
Constable, I.J., Craig, J.E., Kitchner, T.E., Yang, Z., Su, Z., Luo, H., Chen, D., Ouyang, H.,
Flagg, K., Lin, D., Mao, G., Ferreyra, H., Stark, K., von Strachwitz, C.N., Wolf, A., Brandl,
C., Rudolph, G., Olden, M., Morrison, M.A., Morgan, D.J., Schu, M., Ahn, J., Silvestri, G.,
Tsironi, E.E., Park, K.H., Farrer, L.A., Orlin, A., Brucker, A., Li, M., Curcio, C.A., Mohand-
Saïd, S., Sahel, J.-A., Audo, I., Benchaboune, M., Cree, A.J., Rennie, C.A., Goverdhan, S.V.,
Grunin, M., Hagbi-Levi, S., Campochiaro, P., Katsanis, N., Holz, F.G., Blond, F., Blanché,
H., Deleuze, J.-F., Igo, R.P., Truitt, B., Peachey, N.S., Meuer, S.M., Myers, C.E., Moore,
E.L., Klein, R., Hauser, M.A., Postel, E.A., Courtenay, M.D., Schwartz, S.G., Kovach, J.L.,
Scott, W.K., Liew, G., Tan, A.G., Gopinath, B., Merriam, J.C., Smith, R.T., Khan, J.C.,
Shahid, H., Moore, A.T., McGrath, J.A., Laux, R., Brantley, M.A., Agarwal, A., Ersoy, L.,
Caramoy, A., Langmann, T., Saksens, N.T.M., de Jong, E.K., Hoyng, C.B., Cain, M.S.,
Richardson, A.J., Martin, T.M., Blangero, J., Weeks, D.E., Dhillon, B., van Duijn, C.M.,
Doheny, K.F., Romm, J., Klaver, C.C.W., Hayward, C., Gorin, M.B., Klein, M.L., Baird, P.N.,
den Hollander, A.I., Fauser, S., Yates, J.R.W., Allikmets, R., Wang, J.J., Schaumberg, D.A.,
Klein, B.E.K., Hagstrom, S.A., Chowers, I., Lotery, A.J., Léveillard, T., Zhang, K., Brilliant,
M.H., Hewitt, A.W., Swaroop, A., Chew, E.Y., Pericak-Vance, M.A., DeAngelis, M.,
Stambolian, D., Haines, J.L., Iyengar, S.K., Weber, B.H.F., Abecasis, G.R., Heid, I.M., 2016.
A large genome-wide association study of age-related macular degeneration highlights
contributions of rare and common variants. Nat. Genet. 48, 134–143.
https://doi.org/10.1038/ng.3448
Gilchrist, A.S., Partridge, L., 2001. The contrasting genetic architecture of wing size and shape in
Drosophila melanogaster. Heredity 86, 144–152. https://doi.org/10.1046/j.1365-
2540.2001.00779.x
Gompel, N., Prud'homme, B., 2009. The causes of repeated genetic evolution. Dev. Biol. 332,
36–47. https://doi.org/10.1016/j.ydbio.2009.04.040
Gouy, A., Daub, J.T., Excoffier, L., 2017. Detecting gene subnetworks under selection in
biological pathways. Nucleic Acids Res. 45, e149–e149.
https://doi.org/10.1093/nar/gkx626
Green, R.M., Fish, J.L., Young, N.M., Smith, F.J., Roberts, B., Dolan, K., Choi, I., Leach, C.L.,
Gordon, P., Cheverud, J.M., Roseman, C.C., Williams, T.J., Marcucio, R.S., Hallgrímsson,
B., 2017. Developmental nonlinearity drives phenotypic robustness. Nat. Commun. 8,
1970. https://doi.org/10.1038/s41467-017-02037-7
Greene, C.S., Penrod, N.M., Williams, S.M., Moore, J.H., 2009. Failure to Replicate a Genetic
Association May Provide Important Clues About Genetic Architecture. PLoS ONE 4,
e5639. https://doi.org/10.1371/journal.pone.0005639
Houle, D., Mezey, J., Galpern, P., Carter, A., 2003. Automated measurement of *Drosophila*
wings. BMC Evol. Biol. 3, 25. https://doi.org/10.1186/1471-2148-3-25

Huang, W., Massouras, A., Inoue, Y., Peiffer, J., Ramia, M., Tarone, A.M., Turlapati, L., Zichner,
T., Zhu, D., Lyman, R.F., Magwire, M.M., Blankenburg, K., Carbone, M.A., Chang, K., Ellis,
L.L., Fernandez, S., Han, Y., Highnam, G., Hjelmen, C.E., Jack, J.R., Javaid, M., Jayaseelan,
J., Kalra, D., Lee, S., Lewis, L., Munidasa, M., Ongeri, F., Patel, S., Perales, L., Perez, A., Pu,
L., Rollmann, S.M., Ruth, R., Saada, N., Warner, C., Williams, A., Wu, Y.-Q., Yamamoto,
A., Zhang, Y., Zhu, Y., Anholt, R.R.H., Korbel, J.O., Mittelman, D., Muzny, D.M., Gibbs,
R.A., Barbadilla, A., Johnston, J.S., Stone, E.A., Richards, S., Deplancke, B., Mackay, T.F.C.,
2014. Natural variation in genome architecture among 205 *Drosophila melanogaster*
Genetic Reference Panel lines. Genome Res. 24, 1193–1208.
https://doi.org/10.1101/gr.171546.113
Ioannidis, J.P.A., Tarone, R., McLaughlin, J.K., 2011. The False-positive to False-negative Ratio in
Epidemiologic Studies: Epidemiology 22, 450–456.
https://doi.org/10.1097/EDE.0b013e31821b506e
Kelly, J.K., Hughes, K.A., 2019. Pervasive Linked Selection and Intermediate-Frequency Alleles
Are Implicated in an Evolve-and-Resequencing Experiment of *Drosophila simulans.*
Genetics 211, 943–961. https://doi.org/10.1534/genetics.118.301824
King, Elizabeth G., Macdonald, S.J., Long, A.D., 2012. Properties and Power of the *Drosophila*
Synthetic Population Resource for the Routine Dissection of Complex Traits. Genetics
191, 935–949. https://doi.org/10.1534/genetics.112.138537
King, E. G., Merkes, C.M., McNeil, C.L., Hoofer, S.R., Sen, S., Broman, K.W., Long, A.D.,
Macdonald, S.J., 2012. Genetic dissection of a model complex trait using the *Drosophila*
Synthetic Population Resource. Genome Res. 22, 1558–1566.
https://doi.org/10.1101/gr.134031.111
Klingenberg, C.P., 2022. Methods for studying allometry in geometric morphometrics: a
comparison of performance. Evol. Ecol. 36, 439–470. https://doi.org/10.1007/s10682-
022-10170-z
Kofler, Orozco-terWengel, P., De Maio, N., Pandey, R.V., Nolte, V., Futschik, A., Kosiol, C.,
Schlötterer, C., 2011a. PoPoolation: A Toolbox for Population Genetic Analysis of Next
Generation Sequencing Data from Pooled Individuals. PLoS ONE 6, e15925.
https://doi.org/10.1371/journal.pone.0015925
Kofler, Pandey, R.V., Schlotterer, C., 2011b. PoPoolation2: identifying differentiation between
populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics 27,
3435–3436. https://doi.org/10.1093/bioinformatics/btr589
Li, H., Durbin, R., 2010. Fast and accurate long-read alignment with Burrows–Wheeler
transform. Bioinformatics 26, 589–595. https://doi.org/10.1093/bioinformatics/btp698
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352
Madeira, F., Pearce, M., Tivey, A.R.N., Basutkar, P., Lee, J., Edbali, O., Madhusoodanan, N.,
Kolesnikov, A., Lopez, R., 2022. Search and sequence analysis tools services from EMBL-
EBI in 2022. Nucleic Acids Res. 50, W276–W279. https://doi.org/10.1093/nar/gkac240

Marigorta, U.M., Rodríguez, J.A., Gibson, G., Navarro, A., 2018. Replicability and Prediction:
Lessons and Challenges from GWAS. Trends Genet. 34, 504–517.
https://doi.org/10.1016/j.tig.2018.03.005
Marquez, E.J., Houle, D., 2015. Dimensionality and the statistical power of multivariate
genome-wide association studies (preprint). Genomics. https://doi.org/10.1101/016592
Marriage, T.N., King, E.G., Long, A.D., Macdonald, S.J., 2014. Fine-Mapping Nicotine Resistance
Loci in *Drosophila* Using a Multiparent Advanced Generation Inter-Cross Population.
Genetics 198, 45–57. https://doi.org/10.1534/genetics.114.162107
Martin, A., Orgogozo, V., 2013. The Loci of Repeated Evolution: A Catalog of Genetic Hotspots
of Phenotypic Variation. Evolution 67, 1235–1250. https://doi.org/10.1111/evo.12081
McGuigan, K., 2006. Studying phenotypic evolution using multivariate quantitative genetics:
EVOLUTIONARY QUANTITATIVE GENETICS. Mol. Ecol. 15, 883–896.
https://doi.org/10.1111/j.1365-294X.2006.02809.x
Melo, D., Marroig, G., Wolf, J.B., 2019. Genomic Perspective on Multivariate Variation,
Pleiotropy, and Evolution. J. Hered. 110, 479–493.
https://doi.org/10.1093/jhered/esz011
Palsson, A., Dodgson, J., Dworkin, I., Gibson, G., 2005. Tests for the replication of an association
between *Egfr* and natural variation in *Drosophila melanogaster* wing morphology. BMC
Genet. 6, 44. https://doi.org/10.1186/1471-2156-6-44
Palsson, A., Gibson, G., 2004. Association Between Nucleotide Variation in *Egfr* and Wing Shape
in *Drosophila melanogaster*. Genetics 167, 1187–1198.
https://doi.org/10.1534/genetics.103.021766
Pan, D., 2007. Hippo signaling in organ size control. Genes Dev. 21, 886–897.
https://doi.org/10.1101/gad.1536007
Pan, Y., Alégot, H., Rauskolb, C., Irvine, K.D., 2018. The dynamics of Hippo signaling during
Drosophila wing development. Development 145, dev165712.
https://doi.org/10.1242/dev.165712
Pitchers, W., Nye, J., Márquez, E.J., Kowalski, A., Dworkin, I., Houle, D., 2019. A Multivariate
Genome-Wide Association Study of Wing Shape in *Drosophila melanogaster*. Genetics
211, 1429–1447. https://doi.org/10.1534/genetics.118.301342
Porter, H.F., O'Reilly, P.F., 2017. Multivariate simulation framework reveals performance of
multi-trait GWAS methods. Sci. Rep. 7, 38837. https://doi.org/10.1038/srep38837
Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26, 841–842. https://doi.org/10.1093/bioinformatics/btq033
Ray, R.P., Nakata, T., Henningsson, P., Bomphrey, R.J., 2016. Enhanced flight performance by
genetic manipulation of wing shape in *Drosophila*. Nat. Commun. 7, 10851.
https://doi.org/10.1038/ncomms10851
Reddiex, A.J., Chenoweth, S.F., 2021. Integrating genomics and multivariate evolutionary
quantitative genetics: a case study of constraints on sexual selection in *Drosophila*
serrata. Proc. R. Soc. B Biol. Sci. 288, 20211785. https://doi.org/10.1098/rspb.2021.1785
Rohlf, F.J., Slice, D., 1990. Extensions of the Procrustes Method for the Optimal Superimposition
of Landmarks. Syst. Zool. 39, 40. https://doi.org/10.2307/2992207
Schluter, D., 1996. Adaptive Radiation Along Genetic Lines of Least Resistance. Evolution 50,
1766. https://doi.org/10.2307/2410734

Shriner, D., 2012. Moving toward System Genetics through Multiple Trait Analysis in Genome-
Wide Association Studies. Front. Genet. 3. https://doi.org/10.3389/fgene.2012.00001
Spitzer, K., Pelizzola, M., Futschik, A., 2020. Modifying the Chi-square and the CMH test for
population genetic inference: Adapting to overdispersion. Ann Appl Stat 14, 202–220.
https://doi.org/10.1214/19-AOAS1301
Steiner, C.C., Weber, J.N., Hoekstra, H.E., 2007. Adaptive Variation in Beach Mice Produced by
Two Interacting Pigmentation Genes. PLoS Biol. 5, e219.
https://doi.org/10.1371/journal.pbio.0050219
Stephens, M., 2013. A Unified Framework for Association Analysis with Multiple Related
Phenotypes. PLoS ONE 8, e65245. https://doi.org/10.1371/journal.pone.0065245
Stern, D.L., Orgogozo, V., 2008. THE LOCI OF EVOLUTION: HOW PREDICTABLE IS GENETIC
EVOLUTION? Evolution 62, 2155–2177. https://doi.org/10.1111/j.1558-
5646.2008.00450.x
Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., Meyre, D., 2019. Benefits and limitations of
genome-wide association studies. Nat. Rev. Genet. 1. https://doi.org/10.1038/s41576-
019-0127-1
Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A., Yang, J., 2017. 10
Years of GWAS Discovery: Biology, Function, and Translation. Am. J. Hum. Genet. 101,
5–22. https://doi.org/10.1016/j.ajhg.2017.06.005
Wray, N.R., Wijmenga, C., Sullivan, P.F., Yang, J., Visscher, P.M., 2018. Common Disease Is More
Complex Than Implied by the Core Gene Omnigenic Model. Cell 173, 1573–1580.
https://doi.org/10.1016/j.cell.2018.05.051
Wright, S., 1934. Physiological and Evolutionary Theories of Dominance. Am. Nat. 68, 24–53.
https://doi.org/10.1086/280521
915

Figure 1. Projections of data onto RNAi shape change vectors are correlated with major axes
of shape variation among DGRP strains. (A) Shape change vectors from RNAi titration
experiments for *ds, emc* and *neur* were used, and DGRP line means were projected onto these
vectors to calculate shape scores. Eigenvectors for the PCA were estimated based on the same
DGRP line means. Vector correlations between shape change vectors from RNAi knockdown: *ds*
− emc: 0.65, *ds − neur*: 0.03, *emc − neur*: 0.30. (B) Effect of *ds* shape change estimated from
RNAi knockdown, effects not magnified. (C) Landmarks (red) and semi-landmarks (blue) used in
geomorphic morphometric analysis on a *Drosophila* wing. PCs 1–3 account for 22%, 20% and
9% of the overall, among DGRP shape variance.
934
Figure 2. Projections of data onto RNAi shape change vectors are correlated with major axes
of shape variation in wild-caught *Drosophila*. Correlations between projection of shape data
from CMO population onto *ds, emc* and *neur* RNAi shape change vectors, and the first three
eigenvectors from the PCA, calculated from shape data from all samples in the CMO
population. PCs 1–3 account for 24%, 18% and 9% of overall shape variance in the CMO
population.
941
Figure 3. Artificial selection along *ds* shape change vector influences allele frequencies of
variants at *ds*.(A) Phenotypic response to selection based on *ds* shape change vector. Only data
from females is plotted for ease of visualization. Each replicate of up (squares), control (dots)
and down (triangles) selection lineages are plotted (greys). Estimated response to selection
shown along red lines. Wing plots represent the effect of selection on shape change between
generation one and seven (red, effects not magnified). (B) Genomic differentiation (F_{ST})
between up and down selection treatments measured in 5000bp windows. Red line represents
the location of the *ds* locus. Grey line represents 3 standard deviations from genome wide
mean F_{ST}.
951
Figure 4. Artificial selection along *emc* shape change vector has modest influence on allele
frequencies at *emc*, but a greater impact at the *ds locus*. (A) Phenotypic response to selection
based on the *emc* shape change vector. Only data from females is plotted for ease of
visualization. Each replicate of up (squares), control (dots) and down (triangles) selection
lineages are plotted in greys. Estimated response to selection shown along red lines. Shape
change between generation 1 and 7 is indicated on the right. Shape effects have been
magnified 5x. (B) Genomic differentiation between up and down selection lineages (F_{ST})
measured in 5000bp sliding windows. Red and purple vertical lines represent genomic locations
of *ds* and *emc* respectively. Grey line represents 3 standard deviations from genome wide
mean F_{ST}.
962
Figure 5. Genetic differentiation between pools selected based on *ds* shape change among
the wild-caught cohorts. (A) Genome-wide scan for differentiated loci between pools selected
based on *ds* shape change vector using the CMH test implemented in ACER. Points in red
indicate sites with significant differentiation. Position of *ds* gene in blue (B) Genomic
differentiation at *ds* between pools selected based on *ds* shape change vector. No sites are
significantly differentiated in *ds*. The large gap in sites is due to a masked region in the genome

due to repetitive sequence and poor (syntenic) mapping scores. (C) Shape difference between
selected pools of individuals from one representative (CMO) population, with the mean shape
of pools represented in black and red.

993   Table 1. Variants from Pitchers et al. (2019) in *ds* artificial selection experiment. Estimated
994   effect sizes for SNPs are estimated from a GWAS in the DGRP using LASSO regularized
995   coefficients. Average frequency is given with replicate lineage frequencies in brackets.
996   Estimated effect is the l^2-*norm* of shape differences associated with the variant. MAF = minor
997   allele frequency.

Variant	Estimated Effect	DGRP MAF	Estimated MAF in synthetic outcross	Average allele frequency "up" selection	Average allele frequency "down" selection	Average allele frequency "control" selection
2L:655894	0.072	0.44	0.067	0 (0, 0, 0)	0 (0, 0, 0)	0.003 (0, 0, 0.0105)
2L:702560*	**0.159**	**0.056**	**0.06**	**0.995 (1, 0.98, 1)**	**0.446 (0.32, 0.35, 0.67)**	**0.705 (0.69, 0.56, 0.87)**
2L:702798	0.101	0.089	0.1	0.007 (0, 0.0217, 0)	0 (0, 0, 0)	0.005 (0, 0, 0.139)
2L:718623	0.225	0.033	0	0 (0, 0, 0)	0 (0, 0, 0)	0 (0, 0, 0)
2L:718627	0.11	0.033	0	0 (0, 0, 0)	0 (0, 0, 0)	0 (0, 0, 0)

998   * This is a complex polymorphism with linked SNPs and INDELs, in Pitchers *et al* (2019) a SNP in
999   this region was found to be linked. However, the variant calling pipeline used in this work
1000   recognized an INDEL in this region which was used for counting.

1011 Table 2. *ds* Variants from Pitchers et al. (2019) in wild-caught cohorts used in the present study.
1012 Estimated effect sizes for SNPs are estimated from the DGRP GWAS with LASSO regularized
1013 coefficients. MAF in wild cohorts was estimated from sequenced pools of 75 random
1014 individuals.

Variant	Estimated Effect (Pitchers 2019)	DGRP MAF	Estimated MAF CMO	Estimated MAF FVW13	Estimated MAF FVW14	Estimated MAF PHO
2L:655894	0.072	0.445	0	0	0	0
2L:702560*	0.159	0.056	0.375	0.473	0.485	0.336
2L:702798	0.101	0.089	0.077	0.101	0.044	0.034
2L:718623	0.225	0.033	0.051	0.021	0.044	0.100
2L:718627	0.11	0.033	0.055	0.020	0.046	0.099

1015 * This is a complex polymorphism with linked SNPs and INDELs, in Pitchers *et al* (2019) a SNP in
1016 this region was found to be linked. However, the variant calling pipeline used in this work
1017 recognized an INDEL in this region which was used for counting.

1044   Table 3. Significantly differentiated variants for *ds* shape change from the wild-caught cohorts
1045   (BSA).

Location	CMH p-value (FDR corrected)	Gene	FlyBase ID	Distance from ORF (bp)
2R:17491270	0.026	*NT5E-2*	FBgn0050104	0
2R17498059	0.034	*CG30103*	FBgn0050103	2061
2R:17515133	0.022	*CG4853*	FBgn0034230	0
2R: 20537878	0.013	*CG13423*	FBgn0034513	0
2R:23601278	0.005	*CG10332*	FBgn0260455	0
2R:23601278	0.005	*IM18*	FBgn0067903	0
2R:23613785	0.013	*Eglp4*	FBgn0034885	0
2R:23613785	0.013	*Eglp2*	FBgn0034883	0
2R: 23646252	0.016	*retn*	FBgn0004795	0
3L:12831924	0.005	*CG10960*	FBgn0036316	0
3L: 20999119	0.022	*skd*	FBgn0003415	0
3R: 21523866	0.013	*CG7956*	FBgn0038890	0
3R: 2559549	0.011	*Pzl*	FBgn0267430	0
X: 14891220	0.013	*Flo2*	FBgn0264078	0
X: 14891220	0.013	*CG9514*	FBgn0030592	0
X:16039731	0.017	*Muc14a*	FBgn0052580	0
X: 793052	0.011	*CG16989*	FBgn0025621	95
X: 9448676	0.034	*mgl*	FBgn0261260	0

1046
1047

A

ds					
0.69	emc				
0.56	0.83	neur			
-0.56	-0.45	-0.69	PC1		
0.11	0.40	0.15	0.00	PC2	
-0.46	-0.74	-0.64	-0.00	0.00	PC3

B

C