Differential modifications of the C-terminal tails of α-tubulin isoforms and their *in vivo*
importance for kinesin function in *Drosophila*

Mengjing Bao, Ruth Dörig, Dirk Beuchle, Paula Maria Vazquez-Pianzola and Beat Suter
Institute of Cell Biology, University of Bern, Switzerland

Summary

Drosophila TTLL5 is required for:

1. α-tubulin glutamylation in *vivo,* specifically on the C-terminal domain of TBA1 and 3; the
female germline isoform TBA4 is not glutamylated.

2. Kinesin-based localization of Staufen/*osk* mRNA and ooplasmic streaming during
oogenesis.

3. Kinesin-based transport of mitochondria in neurons.

Abstract

Microtubules (MTs) are built from α-/β-tubulin dimers and used as tracks by kinesin and
dynein motors to transport a variety of cargos, such as mRNAs, proteins, and organelles,
within the cell. Tubulins are subjected to several post-translational modifications (PTMs).
Glutamylation is one of them and it is responsible for adding one or more glutamic acids as
branched peptide chains to the C-terminal tails of both α- and β-tubulin. However, very little
is known about the specific modifications found on the different tubulin isoforms *in vivo* and
the role of these PTMs in cargo transport along MTs *in vivo*. In this study, we found that
Drosophila TTLL5 is required specifically for the glutamylation of the α-tubulin isoforms
TBA1 and TBA3 in ovaries, whereas the ovarian isoform TBA4 is not glutamylated.
Furthermore, *TTLL5* is essential for the efficient localization of Staufen/*osk* mRNA and to
give directionality to the fast ooplasmic streaming, two processes known to depend on
kinesin-mediated processes during oogenesis. In the nervous system, the kinesin-dependent
neuronal transport of mitochondria also depends on *TTLL5*. Additionally, α-tubulin
glutamylation plays a role in controlling the pausing of the transport of individual

mitochondria. Our results demonstrate the *in vivo* role of *TTLL5* in differential glutamylation

of α-tubulin isoforms and point to the *in vivo* importance of α-tubulin glutamylation for

kinesin-dependent processes.

Introduction

Microtubules (MTs) are key cytoskeletal filaments that are comprised of heterodimers of α-

and β-tubulins. Despite the high level of conservation in eukaryotes, MTs are still quite

diverse because cells express different isoforms, and these can undergo several different

post-translational modifications (PTMs). Based mainly on *in vitro* assays, these isoform

variations are thought to optimize the MT interaction with different microtubule-associated

proteins (MAPs) and motors.

Drosophila melanogaster possesses four α-tubulin genes. Three of them are expressed

during oogenesis and early embryogenesis, suggesting that they contribute to the formation

of microtubules during these stages. The three genes are *αTub84B*, *αTub84D,* and *αTub67C*

(Kalfayan and Wensink, 1982). *αTub84B* and *αTub84D* encode the TBA1 and TBA3

isoforms of α-tubulin, respectively, which differ in only two amino acids. Their primary

sequence in the C-terminal domain is identical and both are expressed in most tissues and

throughout development. In contrast, *αTub67C* encodes TBA4 and is expressed exclusively

during oogenesis where it is maternally loaded into the egg and embryo. The primary

structure of TBA4 is distinctively different from TBA1/3 (Theurkauf et al., 1986). These

differences are also apparent in the C-terminal domain and include the last residue, which is

phenylalanine (Phe, F) in TBA4 and tyrosine (Tyr, Y) in TBA1/3. *αTub85E,* which encodes

TBA2, is mainly expressed in testes (Kalfayan and Wensink, 1982).

MTs are subjected to several PTMs, including the C-terminal tail glutamylation,

glycylation, and detyrosination / tyrosination (Arce et al., 1975, Eddé et al., 1990, Redeker et

al., 1994, Hallak et al., 1977). Furthermore, acetylation and phosphorylation in the more

central parts of the tubulins have also been identified (Arce et al., 1975; Eddé et al., 1990;

Hallak et al., 1977; L'Hernault and Rosenbaum, 1985; Magiera and Janke, 2014; Redeker et

al., 1994). Glutamylation is a PTM that adds one or more glutamic acids (Glu, E) to the side

chains of glutamic acid residues in the C-terminal tails of both α and β tubulin, causing the

formation of a branched peptide chain (Eddé et al., 1990; Redeker et al., 1992). Glycylation

is another PTM that adds glycines to the γ-carboxyl group of glutamic acid residues in the C-

terminal region (Redeker et al., 1994). At least in vertebrates, tubulin is also subjected to a

special cycle of de-tyrosination-tyrosination (Preston et al., 1979). In this case, the C-

terminal Tyr or Phe of α-tubulin is cyclically removed, resulting in a C-terminal Glu residue,

and a new Tyr can be re-added to the new C-term. The enzymes that catalyze glutamylation,

glycylation, and tyrosination belong to the Tubulin-tyrosine-ligase-like (TTLL) family (Janke

et al., 2005). In *Drosophila melanogaster*, this family encompasses 11 genes (flybase.org).

CG31108 is considered to be the homolog of the mammalian *TTLL5* gene and was

therefore named *DmTTLL5 or TTLL5* (Devambez et al., 2017). It is required for α-tubulin

glutamylation in the *Drosophila* nervous system (Devambez et al., 2017). The murine TTLL5

is composed of an N-terminal core tubulin tyrosination ligase-like (TTLL) domain, a cofactor

interaction domain (CID), and a C-terminal receptor interaction domain (RID) (Lee et al.,

2013). The N-terminal core domain was shown to provide the catalytic activity of TTLL5 and

is highly conserved in *Drosophila* TTLL5 (van Dijk et al., 2007) (**Table 1**). According to the

FlyBase data, *Drosophila TTLL5* mRNA is expressed at high levels in ovaries (**Table S2**),

even higher than in the nervous system where initial studies were performed (Devambez et

al., 2017). We thus focused on *TTLL5*'s possible functions during *Drosophila* oogenesis to

shed light on possible *in vivo* roles of glutamylated α-tubulins.

MTs have an intrinsic polarity with a plus and a minus end. On these tracks, kinesin

motors transport cargo to the plus ends and dynein motors move cargo towards minus ends.

Evidence for the importance of PTMs of tubulin subunits has been reported already *in vitro.*

Artificial tethering of 10E peptides to the side chains at position E445 of α-tubulin and E435

of β-tubulin by maleimide chemistry increased the processivity (the run length) of kinesin

motors on tubulin. For Kinesin-1 this was a 1.5-fold increase (Sirajuddin et al., 2014). In

contrast to Kinesin, Dynein/dynactin motors preferred a Tyr at the C-term of α-tubulin to

initiate processive transport *in vitro*, because the motor favored this isoform as an

attachment point to dock onto the MTs (McKenney et al., 2016). Recent work now also

demonstrated a role for the mouse tubulin glycylation enzymes. *in vitro* fertility assays

showed that the lack of both *TTLL3* and *TTLL8* perturbed sperm motility causing male

subfertility in mice. Structural analyses further suggested that loss of glycylation decreased

axonemal dynein activity (Gadadhar et al., 2021).

A recent *in vivo* mouse study showed that hyperglutamylation of α-tubulin by TTLL1, but

not β-tubulin, increased the overall (both anterograde and retrograde) motility of

mitochondria, perturbed axonal transport, and led to neurodegeneration (Bodakuntla et al.,

2021). Somewhat surprisingly, however, another study showed that the vesicular axonal

transport was not affected by the absence of MT glutamylation in *Drosophila TTLL5* mutant

larvae (Devambez et al., 2017). Given these results and because the *Drosophila* gene is

also highly expressed in ovaries, we studied the *in vivo* role of *TTLL5* with a special focus on

the PTMs of the different α-tubulin isoforms in the ovary. This revealed that *TTLL5* is

essential for the normal glutamylation of the standard TBA1 and 3 isoforms. Surprisingly,

even in wild-type ovaries, the oogenesis-specific TBA4 was not glutamylated, suggesting

that this isoform may have evolved to produce MTs that contain regions that cannot be

glutamylated. We also found that *TTLL5* is required *in vivo* for the localization of mRNAs and

for cytoplasmic streaming, suggesting that these activities indeed depend on

polyglutamylation of α-tubulin. The transport of mitochondria in wing nerves is another

attractive system to study organelle transport (Vagnoni and Bullock, 2016) (Hollenbeck and

Saxton, 2005). We found the initiation of axonal transport of mitochondria in wing nerves

was strongly affected by lack of *TTLL5*. The overall motility and the processivity of the

mitochondrial transport also appeared to depend on α-tubulin glutamylation. Our work,

therefore, reveals the *in vivo* importance of glutamylation of specific α-tubulin isoforms.

Results

Drosophila *TTLL5* is required for MT glutamylation of ovarian TBA1/3.

To study the function of TTLL5, distinct *TTLL5* loss-of-function mutations were created and

characterized (**Fig. 1A**). The *TTLL5^{pBac}* and *TTLL5^{Mi}* mutant strains carry a piggyBac and a

Minos transposon, respectively, within the open reading frame of *TTLL5*. The *TTLL5^{MiEx}*

mutant was generated by imprecise excision of the Minos element, which caused the

formation of a premature stop codon at gene position 8,132 in the open reading frame

(codon position 392). An anti-TTLL5 antibody that cross-reacts with other proteins in ovarian

extracts also recognized a band at the size of TTLL5. This band turned out to be absent in

ovarian extracts prepared from the three *TTLL5* mutants, suggesting that they are protein

nulls for *TTLL5* or make only a truncated protein (**Fig. 1B**). A transgene containing the wild-

type *Drosophila TTLL5* sequence fused to the Venus fluorescent protein sequence under

UAS control (UASP-*Venus::TTLL5*) was also produced as a tool to rescue the phenotypes

produced by the *TTLL5* mutations and for overexpression studies (**Fig. 1A**).

To understand the effect of *TTLL5* on the glutamylation of the C-terminal tail of α-tubulins,

C-terminal peptides of the different α-tubulin isoforms were analyzed for post-translational

modifications by liquid chromatography-mass spectrometry (LC-MS). Proteins extracted from

ovaries were purified through SDS-PAGE. α-tubulin bands were cut out and digested with

the protease trypsin-N for MS. Because the resulting C-terminal peptides are identical

between the TBA1 and TBA3 isoforms, we will refer to them as TBA1/3 (**Fig. 2A**). Analyses

of wild-type ovarian extracts revealed that the general *Drosophila* isoforms TBA1/3, but not

the ovarian specific TBA4 isoform of α-tubulin showed glutamylation on the C-terminal tail of

α-tubulins (**Fig. 2A**, **S1**). For TBA1/3, side chain glutamylation was identified on all 4

glutamic acid residues in the tail region, Glu443, Glu445, Glu448, and Glu449, and the

longest side chains identified consisted of 3Es in wild-type ovaries **(Fig. 2A,2B)**. Determining

the frequency of C-terminal peptides containing the entire primary sequence and 0, 1, 2, and

3 Glu(E) modifications, we found that 62% of the peptides were modified in wild-type ovaries,

41% with 1E, 14% with 2Es, and 7% with 3Es **(Fig. 2C)**. On the other hand, no glutamylation

was observed in *TTLL5^{pBac}* and *TTLL5^{MiEx}* mutants. This clearly shows that *Drosophila*

TTLL5 is essential for the side chain glutamylation of the C-terminal Glu443, Glu445, Glu448,

and Glu449 of TBA1/3.

We also measured the levels of monoglutamylation of α-tubulin by western blotting using

the GT335 antibody. This antibody reacts with the first glutamate on the glutamate side

chain even if it is polygutamylated (**Fig. 3A**) (Wolff et al., 1992). Note that the GT335

antibody does not reliably detect all mono glutamylated residues in *Drosophila* tubulins

because GT335 preferentially recognizes the first Glu attached to a Glu residue in an acidic

environment. In mice, the middle Glu in the sequence -G**E**E- of the C-terminal tail of α-

tubulin is such an "acidic" site. Once glutamylated, the branched Glu modification can be

recognized by GT335 (Bodakuntla et al., 2021). In contrast, in *Drosophila* α-tubulin, both

E443 and E445 in the sequence -G**E**G**E**G- are flanked by Glycines on both sides which does

not provide the acidic environment needed for efficient recognition by GT335. The GT335

signal may therefore mostly stem from the glutamylation on the consecutive Glu residues

E448 and E449 in the sequence -GA**EE**Y. On Western blots from wild-type extracts, the

GT335 antibody produced a band at the expected position and this band was reduced in all

TTLL5 mutants. The GT335 signals were restored and even increased in some *TTLL5*

mutants upon overexpression of the rescue construct *Venus::TTLL5*, showing that the signal

is specific and can be used to monitor *TTLL5* activity **(Fig. 3B)**. We conclude that *TTLL5* is

required for monoglutamylation of α-tubulin.

The 1D5 antibody recognizes α-tubulin with an -XEE sequence at the C-terminal α-

carboxylate group (with X being a variable residue) (Rüdiger et al., 1999) **(Fig. 3A')**.

Therefore, in principle, all the Glu residues on E443, E445, E448, and E449 but also the C-

terminal Glu residues E448/449 can be recognized if they are not followed by another amino

acid (such as Tyr) at the α-carboxylate group. 1D5 does, therefore, not cross-react with C-

terminal Glu residues E448/449 in tyrosinated α-tubulin. Because *w* control ovaries showed

high 1D5 cross-reactivity with a polypeptide of the predicted size, this, therefore, indicates

the presence of glutamate side chains containing at least 2 glutamates and possibly de-

tyrosinated isoforms of α-tubulin (Gadadhar et al., 2017; Preston et al., 1979) **(Fig. 3B',**

3B''). Consistent with the MS analysis, the *TTLL5* mutants showed a strongly reduced signal

with 1D5, indicating that loss of *TTLL5* function prevents normal polyglutamylation. The 1D5

signal was again restored in all *TTLL5* mutants upon expression of the rescue construct.

Overexpression of TTLL5 in *TTLL5^{pBac}* and *TTLL5^{Mi}* even produced a stronger 1D5 signal

and an up-shifted band, suggesting that α-tubulin gets hyper-glutamylated, maybe on

additional sites **(Fig. 3B')**. We also tested whether the overexpression of *TTLL5* in a wild-

type background affected polyglutamylation in the same way. Indeed, *UASP-Venus::TTLL5*

overexpression in ovaries led to hyper glutamylation as evident from the stronger signal and

the additional upshifted band **(Fig. 3C)**. Glutamylation of β-tubulin was also evaluated by

using the antibody β-monoE which specifically recognizes monoglutamylation of the C-

terminal tail of β-tubulin (Bodakuntla et al., 2021) **(Fig. 3B'')**. Results firstly showed a strong

signal of β-monoE in control ovaries, indicating that monoglutamylation occurred also on β-

tubulin in ovaries. However, the β-monoE levels were kept unchanged no matter whether

TTLL5 was overexpressed or mutated. These results reveal that *TTLL5* is essential for the

glutamylation of MTs in the ovary, acting specifically on α-tubulin but not on the β- tubulin

subunit. It also suggests that TTLL5 levels are usually limiting in ovaries.

Most of the murine TTLL5 residues involved in α-tubulin glutamylation (van Dijk et al.,

2007) are conserved in *Drosophila* TTLL5 **(Table 1)**. Two are known to be important for the

general enzymatic activity of TTLLs and two for the interaction with the C-terminal domain of

α-tubulin **(Fig. 3D)**. To find out whether these residues are indeed needed *in vivo* for the

enzymatic activity of *Drosophila TTLL5 towards* TBA1/3, we used CRISPR/Cas9 to mutate

the conserved codons in *Drosophila TTLL5. Drosophila* TTLL5 residues K282 and E517*,*

respectively, correspond to mouse K131 and E336*, respectively.* They directly interact with

ADP/ATP and thus affect general TTLL5 enzymatic activity (van Dijk et al., 2007). TTLL5^{R339}

corresponds to mouse TTLL5^{R188}*,* which is important for TTLL5's glutamylation activity.

MmTTLL5^{R188} resides in the β6-7 loop, which forms a salt-bridge interaction with the C-

terminal tail of α-tubulin and orients it towards the ATP/ADP-binding site (van Dijk et al.,

2007). Glutamylation of α-tubulin was impaired in all these *TTLL5* mutants where the

conserved residue was replaced **(Fig. 3E)**. Since these mutations did not abolish the stability

of the corresponding mutant protein (**Fig. S2**), our results revealed the importance of E517,

K282, and R399 of *Drosophila* TTLL5 for glutamylation. Because almost all active site

residues of human TTL (Prota et al., 2013; van Dijk et al., 2007) are also conserved between

mammalian and *Drosophila* TTLL5 **(Table 2)**, we also mutated the *Drosophila TTLL5^{P522}*

codon corresponding to human *TTL^{P336}* (and mouse *TTLL5^{P371}*), which is required for

anchoring TTL to α-tubulin by forming hydrogen bonds with C-terminal tail residues of α-

tubulin (Prota et al., 2013). However, the importance of *TTLL5^{P371}* has not been reported in

mammalian systems. Somewhat surprisingly, in *Drosophila*, P522 turned out to be essential

for glutamylation as well, revealing P522 as a novel important residue for the glutamylation

function of TTLL5.

Other PTMs of the C-terminal tail of ovarian α-tubulin

Our MS results also detected glycylated C-terminal α-tubulin peptides in ovarian samples.

The glycylations were observed on the C-terminal tail of the *Drosophila* ovarian TBA1/3

isoforms, but not on TBA4 **(Fig. S3A, S3B)**. Up to 2 glycines were detected mostly on

Glu443, Glu445, Glu448, or Glu449 of TBA1/3. The frequency of C-terminal peptides

containing Gly (G) modifications was similar or somewhat higher in *TTLL5* mutants

compared to the wild-type control **(Fig. S3C)**, suggesting that *TTLL5* is not needed for this

modification or even counteracts it.

Western blotting showed no or only minor changes in tyrosinated α-tubulin levels in any of

the null or CRISPR/Cas9 generated point mutants, and *TTLL5* overexpression did also not

affect these levels **(Fig. 3B, 3C, 3E).** TTLL5 levels, therefore, do not appear to affect levels

of tyrosinated α-tubulin in ovaries.

Effect of *TTLL5* on Kinesin-1 dependent refinement of posterior localization of Staufen

osk mRNA localization, combined with translation control, is important for targeting Osk

protein expression to its proper posterior compartment in the cell (Suter, 2018; Weil, 2014).

Staufen is a ribonucleoprotein (RNP) that associates with *osk* mRNA during *osk* mRNA

localization. The localization of Staufen/*osk* mRNA requires the dynein/dynactin mediated

minus-end transport machinery in the early stages of oogenesis. Subsequently, the

Staufen/*osk* mRNA becomes localized to the posterior of the oocyte from mid to late stages

of oogenesis and this particular Staufen/*osk* mRNA localization phase within the oocyte

depends on a kinesin driven process including Kinesin-1 based active transport and

ooplasmic streaming in stage 10B (S10B) oocytes (Kato and Nakamura, 2012) (Lu et al.,

2018) (Brendza et al., 2000) (Zimyanin et al., 2008). To find out whether *TTLL5* contributes

to this process, we studied the role of *TTLL5* in dynein and kinesin transport by analyzing the

localization of Staufen in ovaries lacking functional TTLL5.

An established method to quantify the localization of Staufen to the posterior cortex is to

measure the length of the posterior Staufen crescent along the cortex (Lu et al., 2016). A

shorter crescent is a sign of better, tighter localization. We found that the length of the

posterior Staufen crescent was significantly increased in *TTLL5* null and *TTLL5* point mutant

ovaries compared to control ovaries (**Fig. 4A', 4A''', 4B'-B''**). Furthermore, Z-stack images

revealed a more diffusely spread Staufen signal in all *TTLL5* mutants, although this varied

more between different samples **(Fig. 4A', 4B')**. The increased length of the crescent was

fully reversed to the normal length in all UAS-*Venus::TTLL5* rescued strains **(Fig. 4A''-A''')**.

We also analyzed the distribution of Venus::TTLL5 during oogenesis. Antibody staining

suggested that Venus::TTLL5 strongly accumulated in the oocyte and especially along the

cortex **(Fig. 4A'')**. In conclusion, the posterior localization refinement of Staufen in S10B

oocytes requires *TTLL5* and, because the point mutation specifically blocks the enzymatic

activity of TTLL5, it appears that the proper localization requires glutamylation of tubulin.

To test if *TTLL5* is also required by the dynein/dynactin machinery and the Staufen/*osk*

mRNA transport during previtellogenic stages of oogenesis, we also imaged the localization

of Staufen and Bicaudal-D (BicD) protein in early stages. BicD is the linker that couples

diverse cargos to the dynein/dynactin motor and is thus a good marker to evaluate dynein

transport function (Claussen and Suter, 2005). Both BicD and Staufen showed similar

localization patterns during early oogenesis in *TTLL5* mutant and control ovaries **(Fig. 4C-**

C'). The expression of BicD was not affected by the levels of TTLL5 as assessed by

Western blotting (**Fig. 4D**, **Fig. S4**). These results indicated that in ovaries, kinesin-

dependent localization processes require *TTLL5* whereas the dynein-dependent transport

does not appear to depend on *TTLL5*.

Kinesin-1 dependent ooplasmic streaming in stage 10B oocytes requires *TTLL5*

Ooplasmic streaming is a bulk movement of the oocyte cytoplasm that circulates and

distributes mRNAs and proteins in the oocyte, including the Staufen/*osk* mRNA RNPs. It

allows these mRNA and proteins to reach their intended position where they become

anchored. Slow and nondirectional flows initiate at stages 8–9 and are followed by rapid and

circular streaming at S10B (Lu et al., 2016). Kinesin-1-dependent microtubule sliding is

critical for this fast ooplasmic streaming (Lu et al., 2016). The kinesin-dependent final

posterior localization of Staufen in S10B oocytes depends on this process and *TTLL5*. Aside

from this, the TTLL5 protein was present in the cortical region of the oocyte **(Fig. 4A'')**, the

region where the microtubule sliding occurs (Lu et al., 2018). We thus wanted to find out if
TTLL5 contributes to an efficient kinesin-dependent sliding that drives ooplasmic streaming.

We firstly examined the levels of Kinesin-1 and found that they were neither affected in
TTLL5 mutant ovaries nor in ovaries overexpressing *Venus::TTLL5* (**Fig. 5A, Fig. S4**). To
test if *TTLL5* is needed for ooplasmic streaming, we followed this process in real time by
measuring vesicle movement in DIC time-lapse movies (**Mov. S1-S4**). We then tracked the
movement of the vesicles along the cortex region where microtubule sliding occurs (**Fig. 5B**).
The velocity of the tacked vesicles was not to be affected in *TTLL5* mutant ovaries (**Fig. 5C**).
However, the analysis of the kymographs of these tracked vesicles showed that while control
ovaries have a normal streaming pattern with vesicles moving with a flow direction parallel to
the cortex, *TTLL5^{MiEx}* and *TTLL5^{E517A}* mutant ovaries displayed a disordered streaming
pattern (**Fig. 5D**). The normal streaming pattern was restored in *TTLL5^{MiEx}* ovaries
expressing a *Venus::TTLL5* rescue construct. Because the point mutation specifically blocks
the enzymatic activity of TTLL5, our results suggest that TTLL5 is important to control the
directionality of the fast ooplasmic streaming through its role in MT glutamylation.

Transport of mitochondria in neurons depends on the glutamylation activity of *TTLL5*

Previous studies in *Drosophila* reported that *TTLL5* is required for the initiation of α-tubulin
glutamylation in the nervous system (Devambez et al., 2017). However, the absence of α-
tubulin glutamylation seemed not to be detrimental to the tested nervous system functions in
Drosophila (Devambez et al., 2017). To study neuronal transport in the peripheral nervous
system, the wing nerves are an attractive system to monitor organelle transport such as the
movement of mitochondria (Vagnoni and Bullock, 2016) (Hollenbeck and Saxton, 2005).
Consistent with a previous report, we found that in extracts from a mixture of heads and
wings, α-tubulin was glutamylated and this PTM depended on the enzymatic activity of
TTLL5 (**Fig. 6A**); (Devambez et al., 2017). This suggests that the wing nerve could be a
suitable system for this test.

To study if neuronal transport of mitochondria in wing neurons depends on glutamylation of
α-tubulin, we tracked the movement of GFP-labelled mitochondria (Mito::GFP) in selected
areas of both the L1 and L3 wing veins of freshly eclosed flies (**Fig. 6B**). Time-laps videos of
Mito::GFP movement in the L1 and L3 veins, respectively, are shown in **Mov. S5** and **Mov.**
S6, respectively. Motile mitochondria were manually tracked using the Fiji MTrackJ plugin
(**Fig. 6C**). Movies and still images clearly showed a decrease of motile mitochondria in the
region close to the arch of the L1 vein and no transported mitochondria in the L3 vein of the
TTLL5^{E517A} mutant, pointing to the importance of the enzymatic activity of TTLL5 for proper

transport. For quantitative analysis, the ratio of motile particles was calculated based on the

values divided by total particles from 3 wings of each genotype (**Fig. 6D**). The total particles

include all the static and transporting ones which were counted based on the signal from one

chosen still image. Quantifications showed, firstly, that the total number of particles in the L1

vein (384) was twice as high as in the L3 vein (186). This is likely because the L1 vein

contains more neurons than the L3 (Vagnoni and Bullock, 2016). Secondly, however, in wild-

type wings, the fraction of motile mitochondria is higher (24%) in L3 veins than in L1 veins

(7.8%). This might be because compared to the L3 vein, which consists mainly of axons

(Vagnoni and Bullock, 2016) (**Fig. S5**), the region close to the arch of the L1 vein contained

not only axons but also several cell bodies and dendrites (Vagnoni and Bullock, 2016) (**Fig.**

S5). Mitochondria in dendrites are longer than those in axons (Chang et al., 2006). The

longer mitochondria are less motile than the short ones (Misgeld et al., 2007). The large

number of less motile mitochondria in cell bodies and dendrites "diluted" the motile

mitochondria ratio in L1 veins. Because axons are easily distinguishable in the L3 vein, the

region of L3 seemed to be a better choice for axonal transport studies than the one in the L1

vein. The quantification of the motile ratio in the $TTLL5^{E517A}$ mutant showed no motile

mitochondria in the L3 vein of the $TTLL5^{E517A}$ mutant (**Fig. 6D**), strongly indicating that

$TTLL5$ is required for the initiation of axonal transport of mitochondria. Apart from this, the

quantification of the motile ratio in the L1 showed a drop from 7.8% (plus-end 5.3% and

minus-end 2.5%) in the control to 1.9% in $TTLL5^{E517A}$ mutant wings (plus-end 1.5% and

minus-end 0.4%) (**Fig. 6D** and **Table 3**). These results reveal the importance of the

enzymatic activity of $TTLL5$ for *in vivo* mitochondrial transport in wing nerves.

Because no motile mitochondria were found in the L3 region, we evaluated other

transport parameters in the L1 region. The *Drosophila* sensory neurons contain axons with

plus-end-out MTs and dendrites with minus-end-out MTs (Stone et al., 2008; Zheng et al.,

2008). Though the L1 region is enriched in a mix of axons, dendrites, and cell bodies, the

direction of the organelle transport can still be established. The mitochondria transported

towards the thorax of a fly move by plus-end transport, driven by kinesin, whereas

mitochondria transported away from the thorax are transported by dynein. We examined the

parameters of running patterns, trafficking length, and velocity of mitochondrial movement in

region L1. The start of a run was chosen at the time point when a motile particle began to

move from a stationary position, and the stop of a run was defined when the particle stopped

moving. Pausing of a run was called when the particle moved at a rate lower than 0.2 µm/s.

Particles with a speed higher than 0.2 µm/s were considered as "trafficking". A schematic

running pattern is shown in **Fig. 6E.** Bidirectionally moving particles were seen only very

rarely in our case. In these cases, they usually moved a short distance towards one direction

and suddenly turned back and ran in the opposite direction. For the plus-end transport

(**Table 3)**, the average total duration of a running time showed no significant difference

between controls (92s) and *TTLL5* mutants (98s). As well, no significant difference in the

mean velocity of the tracked mitochondria was observed between controls (0.37 μm/s), and

TTLL5 ^{E517A} mutants (0.36 μm/s). However, the average pausing time ratio in *TTLL5 ^{E517A}*

(63%) showed a decrease compared to the control (74%), suggesting that α-tubulin

glutamylation was required for attaching and holding or stopping mitochondria during their

whole "run" along MTs (see **Fig. 6E**). Consistent with this observation, mitochondria in

TTLL5 ^{E517A} moved longer distances (12μm) than the mitochondria observed in the controls

(7μm). Even though our research focused on the plus-end transport, preliminary numbers for

the minus-end transport suggest that the motile frequency, total duration and pausing time

ratio might decrease in *TTLL5 ^{E517A}*, too (**Table 3**). However, the small numbers for

TTLL5 ^{E517A} do not allow to conclude this.

It appears that two different mechanisms might be involved between the start and the

ongoing transport of individual mitochondria. At the first step, the lack of glutamylation

hinders the initiation of the axonal transport of mitochondria, indicating that initiation of

processive axonal transport heavily depends on glutamylation. Once mitochondria have

started to move, the lack of α-tubulin glutamylation increases the plus-end trafficking time

and trafficking length of the mitochondria in a region of the wing nerve that contains axons,

dendrites and cell bodies, indicating that glutamylation is also needed for holding or

anchoring mitochondria on the MTs during trafficking in neurons. Altogether, these results

demonstrated the role of α-tubulin glutamylation for the efficient mitochondrial transport in

wing nerves.

Discussion

Both western blotting and MS results from wild-type and mutant *Drosophila* revealed that

TTLL5 is essential for polyglutamylation of α-tubulin in ovaries. The complete lack of even

monoglutamylated α-tubulin tails in the presumptive null mutants of *TTLL5* strongly suggests

a requirement for TTLL5 for adding the first branching Glu to the Glu side chains in the C-

term. The requirement for adding the first Glu could also account for the lack of

polyglutamylation in the mutants. This result is also consistent with results from HeLa cells

and the *Drosophila* nervous system (Devambez et al., 2017; van Dijk et al., 2007). Further

MS results indicated that *TTLL5* specifically functions to glutamylate the TBA1/3, but not the

TBA4 isoforms of α-tubulin. Up to three Glu residues were found added to the positions

E443/E445/E448/E449 of TBA1/3 (**Fig. 2A, 2B**). Even though TBA4 is an ovarian-specific

isoform, our MS analysis showed a lower amount of TBA4 peptides than TBA1/3 peptides

(**Table S3**), suggesting that it accumulated at a lower level in ovaries. The fact that we could

not find any glutamylation in TBA4 could indicate that this isoform evolved because the

ovary has a need for non-glutamylated α-tubulins. As we could show here, glutamylation

plays an important role in kinesin-driven processes. The evolutionary appearance of a non-

glutamylatable tubulin isoform may indicate that this tissue has a need for special

microtubules that favour specifically other MT motor transport, and this could be achieved by

setting up MTs or MT regions that are preferentially being used for dynein-driven transport.

Our MS analysis revealed that polyglycylation of α-tubulin also takes place in the ovarian

tissue and that glycylation modifies the same sites in the C-terminal domain as

polyglutamylation does, the side chains of the Glutamic acid residues. Because glycylation is

still seen in the *TTLL5* mutants, TTLL5 does not seem to be the enzyme for this reaction,

and glutamylation of one site is not a prerequisite for glycylation of another site. It would now

be interesting to find out whether glycylation even serves to prevent glutamylation or to

compete with it and whether this is used as another level of control of microtubule activity. A

preliminary analysis of glycylation in the *TTLL5* mutants suggests that this might be possible

(**Fig. S3C**). In *Tetrahymena* such a competition has been described because there, Glutamic

acid and Glycine ligases competed for the same sites or affected each other by sterically

blocking adjacent sites of tubulins (Redeker et al., 2005; Wloga et al., 2009). Only a

surprisingly small fraction of the *TTLL* family is expressed in *Drosophila* ovaries according to

Flybase. Aside from *TTLL5*, we only found evidence for the expression of *TTLL4A* and

TTLL12 (**Table S2**). Our genetic analysis of *TTLL5* showed that it is not required for

glycylation. The predicted mouse homolog of the *Drosophila TTLL4A* was found to

glutamylate β-tubulin (Mahalingan et al., 2020), suggesting that this isoform could be the one

required for β-tubulin glutamylation in ovaries. The function of the predicted homolog of

Drosophila TTLL12 in mammals was poorly studied and the core TTL domain of mouse

TTLL12 is less conserved than other mouse E- and G- ligases of the TTLL family (van Dijk et

al., 2007). It is, therefore, not clear yet which enzyme performs the polyglycylation of α-

tubulin in *Drosophila* ovaries.

The ooplasmic streaming process is required for the posterior localization of Staufen/*osk*

mRNA and is mediated by an MT-sliding mechanism (Lu et al., 2016). This MT-sliding, which

is an essential step for normal oogenesis, involves the sliding between the cortically

anchored microtubules and the free cytoplasmic microtubules that are located close to the

cortex. This sliding between the two classes of microtubules is Kinesin-1 dependent. Our

data show that *TTLL5* is needed for normal oocyte cytoplasmic streaming and for the
posterior localization of Staufen/*osk mRNA*, suggesting that it affects these processes
through a role in kinesin-1 dependent MT sliding.

It is worth pointing out that *in vitro* studies demonstrated that the artificial tethering of
glutamyl peptides to the side chains of both α- and β-tubulin increased the run length of
Kinesin-1 on tubulins (Sirajuddin et al., 2014). *Drosophila TTLL5* is essential for
glutamylation of α-tubulin but not for β-tubulin. These results, first of all, provide the best
evidence yet that *TTLL5* mediated glutamylation of α-tubulin plays an important role in
Kinesin-1 driven processes *in vivo*. However, the phenotypes including reduced ooplasmic
streaming and Staufen localization observed in *TTLL5* mutants were not as severe as the
ones that arose when Kinesin heavy chain (KHC) was not functional (Lu et al., 2016).
Besides, in the wing nerve experiment, though initiation of mitochondria was heavily
dependent on the glutamylation of α-tubulin, there are still some particles which started to be
transported in *TTLL5* mutants. It therefore seems possible that α-tubulin glutamylation is not
absolutely required for transport initiation and this poses the interesting novel question
whether these Kinesin-1 dependent processes can still operate if β-tubulin is glutamylated.

In *KHC* mutants, there is an overall decrease in anterograde as well as dynein-driven
retrograde transport of mitochondria in *Drosophila* larval motor axons (Pilling et al., 2006).
We observed a similar loss of motile mitochondria in the axon-rich region in the *TTLL5*^{E517A}
mutant wing (**Fig. 6C, 6D**). Because both previous studies (Devambez et al., 2017) and our
data (**Fig. 6A**) showed that *TTLL5* was required for α-tubulin glutamylation in the nervous
system, all these results suggested that α-tubulin glutamylation is a crucial factor for the
Kinesin-1 mediated initiation of mitochondrial transported in axons.

By analyzing the dynamics of the few motile mitochondria observed in L1 veins, we found
that in the absence of glutamylation, the plus-end mitochondria can run longer distances with
shorter pausing times. (**Fig. 6G**). Interestingly, a recent study linked the reduced trafficking
time of mitochondria during axonal transport to neurodegeneration in mice and this was
caused by hyperglutamylation of α-tubulin but not β-tubulin (Bodakuntla et al., 2021). It thus
strongly indicated the glutamylation of α-tubulin causes the halting of the ongoing transport
of single mitochondria *in vivo*. Because the orientation of the MTs in the *Drosophila* wing
neurons is unidirectional and the plus-end transport of mitochondria was affected in the
absence of the glutamylation of α-tubulin, we now learned that glutamylation is at least
needed for the Kinesin-1-based transport.

Several mechanisms have been proposed to stop mitochondrial transport in neurons. For

instance, tethering mitochondria to the actin cytoskeleton (Pathak et al., 2010), blocking the

binding of Kinesin-1 to MTs by increasing Ca^{2+} (Wang and Schwarz, 2009), abolishing the

interaction between Kinesin-1 and mitochondria by degrading one of the linker proteins

(Miro; Wang et al., 2011), or possibly traffic jam. We here provided *in vivo* evidence that

glutamylation of MTs is an important regulatory mechanism that contributes to the

sophisticated neuronal mitochondrial transport system.

Materials and methods

Drosophila genetics

The fly stocks used in this study, MattubGal4 (7062), *TTLL5^pBac* (16140), *TTLL5^Mi* (32800), and *Df(3R)BSC679* (26531), were obtained from the Bloomington Drosophila stock center. The *TTLL5^MiEx* mutation was made by imprecise excision of the Minos element, which left a 3 base insertion (CTA) that introduces a premature stop codon at genomic position 8,132 in the open reading frame (polypeptide position 392). The different *TTLL5* point mutations were generated by CRISPR/Cas9. The two Drosophila stocks used for *TTLL5* gene editing, *y[1] M{w[+mC]=nos-Cas9.P}ZH-2A w[*]* and *y[1] v[1] P{y[+t7.7]=nos-phiC31\int.NLS}; P{y[+t7.7]=CaryP} attP40* were obtained from the Bloomington Drosophila Stock Center (54591) (Port et al., 2014) and (25709), respectively. Target sequences near the K131, R339 R376, and E517 codons were selected to introduce INDELs and the following point mutations: K131A, R339A (b6-7 loop), R376A, and E517A. Phosphorylated and annealed primers (Table S1A) needed to construct the corresponding coding sequences for a single guide RNA (sgRNA) were each cloned into the BbsI site of *pCFD5* according to the protocol published on www.crisprflydesign.org. After amplification of the vector in XL1 blue cells, the plasmids were sequenced to verify correct assembly. Subsequently, they were injected into *attP40* embryos. TTLL5-guide stocks with integrated genes encoding sgRNAs were established *(=v; attP40{v^+, TTLL5-guide}/CyO)*. To obtain INDELs for TTLL5, a three-generation crossing schema was set up. Firstly, *nos-cas9* virgins were crossed with males from the different TTLL5-guide stocks. Secondly, the subsequent single male progeny was crossed with *w; Ly/TM3, Sb* virgins. Thirdly, resulting single *TM3, Sb* males were again crossed with *w; Ly/TM3, Sb* virgins to establish the stocks. The nature of the mutated *TTLL5* gene in each stock was determined by sequencing. To introduce specific point mutations in the *TTLL5* gene, embryos from a cross with *y w {w^+, nos-cas9}* virgins and TTLL5-guide males were injected with the appropriate template (**Table S1B).** The resulting individual adult males were crossed with *w; Ly / TM3, Sb* virgins, and stocks were established. The presence of a mutated *TTLL5* gene was confirmed by restriction digestion and subsequent sequencing. All the *TTLL5* alleles were crossed to *Df(3R)BSC679* to test the effect of the *TTLL5* mutations in hemizygous animals. The *UASp-venus::TTLL5* DNA construct was made from the *TTLL5* cDNA amplified by PCR (forward primer: 5′ - GTTCAGATCTATGCCTTCTTCATTGTGTG -3′; reverse primer: 5′ - GAGTCATTCTAGAGCTTCATAGAAATACCTTCTCC- 3′) and inserted into the *pUASP-venus* vector via Xba I/Bgl II ligation. DNA was inserted into the *attP-58A* landing platform

482   (24484) to generate the transgenetic *UASp-venus::TTLL5* fly strain. *ApplGal4* and *UAS-*

483   *mito::GFP* were kindly provided by Simon Bullock's group (Vagnoni et al., 2016).

Western blotting

Western blotting performed with ovarian extracts was carried out with 10-20 pairs of ovaries

for each genotype. Newly hatched females of the indicated genotypes were collected and

crossed with *w* males in fresh vials with ample yeast food for two days. Fully developed

ovaries were then dissected and collected. To analyze neuronal samples, a mixture of head

and wing tissue from 30 flies was collected for each genotype. The samples were directly

lysed in Laemmli buffer, boiled at 95°C for 10 min, and analyzed by SDS-PAGE.

Chemiluminescence detection was done with the Amersham imager 600 (GE healthcare).

The following primary antibodies were used: mouse 1D5 (1:500, Synaptic System), mouse

GT335 (1:200, Adipogen), mouse DM1A (1:1,000, Sigma), mouse tyrosinated α-tubulin

(1:1,000, Sigma), rabbit GFP (1:2,000, Sigma), rabbit GAPDH (1:1,000, GeneTex), rabbit

TTLL5 (1:500, GenScript), mouse BicD (1B11 plus 4C2, 1:5) (Suter and Steward, 1991),

rabbit KHC (1:1,000, cytoskeleton), rabbit β-monoE (1:1,000) (Bodakuntla et al., 2021).

Secondary antibodies were HRP-linked goat anti-mouse (1:1,0000, GE Healthcare) and

HRP-linked goat anti-rabbit (1:10,000, GE Healthcare). Antibodies were incubated in 5%

non-fat milk PBST (PBS with 0.1 % Triton X-100) and washed with PBST.

Immunoprecipitations (IPs) and Mass Spectrometry

Proteins were extracted from the ovaries of 100 3-day-old, well-fed females of each

genotype, crossed to wild-type males, and proteins were immunoprecipitated with an α-

tubulin antibody (Sigma) bound to G-protein coupled magnetic beads. Samples (ligated with

beads) were boiled and run on SDS-PAGE. These were then stained with Coomassie Blue.

The gel pieces containing the α-tubulin region were cut, alkylated, and digested by trypsin N

(ProtiFi) for 3 hours at 55°C. C-terminal peptides of α-tubulin were analyzed by liquid

chromatography (LC)-MS/MS (PROXEON coupled to a QExactive mass spectrometer,

Thermo Fisher Scientific). Samples were searched against the UniProtKB Drosophila

database using Pipeline, PEAKS, Easyprot and MSfragger as search engines. Peptides

were shown by PSM value (peptide spectrum match). To avoid repetition of total PSM

quantification, we searched only for one type of PTM in an analysis per time. Total PSM

indicates the sum of PSMs of primary C-terminal peptides with either glutamylation or

glycylation modifications, or non-modified peptides.

Ooplasmic streaming in live oocytes

Three days old females were kept together with wild-type males under non-crowding
conditions at 25°C on food containing dry yeast. The determination of ooplasmic streaming
was based on the method of Lu and colleagues (Lu et al., 2016). Ovaries were directly
dissected in 10S halocarbon oil (VWR) and eggs were teased apart in the oil drop on a
24×50 mm coverslips (VWR). Living stage 10B oocytes were directly observed and imaged
with the 60×1.4 N.A. oil immersion objective on a Nikon W1 LIPSI spinning disk with a
2x Photometrics Prime BSI CMOS camera. Time-lapse images were acquired every 2s for 2
min under bright-field (BF) controlled by Nikon Elements software. Flow tracks were
generated by kymographs using Fiji (Schindelin et al., 2012) plugin Multi Kymograph (https://
imagej.net/ Multi _Kymograph). The velocity of 100-200 particles/ooplasm was tracked using
Analysis > Distribution of Speed (Track-Based) by DiaTrack 3.05 (Vallotton and Olivier,
2013), with a maximum jumping distance of 3.24 µm/s. The datasets were exported to Excel,
where the mean velocities of each ooplasm were analyzed. Then the velocities of different
ooplasms from different genotypes were shown as means ± S.E.M. by Prism 5 software
(GraphPad).

Immunostaining and confocal microscopy

Ovaries were dissected in 1× PBS and fixed in buffer (200 µl 4% formaldehyde (Sigma), 600
ul Heptane (Merck Millipore), 20 ul DMSO (Sigma) for each group) for 20 min. Samples were
blocked in PBST (PBS with 0.1 % Tween 20) with 5% non-fat milk powder and 0.1 % Triton
X-100 for 1 h. Ovaries were incubated with rabbit anti-Staufen (1:2,000) (St Johnston et al.,
1991), mouse anti-BicD (1B11 plus 4C2, 1:5) (Suter and Steward, 1991) and rabbit anti-GFP
(1:200, Sigma) antibodies at 4°C overnight, and incubated with secondary Alexa Fluor goat
anti-mouse 647 (1:400, Jackson ImmonoResearch) and Alexa Fluor goat anti-rabbit 488
(1:400, Life technologies Thermo Fischer) at room temperature for 3 h. DNA and nuclei were
stained using Hoechst 33528 (Thermo Fischer Scientific) for 10 min. Mounted samples were
imaged with a Leica TCS SP8 confocal microscope equipped with a 63×1.40 oil lens and
controlled by the Las X software (Leica). Images of single focal planes and 12 µm Z
projection were acquired. The crescent length of Staufen was processed by the Fuji
segmented line tool based on the single plane. The data were shown as means ± S.E.M. by
Prism 5 (GraphPad software). The statistical significance was determined by an unpaired
two-tailed non-parametric t-test (Mann-Whitney test).

Mitochondrial transport in Drosophila wing neurons

Mitochondrial transport in Drosophila wing neurons was tracked as described (Vagnoni and
Bullock, 2016). In brief, newly eclosing flies were anesthetized with CO_2 and immobilized in

a custom-built chamber formed of two No.1.5 24×50 mm coverslips (VWR). Imaging of

wings was performed with a Nikon W1 LIPSI spinning disk microscope equipped with a 2x

Photometrics Prime BSI CMOS camera and a 60×1.4 N.A. oil immersion lens. Time-lapse

video of a single focal plane was acquired with an acquisition rate of 0.5 frame/s for 3 min for

Mito::GFP using a 488 nm laser.

Quantification of the movements was performed in Fiji. A region close to the axonal tract of

the wing arch of the L1 vein and a region of the L3 vein were selected and straightened with

the Straighten plugin (Eva Kocsis, NIH, MD). For each "run", the total run distance of more

than 2 µm was defined as "transported or motile mitochondria". To exclude the oscillatory

movements of mitochondria, a minimal instant velocity between each frame was defined at

0.2 µm/s. Transported mitochondria were manually marked with MTrackJ plugin (Meijering

et al., 2012) and tracked by the manual tracking plugin (Fabrice Cordelières, Institute Curie,

France). The total number of mitochondria was counted from a single frame using the

"Analyze particle" function of Fiji. Run length, velocity values, and total running & pausing

times were exported into Excel for analysis. The data were shown as means ± S.E.M. by

Prism 5 (GraphPad). The statistical significance was determined with the student's t-test.

Sequence alignment

Sequences were aligned by CLC Sequence Viewer 8.0 software (Qiagen). The UniProtKB

accession numbers are as follows: *HsTTL*: Q8NG68; *HsTTLL5*: Q6EMB2; *MmTTLL5*:

Q8CHB8; *DmTTLL5*: Q9VBX5;

Acknowledgments

We thank Carsten Janke (Institute Curie, France) for anti-β-monoE antibodies, Simon L.

Bullock (Cambridge, UK) for *ApplGal4* and *UAS-mito::GFP* fly lines, Michel Steinmetz (Paul

Scherrer Institute, Switzerland), and the Janke laboratory members for discussions, and

suggestions. We also thank the Mass Spectrometry center (University of Bern, Switzerland)

members Sophie Braga Lagache and Natasha Buchs for performing the MS experiments,

and Manfred Heller and Anne-Christine Uldry for peptide analysis and discussion of the

results.

Support

This work was financially supported by the project grants 31003A_173188 and

310030_205075 from the Swiss National Science Foundation (SNF; https://www.snf.ch/en)

to B.S. Equipment support was by an equipment grant from SNF (316030_150824) to B.S.

and by the University of Bern (https://www.unibe.ch). Additional salary and material support

were received from the University of Bern (https://www.unibe.ch) to B.S. The funders had no

role in the study design, data collection, and analysis, decision to publish, or preparation of

the manuscript.

The authors declare no competing financial interests.

Author contributions: Mengjing Bao, Beat Suter, and Maria Paula Vazquez-Pianzola

conceived the project and designed the experiments. Mengjing Bao, Beat Suter, Ruth Dörig,

and Dirk Beuchle performed the experiments and analyzed the data; Mengjing Bao and Beat

Suter wrote the manuscript.

Reference

Arce, C.A., J.A. Rodriguez, H.S. Barra, and R. Caputo. 1975. Incorporation of L-tyrosine, L-
phenylalanine and L-3,4-dihydroxyphenylalanine as single units into rat brain tubulin.
Eur J Biochem. 59:145-149.

Bodakuntla, S., X. Yuan, M. Genova, S. Gadadhar, S. Leboucher, M.C. Birling, D. Klein, R.
Martini, C. Janke, and M.M. Magiera. 2021. Distinct roles of α- and β-tubulin
polyglutamylation in controlling axonal transport and in neurodegeneration. *EMBO J*.
40:e108498.

Brendza, R.P., L.R. Serbus, J.B. Duffy, and W.M. Saxton. 2000. A function for kinesin I in
the posterior transport of oskar mRNA and Staufen protein. *Science*. 289:2120-2122.

Chang, D.T., A.S. Honick, and I.J. Reynolds. 2006. Mitochondrial trafficking to synapses in
cultured primary cortical neurons. *J Neurosci*. 26:7035-7045.

Claussen, M., and B. Suter. 2005. BicD-dependent localization processes: from Drosophilia
development to human cell biology. *Ann Anat*. 187:539-553.

Devambez, I., J. van Dijk, S. Benlefki, S. Layalle, Y. Grau, K. Rogowski, M.L. Parmentier,
and L. Soustelle. 2017. Identification of DmTTLL5 as a Major Tubulin Glutamylase in
the Drosophila Nervous System. *Sci Rep*. 7:16254.

Eddé, B., J. Rossier, J.P. Le Caer, E. Desbruyères, F. Gros, and P. Denoulet. 1990.
Posttranslational glutamylation of alpha-tubulin. *Science*. 247:83-85.

Gadadhar, S., G. Alvarez Viar, J.N. Hansen, A. Gong, A. Kostarev, C. Ialy-Radio, S.
Leboucher, M. Whitfield, A. Ziyyat, A. Touré, L. Alvarez, G. Pigino, and C. Janke.
2021. Tubulin glycylation controls axonemal dynein activity, flagellar beat, and male
fertility. *Science*. 371.

Gadadhar, S., S. Bodakuntla, K. Natarajan, and C. Janke. 2017. The tubulin code at a
glance. *J Cell Sci*. 130:1347-1353.

Hallak, M.E., J.A. Rodriguez, H.S. Barra, and R. Caputto. 1977. Release of tyrosine from
tyrosinated tubulin. Some common factors that affect this process and the assembly
of tubulin. *FEBS Lett*. 73:147-150.

Hollenbeck, P.J., and W.M. Saxton. 2005. The axonal transport of mitochondria. *J Cell Sci*.
118:5411-5419.

Janke, C., K. Rogowski, D. Wloga, C. Regnard, A.V. Kajava, J.M. Strub, N. Temurak, J. van
Dijk, D. Boucher, A. van Dorsselaer, S. Suryavanshi, J. Gaertig, and B. Eddé. 2005.
Tubulin polyglutamylase enzymes are members of the TTL domain protein family.
Science. 308:1758-1762.

Kalfayan, L., and P.C. Wensink. 1982. Developmental regulation of Drosophila alpha-tubulin
genes. *Cell*. 29:91-98.

Kato, Y., and A. Nakamura. 2012. Roles of cytoplasmic RNP granules in intracellular RNA
localization and translational control in the Drosophila oocyte. *Dev Growth Differ*.
54:19-31.

L'Hernault, S.W., and J.L. Rosenbaum. 1985. Chlamydomonas alpha-tubulin is
posttranslationally modified by acetylation on the epsilon-amino group of a lysine.
Biochemistry. 24:473-478.

Lee, G.S., Y. He, E.J. Dougherty, M. Jimenez-Movilla, M. Avella, S. Grullon, D.S. Sharlin, C.
Guo, J.A. Blackford, Jr., S. Awasthi, Z. Zhang, S.P. Armstrong, E.C. London, W.
Chen, J. Dean, and S.S. Simons, Jr. 2013. Disruption of Ttll5/stamp gene (tubulin
tyrosine ligase-like protein 5/SRC-1 and TIF2-associated modulatory protein gene) in
male mice causes sperm malformation and infertility. *J Biol Chem*. 288:15167-15180.

Lu, W., M. Lakonishok, A.S. Serpinskaya, D. Kirchenbüechler, S.C. Ling, and V.I. Gelfand.
2018. Ooplasmic flow cooperates with transport and anchorage in. *J Cell Biol*.
217:3497-3511.

Lu, W., M. Winding, M. Lakonishok, J. Wildonger, and V.I. Gelfand. 2016. Microtubule-
microtubule sliding by kinesin-1 is essential for normal cytoplasmic streaming in
Drosophila oocytes. *Proc Natl Acad Sci U S A*. 113:E4995-5004.

Magiera, M.M., and C. Janke. 2014. Post-translational modifications of tubulin. *Curr Biol*.
24:R351-354.

Mahalingan, K.K., E. Keith Keenan, M. Strickland, Y. Li, Y. Liu, H.L. Ball, M.E. Tanner, N.

Tjandra, and A. Roll-Mecak. 2020. Structural basis for polyglutamate chain initiation

and elongation by TTLL family enzymes. *Nat Struct Mol Biol*. 27:802-813.

McKenney, R.J., W. Huynh, R.D. Vale, and M. Sirajuddin. 2016. Tyrosination of α-tubulin

controls the initiation of processive dynein-dynactin motility. *Embo j*. 35:1175-1185.

Meijering, E., O. Dzyubachyk, and I. Smal. 2012. Methods for cell and particle tracking.

Methods Enzymol. 504:183-200.

Misgeld, T., M. Kerschensteiner, F.M. Bareyre, R.W. Burgess, and J.W. Lichtman. 2007.

Imaging axonal transport of mitochondria in vivo. *Nat Methods*. 4:559-561.

Pathak, D., K.J. Sepp, and P.J. Hollenbeck. 2010. Evidence that myosin activity opposes

microtubule-based axonal transport of mitochondria. *J Neurosci*. 30:8984-8992.

Pilling, A.D., D. Horiuchi, C.M. Lively, and W.M. Saxton. 2006. Kinesin-1 and Dynein are the

primary motors for fast transport of mitochondria in Drosophila motor axons. *Mol Biol*

Cell. 17:2057-2068.

Port, F., H.M. Chen, T. Lee, and S.L. Bullock. 2014. Optimized CRISPR/Cas tools for

efficient germline and somatic genome engineering in Drosophila. *Proc Natl Acad Sci*

U S A. 111:E2967-2976.

Preston, S.F., G.G. Deanin, R.K. Hanson, and M.W. Gordon. 1979. The phylogenetic

distribution of tubulin:tyrosine ligase. *J Mol Evol*. 13:233-244.

Prota, A.E., M.M. Magiera, M. Kuijpers, K. Bargsten, D. Frey, M. Wieser, R. Jaussi, C.C.

Hoogenraad, R.A. Kammerer, C. Janke, and M.O. Steinmetz. 2013. Structural basis

of tubulin tyrosination by tubulin tyrosine ligase. *J Cell Biol*. 200:259-270.

Redeker, V., N. Levilliers, J.M. Schmitter, J.P. Le Caer, J. Rossier, A. Adoutte, and M.H. Bré.

1994. Polyglycylation of tubulin: a posttranslational modification in axonemal

microtubules. *Science*. 266:1688-1691.

Redeker, V., N. Levilliers, E. Vinolo, J. Rossier, D. Jaillard, D. Burnette, J. Gaertig, and M.H.

Bré. 2005. Mutations of tubulin glycylation sites reveal cross-talk between the C

termini of alpha- and beta-tubulin and affect the ciliary matrix in Tetrahymena. *J Biol*

Chem. 280:596-606.

Redeker, V., R. Melki, D. Promé, J.P. Le Caer, and J. Rossier. 1992. Structure of tubulin C-

terminal domain obtained by subtilisin treatment. The major alpha and beta tubulin

isotypes from pig brain are glutamylated. *FEBS Lett*. 313:185-192.

Rüdiger, A.H., M. Rüdiger, J. Wehland, and K. Weber. 1999. Monoclonal antibody ID5:

epitope characterization and minimal requirements for the recognition of

polyglutamylated alpha- and beta-tubulin. *Eur J Cell Biol*. 78:15-20.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch,

C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K.

Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for

biological-image analysis. *Nat Methods*. 9:676-682.

Sirajuddin, M., L.M. Rice, and R.D. Vale. 2014. Regulation of microtubule motors by tubulin

isotypes and post-translational modifications. *Nat Cell Biol*. 16:335-344.

St Johnston, D., D. Beuchle, and C. Nüsslein-Volhard. 1991. Staufen, a gene required to

localize maternal RNAs in the Drosophila egg. *Cell*. 66:51-63.

Stone, M.C., F. Roegiers, and M.M. Rolls. 2008. Microtubules have opposite orientation in

axons and dendrites of Drosophila neurons. *Mol Biol Cell*. 19:4122-4129.

Suter, B. 2018. RNA localization and transport. *Biochim Biophys Acta Gene Regul Mech*.

1861:938-951.

Suter, B., and R. Steward. 1991. Requirement for phosphorylation and localization of the

Bicaudal-D protein in Drosophila oocyte differentiation. *Cell*. 67:917-926.

Theurkauf, W.E., H. Baum, J. Bo, and P.C. Wensink. 1986. Tissue-specific and constitutive

alpha-tubulin genes of Drosophila melanogaster code for structurally distinct proteins.

Proc Natl Acad Sci U S A. 83:8477-8481.

Vagnoni, A., and S.L. Bullock. 2016. A simple method for imaging axonal transport in aging

neurons using the adult Drosophila wing. *Nat Protoc*. 11:1711-1723.

Vallotton, P., and S. Olivier. 2013. Tri-track: free software for large-scale particle tracking.

Microsc Microanal. 19:451-460.

van Dijk, J., K. Rogowski, J. Miro, B. Lacroix, B. Eddé, and C. Janke. 2007. A targeted

multienzyme mechanism for selective microtubule polyglutamylation. *Mol Cell*.

26:437-448.

Wang, X., and T.L. Schwarz. 2009. The mechanism of Ca2+ -dependent regulation of

kinesin-mediated mitochondrial motility. *Cell*. 136:163-174.

Wang, X., D. Winter, G. Ashrafi, J. Schlehe, Y.L. Wong, D. Selkoe, S. Rice, J. Steen, M.J.

LaVoie, and T.L. Schwarz. 2011. PINK1 and Parkin target Miro for phosphorylation

and degradation to arrest mitochondrial motility. *Cell*. 147:893-906.

Weil, T.T. 2014. mRNA localization in the Drosophila germline. *RNA Biol*. 11:1010-1018.

Wloga, D., D.M. Webster, K. Rogowski, M.H. Bré, N. Levilliers, M. Jerka-Dziadosz, C. Janke,

S.T. Dougan, and J. Gaertig. 2009. TTLL3 Is a tubulin glycine ligase that regulates

the assembly of cilia. *Dev Cell*. 16:867-876.

Wolff, A., B. de Néchaud, D. Chillet, H. Mazarguil, E. Desbruyères, S. Audebert, B. Eddé, F.

Gros, and P. Denoulet. 1992. Distribution of glutamylated alpha and beta-tubulin in

mouse tissues using a specific monoclonal antibody, GT335. *Eur J Cell Biol*. 59:425-

432.

719 Zheng, Y., J. Wildonger, B. Ye, Y. Zhang, A. Kita, S.H. Younger, S. Zimmerman, L.Y. Jan,

720   and Y.N. Jan. 2008. Dynein is required for polarized dendritic transport and uniform

721   microtubule orientation in axons. *Nat Cell Biol*. 10:1172-1180.

722 Zimyanin, V.L., K. Belaya, J. Pecreaux, M.J. Gilchrist, A. Clark, I. Davis, and D. St Johnston.

723   2008. In vivo imaging of oskar mRNA transport reveals the mechanism of posterior

724   localization. *Cell*. 134:843-853.

726 **Table 1 Conservation between mammalian *TTLL5 and Drosophila TTLL5***
727 Residues labeled in red are critical residues for α-tubulin glutamylation by murine TTLL5 and are
728 conserved in human and *Drosophila* TTLL5 (van Dijk et al., 2007).

	K131 β6-7 loop (R188) R225 E366 N368
Mm TTLL5	TRKDR--KPVASSRGRGVY---VRLYVL---LLEVNLSP
Hs TTLL5	TRKDR--KPVASSRGRGVY---VRLYVL---LLEVNLSP
	K282 β6-7 loop (R339) R376 E517 N519
Dm TTLL5	TRKDR--KPAASSRGRG I F---LRVYVL---LLEI NLSP

731 **Table 2 Conservation between *TTL* and *TTLL5***
732 Many critical **r**esidues of human TTL (red) are conserved in *Drosophila,* mouse and human TTLL5 (Prota et al.,
733 2013; van Dijk et al., 2007).

	R202 R222 D318 E331 N333 P336
Hs TTL	IRSW--VLRTA-----FDFM--LI EVNGAPA
	R376 R398 D504 E517 N519 P522
Dm TTLL5	LRVY---I VRLA-----FDIL----LLEI NL SPS
	R225 R247 D353 E366 N368 P371
Mm TTLL5	VRLY---LARFA-----FDVL---LLEVNL SPS
Hs TTLL5	VRLY---LARFA-----FDVL---LLEVNL SPS

736 **Table 3 Transport parameters for mitochondria in the selected region of L1 veins.**

Genotype	*Control*		*TTLL5[E517A]*	
Direction	Plus-end	Minus-end	Plus-end	Minus-end
Number of wings (motile mito[1])	3 (20)	3 (11)	3 (8)	3 (2)
Motile Mito[1] (% total mito)	5.3±1.7	2.5±3.2	**1.5±1.3***	0.4±0.4
Total time (s)	92±45	100±43	98±48	64±37
Pausing (% total time)[2]	74±14	77±15	**63±13***	61±20
Length (µm)	7.2±4.5	6.7±1.6	**12.3±5.3***	7.5±2.8
Velocity (µm/s) [3]	0.37±0.08	0.38±0.14	0.36±0.08	0.36±0.16

[1] Motile mitochondria showed a run distance ≥ 2 µm.
[2] The pausing time ratio indicates the pausing time divided by the total duration of each motile mitochondria.
[3] Mean velocity corresponds to the mean values of instant trafficking velocity. Significant differences for
mitochondria of plus-end transport between control and *TTLL5[E517A]* is shown in bold face followed by * (p<0.05).
Only two mitochondria transported toward the MT minus-end were observed in *TTLL5[E517A]*. A comparison of the

minus-end transport between control and $TTLL5^{E517A}$ is not warranted, and these parts are shaded in grey to
indicate this.

 Figure legends

Fig. 1 Schematic representation of the *Drosophila TTLL5* gene, TTLL5, TTLL5^{MiEx}, and
Venus::TTLL5 protein. A) The *TTLL5* gene structure is based on the Flybase data for the
CG31108-RA transcript. The positions of the loss-of-function mutations are marked on the
TTLL5 gene. Grey triangles: transposon insertions. Grey asterisks: codons targeted for
generating InDels and point mutations. All *TTLL5* alleles were analyzed as hemizygous
animals over *Df(3R)BSC679*, which removes the region of the *TTLL5* gene. B) Western blot
from the soluble fraction of ovarian extracts. *w* (*TTLL5^{+}/$^{+}$),* which contains 2 wild-type copies
of *TTLL5*, expressed the highest levels of TTLL5. The TTLL5 signal is strongly reduced or
abolished in the null mutants *TTLL5^{pBac}/, TTLL5^{Mi}/* and *TTLL5^{MiEx}/* compared to *w*. The
hemizygous *TTLL5^{+}* ovaries (*TTLL5^{+}/*) expressed less TTLL5 than *w*. The signal of an
unspecific band (labeled with *), produced after a short exposure, was used as loading
control. In the following we will refer to the genotypes as *w* for the control (*TTLL5^{+}/$^{+}$*) and as
TTLL5^{allele} for the hemizygous mutants *(TTLL5^{allele}/ $^{-}$).*

Fig. 2 Mass spectrometry analysis of the glutamylation of the C-terminal domains of
the α-tubulin isoforms. A) In *Drosophila* ovaries**,** the C-terminal peptides of TBA1/3 can be
found modified by glutamic acids residue added to the side chains of E443, E445, E448, and
E449**.** B) The C-terminal peptides containing the full primary sequence were analyzed. The
structure of the different C-terminal tail peptides of TBA1/3 modified by glutamylation are
shown. Different combinations of Glu side chain modifications were found added at the
positions E443, E445, E448, and E449 of TBA1/3. C) Frequency of glutamylated C-terminal
peptides (containing the entire primary sequence) in *w* controls, *TTLL5^{pBac}* and *TTLL5^{MiEx}*
mutants containing modifications of 1E, 2E or 3Es. "Total Glu modification" shows the sum of
the 3 frequencies. The ratio was quantified based on the values of glutamylated C-terminal
TBA1/3 PSM (peptide spectrum match) divided by total PSM of the full primary C-terminal
TBA1/3 sequence.

Fig. 3 *TTLL5* is required for the glutamylation of the TBA1/3 isoforms of ovarian α-
tubulin. A) A') Sites on TBA1/3 recognized by the GT335 and 1D5 antibodies, respectively.
B) B') B'') The glutamylation signal produced by western blotting with the GT335 and 1D5

antibodies is lower in *TTLL5^{pBac}, TTLL5^{Mi}*, and *TTLL5^{MiEx}* than in the *w* control, but is restored

and even elevated in mutant ovaries that were rescued by *MattubGal4>Venus::TTLL5*. The

up-shifted bands recognized by 1D5 can be seen in the rescued *TTLL5^{pBac}* and *TTLL5^{Mi}*

mutants. The total tyrosinated α-tubulin levels seemed unaffected by the TTLL5 levels. B'')

The anti-βmono-E levels were similar between *w* control, *TTLL5^{MiEx}*, and rescue of *TTLL5^{MiEx}*.

α-tubulin and GAPDH served as loading controls. C) Stronger up-shifted bands were

observed upon *MattubGal4>Venus::TTLL5* overexpression in wild-type ovaries (anti-1D5

antibody). Total α-tubulin remained at similar levels in samples with excessive expression of

TTLL5. GAPDH was a loading control. D) The genotypes and rational of the *TTLL5* alleles

generated by Crispr/Cas9 are listed. The selected residues are predicted to be either

important for the glutamylation or tyrosination function based on the alignment shown in

Tables 1 and **2**. E) Western blotting showed that the polyglutamylation signal is absent in all

point mutants. *w* with 2 copies of *TTLL5^{+}* is the wild-type control and *TTLL5^{K282}* indel

(*TTLL5^{K282 indel3}*) that with only one copy of *TTLL5* still kept K282 is therefore an additional

control. Relative 1D5 level was observed slightly decreased in the hemizygous situation with

only one *TTLL5* copy. All *TTLL5* alleles were analyzed as hemizygous animals over

Df(3R)BSC679, which removes the region of the *TTLL5* gene. The genotypes for rescues

were *MattubGal4>UAS-Venus::TTLL5;TTLL5^{alleles}/Df(3R)BSC679.* The genotypes for

Venus::TTLL5 overexpression were *MattubGal4>UAS-Venus::TTLL5 ;*

Fig. 4 Effect of *TTLL5* on Staufen posterior localization refinement in oocytes.

A) A') Confocal micrographs showing stage 10B oocytes of *w* controls and *TTLL5^{pBac},*

TTLL5^{Mi}, and *TTLL5^{MiEx}* mutants. A'') Confocal micrographs showing stage 10B oocytes of

TTLL5^{pBac}, TTLL5^{Mi}, and *TTLL5^{MiEx}* mutants and their rescue with *Mattub4>UASP-*

Venus::TTLL5. Anti-Staufen is shown in green, anti-GFP in red, and Hochest in blue. A''')

Quantification of the Staufen crescent length along the posterior cortex based on A, A' and

A''. B) B') Confocal micrographs showing stage 10B oocytes of *w* controls, *TTLL5^{E517A},*

TTLL5^{P522A}, and *TTLL5^{R339A}* mutants. B'') Quantification of Staufen's crescent length along

the posterior cortex based on B and B'', respectively. C) C') Confocal micrographs showing

previtellogenic egg chambers of *w* controls, *TTLL5^{E517A}, TTLL5^{P522A},* and *TTLL5^{R339A}* mutants.

anti-BicD is shown in red, anti-Staufen in green, and Hochest in blue. D) The BicD levels

remained unchanged between *w* control, *TTLL5^{MiEx}*, and *TTLL5^{MiEx}* rescued by

Mattub4>Venus::TTLL5. GAPDH served as loading control. *** p<0.005; ** <0.01. Scale Bar:

25 μm. All *TTLL5* alleles were hemizygous over *Df(3R)BSC679*. The genotypes for rescued

animals were *MattubGal4/UAS-Venus::TTLL5;TTLL5 ^{alleles}/Df(3R)BSC679.*

Fig. 5 Role of *TTLL5* in ooplasmic streaming. A) KHC levels remained unchanged
between *w* control, *TTLL5^{MiEx}*, and *TTLL5^{MiEx}* rescued by *Mattub4>Venus::TTLL5.* GAPDH
served as loading control. B) Example picture of a stage 10B oocyte used for particle flow
measurements based on time-lapse movies of ooplasmic streaming (Mov.S1-S4). Scale Bar:
25 μm. C) 100-200 particles were selected along the cortex line from two sides of each
ooplasm to determine the mean velocities of the particles from the different genotypes. One
stage 10B oocyte was chosen randomly for velocity quantification for each fly. The fly
numbers for *w* control, *TTLL5^{MiEx}* mutant, *TTLL5^{E517A}* mutant, and *UASP-Venus::TTLL5*
transgene rescuing the *TTLL5^{MiEx}* mutant were 7, 9, 5, and 5, respectively. No significant
changes of particle velocity were found between different genotypes. D) Representative
kymographs of particles in oocytes of a *w* control, a *TTLL5^{MiEx}* mutant, the *TTLL5^{E517A}* mutant
and the *TTLL5^{MiEx}* mutant rescued by overexpression of the *UASP-Venus::TTLL5* transgene.
Kymographs were generated along a line (the yellow line in B) close to the cortex on each
side of the ooplasm from 3 min time-lapse movies. Representative kymographs showed that
the lines generated along the cortex were parallel to the particle flow in both control and
Venus::TTLL5 rescue. The moving particles revealed different flow patterns in the case
when TTLL5 activity was low. Genotypes for *TTLL5^{MiEx}* and *TTLL5^{E517A}* were
TTLL5^{MiEx}/Df(3R)BSC679 and *TTLL5^{E517A}/Df(3R)BSC679*, respectively. The genotype of the
rescued flies was *MattubGal4/UAS-Venus::TTLL5; TTLL5^{MiEx}/Df(3R)BSC679.*

Fig. 6 Effect of *TTLL5* on mitochondria transport in the wing nerve. A) In head and wing
extracts, the relative glutamylation signal of 1D5 is lower in *TTLL5^{E517A}* (genotype: *ApplGal4;
UAS-Mito::GFP; TTLL5^{E517A}/Df(3R)BSC679*) than in the control (genotype: *ApplGal4;UAS-
Mito::GFP;+/Df(3R)BSC679*). B) Cartoon of the *Drosophila* wing showing the L1 and L3 vein
and the neuron rich regions analyzed for tracking mitochondria (red and green boxes). C)
Representative still image of GFP-labelled mitochondria in axons of the L1 and L3 vein (from
Mov. S5 and **Mov. S6**, respectively) of both control and *TTLL5^{E517A}* wings. Transported
mitochondria (run distance ≥ 2 μm) were manually labeled with the Fiji MTrackJ plugin.
Some circles only mark a mitochondrial signal that appeared in another time frame. Scale
Bar: 10 μm. D) Quantification of the fraction of transported mitochondria. Note that in control
wings a much higher fraction of mitochondria is transported in the L3 vein than in L1. The
results from L1 show a dramatic decrease of motile mitochondria in the *TTLL5^{E517A}* mutant.
In the L3 vein, no transported mitochondria were found in the *TTLL5^{E517A}* mutant. The total

particles (indicated by "n") with the diameter from 0.1μm-1.6μm were analyzed by Fiji from

the signals of representative still images from three wings for each genotype. The motile

mitochondria ratio indicates the value of transported mitochondria divided by total

mitochondria particles. E) Scheme of the transport behavior of the mitochondria in the wing

neuron. Start of a run was either considered the time when a motile particle began to move

from a stationary position or entered into the focal plane. Particles were defined as

"trafficking" when their instant velocity was ≥ 0.2 μm/s and they were considered as "pausing"

when they moved at less than 0.2 μm/s, but later resumed their movement. As stop of a run

we considered particles that entirely terminated or moved out of the focal plane. For each

run, the total run distance of more than 2 μm was considered as "transported or motile

mitochondria". *** $p<0.005$; * <0.05.

Fig. 1

A

B

Fig. 2

A

α-tubulin isoforms in *Drosophila* ovaries

E-residues were found added at the TBA1/3 positions: E443, E445, E448 and E449.

B

Glutamylation on Glu residues in the C-terminal domain of TBA1/3

```
GEGEGAEEY      GEGEGAEEY      GEGEGAEEY      GEGEGAEEY
    |              |              |              |
    E              E              E              E

GEGEGAEEY      GEGEGAEEY      GEGEGAEEY      GEGEGAEEY
    |              |              |              |
    EE             EE             EE             EE

GEGEGAEEY      GEGEGAEEY      GEGEGAEEY      GEGEGAEEY
    |              |              |              |
    EEE            EEE            EEE            EEE

GEGEGAEEY      GEGEGAEEY      GEGEGAEEY      GEGEGAEEY
    |  |           |  |           |  |           |  |
    E  E           E  EE          E    E         E   EE

GEGEGAEEY      GEGEGAEEY
      |  |        |  |  |
      EEE         E  E  E
```

C

Role of *TTLL5* for the Glutamylation of the C-terminal domain of TBA1/3			
	w (n=56)	***TTLL5*** [pBac] (n=29) (*%E mod*)	***TTLL5*** [MiEx] (n=7)
+1	41	0	0
+2	14	0	0
+3	7	0	0
Total Glu modification	**62**	**0**	**0**

Fig. 3

A

443　445　　448　449

G D G E G E G A E E Y

anti-GT335　　anti-GT335

A'

443　445　　448　449

anti-1D5

G D G E G E G A E E Y

anti-1D5　anti-1D5　anti-1D5　anti-1D5

B

w (control) │ TTLL5^pBac │ Venus::TTLL5;TTLL5^pBac │ w (control) │ TTLL5^Mi │ Venus::TTLL5;TTLL5^Mi │ w (control) │ TTLL5^MiEx │ Venus::TTLL5;TTLL5^MiEx

130- ── GFP

55- ── GT335

55- ── tyr-α-tubulin

35- ── GAPDH

B'

w (control) │ TTLL5^pBac │ Venus::TTLL5;TTLL5^pBac │ w (control) │ TTLL5^Mi │ Venus::TTLL5;TTLL5^Mi

130- ── GFP

55- ── 1D5

55- ── α-tubulin

35- ── GAPDH

B''

w (control) │ TTLL5^MiEx │ Venus::TTLL5;TTLL5^MiEx

130- ── GFP

55- ── 1D5

55- ── β-monoE

55- ── α-tubulin

35- ── GAPDH

C

w (control) │ Venus::TTLL5 o.e.

130- ── GFP

55- ── 1D5

55- ── α-tubulin

55- ── tyr-α-tubulin

35- ── GAPDH

D

Residues of *Drosophila TTLL5*	Function	Corresponding residues to mammalian *TTLL5* or *TTL*	Mutations
E517	ATP/ADP binding	*MmE336 of TTLL5 HsE331 of TTL*	*E517 indel* : WLLE517 → S E517A
K282	ATP/ADP binding	*MmK131 of TTLL5*	*K282 indel 1* : K282 → stop codon *K282 indel 2* : TRKD283 → N *K282 indel 3* : TRK282 → KGK282
R339	(In β6-7 loop) Stabilization tubulin with TTLL5 active site	*MmR188 of TTLL5*	R339A
P522	(Conserved residue of human TTL) Anchor enzyme to α-tubulin	*HsP336 of TTL*	P522A

E

w (positive control) │ TTLL5^E517 indel │ TTLL5^E517A │ TTLL5^K282 indel 1 │ TTLL5^K282 indel 2 │ TTLL5^R339A │ TTLL5^P522A │ TTLL5^K282 indel 3 (positive control)

ATP/ADP │ ATP/ADP │ Glu │ Tyr

55- ── 1D5

55- ── tyr-α-tubulin

35- ── GAPDH

Fig. 4

A

w (control)

12 µm

DNA   Staufen   GFP

A'

TTLL5^pBac   *TTLL5^MiEx*   *TTLL5^Mi*

12 µm   12 µm   12 µm

A''

Venus::TTLL5

DNA   Staufen   GFP

A'''

Cresent length (µm)

n: 50 19 17 26 24 20 20

w (control)
TTLL5^pBac
Venus::TTLL5;TTLL5^pBac
TTLL5^Mi
Venus::TTLL5;TTLL5^Mi
TTLL5^MiEx
Venus::TTLL5;TTLL5^MiEx

B

S10B

w (control)

12 µm

DNA   Staufen

B'

S10B

TTLL5^E517A

TTLL5^P522A

TTLL5^R339A

12 µm

DNA   Staufen

B''

Cresent length (µm)

n: 20   21   15   16

w (control)
TTLL5^E517A
TTLL5^P522A
TTLL5^R339A

C

Previtellogenic stages

w (control)

BicD   Staufen

C'

Previtellogenic stages

TTLL5^E517A

TTLL5^P522A

TTLL5^R339A

BicD   Staufen

D

w (control)   *TTLL5^MiEx*   *Venus::TTLL5;TTLL5^MiEx*

130-   GFP

97-   BicD

35-   GAPDH

Fig. 5

Fig. 6

Fig. S1

A) Example_Es on *w* TBA1/3: GEGEGAEEY
EE

PEAKS

Easyprot

B) Example_*w* TBA4: DEDFDEF

MSfragger

Fig. S2

Fig. S3

A

α-tubulin isoforms in *Drosophila* ovaries

G-residues were found added at the TBA1/3 positions E443, E445, E448 and E449.

B

Glycylation on Glu residues in the C-terminal domain of TBA1/3

C

Role of *TTLL5* for the Glycylation of the C-terminal domain of TBA1/3

	w (n=56)	*TTLL5* [pBac] (n=29)	*TTLL5* [MiEx] (n=7)
		(%G mod)	
+1	3.5	10	14
+2	3.5	0	0
Total Gly modification	**7**	**10**	**14**

Fig. S4

w (Control)

Venus::TTLL5 o.e.

130- GFP

130-
97- KHC

97- BicD

55- α-tubulin

Fig. S5

Appl-Gal4>Mito::GFP

Control

L1

10 μm

L3

10 μm

➡ Thorax

Supplementary Information

S-Tables

Table S1A: primers for cloning the sequences encoding the sgRNAs into the *Drosophila* transformation vector pCFD5

for guide 2	TTLL5 R339 top	$^{5'}$TGC AGC GGC ATC CAG TCG AGG AAG
	TTLL5 R339 rev	$^{5'}$ AAA CCT TCC TCG ACT GGA TGC CGC
for guide 3	TTLL5 K282 top	$^{5'}$ TGC ATC CTA CGA AAT GAC ACG AA
	TTLL5 K282 rev	$^{5'}$ AAA CTT CGT GTC ATT TCG TAG GA
for guide 4	TTLL5 E517 top	$^{5'}$ TGC AAG ATT GAT TTC CAG TAG CCA
	TTLL5 E517 rev	$^{5'}$ AAA CTG GCT ACT GGA AAT CAA TCT
for guide 5	TTLL5 R376 top	$^{5'}$ TGC ATG GTC ACA AAT GTG ATT TGA
	TTLL5 R376 rev	$^{5'}$ AAA CTC AAA TCA CAT TTG TGA CCA

Table S1B: templates to introduce point mutations

TTLL5^{R339A}
= Arg (AGG) -> Ala (**GCT**) on the reverse strand; introducing an AluI site (AGCT) at the same time; PAM (underlined) on the other strand (5'TGG) $^{5'}$GAATTATAGCAAGTTGTTGGTTTAGTTACCGAATTAACAATAAATATT<u>CC**AGC**</u>T'CCTCGACTGGATGCCGCTGGCTTAACTATCCATGGTCCGCGATGCTTGTT
TTLL5^{E517A}
= Glu (GAA) -> Ala (**GCC**) on the reverse strand; introducing a HaeIII site (GGCC) at the same time; PAM (underlined) on this strand (5'TGG) $^{5'}$ACCTTAGTGTCCAAAGGACTATCCACACCCATTGACGGCGATAGATTGAT**GGC**CAGTAG'CCA<u>TGG</u>CTTAAGTGCGTTGTCTATTAATATATCGAATCCATACAGCTCAA
TTLL5^{P522A}
= Pro (CCG) -> Ala (GCG) on the reverse strand; introducing a BsaHI site (GRCGYC) at the same time; PAM (underlined) on this strand (5'TGG); grey nucleotide changes that are silent mutations to prevent multiple recognition after DNA was cut $^{5'}$GTCTGCCATCAGACATGACTTAACCTTAGTGTCCAAAGGACTATCCACACCCATTGA**CGC**CGATAGATTGATTTC**G**AG**A**AG'CCA<u>TGG</u>CTTAAGTGCGTTGTCTATTAATATATCGAATCCATACAGCTCAA

Table S2: Tissue specific expression of the 11 *Drosophila TTLL* genes

TTLL1A	testis	imaginal disc			
TTLL1B	testis	imaginal disc			
TTLL3A					
TTLL3B	testis	imaginal disc	accessory gland	central nervous system	
TTLL4A	testis	imaginal disc	ovary	central nervous system	
TTLL4B	testis	imaginal disc	accessory gland	fat body	
TTLL5	ovary	imaginal disc	testis	accessory gland	central nervous system
TTLL6A	testis	imaginal disc	accessory gland		
TTLL6B	testis	imaginal disc	accessory gland	fat body	
TTLL12	central nervous system	carcass	ovary/head/ imaginal disc	fat body	
TTLL15					

Table S3 Number of peptides of TBA1/3/4 found by MS

Isoform	*w*	*TTLL5^{pBac}*	*TTLL5^{MiEX}*
		(PSM value)	

TBA1/3	56	29	7
TBA4	12	12	4

Supplementary Information

Legends for **Fig. S1-S5 & Mov. S1-S5**

Fig. S1: Example mass spectrum for C-terminal peptides. A) Example of a mass spectrum of Glutamylation of the C-terminal domain of the TBA1/3 isoform by PEAKS and Easyprot search engines. B) Example mass spectrum of the C-terminal domain of the TBA4 isoform by the MSfragger search engine.

Fig. S2: TTLL5 was stably expressed in all Crispr/Cas9 generated *TTLL5* mutants. TR**K**D283N refers to a more complex mutation in which the codons TR**K**D283 were replaced by a N. All Crispr/Cas9 generated *TTLL5* mutations were analyzed hemizygously over *Df(3R)BSC679.* We, therefore, expect extracts from these animals to contain at most half the amount of TTLL5*. *w* (*TTLL5$^+$/TTLL5$^+$*) was the positive control and the null mutation *TTLL5^{MiEx}/Df(3R)BSC679* was the negative control. The signal of an unspecific band (labeled with *) under short exposure was treated as a loading control. *w,* which contains 2 copies of the *TTLL5* allele expressed highest levels of TTLL5. Crispr/Cas9 generated *TTLL5* mutations with a single allele of *TTLL5* expressed less TTLL5 compared to *w,* but more compared to the null mutation *TTLL5^{MiEx}/Df(3R)BSC679*. The TTLL5 levels of the hemizygous point mutants were similar to the ones of the hemizygous wild type (refer to **Fig 1B**). The faint band seen in *TTLL5^{MiEx}/Df(3R)BSC679* is probably a background band.

Fig. S3: Mass spectrometry analysis of the glycylation of the C-terminal end of the α-tubulin isoforms. A) The following Glu residues in the C-terminal peptides were found to be modified in *Drosophila* ovaries. E443, E445, E448, and E449 of TBA 1/3 were found modified by the addition of 1-2 glycines. B) The C-terminal peptides containing the full primary sequence were analyzed. The C-terminal tail of the TBA1/3 peptides with the Glycylation modifications found by MS are shown. Gly side chains were found added to E443, E445, E448, and E449 of TBA 1/3. C) The Glycylation frequency for the C-terminal peptides of *w* controls, *TTLL5^{pBac},* and *TTLL5^{MiEx}* mutants is shown. +1=frequency of peptides contained 1E modification. +2=frequency of peptides contained 2E modifications. Total Gly modification shows the sum of the 1E and 2E frequencies. The frequency was quantified based on the values of glycylated C-terminal TBA1/3 PSM (peptide spectrum match) divided by total PSM of the full primary C-terminal TBA1/3 sequence.

Fig. S4: *TTLL5* overexpression in ovaries neither alters Kinesin-1 heavy chain nor BicD levels. The KHC and BicD levels kept unchanged between *w* control and *Venus::TTLL5* overexpressing ovaries. α-tubulin was used as loading control.

Fig. S5: Still picture of Mito::GFP movie showing a selected region of L1 and L3 neurons in control wing nerves. Yellow triangle: cell bodies. Yellow bracket: axons.

Mov. S1 Ooplasmic streaming in *w* (control; 2 ooplasms)

Mov. S2 Ooplasmic streaming in *TTLL5^MiEx^* (2 ooplasms)

Mov. S3 Ooplasmic streaming in *TTLL5^E517A^* (2 ooplasms)

Mov. S4 Ooplasmic streaming in *Venus::TTLL5; TTLL5^MiEx^* (2 ooplasms)

Mov. S5 Appl-Gal4>Mito::GFP trafficking in the L1 vein. Top movie: control; bottom movie: *TTLL5^E517A^*.

Mov. S6 Appl-Gal4>Mito::GFP trafficking in the L3 vein. Top movie: control; bottom movie: *TTLL5^E517A^*.