1

Haploidy-linked cell proliferation defects limit larval growth in Zebrafish

Kan Yaguchi[1,2], Daiki Saito[1], Triveni Menon[3], Akira Matsura[1], Takeomi Mizutani[4],
Tomoya Kotani[5], Sreelaja Nair[6], and Ryota Uehara[1,2]

[1]Graduate School of Life Science, Hokkaido University, Kita 21, Nishi 11, Kita-Ku,
Sapporo, 001-0021, Japan
[2]Faculty of Advanced Life Science, Hokkaido University, Kita 21, Nishi 11, Kita-Ku,
Sapporo, 001-0021, Japan
[3]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, Virginia, USA
[4]Department of Life Science and Technology, Faculty of Engineering, Hokkai-Gakuen
University, Minami 26, Nishi 11, Chuo-ku, Sapporo, 064-0926, Japan
[5]Department of Biological Sciences, Faculty of Science, Hokkaido University, Kita 10,
Nishi 8, Kita-Ku, Sapporo 060-0810, Japan
[6]Department of Biosciences and Bioengineering, Indian Institute of Technology Bombay,
Powai, Mumbai, 400076, India

Corresponding authors:

Sreelaja Nair, Department of Biosciences and Bioengineering, Indian Institute of
Technology Bombay, Powai, Mumbai, 400076, India
E-mail: sreelaja@iitb.ac.in
Phone: 91-22-25766758

Ryota Uehara, Faculty of Advanced Life Science, Hokkaido University, Kita 21, Nishi 11,
Kita-Ku, Sapporo 001-0021, Japan
E-mail: ruehara@sci.hokudai.ac.jp
Phone: 81-11-706-9238

Keywords: Ploidy, Centrosome, Zebrafish

Abstract

Haploid larvae in non-mammalian vertebrates are lethal with characteristic organ growth retardation collectively called "haploid syndrome." In contrast to mammals whose haploid intolerance is attributed to imprinting misregulation, the cellular principle of haploidy-linked defects in non-mammalian vertebrates remains unknown. Here, we investigated cellular defects that disrupt the ontogeny of gynogenetic haploid zebrafish larvae. Unlike diploid control, haploid larvae manifested unscheduled cell death at the organogenesis stage, attributed to haploidy-linked p53 upregulation. Moreover, we found that haploid larvae specifically suffered the gradual aggravation of mitotic spindle monopolarization during 1-3 days post fertilization, causing spindle assembly checkpoint-mediated mitotic arrest throughout the entire body. High-resolution imaging revealed that this mitotic defect accompanied the haploidy-linked centrosome loss occurring concomitantly with the gradual decrease in larval cell size. Either resolution of mitotic arrest or depletion of p53 significantly improved organ growth in haploid larvae. Based on these results, we propose that haploidy-linked mitotic defects and cell death are critical cellular causes that limit the larval growth in the haploid state, potentially placing an evolutionary constraint on allowable ploidy status in the non-mammalian vertebrate life cycle.

Introduction

Compared to plants or fungi, the life cycle in animals is restricted regarding ploidy. In plant and fungal life cycles, cells undergo mitotic divisions and develop multicellular bodies in the haploid and diploid generations (Mable and Otto, 1998). In contrast, the multicellular stage in animals is restricted to the diploid generation, except in some orders of haplodiplontic invertebrates (Mable and Otto, 1998; Otto and Jarne, 2001). The ploidy restriction is especially stringent in vertebrates: Haploid embryos generated through egg activation without the contribution of one parent's genome (e.g., parthenogenesis, gynogenesis, or androgenesis) are almost invariably lethal in vertebrates (Wutz, 2014).

In mammals, haploid embryonic lethality is mainly attributed to misregulation of imprinted genes. Differential expression of parental alleles of imprinted genes disables the development of mammalian uniparental haploid embryos beyond the preimplantation blastocyst stage (Leeb and Wutz, 2013; Tilghman, 1999). On the contrary, non-mammalian vertebrates are devoid of parent-specific genomic imprinting and, hence, free from imprinting-associated developmental defects. However, despite the lack of the detrimental feature of imprinting misregulation, haploid larvae are lethal even in non-mammalian vertebrates.

Compared to mammals, ontogenetic defects become evident at later developmental stages

in haploid non-mammalian vertebrates. For example, haploid larvae in zebrafish undergo

gastrulation and somite formation and initiate organogenesis by 24 hours post fertilization

(hpf) with only a modest delay from diploid counterparts (Menon et al., 2020). Haploid

larvae in other fish and amphibians also reach the organogenesis stage (Fankhauser, 1938;

Hamilton, 1966; Uwa, 1965). However, after the onset of organogenesis, haploid larvae in

these non-mammalian species manifest pleiotropic developmental defects and succumb to

lethality after hatching. For example, haploid embryos and larvae suffer abnormalities in

collective cell migration during gastrulation (Menon et al., 2020), water inflow control

through ectoderm in gastrulae (Hamilton and Tuft, 1972), prechordal plate function (Araki

et al., 2001), epidermal cell arrangements during post-gastrula stages (Ellinger, 1979;

Ellinger and Murphy, 1980), melanophore pigmentation (Uwa, 1965), and blood circulation

(Uwa, 1965). These developmental abnormalities have been collectively called "haploid

syndrome" (Dasgupta and Matsumoto, 1972; Hamilton, 1966; Luo and Li, 2003).

In the complex profile of haploid syndrome, a particularly common abnormality across

non-mammalian vertebrate species is severe growth retardation of organs, such as the brain

and eyes (Dasgupta and Matsumoto, 1972; Fankhauser and Griffiths, 1939; Hertwig, 1911;

Oppermann, 1913; Purdom, 1969; Uwa, 1965). Forced diploidization of haploid larvae by

artificial induction of whole-genome duplication in the early cleavage stages resolves organ

retardations, suggesting that the haploid state per se, rather than loss of heterozygosity of

deleterious recessive alleles, causes these defects (Menon and Nair, 2018; Nagy et al.,

1978; Streisinger et al., 1981; Subtelny, 1958). Though previous studies identify global

developmental errors associated with haploidy, the specific cell biological triggers of

haploid organ retardations remain largely unknown. Elucidation of a fundamental cellular

cause of the haploidy-linked organ retardations in non-mammalian vertebrates would

provide a profound insight into evolutionary constraints that limit the multicellular stage of

vertebrate life cycles exclusively to the diploid generation.

In this study, we investigated cellular defects in haploid zebrafish during the larval stages,

when morphological abnormalities became evident (Kroeger et al., 2014). Haploid larvae

suffered unscheduled cell death associated with p53 upregulation. High-resolution imaging

revealed that, as cell size gradually decreased after 12 hpf, drastic centrosome loss occurred

almost exclusively in haploid cells below a certain size threshold. This cell size-linked

centrosome loss led to the gradual accumulation of mitotic defects in haploid larvae, which

coincided with the characteristic temporal pattern of organ growth retardation at the

developmental stage. Alleviating mitotic arrest or artificially reducing p53 levels

significantly improved organ growth, indicating that haploidy-linked poor organ growth

stems at least in part from these cellular defects. Our results revealed haploidy-linked

incoordination of cell proliferation, whose progression pattern may shape the characteristic

profile of "haploid syndrome" in non-mammalian vertebrates.

Results

A drastic increase in unscheduled apoptosis in haploid larvae

We investigated the developmental processes of haploid and diploid zebrafish larvae

generated by *in vitro* fertilization using UV-irradiated and non-irradiated spermatozoa,

respectively (Fig. S1). At 3.5 days post fertilization (dpf), by when most organs have

formed, haploid larvae manifested typical "haploid syndrome" defects, such as curled, short

body axis and reduced brain and eye sizes compared with diploid counterparts (Fig. 1A and

B) (Kroeger et al., 2014; Luo and Li, 2003; Purdom, 1969; Uwa, 1965). The morphological

defects of haploid syndrome occurred in varying phenotypic grades, with some larvae

manifesting severe defects and others with milder defects within a clutch (Fig. 1A).

Consistent with this, we observed a larger variance of body and organ size distributions in

haploid larval groups than in diploids (Fig. 1B). DNA content analysis using flow

cytometry showed that the 1C and 2C populations (possessing one and two genome copies,

respectively) predominated in the haploid larval groups at 3 and 5 dpf, confirming that cells

in haploid zebrafish larvae retained the haploid state throughout the time duration (Fig. S1).

This is in sharp contrast to haploid mammalian embryonic cells that quickly convert to

diploids during early embryogenesis (Leeb and Wutz, 2011; Sagi et al., 2016).

The drastically poor organ growth in haploid larvae indicated a possibility that haploid cells

suffered defects in cell proliferation. To test this idea, we visualized apoptotic and mitotic

cells by immunostaining cleaved caspase-3 and phospho-histone H3 (pH3), respectively

(Mendieta-Serrano et al., 2013; Sorrells et al., 2013), in 3 dpf haploid and diploid larvae

(Fig. 2A). In diploid larvae, cleaved caspase-3-positive cells were infrequent and found at

specific sites, such as the optic tectum, mid-brain, and the inner retinal layers (Fig. 2A, B,

and S2), reflecting programmed apoptosis underlying tissue organization (Yamashita,

2003). On the other hand, in haploid larvae, cleaved caspase-3-positive cells were

frequently detected throughout the whole body (Fig. 2A, B, and S2). We also observed

clusters of apoptotic cells in the retinal periphery and brain ventricles in haploid larvae (Fig.

2A and S2, see also Fig. 2B). Moreover, haploid larvae showed a drastic increase in pH3-

positive mitotic cells compared to diploids throughout the larval body. These results

demonstrate that haploid larvae suffer irregularly patterned apoptosis and cell division

throughout the body, potentially impairing organ growth during development. Therefore,

we decided to address the molecular and cellular bases of these haploidy-linked defects in

cell survival and proliferation.

Irregular p53 upregulation limits organ growth in haploid larvae

We next sought to determine the cause of the haploidy-linked unscheduled apoptosis. As a

candidate apoptosis inducer, we investigated expression levels of p53 protein in haploid

and diploid larvae at 3 dpf by immunoblotting (Fig. 3A). p53 expression was significantly

higher in haploid larvae than in diploids (Fig. 3A and B). p53 level in haploids was even

higher than that in diploid larvae irradiated with UV to induce DNA damage-associated p53

upregulation (Fig. 3B).

To investigate the causality between irregular p53 upregulation and haploidy-linked defects,

we next tested the effects of morpholino-mediated p53 depletion on apoptosis in haploid

larvae. p53 expression was substantially reduced by a p53-specific morpholino (Fig. 3C and

D). Cleaved caspase-3-positive cells were less frequently observed in haploid p53

morphants than in control haploid larvae injected with a 4-base mismatch morpholino or

uninjected haploid larvae (Fig. 3E, F, and S3A; see also Fig. 2). Interestingly, the apoptotic

cell clusters frequently observed in haploid larvae were absent in haploid p53 morphants

(Fig. 3F and S3A). DNA content in haploid p53 morphants was equivalent to that in control

haploids, demonstrating that the ploidy level did not change upon p53 depletion in haploid

larvae (Fig. S3B). These results suggest that p53 upregulation contributes to enhanced

apoptosis in haploid larvae.

We also tested the effect of p53 depletion on organ size in haploid larvae (Fig. 3G). p53

depletion significantly increased body-axis length and brain width assessed at the mid-brain

in haploid larvae (Fig. 3G and H). p53 depletion notably eliminated the population of

haploid larvae that suffered particularly severe organ size reduction, suggesting efficient

alleviation of the haploidy-linked poor organ growth (Fig. 3H). The above data indicate

p53-induced apoptosis as a cause of haploidy-linked organ growth retardation.

Haploidy larvae suffer severe mitotic arrest

Next, we sought to understand the reason for the irregular cell proliferation in haploid

larvae and its possible influence on developmental defects in the haploid state. To

quantitively analyze the haploidy-linked change in cell proliferation pattern, we conducted

flow cytometry for DNA content and pH3-positive cell proportion in 1-3 dpf haploid and

diploid whole-larval cells (Fig. 4A and B). In diploid larvae, the proportion of pH3-positive

cells was 1.6 % at 1 dpf, which decreased to 0.4% at 3 dpf (Fig. 4A and B), possibly

reflecting the transition from proliferative to postmitotic state in different cell lineages

during larval stages (Li et al., 2000; Sugiyama et al., 2009). On the other hand, the

proportion of pH3-positive cells in haploid larvae was 1.6% at 1 dpf and 1.2% at 3 dpf (Fig.

4A and B). The mitotic index was significantly higher in haploid larvae than in diploids at

2-3 dpf (Fig. 4B), consistent with the finding in the whole-mount larval imaging (Fig. 2A).

To address the cause of the mitotic index increase in haploids, we conducted live imaging

and analyzed the mitotic progression of endothelial cells in 1.5-3 dpf haploid and diploid

larvae stably expressing histone-H2B-mCherry (Tg(fli1:h2b-mCherry)/ncv31Tg) (Yokota

et al., 2015). In diploid larvae, almost all cells that entered mitosis (marked by nuclear

envelope breakdown (NEBD)) completed chromosome alignment and segregation within

30 min after mitotic entry (Fig. 4C and D). On the other hand, 15% of haploid cells spent >

60 minutes in the mitotic phase, revealing a severe mitotic delay in haploid larvae. Such

mitotically delayed haploid cells often underwent mitotic death or mitotic slippage (exiting

mitosis without proper chromosome segregation) (Fig. 4C and E). Mitotic progression was

slower in haploid larvae than in diploids for the duration of assessment from 1.5-3 dpf, with

more frequent mitotic defects later in the observation (Fig. S4A, B, and D). We also

observed severe chromosome misalignment in haploid cells delayed in mitosis (Fig. 4C),

suggesting that the increased mitotic index and mitotic delay seen in haploid larvae could

be due to the activation of the spindle assembly checkpoint (SAC).

Gradual aggravation of mitotic spindle monopolarization with centriole loss in

haploid larvae during 1-3 dpf

To understand the cause of the frequent mitotic arrest with chromosome misalignment in

haploid larvae, we investigated mitotic spindle organization by immunostaining of α-

tubulin and centrin, which mark microtubules and the centrioles, respectively, in the head

region of haploid and diploid larvae from 0.5 to 3 dpf (Fig. 5 and S5). Almost all mitotic

cells in diploids possessed bipolar spindles with the normal number of 4 centrioles in all

stages tested. At 0.5 dpf, haploid embryonic cells also possessed bipolar spindles with 4

centrioles (Fig. 5A-C, and S5A-D). However, as development progressed, cells in haploid

larvae had monopolar spindles with reduced centriole number and severe chromosome

misalignment (Fig. 5B and C). In the head region of haploid larvae, including eyes, brain,

skin, and olfactory organ, 14.5% of mitotic cells had monopolar spindles at 1 dpf, which

increased to 32.6% and 61.5% at 2 and 3 dpf, respectively (Fig. 5B). The timing of the

gradual aggravation of spindle monopolarization and centrosome loss corresponded well

with the time when mitotic index became significantly higher in haploids than in diploids

with increased cell death (Fig. 4A, B and S4D). The trajectory of centriole loss and spindle

monopolarization was organ-specific: these defects commenced earlier in the eyes and

brain than in the skin (Fig. S5A-D). However, the frequency of monopolar mitotic cells

with centriole loss reached over 50% in all these organs by 3 dpf, revealing the general

nature of the cellular defects in haploid larvae.

The developmental stage when haploid larvae suffered the gradual aggravation of

centrosome loss corresponded to the stage when larval cell size gradually decreased

through successive cell divisions. Therefore, we tested the relationship between cell size

and centriole number in mitotic cells in subepidermal cell layers in the head region of 0.5-3

dpf haploid or diploid larvae (Fig. 5D and E; *see Material and methods*). During 0.5-3 dpf,

the largest slice area of mitotic cells reduced from 43 μm^2 to 16 μm^2 or from 61 μm^2 to 29

μm^2 in haploids or diploids, respectively (Fig. 5D). Interestingly, monopolar spindles with

reduced number of the centrioles were observed almost exclusively in the cells whose

largest slice area was less than 35 μm^2, suggesting a lower limit of cell size for supporting

centrosome number homeostasis in haploid larvae (Fig. 5D and E). Below the limit,

frequency of monopolar spindles monotonically increased as cell size reduced in haploids

(Fig. 5E). In contrast, diploid larval cells whose size became lower than this limit still

formed bipolar spindles. The above data indicates that the stage-specific aggravation of

mitotic spindle monopolarization through the haploidy-specific centrosome loss results in

severe mitotic arrest in haploid larvae.

Resolution of SAC-dependent mitotic arrest improves larval growth in the haploid state

Finally, we sought to address the causality between haploidy-linked mitotic arrest and larvae growth defects. For this, we attempted to resolve the haploidy-linked mitotic arrest by suppressing SAC using reversine, an inhibitor of Mps1 kinase required for SAC activation (Santaguida et al., 2010). In live imaging of histone-H2B-mCherry, 97% of reversine-treated haploid larval cells underwent chromosome segregation or exited mitosis within 30 min after NEBD (Fig. 6A, B, and S4C). Notably, mitotic cell death was substantially suppressed by reversine (Fig. 6C, see also Fig. 4E). Therefore, the SAC inactivation resolved the severe mitotic delay and mitigated mitotic cell death in haploid larvae.

We next tested whether SAC inactivation resolved the abnormal cell proliferation pattern in haploid larvae. For this, we analyzed the mitotic index in 3 dpf haploid larvae treated with reversine using flow cytometry (Fig. 6D and E). Reversine significantly reduced the proportion of pH3-positive cells in haploid larvae to levels equivalent to that in diploids (Fig. 6D and E, see also Fig. 4A and B). In immunostained haploid larvae, we also observed a reduction in pH3-positive cells across tissues upon reversine treatment (Fig. 6F). These results indicate that the abnormal cell proliferation pattern with the high mitotic index in haploid larvae is mainly due to the SAC-dependent mitotic arrest.

Based on the finding that SAC inactivation mitigated abnormal cell proliferation in haploid

larvae, we tested the effect of reversine treatment on organ growth in haploid larvae.

Reversine treatment significantly increased body axis length and eye size in haploid larvae

compared to DMSO-treated control (Fig. 6G and H). These results indicate that the

haploidy-linked mitotic stress with SAC activation is a primary constraint for organ growth

in haploid larvae.

Discussion

Cell proliferation defects limit organ growth in haploid zebrafish larvae

Poor organ growth is a common feature of haploid syndrome in non-mammalian

vertebrates. Since haploid somatic cells are generally smaller than isogenic diploids in

animals (Gibeaux et al., 2018; Menon et al., 2020; Yaguchi et al., 2018), haploid embryos

have a higher demand for cell proliferation (i.e., need more cells) than diploids for

achieving normal organ size. Indeed, such compensatory cell number increase occurs in

some cases, such as pronephric tubules and ducts in haploid newt larvae or cellular

blastoderms of haploid fry (Edgar et al., 1986; Fankhauser, 1945). However, we found that

haploid zebrafish larvae manifested frequent mitotic arrest and cell death across various

organs, which blocked efficient cell proliferation during organogenesis. Resolution of

mitotic arrest by SAC inactivation or suppression of apoptosis by p53 depletion

significantly improved organ growth in haploid larvae. These results support the idea that

poor organ growth in haploid larvae stems, at least in part, from haploidy-linked defects in

cell proliferation control, which abolishes the compensatory cell number increase required

for ontogenesis in the haploid state.

Developmental stage-dependent aggravation of mitotic defects

Haploid larvae suffered severe mitotic arrest with the abnormally high mitotic index after 1

dpf (Fig. 4). Corresponding to these defects, mitotic chromosome misalignment associated

with spindle monopolarization became evident and gradually aggravated after 1 dpf in

haploid larvae (Fig. 5). A previous study reported that normal zebrafish embryos acquire

SAC function by 4 hpf (Zhang et al., 2015), suggesting that spindle monopolarization after

1 dpf activated SAC and arrest mitotic progression. Indeed, SAC inhibition by reversine

sufficiently restored mitotic progression and cell proliferation patterns in haploid larvae

(Fig. 6), demonstrating that monopolar spindle-driven SAC is a cause of haploidy-linked

abnormality in cell proliferation. This data also shows that SAC acquisition is normal even

in the haploid state in zebrafish larvae.

The correlation between centrosome loss and spindle monopolarization indicates that

haploid larval cells fail to form bipolar spindle because of the haploidy-linked centrosome

loss. We found that the centrosome loss progressed from 1 to 3 dpf in haploid larvae,

concomitantly with the gradual reduction in cell size through continuous cell divisions

without full cell growth (Menon et al., 2020). Interestingly, centrosome loss occurred

almost exclusively in haploid cells whose size became smaller than a certain border (Fig. 5),

indicating that cell size is a key determinant of centrosome number homeostasis in the

haploid state. Because of the cell size-coupled progression of centrosome loss, severe

mitotic defects and size retardation took place mainly in the organs that remained

mitotically active even after 1 dpf. Based on these results, we propose that the stage-

dependent propagation of centrosome loss shapes the characteristic profile of "haploid

syndrome" in zebrafish larvae.

While it is currently unknown why haploid cells lose their centrosomes only below a

certain cell size threshold, a possible explanation could be the depletion of centrosomal

protein pool in the cytoplasm with drastic cell size reduction. Consistent with this idea, a

previous study in early C. elegans embryos proposed that the total centrosomal protein

amount proportional to embryonic cell size determined the extent of protein accumulation

at the mitotic centrosomes (Decker et al., 2011). It is important to note that diploid larval

cells also became smaller than the size border found in haploids, though they reached the

border 1 or 2 days later than haploids (Fig. 5D). However, diploid larval cells remained free

from centrosome loss even when smaller than the border. Therefore, interesting future

perspectives are to address the mechanism that secures centrosome number stability upon

cell size changes in diploids and to specify the cause that abolishes the mechanism in

haploids.

In contrast to the later organogenetic stages, haploid embryos at 0.5 dpf possessed a normal

number of centrosomes. This result demonstrates that centrioles provided by the UV-

irradiated sperms are functional to fully support centriole duplication during early

developmental stages before 0.5 dpf. This finding excludes the possibility that centrosome

loss in later haploid larvae is merely a side effect of sperm UV irradiation for gynogenesis.

Generality of haploidy-linked cellular defects across vertebrates

In mammalian cultured cells and embryos, centrosome loss causes chromosome

missegregation and chronic mitotic delay, resulting in gradual p53 accumulation that

eventually blocks cell proliferation or viability (Bazzi and Anderson, 2014; Fong et al.,

2016; Lambrus et al., 2016; Meitinger et al., 2016). Therefore, it is intriguing to speculate

that the haploidy-linked p53 upregulation in zebrafish larvae stems from severe centrosome

loss. Interestingly, haploid mammalian cultured cells also suffer centrosome loss and

chronic p53 upregulation, limiting their proliferative capacity (Olbrich et al., 2017; Yaguchi

et al., 2018). Our findings demonstrate a striking commonality of the haploidy-linked

cellular defects between mammalian and non-mammalian species while manifesting

diverse organismal defects in the haploid state.

A limitation of our current approach to alleviating mitotic arrest using reversine is that SAC

inactivation does not rescue chromosome missegregation caused by premature mitotic exit

with erroneous kinetochore-microtubule attachments (Santaguida et al., 2010). This

precludes us from investigating the relationship between the mitotic fidelity and viability of

haploid larvae. For further investigating the causality between mitotic defects and haploid

developmental abnormalities, it would be ideal to have an experimental condition that

restores intact centrosomal control in haploid larvae. Previously, we found that artificial re-

coupling of the DNA replication cycle and centrosome duplication cycle by delaying the

progression of DNA replication resolved chronic centrosome loss and mitotic defects in

human haploid cultured cells (Yoshizawa et al., 2020). Though we tried to restore

centrosome loss by treating aphidicolin in haploid larvae, severe toxicity of the compound

on haploid larvae after 1 dpf precluded us from testing its effect on centrosome control.

Genetic manipulation of centrosome duplication control in haploid larvae may provide an

excellent opportunity to investigate the causality of centrosome loss in the haploid

syndrome in future studies.

Developmental incompetence of haploid larvae is likely a crucial evolutionary constraint of

the diplontic life cycle (Sagi and Benvenisty, 2017). Though parent-specific genome

imprinting would serve as the primary mechanism for blocking the development of haploid

embryos in mammals, what precludes haploid development in non-mammalian species has

been unknown. Based on our results, we propose that the ploidy-centrosome link, as a

broadly conserved mechanism, limits the developmental capacity of haploid embryos in

non-mammalian vertebrates. It is an intriguing future perspective to address how the

ploidy-centrosome link is preserved or modulated in invertebrate animal species, especially

those with a haplodiplontic life cycle.

Material and Methods

Zebrafish strain and embryos

Wild-type zebrafish were obtained from the National BioResource Project Zebrafish Core

Institution (NZC, Japan) or a local aquarium shop (Homac, Japan). The *Tg(fli1:h2b-*

mCherry)/ncv31Tg line (Yokota et al., 2015) was provided by NZC. Transgenic animal

experiments in this study were approved by the Committee on Genetic Recombination

Experiment, Hokkaido University. Fish were maintained at 28.5°C under a 14 h light and

10 h dark cycle. For collecting sperm for *in vitro* fertilization, whole testes from a single

male were dissected into 1 mL cold Hank's buffer (0.137 M NaCl, 5.4 mM KCl, 0.25 mM

Na_2HPO_4, 0.44 mM KH_2PO_4, 1.3 mM $CaCl_2$, 1.0 mM $MgSO_4$ and 4.2 mM $NaHCO_3$). For

sperm DNA inactivation, sperm solution was irradiated with 254 nm UV (LUV-6, AS ONE)

at a distance of 30 cm for 1 min with gentle pipetting every 30 seconds (s). For

insemination, we added 500 µL sperm solution to ~200 eggs extruded from females

immobilized by anesthesia (A5040, Sigma-Aldrich; 0.08% ethyl 3-aminobenzoate

methanesulfonate salt, pH 7.2). After 10 s, we added 500 mL Embryo medium (5 mM NaCl,

0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$ and 10^{-5}% Methylene Blue). After the

chorion inflation, embryos were grown in Embryo medium at 28.5°C until use.

To inhibit pigmentation, we treated embryos with 0.03 g/L N-phenylthiourea (P7629,

Sigma Aldrich) at 0.5 dpf. Dechorionation and deyolking were done manually in the cold

fish ringer's buffer without Ca^{2+} (55 mM NaCl, 1.8 mM KCl, 12.5 mM NaHCO$_3$, pH 7.2).

In the case of inducing p53 upregulation for checking the specificity of the anti-p53

antibody, diploid larvae were irradiated with 254 nm UV at a distance of 10 cm for 3 min at

66 hours post-fertilization (hpf). For SAC inactivation, larvae were treated with 5 μM

reversine (10004412, Cayman Chemical) from the time points described elsewhere.

Treatment with 0.5% DMSO was used as vehicle control for reversine treatment.

Antibodies

Antibodies were purchased from suppliers and used at the following dilutions: mouse

monoclonal anti-α-tubulin (1:800 for Immunofluorescence staining (IF); YOL1/34; EMD

Millipore); mouse monoclonal anti-β-tubulin (1:1000 for Immunoblotting (IB); 10G10;

Wako); mouse monoclonal anti-centrin (1:400 for IF; 20H5; Millipore); rabbit monoclonal

anti-active caspase-3 (1:500 for IF; C92-605; BD Pharmingen); rabbit polyclonal anti-p53

(1:1000 for IB; GTX128135; Gene Tex); Alexa Fluor 488-conjugated rabbit monoclonal

anti-phospho-histone H3 (pH3) (1:200 for IF and 1:50 for flow cytometry; D2C8; Cell

Signaling Technology); and fluorescence (Alexa Fluor 488 or Alexa Fluor 594) or

horseradish peroxidase-conjugated secondaries (1:100 for IF and 1:1000 for IB; Abcam or

Jackson ImmunoResearch Laboratories). Hoechst 33342 was purchased from Dojinjo (1

mg/mL solution; H342) and used at 1:100.

Flow cytometry

For isolating whole-larval cells, ~ 10 deyolked embryos were suspended in a cold trypsin

mixture (27250-018, Gibco; 0.25% trypsin in 0.14 M NaCl, 5 mM KCl, 5 mM glucose, 7

mM NaHCO$_3$, 0.7 mM EDTA buffer, pH 7.2) for ~15 min on ice with continuous pipetting.

For 2 dpf or older larvae, 8 mg/mL collagenase P (Roche) was added to the trypsin mixture

for thorough digestion. The isolated cells were collected by centrifugation at 1,300 rpm for

15 min at 4°C, fixed with 8% PFA in Dulbecco's phosphate-buffered saline (DPBS, Wako)

for 5 min at 25°C, permeabilized by adding an equal amount of 0.5% Triton X-100 in

DPBS supplemented with 100 mM glycine (DPBS-G), and collected by centrifugation as

above for removing the fixative. For DNA content and mitotic index analyses, cells were

stained with Hoechst 33342 and Alexa Fluor 488-conjugated anti-pH3, respectively, for 30

min at 25°C, washed once with DPBS, and analyzed using a JSAN desktop cell sorter (Bay

bioscience).

Immunofluorescence staining

For staining active caspase-3, or pH3, larvae were fixed with 4% PFA in DPBS for at least

2 h at 25°C, followed by partial digestion with cold trypsin mixture for 3 min. For staining

centrin and α-tubulin, larvae were fixed with 100% methanol for 10 min at −20°C. Fixed

larvae were manually deyolked in 0.1% Triton X-100 in DPBS and permeabilized with 0.5%

Triton X-100 in DPBS overnight at 4°C, followed by treatment with BSA blocking buffer

(150 mM NaCl, 10 mM Tris-HCl, pH 7.5, 5% BSA, and 0.1% Tween 20) for >30 min at

4°C. Larvae were subsequently incubated with primary antibodies for >24 h at 4°C and

with secondary antibodies overnight at 4°C. Following each antibody incubation, larvae

were washed three times with 0.1% Triton X-100 in DPBS. Stained larvae were mounted in

Fluoromount (K024, Diagnostic BioSystems).

To estimate mitotic cell size in 1-3 dpf larvae, we measured the area of the largest confocal

sections of subepidermal mitotic cells (cells located below the top two epidermal cell

layers) using median 3D filters and the polygon selection tool of ImageJ. The cell contour

was judged based on microtubule staining.

Microscopy

Immunostainings of active caspase-3 or pH3 were observed on an A1Rsi microscope

equipped with a 60× 1.4 NA Apochromatic oil immersion objective lens, an LU-N4S

405/488/561/640 laser unit, and an A1-DUG detector unit with a large-image acquisition

tool of NIS-Elements (Nikon). For live imaging of histone H2B-mCherry-expressing larvae,

the larvae were embedded in agarose gel (5805A, Takara) in E3 buffer supplemented with

anesthesia and N-phenylthiourea and observed using a TE2000 microscope (Nikon)

equipped with a Thermo Plate (TP-CHSQ-C, Tokai Hit; set at 30°C), a 60× 1.4 NA Plan-

Apochromatic oil immersion objective lens (Nikon), a CSU-X1 confocal unit (Yokogawa),

and iXon3 electron multiplier-charge-coupled device camera (Andor). Immunostaining of

centrin and α-tubulin was observed on a C2si microscope equipped with a 100× 1.49 NA

Plan-Apochromatic oil immersion objective lens, an LU-N4 405/488/561/640 laser unit,

and a C2-DU3 detector unit (Nikon).

Immunoblotting

Embryos were deyolked in cold-DPBS supplemented with cOmplete proteinase inhibitor

cocktail (Roche, used at 2× concentration), extracted with RIPA buffer (50 mM Tris, 150

mM NaCl, 1% NP-40, 0.5% Sodium Deoxycholate and 0.1% SDS) supplemented with 2×

cOmplete and centrifuged at 15,000 rpm for 15 min at 4°C to obtain the supernatant.

Proteins separated by SDS-PAGE were transferred to Immun-Blot PVDF membrane (Bio-

Rad). Membranes were blocked with 0.3% skim milk in TTBS (50 mM Tris, 138 mM NaCl,

2.7 mM KCl, and 0.1% Tween 20) and incubated with primary antibodies overnight at 4°C

or for 1 h at 25°C and with secondary antibodies overnight at 4°C or 30 min at 25°C. Each

step was followed by 3 washes with TTBS. For signal detection, the ezWestLumi plus ECL

Substrate (ATTO) and a LuminoGraph II chemiluminescent imaging system (ATTO) were

used. Signal quantification was performed using the Gels tool of ImageJ/Fiji software

(NIH).

Measurement of larval body size

Larvae were anesthetized, mounted in 3% methylcellulose (M0387, Sigma), and observed

under a BX51 transparent light microscope (Olympus) equipped with a 10× 0.25 NA

achromatic objective lens (Olympus) and a 20× 0.70 NA Plan-Apochromatic lens

(Olympus). We measured the body length, brain width, and lateral eye area of larvae using

the segmented line tool of ImageJ. Since haploid larvae were often three-dimensionally

curled or bent, we measured lengths of body axes viewed from lateral and dorsal sides and

used longer ones for statistical analyses. We realized that the severeness of the haploidy-

linked morphological defects tended to differ among larvae from different female parents.

Therefore, we used clutches of larvae obtained from the same female parents for

comparative analyses of the experimental conditions.

Morpholino injection

The morpholino used in this study were 5' -GCG CCA TTG CTT TGC AAG AAT TG- 3'

(p53 antisense) and 5' -GCa CCA TcG CTT gGC AAG cAT TG- 3' (4 base-mismatch p53

antisense) (Langheinrich et al., 2002). One nL morpholino (dissolved in 0.2 M KCl with

0.05% phenol red at 3 mg/mL) was microinjected into haploid embryos at the 1 or 2-cell

stage using FemtoJet and InjectMan NI2 (Eppendorf).

Statistical analysis

Analyses for significant differences between the two groups were conducted using a two-

tailed Student's *t*-test in Excel (Microsoft). Multiple group analysis in Fig. 2B was

conducted using one-way ANOVA with Tukey post-hoc test in R software (The R

Foundation). Statistical significance was set at $p < 0.05$. p-values are indicated in figures or

the corresponding figure legends.

Acknowledgment

We are grateful to Kentaro Kobayashi and other members of the Nikon Imaging Center at Hokkaido University for imaging technical support, Yoshimitsu Sagara, Kuniharu Ijiro, Hiroshi Hinou, and Shin-Ichiro Nishimura for microinjectors and microscopes, Mithilesh Mishra for kind supports, and the Open Facility, Global Facility Center, Creative Research Institution, Hokkaido University for the flow cytometer. This work was supported by JSPS KAKENHI (Grant Numbers JP19J12210 and JP21K20737 to K.Y., and JP19KK0181, JP19H05413, JP19H03219, JPJSBP120193801, and JP21K19244 to R.U.), the India Alliance Wellcome Trust/Department of Biotechnology Intermediate Fellowship IA/I/13/2/501042 to S.N., the Princess Takamatsu Cancer Research Fund, the Kato Memorial Bioscience Foundation, the Orange Foundation, the Smoking Research Foundation, Daiichi Sankyo Foundation of Life Science, the Terumo Life Science Foundation, and the Nakatani Foundation to R.U. The authors declare no competing financial interests.

Author Contributions

Conceptualization, K.Y., and R.U.; Methodology, K.Y., D.S., T.Me., T.Mi., T.K., S.N., and R.U.; Investigation, K.Y., and D.S; Formal Analysis, K.Y., and D.S.; Resources, K.Y., D.S., T.Me., A.M., T.K., S.N., and R.U.; Writing – Original Draft, K.Y., and R.U.; Writing – Review & Editing, K.Y, T.Me., A.M., S.N., and R.U.; Funding Acquisition, K.Y., S.N., and R.U.

Figure legend

Figure 1. Morphological comparison between haploid and diploid larvae

(A) Transparent microscopy of haploid or diploid larvae at 3.5 dpf. Broken line: body axis

(left panels), brain width (middle panels), or lateral eye contour (right panels). Arrowheads:

curling or bending of body axis. Arrows: Edema. **(B)** Quantification of body axis length,

brain width, or lateral eye area of haploid or diploid larvae in A. Box plots and beeswarm

plots of ≥17 larvae (≥34 eyes) from three independent experiments (**$p < 0.01$, two-tailed

t-test).

Figure 2. Haploid zebrafish larvae suffer irregularly increased apoptosis

(A) Immunostaining of active caspase-3 and phospho-histone H3 (pH3) in whole-mount

haploid or diploid larvae at 3 dpf. DNA was stained by DAPI. Z-projected images of

confocal sections containing peripheral brain surface are shown. Broken lines indicate brain

area. The arrowhead indicates clusters of apoptotic cells in the haploid larvae. **(B)**

Classification of the level of apoptosis in haploid and diploid larvae. Seven haploid and 7

diploid larvae from 2 independent experiments were analyzed. All larvae observed in this

experiment are shown in Fig. S2.

Figure 3. p53 upregulation limits organ growth in haploid larvae

(A, C) Immunoblotting of p53 in haploid, diploid, or UV-irradiated diploid larvae (A) or

haploid larvae treated with control or p53 antisense morpholino (C) at 3 dpf. Arrows and

open arrowheads indicate full-length and shorter p53 isoforms, respectively. β-tubulin was

detected as a loading control. **(B, D)** Quantification of relative expression of p53 proteins in

A (B) or C (D). Mean ± standard error (SE) of 3 independent experiments (*$p < 0.05$, **$p <$

0.01, one-way ANOVA with Tukey post-hoc test in B, and two-tailed t-test in D). **(E)**

Immunostaining of active caspase-3 in whole-mount haploid control or p53 morphant at 3

dpf. DNA was stained by DAPI. Z-projected images of confocal sections containing

peripheral brain surfaces are shown. Broken lines indicate brain area. The arrowhead

indicates a cluster of apoptotic cells. **(F)** Classification of the level of apoptosis in haploid

morphants. Six control morphants and 4 p53 morphants from 2 independent experiments

were analyzed. All larvae observed in this experiment are shown in Fig. S3A. **(G)**

Transparent microscopy of haploid control and p53 morphants at 3.5 dpf. Broken line: body

axis (left panels), brain width (middle panels), and lateral eye contour (right panels). **(H)**

Quantification of the body axis, brain width, and lateral eye area in G. Box plots and

beeswarm plots of 29 larvae (58 eyes) from 3 independent experiments (**$p < 0.01$, two-

tailed *t*-test).

Figure 4. Frequent mitotic delay and failures in haploid larvae

(A) Flow cytometric analysis of DNA content (Hoechst signal) and mitotic proportion

(marked by anti-pH3) in isolated haploid or diploid larval cells at 1, 2, or 3 dpf. Magenta

boxes indicate the pH3-positive mitotic populations. **(B)** Quantification of mitotic index in

A. Mean ± SE of ≥3 independent experiments (**p < 0.01, two-tailed *t*-test). **(C)** Live

images of endothelial cells expressing histone H2B-mCherry in haploid or diploid larvae.

Images were taken at a 7.5 min interval from 1.5 to 3 dpf. Asterisks indicate neighbor cells.

(D, **E)** Distribution of mitotic length (time duration from NEBD to anaphase onset; D) or

frequency of mitotic fates (E) in C. Data were sorted into separated graphs by mitotic fates

(completion, mitotic death, or mitotic slippage) in E. At least 78 cells of 8 larvae from 8

independent experiments were analyzed.

Figure 5. Centrosome loss and spindle monopolarization in haploid larvae

(A) Immunostaining of α-tubulin and centrin in haploid and diploid larvae at 0.5, 1, 2, and

3 dpf. Magenta boxes in the left or middle panels indicate the enlarged regions of eyes

(shown in the middle panels) or mitotic cells (shown in the right panels), respectively.

Insets in the right panels show 3× enlarged images of centrioles. **(B**, **C)** Frequency of

spindle polarity and centrin foci number in mitotic cells in the whole-head region

(including eyes, brain, skin epithelia, and olfactory organ) at different developmental stages.

Mean ± SE of ≥4 larvae from ≥2 independent experiments (≥98 cells were analyzed for

each condition; asterisks indicate statistically significant differences from diploids at the

corresponding time points; *p < 0.05, **p < 0.01, two-tailed *t*-test). Data points taken from

the eyes, the brain, or skin epithelia are also shown separately in Fig. S5. **(D**, **E)** Areas of

the largest confocal section of subepidermal mitotic cells in haploid and diploid larvae

during 0.5-3 dpf (D), or frequency of monopolar spindle in haploid and diploid larval cells

sorted by their size (E). Centriole number and spindle polarity in each cell are indicated as

depicted in the graph legends in D. At least 27 cells from ≥4 larvae from ≥2 independent

experiments were analyzed for each condition in D. At least 13 cells were analyzed for each

cell size bin in E.

Figure 6. SAC inactivation mitigates abnormal mitotic patterns and organ growth

defects in haploid larvae

(A) Live images of endothelial histone H2B-mCherry cells in haploid larvae treated with

reversine approximately from 1.5 dpf. Images were taken at a 7.5 min interval from 1.5 to 3

dpf. **(B**, **C)** Distribution of mitotic length (B) or frequency of mitotic fates (C) in A. Data

were sorted into separate graphs by mitotic fates (completion, mitotic death, or mitotic

slippage) in C. At least 75 cells of 6 larvae from 6 independent experiments were analyzed.

(D) Flow cytometric analysis of DNA content (Hoechst signal) and mitotic proportion

(marked by anti-pH3) in the cells isolated at 3 dpf from haploid larvae treated with DMSO

or reversine from 0.5 to 3 dpf. Magenta boxes indicate the pH3-positive mitotic populations.

(E) Quantification of mitotic index in D. Mean ± SE of 4 independent experiments (*$p <$

0.05, two-tailed *t*-test). **(F)** Immunostaining of pH3 in haploid 3-dpf larvae treated with

DMSO or reversine from 1.5 to 3 dpf. Z-projected images of confocal sections containing

peripheral brain surfaces are shown. Broken lines indicate brain area. **(G)** Transparent

microscopy of haploid 3.5-dpf larvae treated with DMSO or reversine from 1.5 to 3.5 dpf.

Broken line: body axis (left panels), brain width (middle panels), and lateral eye contour

(right panels). **(H)** Measurement of the body axis, brain width, and lateral eye area in G.

Box plots and beeswarm plots of 24 larvae (48 eyes) from 3 independent experiments (**p

< 0.01, two-tailed t-test).

Supplemental Figure 1. Generation of haploid and diploid larvae

Experimental scheme of in vitro fertilization for generating haploid and diploid larvae (left)

and flow cytometric DNA content analysis in Hoechst-stained larval cells (right; isolated

from 3- or 5-dpf larvae). The labels on the plot (1c, 2c, and 4c) indicate the relative DNA

amount (c-value). Representative data from 2 independent experiments are shown.

Supplemental Figure 2. Visualization of apoptotic cells in haploid and diploid larvae

Immunostaining of active caspase-3 in whole-mount haploid and diploid larvae at 3 dpf. Z-

projected images of confocal sections containing the peripheral brain surface and inner

brain area. We classified the severity of apoptosis according to the degree of active

caspase-3 staining. Arrowheads indicate clusters of apoptotic cells. Broken lines indicate

brain area. All larvae analyzed in Fig. 2B are shown. The panel includes the larvae identical

to those shown in Fig. 2A (marked by asterisks) to indicate the categories of their apoptotic

levels.

Supplemental Figure 3. Visualization of apoptotic cells in haploid control and p53

morphants

(A) Immunostaining of active caspase-3 in haploid control and p53 morphants at 3 dpf. Z-

projected images of confocal sections containing the peripheral brain surface and inner

brain area. Arrowheads indicate clusters of apoptotic cells. Broken lines indicate brain area.

All larvae analyzed in Fig. 3F are shown. The panel includes the larvae identical to those

shown in Fig. 3E (marked by asterisks) to indicate the categories of their apoptotic levels.

(B) Flow cytometric DNA content analysis in Hoechst-stained larval cells isolated from

haploid control and p53 morphant at 3 dpf. Representative data from 2 independent

experiments are shown.

Supplemental Figure 4. Mitotic progression in haploid, diploid, or reveresine-treated

haploid larvae

(A-C) Mitotic length plot against NEBD time point (dpf) in haploid or diploid larvae in Fig.

4C (A, B) or the reveresine-treated haploid larvae in Fig. 5A (C). At least 74 cells of 6

larvae from 6 independent experiments were analyzed. **(D)** The frequency of mitotic fates

of haploid larval cells arrested at mitosis for > 60 min (shown as percentages in the total

mitotic events analyzed in A; 91 haploid larval cells of 8 larvae from 8 independent

experiments). Data were sorted by NEBD time point (dpf).

Supplemental Figure 5. Haploidy-linked centrosome loss in different organs

(A, **B)** Immunostaining of centrin and α-tubulin in haploid and diploid larvae.

Representative data of skin (A) and brain (B) of haploid and diploid larvae at 0.5, 1, 2, and

3 dpf. Magenta boxes in the left or middle panels indicate the enlarged regions of each

organ (shown in middle panels) or mitotic cells (shown in right panels), respectively. Insets

in the right panels show 3× enlarged images of centrioles. **(C, D)** Spindle polarity and

centrin foci number in mitotic cells in each organ at different developmental stages. At least

11 cells of 4 larvae from 2 independent experiments were analyzed for each condition.

References

Araki, K., H. Okamoto, A.C. Graveson, I. Nakayama, and H. Nagoya. 2001. Analysis of haploid development based on expression patterns of developmental genes in the medaka Oryzias latipes. *Dev Growth Differ*. 43:591-599.

Bazzi, H., and K.V. Anderson. 2014. Acentriolar mitosis activates a p53-dependent apoptosis pathway in the mouse embryo. *Proceedings of the National Academy of Sciences of the United States of America*. 111:E1491-1500.

Dasgupta, S., and L. Matsumoto. 1972. The haploid syndrome in isogenic haploid frog embryos of Rana pipiens derived by nuclear transplantation. *Journal of Experimental Zoology*. 180:413-419.

Decker, M., S. Jaensch, A. Pozniakovsky, A. Zinke, K.F. O'Connell, W. Zachariae, E. Myers, and A.A. Hyman. 2011. Limiting amounts of centrosome material set centrosome size in C. elegans embryos. *Current biology : CB*. 21:1259-1267.

Edgar, B.A., C.P. Kiehle, and G. Schubiger. 1986. Cell cycle control by the nucleo-cytoplasmic ratio in early Drosophila development. *Cell*. 44:365-372.

Ellinger, M.S. 1979. Ontogeny of melanophore patterns in haploid and diploid embryos of the frog, Bombina orientalis. *Journal of Morphology*. 162:77-91.

Ellinger, M.S., and J.A. Murphy. 1980. Cellular morphology in haploid amphibian embryos. *Journal of embryology and experimental morphology*. 59:249-261.

Fankhauser, G. 1938. The microscopical anatomy of metamorphosis in a haploid salamander, triton taeniatus laur. *Journal of Morphology*. 62:393-413.

Fankhauser, G. 1945. Maintenance of normal structure in heteroploid salamander larvae, through compensation of changes in cell size by adjustment of cell number and cell shape. *Journal of Experimental Zoology*. 100:445-455.

Fankhauser, G., and R.B. Griffiths. 1939. Induction of Triploidy and Haploidy in the Newt, Triturus Viridescens, by Cold Treatment of Unsegmented Eggs. *Proceedings of the National Academy of Sciences of the United States of America*. 25:233-238.

Fong, C.S., G. Mazo, T. Das, J. Goodman, M. Kim, B.P. O'Rourke, D. Izquierdo, and M.F. Tsou. 2016. 53BP1 and USP28 mediate p53-dependent cell cycle arrest in response to centrosome loss and prolonged mitosis. *eLife*. 5.

Gibeaux, R., K. Miller, R. Acker, T. Kwon, and R. Heald. 2018. Xenopus Hybrids Provide Insight Into Cell and Organism Size Control. *Frontiers in physiology*. 9:1758.

Hamilton, L. 1966. The role of the genome in the development of the haploid syndrome in Anura. *Development*. 16:559-568.

Hamilton, L., and P.H. Tuft. 1972. The role of water-regulating mechanisms in the development of the haploid syndrome in Xenopus laevis. *Development*. 28:449-462.

Hertwig, O. 1911. Die Radiumkrankheit tierischer Keimzellen, Ein Beitrag zur experimentellen Zeugungs und Vererbungslehre. . *Arch. Mikrosk. Anat.* 77:97-164.

Kroeger, P.T., Jr., S.J. Poureetezadi, R. McKee, J. Jou, R. Miceli, and R.A. Wingert. 2014. Production of haploid zebrafish embryos by in vitro fertilization. *Journal of visualized experiments : JoVE*.

Lambrus, B.G., V. Daggubati, Y. Uetake, P.M. Scott, K.M. Clutario, G. Sluder, and A.J. Holland. 2016. A USP28-53BP1-p53-p21 signaling axis arrests growth after centrosome loss or prolonged mitosis. *The Journal of cell biology*. 214:143-153.

Langheinrich, U., E. Hennen, G. Stott, and G. Vacun. 2002. Zebrafish as a model organism for the identification and characterization of drugs and genes affecting p53 signaling. *Current biology : CB*. 12:2023-2028.

Leeb, M., and A. Wutz. 2011. Derivation of haploid embryonic stem cells from mouse embryos. *Nature*. 479:131-134.

Leeb, M., and A. Wutz. 2013. Haploid genomes illustrate epigenetic constraints and gene dosage effects in mammals. *Epigenetics & Chromatin*. 6:41.

Li, Z., M. Hu, M.J. Ochocinska, N.M. Joseph, and S.S. Easter, Jr. 2000. Modulation of cell proliferation in the embryonic retina of zebrafish (Danio rerio). *Developmental dynamics : an official publication of the American Association of Anatomists*. 219:391-401.

Luo, C., and B. Li. 2003. Diploid-dependent regulation of gene expression: a genetic cause of abnormal development in fish haploid embryos. *Heredity*. 90:405-409.

Mable, B.K., and S.P. Otto. 1998. The evolution of life cycles with haploid and diploid phases. *BioEssays*. 20:453-462.

Meitinger, F., J.V. Anzola, M. Kaulich, A. Richardson, J.D. Stender, C. Benner, C.K. Glass, S.F. Dowdy, A. Desai, A.K. Shiau, and K. Oegema. 2016. 53BP1 and USP28 mediate p53 activation and G1 arrest after centrosome loss or extended mitotic duration. *The Journal of cell biology*. 214:155-166.

Mendieta-Serrano, M.A., D. Schnabel, H. Lomelí, and E. Salas-Vidal. 2013. Cell proliferation patterns in early zebrafish development. *Anat Rec (Hoboken)*. 296:759-773.

Menon, T., A.S. Borbora, R. Kumar, and S. Nair. 2020. Dynamic optima in cell sizes during early development enable normal gastrulation in zebrafish embryos. *Developmental biology*. 468:26-40.

Menon, T., and S. Nair. 2018. Transient window of resilience during early development minimizes teratogenic effects of heat in zebrafish embryos. *Developmental dynamics : an official publication of the American Association of Anatomists*. 247:992-1004.

Nagy, A., K. Rajki, L. Horvárth, and V. Csárnyi. 1978. Investigation on carp, Cyprinus carpio L. gynogenesis. *Journal of Fish Biology*. 13:215-224.

Olbrich, T., C. Mayor-Ruiz, M. Vega-Sendino, C. Gomez, S. Ortega, S. Ruiz, and O. Fernandez-Capetillo. 2017. A p53-dependent response limits the viability of mammalian haploid cells. *Proceedings of the National Academy of Sciences of the United States of America*. 114:9367-9372.

Oppermann, K. 1913. Die Entwicklung von Forelieneiern nach Befruchtung mit radium bestrahlten Samemfaden. . *Arch. Mikrosk. Anat.* 83:141-189.

Otto, S.P., and P. Jarne. 2001. Haploids--Hapless or Happening? *Science*. 292:2441-2443.

Purdom, C.E. 1969. Radiation-induced gynogenesis and androgenesis in fish. *Heredity*. 24:431-444.

Sagi, I., and N. Benvenisty. 2017. Haploidy in Humans: An Evolutionary and
Developmental Perspective. *Developmental Cell*. 41:581-589.
Sagi, I., G. Chia, T. Golan-Lev, M. Peretz, U. Weissbein, L. Sui, M.V. Sauer, O. Yanuka,
D. Egli, and N. Benvenisty. 2016. Derivation and differentiation of haploid human
embryonic stem cells. *Nature*. 532:107-111.
Santaguida, S., A. Tighe, A.M. D'Alise, S.S. Taylor, and A. Musacchio. 2010. Dissecting
the role of MPS1 in chromosome biorientation and the spindle checkpoint through
the small molecule inhibitor reversine. *Journal of Cell Biology*. 190:73-87.
Sorrells, S., C. Toruno, R.A. Stewart, and C. Jette. 2013. Analysis of apoptosis in zebrafish
embryos by whole-mount immunofluorescence to detect activated Caspase 3.
Journal of visualized experiments : JoVE:e51060.
Streisinger, G., C. Walker, N. Dower, D. Knauber, and F. Singer. 1981. Production of
clones of homozygous diploid zebra fish (Brachydanio rerio). *Nature*. 291:293-296.
Subtelny, S. 1958. The development of haploid and homozygous diploid frog embryos
obtained from transplantations of haploid nuclei. *Journal of Experimental Zoology*.
139:263-305.
Sugiyama, M., A. Sakaue-Sawano, T. Iimura, K. Fukami, T. Kitaguchi, K. Kawakami, H.
Okamoto, S.-i. Higashijima, and A. Miyawaki. 2009. Illuminating cell-cycle
progression in the developing zebrafish embryo. *Proceedings of the National
Academy of Sciences*. 106:20812-20817.
Tilghman, S.M. 1999. The Sins of the Fathers and Mothers: Genomic Imprinting in
Mammalian Development. *Cell*. 96:185-193.
Uwa, H. 1965. Gynogenetic haploid embryos of the Medaka (Oryzias Latipes).
Embryologia. 9:40-48.
Wutz, A. 2014. Haploid animal cells. *Development*. 141:1423-1426.
Yaguchi, K., T. Yamamoto, R. Matsui, Y. Tsukada, A. Shibanuma, K. Kamimura, T. Koda,
and R. Uehara. 2018. Uncoordinated centrosome cycle underlies the instability of
non-diploid somatic cells in mammals. *The Journal of cell biology*. 217:2463-2483.
Yamashita, M. 2003. Apoptosis in zebrafish development. *Comparative Biochemistry and
Physiology Part B: Biochemistry and Molecular Biology*. 136:731-742.
Yokota, Y., H. Nakajima, Y. Wakayama, A. Muto, K. Kawakami, S. Fukuhara, and N.
Mochizuki. 2015. Endothelial Ca 2+ oscillations reflect VEGFR signaling-regulated
angiogenic capacity in vivo. *eLife*. 4.
Yoshizawa, K., K. Yaguchi, and R. Uehara. 2020. Uncoupling of DNA Replication and
Centrosome Duplication Cycles Is a Primary Cause of Haploid Instability in
Mammalian Somatic Cells. *Frontiers in cell and developmental biology*. 8:721.
Zhang, M., P. Kothari, and M.A. Lampson. 2015. Spindle assembly checkpoint acquisition
at the mid-blastula transition. *PLoS One*. 10:e0119285.

Figure 1

A
Whole body | Brain | Eye

3.5 dpf

Haploid

Diploid

500 μm   200 μm   200 μm

B

Body axis length
Length (μm)
**
Haploid   Diploid

Brain width
Width (μm)
**
Haploid   Diploid

Eye surface area
Area (μm²)
**
Haploid   Diploid

Figure 2

A

| | Active caspase-3 | DAPI/ Active caspase-3 | pH3 | DAPI/ pH3 |

3 dpf — Haploid, Diploid

B

Apoptosis level	Haploid	Diploid
Scarce	None	2/7
Mild	2/7	5/7
Severe (w/ clusters)	5/7 (3 out of 5)	None (None)

Figure 3

A 3 dpf

α-p53

α-β-tub

(kDa)
— 50
— 37
— 50

B p53 intensity:
■ Full-length
□ Shorter isoform

Normalized intensity (a.u.)

**

*

Haploid Diploid UV-irradiated diploid

C 3 dpf haploid

Control p53 MO

α-p53

α-β-tub

(kDa)
— 50
— 37
— 50

D p53 intensity:
■ Full-length
□ Shorter isoform

Normalized intensity (a.u.)

** *

Control p53 MO

E Active caspase-3   DAPI/ Active caspase-3

3 dpf haploid

Control

p53 MO

200 μm

F

Apoptosis level	Control	p53 MO
Scarce	None	None
Mild	None	3/4
Severe (w/ clusters)	6/6 (5 out of 6)	1/4 (None)

G 3.5 dpf haploid

Control

p53 MO

500 μm   200 μm   200 μm

H Body axis length

Length (μm)

**

Control p53 MO

Brain width

Distance (μm)

**

Control p53 MO

Eye surface area

Area (μm²)

Control p53 MO

Figure 4

Figure 5

Figure 6