An atlas of epithelial cell states and plasticity in lung adenocarcinoma

Guangchun Han*[1], Ansam Sinjab*[2], Warapen Treekitkarnmongkol[2], Wei Lu[2], Alejandra G. Serrano[2], Sharia D. Hernandez[2], Zahraa Rahal[2], Xuanye Cao[1], Enyu Dai[1], Lorena I. Gomez-Bolanos[2], Edwin R. Parra[2], Marcelo Vailati Negrao[3], Tina Cascone[3], Boris Sepesi[4], Seyed Javad Moghaddam[5], Jichao Chen[5], Steven M. Dubinett[6], Paul Scheet[7], Junya Fujimoto[2], Luisa M. Solis[2], Ignacio I. Wistuba[2], Christopher S. Stevenson[8], Avrum E. Spira[8,9], Linghua Wang[#1,10], Humam Kadara[#2,10]

Departments of [1]Genomic Medicine, [2]Translational Molecular Pathology, [3]Thoracic, Head and Neck Medical Oncology, [4]Cardiovascular and Thoracic Surgery, and [5]Pulmonary Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX. [6]Department of Medicine, The University of California Los Angeles, Los Angeles, CA. [7]Department of Epidemiology, The University of Texas MD Anderson Cancer Center, Houston, TX. [8]Lung Cancer Initiative at Johnson and Johnson, Boston, MA. [9]Section of Computational Biomedicine, School of Medicine, Boston University, Boston, MA. [10]The University of Texas Graduate School of Biomedical Sciences, Houston, TX.

* These authors contributed equally

[#] **Correspondence**: Humam Kadara, PhD, Email: hkadara@mdanderson.org ; Linghua Wang, MD, PhD: lwang22@mdanderson.org.

Abstract

Understanding cellular processes underlying early lung adenocarcinoma (LUAD) development is needed to devise intervention strategies[1]. Here, we studied 246,102 single epithelial cells from 16 early-stage LUADs and 47 matched normal lung samples. Epithelial cells comprised diverse normal alveolar and airway lineages as well as cancer cell populations. Diversity among cancer cells was strongly linked to LUAD patient-specific oncogenic drivers. *KRAS*-mutant cancer cells were unique in their transcriptional features, strikingly reduced differentiation, low levels of DNA copy number changes, and increased variability amongst the cells themselves. The local niche of LUADs, relative to that of normal lungs, was enriched with intermediary cells in lung alveolar differentiation with potential to transition to *KRAS*-mutant cancer cells. A subset of alveolar intermediate cells with elevated *KRT8* expression (*KRT8*+ alveolar cells; KACs) showed increased plasticity, and their gene expression profiles were enriched in lung precancer and LUAD and signified poor survival. Murine KACs were evident in lungs of tobacco carcinogen-exposed mice that develop *KRAS*-mutant LUADs but not in the saline-treated control group. While murine KACs emerged prior to tumor onset, they persisted for months after carcinogen cessation, acquired driver *Kras* mutations and, like their human counterparts, were poorly differentiated and harbored *KRAS*-specific transcriptional programs. This study provides new insights into early LUAD development and identifies potential targets for treatment.

Main

Lung adenocarcinomas (LUAD) represent the most common subtype of lung cancer. Thanks to enhanced screening by low dose CT, more LUADs are being detected at early pathological stages[2-4]. Yet, patient prognosis remains moderate to poor thus posing a growing public health burden that warrants improved early treatment strategies. Decoding the earliest events driving LUADs can inform of ideal targets for its interception. Previous work showed that smoking leads to a pervasive molecular (e.g., *KRAS* mutations) and immune changes that are shared between LUADs and their adjacent normal-appearing ecosystems and that are strongly associated with development of lung premalignant lesions (PMLs) and LUAD[1,5-12]. These earlier studies have shed light on events that are likely implicated in early development of lung tumors driven by somatic mutations in a diverse set of oncogenes (e.g. *KRAS*, *EGFR*, *MET*). Yet, most of these reports were based on bulk approaches and were designed to focus on tumor and distant normal sites in the lung, thereby discounting the cellular and transcriptional phenotypes of the expanded LUAD landscape. Furthermore, while many lung single-cell studies provide details on immune and stromal states[13,14], little insight is drawn to epithelial cells given their paucity (~4%) when performing unbiased scRNA-seq analysis (i.e., without prior enrichment). Therefore, little is known of the identities of specific epithelial subsets or how they promote a field of injury, trigger progression of normal lung to PML, and inspire LUAD pathogenesis. Understanding cell type-specific changes at the root of LUAD inception will help delineate tumor heterogeneity and identify actionable targets and strategies for prevention of this morbid disease. Here, our efforts were focused on in-depth single-cell interrogation of normal and malignant epithelial cells from early-stage LUADs and from an experimental mouse model that faithfully recapitulates the

disease, with attention to how specific populations evolve to give rise to malignant tumors.

Epithelial transcriptional landscape

Our study design converges in-depth single-cell RNA-sequencing (scRNA-seq) of early-stage

LUADs clinical specimens, and, for validation, interrogation of murine lungs from a human-

relevant mouse model of LUAD development following exposure to tobacco carcinogen (**Fig.**

1a). In the human cohort, we studied by scRNA-seq enriched (by sorting for EPCAM+)

epithelial cell subsets from early-stage LUADs of 16 patients and 47 paired NL samples (**Fig.**

1a). As previously described[15], NL tissues were collected at defined spatial proximities from the

LUADs – namely, tumor-adjacent, -intermediate, and -distant locations - such that the samples

span a topographic continuum (**Fig. 1a**). Our design was purposefully outlined to interrogate

normal-appearing lung tissues with variable proximities from the tumors in order to determine

cell subsets that are tightly linked with spatial location of the primary tumor. Based on our prior

experience as well as others' showing a relatively small fraction of epithelial cells (~4%) in

unbiased approaches[15,16], we chose to enrich for epithelial cells, including non-malignant cells in

tumor tissues, prior to scRNA-seq. In addition, we collected, from the same tumor and normal

regions, a set of tissues for genomic profiling using whole exome sequencing (WES), and

another set for high-resolution imaging analysis (**Fig. 1a**).

Following quality control, we retained 246,102 high-quality epithelial cells for analysis

(**Extended Data Fig. 1a, b, Supplementary Table 1**). We identified malignant cells (n =

17,064) from otherwise non-malignant "normal" cells (n = 229,038) by integrating information

from inferred copy number variations (inferCNV[17]), clustering distribution, lineage-specific gene

expression, and the presence of reads carrying $KRAS^{G12D}$ somatic mutations (**Fig. 1b-e,**

Extended Data Fig. 1c). We did not detect any malignant cells from P12 or P16, possibly due to

low number of epithelial cells sequenced in these LUADs (Supplementary Data Table 1).

Normal epithelial cells we studied also include those that are within the local vicinity of the

tumor, i.e., from the LUAD tissues themselves. We identified two main clusters of normal cells,

namely alveolar and airway, as well as a small cluster of proliferative cells (**Fig. 1b,**

Supplementary Table 2). Normal airway epithelial cells (n = 40,607) included diverse lineages

such as basal ($KRT17^+$), ciliated ($FOXJ1^+$), and club and secretory ($SCGB1A1^+$) populations, as

well as rare types such as ionocytes ($ASCL3^+$), neuroendocrine ($ASCL1^+$), and tuft cells

($GNAT3^+$). Hierarchical clustering showed that club and basal cells were more closely

associated, pointing towards overall transcriptomic similarities, in contrast to tuft and ciliated

cells which clustered at further distances from the rest of airway cells (**Fig. 1c**). Alveolar cells (n

=187,768) consisted of alveolar type 1 (AT1; $AGER1^+$, $ETV5^+$), alveolar type 2 (AT2; $SFTPB^+$,

$SFTPC^+$), bronchioalveolar (BA) that are $SFTPC/SFTPB+$ but with club cell markers, as well as

a cluster of alveolar intermediate cells (AICs) that was closely tucked between AT1 and AT2

clusters and that shared gene expression features with both major types of alveolar cells (**Fig. 1b,**

c). Relative to AT1 and AT2 cells, AICs and BA cells were more closely associated with each

other - in line with their position on the UMAP and their lineage-intermediate expression

features (**Fig. 1c**).

In contrast to normal cells, malignant cells had low/no expression of lineage-specific markers

and showed, overall, loss of lineage identity (**Fig. 1c**). Malignant cells formed 14 clusters (**Fig.**

1d) that were for the most part patient-specific (**Fig. 1e, left**) signifying strong interpatient

heterogeneity. Malignant cells overall showed high levels of aneuploidy based on inferred CNV

scores (**Fig. 1e, middle**). We did not detect any unique clustering pattern with respect to smoking

status (**Extended Data Fig. 1d**). Annotation based on genomic profiling (by WES) showed that

malignant cells from 3 *KRAS*-mutant (KM-) LUAD patients in our cohort (P2, P10, P14)

clustered closely together and in comparison to malignant cells from 5 EGFR-mutant, 2 MET-

mutant, or other LUADs, which showed a more dispersed clustering pattern (**Fig. 1e, right,**

Extended Data Fig. 1e). We further confirmed expression of somatic $KRAS^{G12D}$ mutation

(inferred from our single-cell data) in malignant cells from the KM-LUAD tumors (**Fig. 1e,**

right).,

LUAD malignant transcriptional programs

We then performed deeper analysis of malignant cells in the context of recurrent driver

mutations. Principal component analysis showed that cells from KM-LUADs clustered together

and distinctively from cells of *EGFR*-mutant LUADs (EM-LUADs) or *MET*-mutant LUADs

(**Fig. 2a**). To further interrogate tumors driven by the same oncogenic drivers, we applied

Harmony[18] to adjust for patient-specific effects (see Methods). Among the 7 distinct malignant

cell sub-clusters, cluster 5 was primarily enriched with malignant cells from KM-LUAD patients

(P2, P10 and P14), (**Extended Data Fig. 1e, Extended Data Fig. 2a**). The vast majority of the

KRAS-mutant malignant cells clustered together in dimension reduction space and separated

from the rest of malignant cells, indicating unique transcriptional reprogramming in *KRAS*-

mutant cells (**Fig. 2b**). In line with previous reports[15] showing low levels of chromosomal

instability in LUADs driven by strong driver mutations *in vivo*[19], malignant cells from KM-

LUADs were chromosomally more stable (indicated by significantly lower CNV scores) in

contrast to those from EM-LUADs - which exhibited highest levels of aneuploidy (**Fig. 2b,**

middle, $p < 0.001$). CNV events were significantly more prominent in malignant cells from

smoker relative to never-smoker patients (**Fig. 2b, right**, $p < 0.001$). We noted high degree of

inter-patient heterogeneity in the differentiation states of malignant cells, whereby, irrespective

of tumor mutation load, cells from KM-LUADs were the least differentiated as indicated by their

highest CytoTRACE[20] scores, followed by EM-LUADs (**Fig. 2c**, $p < 0.001$). We also noted

intra-tumor heterogeneity (ITH) with respect to differentiation states (e.g. P2, P9, P14, P15),

whereby malignant cells from 7 out of the 14 patients with detectable malignant cells (50%)

exhibited a broad distribution of CytoTRACE scores, independent of oncogenic mutations (**Fig.**

2c, right). These results point towards extensive heterogeneity within and across tumors, with

malignant cells from KM-LUADs exhibiting distinctively reduced differentiation.

To better understand malignant cell transcriptional programs that are shared across tumors and

govern tumor cell heterogeneity, we measured, in malignant cells from our cohort, the expression

levels of 23 recurrent malignant meta-programs (MPs) that were recently defined in the

comprehensive study by Gavish and colleagues[21] and which interrogated >1,000 human tumor

samples across a large number of scRNA-seq datasets. Unsupervised clustering identified 5 MP-

driven cell clusters (Meta_C1- Meta_C5) based on expression [21] (**Fig. 2d, Supplementary Table**

3). Notably, 96.8% of malignant cells from KM-LUADs were clustered in Meta_C1 which

showed distinct expression of these MPs compared to malignant cells in other clusters (**Fig. 2d,**

e, Extended Data Fig. 2b), in line with these cells exhibiting unique transcriptome profiles (**Fig.**

2a, b). Consistently, cells from Meta_C1 exhibited the highest expression of a gene module

associated with $KRAS^{G12D}$ present in pancreatic ductal adenocarcinoma (MP30)[21], epithelial to

mesenchymal transition (EMT-III; MP14) and epithelial senescence (MP19), lowest levels of

alveolar differentiation (MP31), in line with the findings above (**Fig. 2c**), overall increased

cellular respiration (MP21) particularly in cells with reduced differentiation (**Fig. 2d, f,**

Extended Data Fig. 2c). Meta_C2 was predominantly composed of malignant cells from EM-

LUADs (**Fig. 2d, e**). Analysis of Meta_C2 malignant cells pointed towards MP patterns unique

to highly differentiated EM-LUADs such as high expression of alveolar MP31, and the lowest

expression of EMT-II (MP13), a glioma-pertinent mesenchymal signature MES (MP16), and

hypoxia (MP6) relative to other MP clusters (**Fig. 2d-f, Extended Data Fig. 2c**). In contrast to

Meta_C2, Meta_C5 had the highest signatures scores for hypoxia (MP6) and the glioma-

pertinent mesenchymal signature (MES, MP16), as well as cell cycle signature enriched with

chromatin regulator genes (MP4) (**Fig. 2d, f, Extended Data Fig. 2c**). Notably, Meta_C5 was

enriched with the most differentiated malignant cells and almost entirely derived from LUADs

from never-smokers and that are neither KM- nor EM-LUADs (**Fig. 2d-f, Extended Data Fig.**

2b, c). MP analysis uncovered a high degree of malignant transcriptional heterogeneity at the

single cell level.

Given the stark contrast of transcriptional programs in KM-LUADs relative to other LUADs, we

further analyzed malignant cell states and heterogeneity in KM-LUADs. We examined cell

differentiation states, expression of MPs, as well as presence of *KRAS* mutations and CNVs in

malignant cells from P14 KM-LUAD (**Fig. 2g, left**). We found increased activation of MP30

signature in a subset of $KRAS^{G12D}$ cells (**Fig. 2g, middle)**, and similarly, diverse activation

patterns of other MPs (e.g. cellular respiration, EMT-III, cell cycle) across the mutant cells

(**Extended Data Fig. 2d**), pointing to marked heterogeneity in transcriptional programs among

$KRAS^{G12D}$ cells. Overall, $KRAS^{G12D}$ cells showed reduced differentiation (higher CytoTRACE

scores) relative to malignant cells without the driver mutation (**Fig. 2g**). Trajectory analysis

showed strong correlation between reduced differentiation (increased CytoTRACE scores) and

$KRAS^{G12D}$ cells (**Fig. 2g, right**) which is concordant with loss of alveolar differentiation (MP31)

in KM-LUADs (**Fig. 2d-f**). To further interrogate malignant cell genomic heterogeneity, we

performed unsupervised CNV clustering analysis and identified three clusters with differential

CNV profiles (**Extended Data Fig. 2e, left**). $KRAS^{G12D}$ cells were enriched in cells harboring

late CNV events (e.g. Chr 1p loss, Chr 8 and Chr 12 gains, identified by clustering analysis of

inferred CNVs, as previously described[15]), and reduced alveolar signature scores in line with

attenuated differentiation (**Extended Data Fig. 2e, f**). These results suggest that signaling by

mutant *KRAS* likely drives transcriptional reprogramming events in KM-LUAD evolution.

To better characterize malignant transcriptional programs unique to early-stage KM-LUADs, we

derived a *KRAS* activation signature based on unique expression patterns of *KRAS*-mutant

malignant cells from our cohort (C5; **Extended Data Fig. 2a**). This signature correlated strongly

and significantly with *KRAS* MP30 present in *KRAS*-mutant pancreatic ductal adenocarcinomas

(**Extended Data Fig. 2g**; $p < 0.0001$). Further analysis of TCGA LUAD cohort showed that

KM-LUADs with relatively high expression of our *KRAS* signature were enriched with activated

KRAS MP30 as well as other MPs we had found to be increased in Meta_C1 (e.g. epithelial

senescence MP19, EMT-III MP14) (n = 498; **Fig. 2h, left**). Relative to *KRAS* signature-low

LUADs, LUADs with high expression of the signature displayed significantly reduced overall

survival (OS) irrespective of *KRAS* mutation status (**Extended Data Fig. 2h**, **left**; $p = 0.005$). A

similar survival disadvantage was observed in *KRAS* signature-high tumors with wild-type

KRAS, and in *KRAS*[G12D]-LUADs despite the small cohort size (**Fig. 2i**; $p = 0.02$). Notably, and

compared to other more frequently observed *KRAS* codon 12 driver mutations (e.g. *KRAS*[G12C],

KRAS[G12V]), *KRAS*[G12D]-LUADs clustered closer to KM-LUADs with high *KRAS* signature

expression and showed increased activation of associated MPs: *KRAS* (MP30), epithelial

senescence (MP19), and EMT-III (MP14) (**Extended Data Fig. 2h, right**; $p < 0.05$). Taken

together, our analysis of a large number of malignant cells from early-stage LUADs identified

neoplastic transcriptomic heterogeneity, cell-level ITH within a single LUAD region, and a

malignant signature that is biologically- and possibly- clinically-relevant to KM-LUADs.

Alveolar intermediate cells in LUADs

Our data suggesting epithelial cell plasticity and heterogeneity in early LUADs, prompted us to

investigate specific cell subsets and states that are implicated in LUAD development. We

analyzed proportions of epithelial lineages (e.g. alveolar and airway) among epithelial cells in

LUAD tissues and the multi-region uninvolved normal lung (NL) samples and found no

significant changes in fractions of AT1, basal, ciliated, or club and secretory cell populations

across the different tissues (**Extended Data Fig. 3a, b**). Notably, the fraction of AT2 cells was

significantly reduced with closer proximity to the LUADs ($p = 0.002$) (**Fig. 3a, b**). Interestingly,

we noted a distinctively significant enrichment of alveolar intermediate cell (AICs) in the LUAD

tissues relative to all NLs ($p = 0.02$) (**Fig. 3a, b**) possibly signifying alveolar changes that are

pertinent to the local niche of LUADs. These observations were overall consistent across

individual patient samples (**Extended Data Fig. 3a**). Recent reports have identified an alveolar

intermediate state which emerges in response to acute lung injury in mice and along the AT2-to-

AT1 differentiation axis, a process which serves to replenish damaged lung tissues[22]. Yet, the

role of these repair mechanisms in the context of LUAD development in humans are poorly

understood. We interrogated developmental and differentiation trajectories among alveolar cells

in our cohort and found that AICs represented an intermediary state (intermediate pseudotime

and CytoTRACE scores) along the differentiation trajectory of AT2 cells to the relatively highly

differentiated AT1 cells (lowest CytoTRACE score) (**Fig. 3c, d**). The proportion of least

differentiated AICs in LUAD tissues was higher than that of their more differentiated

counterparts (low CytoTRACE scores) (29% versus 11%, respectively, **Extended Data Fig. 3c**).

Notably, trajectory analysis suggests that AICs could also transition to malignant cells (**Fig. 3c,**

top branch of the trajectory). A fraction of these cells harbored driver *KRAS* mutations, which

were overall significantly more developmentally late relative to *EGFR*-mutant malignant cells

(**Fig. 3d, bottom**). This implies that AICs could be associated with development of LUADs in

these patients. Next, we took a deeper dive into AICs and identified two transcriptionally distinct

sub-populations whereby one subset featured high expression of *KRT8* (**Fig. 3e**). We refer to

these AICs as "*KRT8*-expressing Alveolar Cells" or KACs and found that they had uniquely

increased expression of *CDKN1A/2A, PLAUR, CLDN4, LCN2, UC5B* and *TTF3* relative to other

AICs (**Fig. 3e, Extended Data Fig. 3d, Supplementary Table 4**). KACs were also significantly

less differentiated (higher CytoTRACE score) relative to other AICs (**Fig. 3f**). The proportion of

KACs among non-malignant epithelial cells detected in LUAD samples was strongly and

significantly increased relative to that among non-malignant epithelial cells of multi-region NL

tissues (**Fig. 3g**). Furthermore, the fraction of KACs in LUAD samples was significantly higher

than that of AT1, AT2, or other AICs (**Fig. 3h, right**). Of note, KACs were also evident in NL

samples and they clustered closer to other AICs relative to tumor derived KACs (**Fig. 3h, left**).

High-resolution, multi-plex, and spatial imaging analysis of KRT8, CLDN4, and pan-cytokeratin

(PanCK) validated the presence of KACs *in situ* in LUAD tissues from multiple patients (**Fig. 3i,**

Extended Data Fig. 3e). KACs were not only enriched in tumor-adjacent regions (TANs) but

were found right next to malignant cells, the latter which highly expressed KRT8 and CLDN4

(**Fig. 3i, Extended Data Fig. 3e**). While KACs were also found in the uninvolved NLs,

consistent with our scRNAseq-analysis, only in the TANs did they display features of "reactive"

or hyperplastic epithelial cells (**Fig. 3i, Extended Data Fig. 3e**). Intriguingly, we found that

expression of *KRAS* signature (see **Fig. 2**) was the highest in KACs relative to AT1, AT2, or

other AICs, and in stark contrast to alveolar signature (MP31) and AT1 signature (**Fig. 3j, left,**

Extended Data Fig. 3f). We derived a KAC signature and found that it significantly and

positively correlated with *KRAS* signature (r = 0.45; $p < 2.2 \times 10^{-16}$) and inversely correlated with

alveolar signature (r = -0.77; $p < 2.2 \times 10^{-16}$; **Fig. 3j, right**). KAC signature was also enriched in

malignant cells from KM-LUADs relative to LUADs with wild-type *KRAS* ($p < 0.001$;

Extended Data Fig. 3g). We then found that the KAC signature was significantly enriched in

bulk transcriptomes of human TCGA LUADs compared to uninvolved NL tissues (**Fig. 3k, top**;

$p = 1.9 \times 10^{-15}$), and in contrast to alveolar signature which was significantly reduced in LUADs

($p < 0.001$; **Extended Data Fig. 3h**). By analysis of our independent cohort of preneoplasia, we

also found that the KAC signature was significantly and progressively increased along the

pathologic continuum of NL, atypical adenomatous hyperplasia (AAH; the earliest precursor of

LUAD), and matching invasive LUAD (**Fig. 3k, bottom**; $p < 0.05$). The fraction of KM-LUADs

among TCGA samples with relatively high KAC signature score was significantly higher than

those with low expression of the signature (n = 498, $p < 0.05$, **Fig. 3l, left**). Also, patients with

KAC signature-high LUADs (n = 243) showed significantly reduced OS ($p = 0.005$) relative to

those with low expression of the signature (n = 246; **Fig. 3l, right**). In a separate cohort, KAC

signature score significantly and positively correlated with expression of *LCN2*, a gene we

previously found to be strongly associated with KM-LUAD (n = 150; r = 0.6; $p < 10^{-15}$)[23], and

was associated with poor OS ($p = 0.04$; **Fig. 3m**). These data describe an intermediate alveolar

cell state that is linked to LUAD pathogenesis in humans.

A KAC state is linked to murine KM-LUADs

Since the association of KACs with human LUAD was not previously reported and their

implication in lung tumor development is not known, we thus sought to validate the association

of these cells with LUAD development *in vivo*, and if confirmed, determine their temporal

dynamics in pathogenesis of the malignancy. We performed scRNA-seq analysis of lung

epithelial cells from mice with knockout of the airway lineage-specific G-protein coupled

receptor (*Gprc5a$^{-/-}$*)[24,25] in the presence or absence of tobacco-carcinogen exposure. LUADs

form in *Gprc5a$^{-/-}$* mice but not in their wild-type littermates, are accelerated by tobacco-

carcinogen exposure, and acquired somatic *Kras^{G12D}* mutations – we surmised that these features

are highly pertinent to KM-LUAD development[24,26,27] and thus to exploring KACs in this setting.

We studied by scRNA-seq lungs from *Gprc5a$^{-/-}$* mice treated with nicotine-derived nitrosamine

ketone (NNK) or saline at the end of exposure (EOE) and at 7 months post-exposure, the time

point of KM-LUAD onset in all mice (n = 4 mice per group and time point) (**Fig 4a**). Clustering

analysis of 9,272 high-quality epithelial cells revealed distinct lineages that were similarly

observed in the human lungs including KACs that clustered in between AT1 and AT2 cell

subsets and close to tumor cells (**Fig. 4b, c**). Like their human counterparts, malignant cells

displayed low expression of lineage-specific genes (**Fig. 4c**). Cells from the malignant cluster

harbored high CNV scores, expressed $Kras^{G12D}$ mutation, and showed increased expression of

markers associated with loss of alveolar differentiation (*Kng2* and *Meg3*) and the epithelial-

specific immune checkpoint *Cd24a* which we had previously shown to be associated with LUAD

pathogenesis[28] (**Fig. 4d, Extended Data Fig. 4a**). Intriguingly, relative to AT1 and AT2 cells,

KACs displayed markedly increased prevalence of $Kras^{G12D}$ mutation, more so than CNV

burden, and increased expression of gastric differentiation gene *Gnk2*- in line with previous

reports[29] and suggestive of reduced alveolar differentiation - albeit to lesser extents compared to

malignant cells (**Fig. 4e, Extended Data Fig. 4b**). We then probed temporal dynamics of

alveolar cell subsets and malignant cells. Malignant cells were present only at 7 months post-

NNK and were absent from samples collected at EOE to carcinogen or in saline-treated animals

(**Fig. 4f, g**). In contrast, fractions of KACs were markedly increased early on in NNK-exposed

mice relative to control littermates treated with saline at EOE to the carcinogen and prior to

tumor formation and, were for the most part, maintained at 7 months post exposure ($p < 0.01$;

Fig. 4f, g; Extended Data Fig. 4c, d). Of note, fractions of AT2 cells were reduced with time

(**Extended Data Fig. 4d, e**) in line with our observations in human LUADs (**Fig. 3b**).

Immunofluorescence (IF) analysis of Krt8, Lamp3 (AT2), and Pdpn (AT1) expression in mouse

lung tissues showed that KACs were present in NNK-exposed normal lungs and were nearly

absent in uninvolved normal lung regions of mice exposed to saline (**Fig. 4h**). Of note, LUADs

themselves also displayed prominent expression of Krt8 (**Fig. 4h**).

Developmental trajectory analysis revealed that while KACs are likely AT2-derived and act as

intermediate states in AT2 to AT1 differentiation (from left to right), in close agreement with

recent reports in the context of acute lung injury[30,31] and our analysis of human LUADs (**Fig. 3**),

they also likely transitioned to malignant cells (**Fig. 5a, top row**). KACs assumed an

intermediate differentiation state (resembling malignant cells) along the path from least

differentiated AT2 cells and up to the most differentiated AT1 cells (highest to lowest

CytoTRACE scores), thereby suggesting that KACs may well represent tumor progenitors (**Fig.**

5a). Intriguingly, a closer look showed that KACs at EOE were-less differentiated (higher

CytoTRACE scores) relative to the more differentiated KACs at 7 months post-exposure (**Fig.**

5a, bottom right). Indeed, we found that KACs were enriched in organoids derived from AT2

cells of NNK- but not saline-exposed $Gprc5a^{-/-}$ mice with reporter labelled-AT2 cells

($Sftpc^{CreER/+}$; $Rosa^{Sun1GFP/+}$; $Gprc5a^{-/-}$) (**Extended Data Fig. 4f**). Analysis of differential gene

expression among KACs, AT1, AT2, and malignant cells, showed that KACs clustered most

closely with tumor cells (**Fig. 5b**). Very much like their human counterparts (**Fig. 3**), KACs

displayed markedly upregulated expression of $Krt8$, $Cldn4$, key players in p53 signaling

including $Trp53$ itself, and $Cdkn2a$, the negative regulator of cell cycle(**Fig. 5c**). Single-cell

pathway analysis showed that NF-κB, p53 signaling, and EMT were the top pathways enriched

or activated in KACs (**Fig. 5d**). We derived a murine KAC signature and found that it was

gradually and significantly enriched in AT2 cells and up to KACs and malignant cells ($p < 0.05$;

Fig. 5e). The murine KAC signature was also enriched in human LUADs, as well as gradually

increased in transcriptomes of premalignant AAHs and invasive LUADs, and relative to matched

NL tissues (**Extended Data Fig. 5g**). The mouse KAC signature partially overlaps (21%) with

the human KAC counterpart (**Fig. 3**), and significantly so when compared to overlap by chance

(3%, $p < 0.001$; **Fig. 5f**). Expression of human KAC marker gene orthologs was highly enriched

in mouse KACs relative to other alveolar cells (**Fig. 5g**). Our *in vivo* analyses identify KACs as

an intermediate cell state in early development of KM-LUAD and following tobacco carcinogen

exposure.

Discussion

Our high-resolution analysis of epithelial cells from early-stage LUADs and the peripheral lung

uncovered diverse malignant states, patterns of ITH, and cell plasticity programs that are linked

to KM-LUAD pathogenesis. Our analysis of "normal" epithelial cell subsets from LUADs and

the uninvolved normal lungs describe a scenario whereby intermediary cells that arise upon

activation of alveolar differentiation programs are linked to transformation of normal lung

epithelial cells to malignant states (**Fig. 5h**). AICs can transition to KACs, which themselves

likely comprise LUAD progenitors, and emerge early on and acquire unique transcriptional

programs, disrupted differentiation, and activation of epithelial plasticity programs - phenotypes

that are important for LUAD development. We also show that AICs temporally evolve early on

post tobacco carcinogen exposure *in vivo* and prior to tumor onset, thereby lending support to the

notion that these cells could be a discrete phenotypic attribute of LUAD *field of injury*. It is

plausible that the unique and evolutionarily conserved AIC states are at the root of LUAD

inception and therefore, comprise a feature of a primed normal-appearing or premalignant

injured/mutated cells[28] (**Fig. 5h**). Thus, our discoveries inspire the derivation of targets to

prevent initiation and development of this deadly disease.

Our study catalogs a rich cellular and transcriptional landscape of lung epithelial cells from

early-stage LUAD patients. We describe remarkable inter-patient heterogeneity of malignant cell

subsets that is likely based, and possibly partially driven by, different driver mutations. Of note,

we uncover unique transcriptomic aspects of *KRAS*-mutant LUADs, whereby loss of epithelial

lineage identity was a hallmark of malignant cells from these tumors. Our deep single-cell

profiling of *KRAS*-mutant cells was not only reminiscent of previous reports showing loss of

AT2 identity in organoid models of KM-LUAD[32], and lineage promiscuity in metastatic

LUADs[16], but also pinpointed marked heterogeneity in transcriptional programs among these

cells. In-depth interrogation of a large number of malignant cells also enabled us to derive a

KRAS signature with potential translational applications - this signature marks *KRAS*-mutant

tumors when examined in public datasets and is significantly associated with poor survival.

These findings point towards possible treatment targets for KM-LUAD that may be contextually

relevant to emerging therapeutic strategies that target *KRAS*[33]. Our results also motivate a timely

interrogation of potential combinatorial approaches that could amplify the benefits of next-

generation *KRAS*[G12D] or pan-*KRAS* inhibitors for treatment of this highly disadvantaged

population of patients.

We also had the capacity to study our large cohort of non-malignant epithelial cells in more

detail. In human LUADs, we found AICs which, for the most part, had been previously

described in animal models as intermediate cells in a transitionary state between AT1 and AT2

cells[30,31,34]. This led us to identify tumor-associated KACs, a subset of AICs that arises from AT2

cells and that is vastly enriched in LUADs. Our results support the existence of alveolar

plasticity programs that are active early on in LUAD development. Recent studies have shown

that following acute lung injury, AT2 cells differentiate into AICs that are characterized by high

expression of *Krt8* and that are crucial for regeneration of AT1 cells[30,31,34]. In parallel, few

studies have described cellular disruptive contexts whereby regeneration programs constitute

mechanisms that not only could underlie cancer development, but that may also drive malignant

cell heterogeneity[14,35]. Despite these findings, the implications of AT2-mediated repair

mechanisms in lung tumorigenesis have been for the most part, largely elusive. We propose that

KACs could short-circuit the AT2-to-AT1 differentiation trajectory, thereby transitioning to

malignant cells including those harboring driver *KRAS*^{G12D} mutations (**Fig. 5h**). We also

characterized a KAC signature that associated with poor cell differentiation status and

augmented epithelial plasticity, a functionally-discrete hallmark of cancer[36]. Congruent with

these findings, we showed that tobacco carcinogen exposure *in vivo* activated cellular programs

that enabled the phenotypic switching of AT2 cells to KACs and which were closely linked to

emergence of malignant cells in KM-LUADs, suggesting that the early appearance of these cells

following tobacco exposure may represent development of a *field of injury*[11]. Additional studies

can further unravel the role of plastic cell states in the continuum of lung malignancy and

identify potential KAC signals that could be harnessed as targets for prevention of PML

inception and/or progression at the earliest possible stages.

From our downstream analysis of KACs, we uncovered multiple lines of evidence pointing

towards a strong link between KACs and LUADs that specifically harbor *KRAS* mutations. First,

we found that KACs, much like *KRAS*-mutant malignant cells, are less differentiated relative to

AT1 and AT2 subsets. This is in line with our analysis of malignant cells in KM-LUAD and

whereby loss of alveolar differentiation was a unique characteristic of these cells. Furthermore,

gene expression in both KACs and malignant *KRAS* mutant tumors showed activation of EMT

and mutant p53 programs. Intriungly, some of these transcriptional modules were recently

reported in mice undergoing lung repair in respose to acute injury[30,31,34]. Furthemore, while our

in vivo findings are consistent with previous independent reports showing that AT2 cells are the

preferential cells-of-origin in *Kras*-driven murine LUADs in aimals[37-39], our results enable a

deeper scrutiny of the unique attributes and states of alveolar intermediary cells that have

potential to constitute progenitors for KM-LUADs. Future analysis of additional LUAD tissues

as well as mouse models of LUAD driven by other oncogenes (such as *EGFR*) can serve to

determine whether KACs are specific to KM-LUADs.

Our study design converges an in-depth interrogation of early-stage LUADs as well as a human-

relevant mouse model of KM-LUAD which enabled the analysis of an expansive number of

epithelial cells. This atlas of epithelial and malignant cell states in human and murine LUADs

underscores new cell-specific states that underlie inception of LUADs, and that can thus guide

novel strategies to prevent the initiation and development of this morbid disease.

References

1   Kadara, H., Scheet, P., Wistuba, II & Spira, A. E. Early Events in the Molecular

Pathogenesis of Lung Cancer. *Cancer Prev Res (Phila)* **9**, 518-527, doi:10.1158/1940-

6207.CAPR-15-0400 (2016).

2   Cardarella, S. & Johnson, B. E. The impact of genomic changes on treatment of lung

cancer. *American journal of respiratory and critical care medicine* **188**, 770-775,

doi:10.1164/rccm.201305-0843PP (2013).

3   Timar, J. The clinical relevance of KRAS gene mutation in non-small-cell lung cancer.

Current opinion in oncology **26**, 138-144, doi:10.1097/CCO.0000000000000051 (2014).

4   Tomasini, P., Walia, P., Labbe, C., Jao, K. & Leighl, N. B. Targeting the KRAS Pathway

in Non-Small Cell Lung Cancer. *The oncologist* **21**, 1450-1460,

doi:10.1634/theoncologist.2015-0084 (2016).

5   Kadara, H. *et al.* Transcriptomic architecture of the adjacent airway field cancerization in

non-small cell lung cancer. *J Natl Cancer Inst* **106**, dju004, doi:10.1093/jnci/dju004

(2014).

6   Belinsky, S. A. *et al.* Aberrant methylation of p16(INK4a) is an early event in lung

cancer and a potential biomarker for early diagnosis. *Proc Natl Acad Sci U S A* **95**,

11891-11896 (1998).

7   Belinsky, S. A. *et al.* Aberrant promoter methylation in bronchial epithelium and sputum

from current and former smokers. *Cancer Res* **62**, 2370-2377 (2002).

8   Franklin, W. A. *et al.* Widely dispersed p53 mutation in respiratory epithelium. A novel

mechanism for field carcinogenesis. *J Clin Invest* **100**, 2133-2137,

doi:10.1172/JCI119748 (1997).

9   Spira, A. *et al.* Airway epithelial gene expression in the diagnostic evaluation of smokers

with suspect lung cancer. *Nat Med* **13**, 361-366, doi:nm1556 [pii] 10.1038/nm1556

(2007).

10 Spira, A., Halmos, B. & Powell, C. A. Update in Lung Cancer 2014. *American journal of*

respiratory and critical care medicine **192**, 283-294, doi:10.1164/rccm.201504-0756UP

(2015).

11　Steiling, K., Ryan, J., Brody, J. S. & Spira, A. The field of tissue injury in the lung and

airway. *Cancer Prev Res (Phila Pa)* **1**, 396-403, doi:1/6/396 [pii] 10.1158/1940-

6207.CAPR-08-0174 (2008).

12　Wistuba, II *et al.* Molecular damage in the bronchial epithelium of current and former

smokers. *J Natl Cancer Inst* **89**, 1366-1373 (1997).

13　Lambrechts, D. *et al.* Phenotype molding of stromal cells in the lung tumor

microenvironment. *Nat Med* **24**, 1277-1289, doi:10.1038/s41591-018-0096-5 (2018).

14　Maynard, A. *et al.* Therapy-Induced Evolution of Human Lung Cancer Revealed by

Single-Cell RNA Sequencing. *Cell* **182**, 1232-1251.e1222,

doi:10.1016/j.cell.2020.07.017 (2020).

15　Sinjab, A. *et al.* Resolving the Spatial and Cellular Architecture of Lung

Adenocarcinoma by Multiregion Single-Cell Sequencing. *Cancer discovery* **11**, 2506,

doi:10.1158/2159-8290.CD-20-1285 (2021).

16　Laughney, A. M. *et al.* Regenerative lineages and immune-mediated pruning in lung

cancer metastasis. *Nat Med* **26**, 259-269, doi:10.1038/s41591-019-0750-6 (2020).

17　Tirosh, I. *et al.* Single-cell RNA-seq supports a developmental hierarchy in human

oligodendroglioma. *Nature* **539**, 309-313, doi:10.1038/nature20123 (2016).

18　Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nature methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).

19　McFadden, D. G. *et al.* Mutational landscape of EGFR-, MYC-, and Kras-driven

genetically engineered mouse models of lung adenocarcinoma. *Proc Natl Acad Sci U S A*

113, E6409-e6417, doi:10.1073/pnas.1613601113 (2016).

20   Gulati, G. S. *et al.* Single-cell transcriptional diversity is a hallmark of developmental

481   potential. *Science* **367**, 405-411, doi:10.1126/science.aax0249 (2020).

482   21   Gavish, A. *et al.* The transcriptional hallmarks of intra-tumor heterogeneity across a

483   thousand tumors. *bioRxiv*, 2021.2012.2019.473368, doi:10.1101/2021.12.19.473368

484   (2021).

485   22   Treutlein, B. *et al.* Reconstructing lineage hierarchies of the distal lung epithelium using

486   single-cell RNA-seq. *Nature* **509**, 371-375, doi:10.1038/nature13173 (2014).

487   23   Treekitkarnmongkol, W. *et al.* Augmented Lipocalin-2 Is Associated with Chronic

488   Obstructive Pulmonary Disease and Counteracts Lung Adenocarcinoma Development.

489   *American journal of respiratory and critical care medicine* **203**, 90-101,

490   doi:10.1164/rccm.202004-1079OC (2021).

491   24   Fujimoto, J. *et al.* Development of Kras mutant lung adenocarcinoma in mice with

492   knockout of the airway lineage-specific gene Gprc5a. *International journal of cancer*

493   **141**, 1589-1599, doi:10.1002/ijc.30851 (2017).

494   25   Westcott, P. M. *et al.* The mutational landscapes of genetic and chemical models of Kras-

495   driven lung cancer. *Nature* **517**, 489-492, doi:10.1038/nature13898 (2015).

496   26   Cancer Genome Atlas Research, N. Comprehensive molecular profiling of lung

497   adenocarcinoma. *Nature* **511**, 543-550, doi:10.1038/nature13385 (2014).

498   27   Kantrowitz, J. *et al.* Genome-Wide Gene Expression Changes in the Normal-Appearing

499   Airway during the Evolution of Smoking-Associated Lung Adenocarcinoma. *Cancer*

500   *Prev Res (Phila)* **11**, 237-248, doi:10.1158/1940-6207.CAPR-17-0295 (2018).

28   Sinjab, A., Han, G., Wang, L. & Kadara, H. Field Carcinogenesis in Cancer Evolution:

What the Cell Is Going On? *Cancer Research* **80**, 4888, doi:10.1158/0008-5472.CAN-

20-1956 (2020).

29   Snyder, E. L. *et al.* Nkx2-1 represses a latent gastric differentiation program in lung

adenocarcinoma. *Mol Cell* **50**, 185-199, doi:10.1016/j.molcel.2013.02.018 (2013).

30   Choi, J. *et al.* Inflammatory Signals Induce AT2 Cell-Derived Damage-Associated

Transient Progenitors that Mediate Alveolar Regeneration. *Cell Stem Cell* **27**, 366-

382.e367, doi:10.1016/j.stem.2020.06.020 (2020).

31   Strunz, M. *et al.* Alveolar regeneration through a Krt8+ transitional stem cell state that

persists in human lung fibrosis. *Nature Communications* **11**, 3559, doi:10.1038/s41467-

020-17358-3 (2020).

32   Dost, A. F. M. *et al.* Organoids Model Transcriptional Hallmarks of Oncogenic KRAS

Activation in Lung Epithelial Progenitor Cells. *Cell Stem Cell* **27**, 663-678.e668,

doi:10.1016/j.stem.2020.07.022 (2020).

33   Salgia, R., Pharaon, R., Mambetsariev, I., Nam, A. & Sattler, M. The improbable targeted

therapy: KRAS as an emerging target in non-small cell lung cancer (NSCLC). *Cell*

Reports Medicine **2**, 100186, doi:https://doi.org/10.1016/j.xcrm.2020.100186 (2021).

34   Kobayashi, Y. *et al.* Persistence of a regeneration-associated, transitional alveolar

epithelial cell state in pulmonary fibrosis. *Nat Cell Biol* **22**, 934-946,

doi:10.1038/s41556-020-0542-8 (2020).

35   Kadur Lakshminarasimha Murthy, P. *et al.* Human distal lung maps and lineage

hierarchies reveal a bipotent progenitor. *Nature* **604**, 111-119, doi:10.1038/s41586-022-

04541-3 (2022).

36   Hanahan, D. Hallmarks of Cancer: New Dimensions. *Cancer discovery* **12**, 31-46,

doi:10.1158/2159-8290.Cd-21-1059 (2022).

37   Lin, C. *et al.* Alveolar type II cells possess the capability of initiating lung tumor

development. *PloS one* **7**, e53817, doi:10.1371/journal.pone.0053817 (2012).

38   Liu, B. *et al.* An entropy-based metric for assessing the purity of single cell populations.

Nature Communications **11**, 3155, doi:10.1038/s41467-020-16904-3 (2020).

39   Xu, X. *et al.* Evidence for type II cells as cells of origin of K-Ras-induced distal lung

adenocarcinoma. *Proc Natl Acad Sci U S A* **109**, 4910-4915,

doi:10.1073/pnas.1112499109 (2012).

Figure legends

Fig. 1. Transcriptional landscape of lung epithelial cells in early-stage LUAD patients. a,

schematic overview of the experimental design and the analysis workflow. State: cellular

transcriptional state. Composition: composition of cell subsets. Program: transcriptional

programs in malignant cells. **b,** UMAP plots of 229,038 normal epithelial cells from 63 samples.

Each dot represents a single cell colored by major cell lineages (left), airway sub lineages (top)

and alveolar sub lineages (bottom). AIC: alveolar intermediate cell. **c,** Sub-lineage hierarchy

within airways cells and alveolar cells (left) and expression (proportions and average expression

levels) of selected marker genes for 10 normal epithelial cell types as defined in panel b (right)

and malignant cells. **d,** Unsupervised clustering of 17,064 malignant cells colored by cluster ID.

The zoom-in view (top left) showing malignant cells colored by *KRAS*^{G12D} mutation status

identified using scRNA-seq data. **e,** UMAP plots of malignant cells as shown in **d,** but colored

by patient ID (left), inferred CNV score (middle), and driver mutations identified in each tumor

sample using whole-exome sequencing (right).

Fig. 2. Deep single-cell analysis of malignant cells reveals unique signatures, states, and

malignant programs. a, Principal component analysis (PCA) plot of malignant cells colored by

driver mutations identified in each tumor sample using whole-exome sequencing. **b,** Harmony-

corrected UMAP plots of malignant cells colored by patient ID (left, which were assigned to

color palette groups based on driver mutation status), CNV scores (top middle), smoking status

(top right). Comparison of CNV scores between malignant cells from samples carrying different

driver mutations (bottom left) or between smokers and never smokers (bottom right). ***: FDR

adjusted p-value < 0.001. Box, median ± interquartile range; whiskers, 1.5× interquartile range;

center line: median. **c,** Landscape of malignant cell differentiation status (top left). Comparison

of differentiation states (inferred by CytoTRACE) between malignant cells from samples with

different driver mutations (bottom left). Per sample distribution of cell CytoTRACE score

(right). TMB: tumor mutation burden. Box definitions are the same as in panel **b**. **d,**

Unsupervised clustering of malignant cells based on expression of 23 previously defined

consensus intratumor heterogeneity meta-programs (MPs). **e,** Enrichment of clusters (C1-C5)

from panel **d** in samples with different driver mutations. Distribution of CNV scores (top right)

and CytoTRACE scores (bottom right) across the 5 clusters defined in panel **d**. Box definitions

are the same as in panel **b**. **f,** Distribution of signature scores of 4 representative MPS: across

clusters defined in panel **d** in panel d. Box definitions are the same as in panel **b**. **g,** Profiling of

intratumoral heterogeneity in malignant cells from P14. UMAP plots show P14 malignant cells

colored by (left to right) $KRAS^{G12D}$ mutation status, $KRAS$ signature expression, and cell

differentiation status (CytoTRACE), respectively. Developmental trajectory of P14 malignant

cells colored by inferred pseudotime, differentiation status, and the presence of $KRAS^{G12D}$

mutations. **h,** $KRAS$ signature expression in TCGA LUAD cohort. Enrichment of significantly

correlated meta-programs with $KRAS$ signature in TCGA LUAD cohort are shown. Samples

harboring somatic $KRAS^{G12D}$ mutations are annotated on the top. **i,** Survival probability

differences between samples carrying somatic $KRAS^{G12D}$ mutation, and those with high or low

level of $KRAS$ signature expression. $KRAS$ sig. high: samples showing high $KRAS$ signature

expression (the top quartile) . $KRAS$ sig. low: samples with low $KRAS$ signature expression

(below the third quartile). $KRAS^{G12D}$: samples with $KRAS^{G12D}$ mutation.

Fig. 3. Identification and characterization of *KRT8*+ alveolar intermediate cells that are

implicated in human LUAD development. a, Composition of normal epithelial lineages across

spatial regions as defined in **Fig. 1a**. Dis: distant normal. Int: intermediate normal. Adj: adjacent

normal. AIC: alveolar intermediate cells. AT1: alveolar type 1. AT2: alveolar type 2. BA:

bronchioalveolar. NE: neuroendocrine. **b,** Changes in cellular fractional of AT2 cells (left) and

AICs (right) across the spatial samples. *P*-values were calculated with Kruskal-Wallis test. Box,

median ± interquartile range; whiskers, 1.5× interquartile range; center line: median. **c,**

Pseudotime trajectory analysis of alveolar and malignant cells identified in this study. **d,**

Distribution of CytoTRACE scores in AICs, AT1 and AT2 cells (top). Distribution of

pseudotime scores in malignant cells of *EGFR*- and *KRAS*-mutant tumors. ***: *p*-value < 0.001.

P-value was calculated with Mann-Whitney U test. Box definitions are the same as in panel **b. e,**

Sub-clustering analysis of AICs (left). KACs: $KRT8^{+}$ alveolar intermediate cells. Expression

levels and frequencies of representative marker genes of KACs (right). **f,** CytoTRACE score in

KACs versus other AICs. Box definitions are the same as in panel **b. g,** Proportion of KACs

among non-malignant epithelial cells across the spatial samples. *P*-value was calculated using

Kruskal-Wallis test. Box definitions are the same as in panel **b. h,** Differences in cell density

between LUAD (left) and normal lung tissues (middle). NL: normal lung. Barplot (right) shows

fraction of alveolar cell subsets colored by sample type. ***: FDR adjusted *p*-value < 0.001. *P*-

values were calculated using Fisher's exact tests. **i,** Microphotographs of P14 LUAD (1-5) and

paired uninvolved NL tissue (6-10). Top panels: Hematoxylin & Eosin staining showing LUAD

tumor area (T) and tumor adjacent normal tissue (TAN) displaying reactive hyperplasia of AT2

cells and presence of intra-alveolar macrophages (1, top left), while 6 shows TAN.

Immunofluorescence staining with KRT8 (2 and 7, red), CLDN4 (3 and 8, yellow), PanCK (4

and 9, green), and composite image of all three antibodies (5 and 10). Blue nuclear staining was

done using Syto13. Magnification 20x. Bar 200 μm. **j,** Enrichment of *KRAS* signature score (top

left) and the alveolar lineage signature score (bottom left) across different subsets of alveolar

cells. Correlation analysis between KAC signature scores and *KRAS* signature scores (top right)

or alveolar lineage signature scores (bottom right). *P*-values were calculated with Spearman

correlation test. R denotes the Spearman correlation coefficients. Box definitions are the same as

in panel **b. k,** Enrichment of KAC signature score across samples of TCGA LUAD (top) and

premalignancy (bottom) cohorts. AAH: atypical adenomatous hyperplasia. *P*-values were

calculated using Wilcoxon signed-rank test. Box definitions are the same as in panel **b. l,**

Fraction of *KRAS*-mutant samples among TCGA samples with high or low KAC signature score

(left). MUT: *KRAS* mutant. WT: *KRAS* wild type. Low: LUAD samples with signature scores

lower than the group median value. High: LUAD samples with signature scores higher than the

group median value. Differences in overall survival probability between samples with high

versus low KAC signature scores (right). Mo: months. **m,** Correlation between the expression

levels of *LCN2* and KAC signature scores in the PROSPECT cohort (left). Overall survival

probability differences between samples with high *versus* low KAC signature scores in

PROSPECT cohort (right). Group median value of KAC signature scores was used dichotomize

the high and low groups.

Fig. 4. Krt8+ alveolar intermediate cells evolve early and prior to tumor onset *in vivo*

during tobacco-associated KM-LUAD pathogenesis. a, Schematic view of the experimental

design. NNK: nicotine-derived nitrosamine ketone. **b,** UMAP view of mouse epithelial cell

subsets. **c,** Proportions and average expression levels of selected marker genes for mouse normal

epithelial cell lineages and malignant cells as defined in panel **a**. NE: neuroendocrine. KACs:

Krt8+ alveolar cells. **d,** UMAP distribution of alveolar and malignant cells colored by CNV

score (top left), $Kras^{G12D}$ mutation status (top right), and expression level of oncogenic markers

Kng2 and *Meg3*. **e,** Distribution of CNV scores (left) among alveolar and malignant cells and

fraction of $Kras^{G12D}$ mutant cells in KACs, malignant, AT1 and AT2 subsets (right). Absolute

numbers of $Kras^{G12D}$ mutant cells are indicated on top of each bar. **f,** Enlarged UMAPs of

alveolar and malignant cells colored by cell subsets (left), $Kras^{G12D}$ mutation status (middle) and

CNV level (right), both at end-of-exposure to NNK (top row) and at 7 months post-NNK

exposure (bottom row). EOE: end of exposure. Mo: months. **g,** Fraction of malignant cells (left)

and KACs (right) across treatment groups and time points. *: FDR adjusted *p*-value < 0.05. *P*-

values were calculated using Mann-Whitney U test. Box, median ± interquartile range; whiskers,

1.5× interquartile range; center line: median. **h,** Immunofluorescence of Krt8, Lamp3 and Pdpn

level in mouse lung tissues. NL: normal lung. Scale bars: 10 μm.

Fig. 5. *Krt8+* **alveolar cells are implicated in transition of alveolar cells to** *Kras***-mutant**

malignant cells. a, Developmental trajectory of alveolar and malignant cells colored by inferred

pseudotime (top left), cell differentiation status (top middle), and cell type (top right).

Distribution of inferred pseudotime (bottom left) and CytoTRACE (bottom middle) across the

indicated cell subsets. CytoTRACE score distribution in KACs at the two time points. EOE: end

of exposure (bottom right). 7 mo: 7 months post exposure to NNK. Box, median ± interquartile

range; whiskers, 1.5× interquartile range; center line: median. **b,** Normalized expression level of

differentially expressed genes in AT1, KACs, malignant cells and AT2 cells. **c,** Expression levels

of representative KAC marker genes across 4 indicated cell subsets. Box, median ± interquartile

range; whiskers, 1.5× interquartile range; center line: median. **d,** Significantly enriched

functional pathways of marker genes of KACs. **e,** Enrichment of KAC signature score, defined

using marker genes of KACs, across the four cell subsets. ***: FDR q-value < 0.001; *: FDR q-

value < 0.05. *P*-values were calculated using Mann-Whitney U test. **f,** Fraction of genes

belonging to human KAC signature among mouse KAC genes *versus* in genome background. *P*-

value was calculated using Fisher's exact test. **g,** Proportions and average expression levels of

mouse orthologs of selected human KAC marker genes in murine KACs and other alveolar cells.

h, Proposed model for alveolar plasticity dynamics in lung cancer whereby a subset of the AICs

in the intermediate AT2-to-AT1 differentiation state can evolve into KACs and later, lead to

LUAD development.

METHODS

Multiregional sampling of surgically resected LUADs and normal lung tissues

Study subjects were evaluated at MD Anderson Cancer Center and underwent standard-of-care

surgical resection of early-stage LUAD (I–IIIA). All samples in the study were obtained under

waiver of consent from banked or residual tissues approved by MD Anderson institutional

review board protocols. Residual surgical specimens were then used for derivation of

multiregional samples for single-cell analysis (**Supplementary Table 1**). Immediately following

surgery, resected tissues were processed by an experienced pathologist assistant. One side of the

specimen was documented and measured, followed by tumor margin identification. On the basis

of the placement of the tumor within the specimen, incisions were made at defined collection

sites in one direction along the length of the specimen and spanning the entire lobe: tumor-

adjacent and tumor-distant normal parenchyma at 0.5 cm from the tumor edge and from the

periphery of the overall specimen/lobe, respectively. An additional tumor-intermediate normal

tissue was selected for P2–P16 and ranged between 3 and 5 cm from the edge of the tumor.

Sample collection was initiated at normal lung tissues that are farthest from the tumor moving

inward toward the tumor to minimize cross-contamination during collection.

Single-cell isolation from LUADs and normal spatial samples

Fresh tissues were collected in RPMI medium supplemented with 2% fetal bovine serum (FBS)

and maintained on ice for immediate processing. Tissues were placed in a cell culture dish

containing Hank's balanced salt solution (HBSS) on ice and extra-pulmonary airways and

connective tissue were removed with scissors. Samples were transferred to a new dish on ice and

minced into tiny pieces (approximately 1mm^3) followed by enzymatic digestion using a cocktail

containing Collagenase A (10103578001, Sigma Aldrich), Collagenase IV (NC9836075, Thermo

Fisher Scientific), DNase I, (11284932001, Sigma Aldrich), Dispase II (4942078001, Sigma

Aldrich), Elastase 1 (NC9301601, Thermo Fisher Scientific) and Pronase (10165921001, Sigma

Aldrich), as previously described [40]. Samples were incubated in a 37 °C oven for 10 minutes

with gentle rotation, pipet mixed at room temperature for further disaggregation, and then

incubated at 37 °C for an additional 10 minutes. Samples were then filtered through 70 μm

strainers (Miltenyi biotech, 130-098-462) and washed with ice-cold HBSS. Filtrates were then

centrifuged and re-suspended in ice-cold ACK lysis buffer (A1049201, Thermo Fisher

Scientific) for red blood cell (RBC) lysis. Following RBC lysis, samples were centrifuged and

re-suspended in ice-cold FBS, filtered (using 40 μm FlowMi tip filters; H13680-0040,

Millipore), and an aliquot was taken to enumerate cells and check for viability by Trypan blue

exclusion analysis.

Sorting and enrichment of viable lung EPCAM+ epithelial singlets

Single cells from patient 1 (P1) were stained with SYTOX Blue viability dye (S34857, Life

Technologies), and processed on a fluorescence-activated cell sorting (FACS) Aria I instrument.

Cells from patients 2 through 16 (P2-P16) were stained with anti-EPCAM-PE (347198, BD

Biosciences;1:50 dilution in ice-cold phosphate-buffered saline, PBS, containing 2% FBS) for 30

minutes with gentle rotation at 4°C. EPCAM-stained cells were then washed, filtered using 40

μm tip filters, stained with SYTOX Blue and processed on a FACS Aria II instrument. Doublets

and dead cells were eliminated, and f viable (SYTOX-negative) EPCAM-positive (EPCAM+,

epithelial) singlets were collected in PBS containing 2% FBS. Cells were washed again to

eliminate ambient RNA, and a sample was taken for counting by Trypan Blue exclusion (T8154,

Sigma Aldrich) before loading on 10X Chromium microfluidic chips.

Preparation of single-cell 5' gene expression libraries

Up to 10,000 cells per sample were partitioned into nanoliter-scale Gel beads-in-emulsion

(GEMs) using Chromium Next GEM Single Cell 5' Gel Bead Kit v1.1 (1000169, 10XGenomics)

and by loading onto Chromium Next GEM Chips G (1000127, 10XGenomics). GEMs were then

recovered to construct single-cell 5' gene expression libraries using the Chromium Next GEM

Single Cell 5' Library kit (1000166, 10XGenomics) according to the manufacturer's protocol.

Briefly, recovered barcoded GEMs were broken and pooled, followed by magnetic bead clean-up

(Dynabeads MyOne Silane, 37002D, Thermo Fisher Scientific). 10X-barcoded full-length cDNA

was then amplified by PCR and analyzed using Bioanalyzer High Sensitivity DNA kit (5067-

4626, Agilent). Up to 50 ng of cDNA was carried over to construct gene expression libraries and

was enzymatically fragmented and size-selected to optimize the cDNA amplicon size prior to 5'

gene expression library construction. Further, samples were subject to end-repair, A-tailing,

adaptor ligation, and sample index PCR using Single Index Kit T Set A (2000240,

10XGenomics) to generate Illumina-ready barcoded gene expression libraries. Library quality

and yield was measured using Bioanalyzer High Sensitivity DNA (5067-4626, Agilent) and

Qubit dsDNA High Sensitivity Assay (Q32854, Thermo Fisher Scientific) kits. Indexed libraries

were normalized by adjusting for the ratio of the targeted cells per library as well as individual

library concentration and then pooled to a final concentration of 10 nM. Library pools were then

denatured and diluted as recommended for sequencing on the Illumina NovaSeq 6000 platform.

Single-cell RNA-seq quality control

The raw scRNA-seq data were pre-processed (demultiplex cellular barcodes, read alignment, and

generation of gene count matrix) using Cell Ranger Single Cell Software Suite (version 3.0.1)

provided by 10X Genomics. For reads alignment of human and mouse scRNA-seq data, the

human reference, GRCh38 (hg38) and mouse reference, GRCm38 (mm10) were used. Detailed

quality control (QC) metrics were generated and evaluated, samples and cells were carefully and

rigorously filtered to obtain high-quality data for downstream analyses[15]. Briefly, for basic

quality filtering, cells with low complexity libraries (in which detected transcripts are aligned to

< 200 genes) were filtered out and excluded from subsequent analyses. This filter will help

remove cell debris, empty drops, and low-quality cells. Likely dying or apoptotic cells where >

15% of transcripts derived from the mitochondrial genome were also excluded.

Doublets detection and removal and batch effect evaluation and correction

Likely doublets or multiplets were detected and carefully removed through a multi-step approach

as described in our recent studies[15,41]. Briefly, doublets or multiplets were identified by the

following methods: 1) Library complexity: cells with high-complexity libraries in which detected

transcripts are aligned to > 6,500 genes were removed. 2) Cluster distribution and marker gene

expression: some doublets or multiplets can form distinct clusters with hybrid expression

features, exhibiting an aberrantly high gene count. We reviewed the expression levels and

proportions of canonical lineage-related marker genes in each identified cluster. Clusters co-

expressing discrepant lineage markers were identified and removed. 3) Doublet detection

algorithms: we applied DoubletFinder[42], an algorithm to predict doublets in scRNA-seq data, to

detect doublets. The proportion of expected doublets was estimated based on the number of cells

used for scRNA-seq library construction. Library size normalization was then performed using

Seurat[43] on the filtered gene-cell matrix to obtain normalized UMI count. Statistical assessment

of possible batch effects was performed using the R package k-BET[44], as described in our

previous studies[15,41] and Harmony[18] was run with default parameters to remove batch effects

present in the PCA space.

Unsupervised clustering and subclustering analysis

Seurat[43] was applied to the normalized gene-cell matrix to identify highly variable genes (HVGs)

for unsupervised cell clustering. Principal component analysis (PCA) was performed on the top

2,000 HVGs. The elbow plot was generated with the *ElbowPlot* function of Seurat and, based on

which, the number of significant principal components (PCs) was determined. The

FindNeighbors function of Seurat was used to construct the Shared Nearest Neighbor (SNN)

Graph, based on unsupervised clustering performed with Seurat function *FindClusters*. Multiple

rounds of clustering and subclustering analysis were performed to identify major cell types and

distinct cell transcriptional states. Dimensionality reduction and 2-D visualization of the single

cell clusters was performed using Uniform Manifold Approximation and Projection (UMAP)[45]

with Seurat function *RunUMAP*. The number of PCs used to calculate the embedding was the

same as that used for clustering. To determine whether cells are over-clustered, we checked for

each cell type, whether multiple subclusters represent the same cell state without showing any

unique features. In addition, we applied ROGUE[38], an entropy-based statistic, to quantify the

cellular transcriptional heterogeneity of each cluster. Subclustering analysis was then performed

for low-purity clusters suggested by ROGUE. Hierarchical clustering of major epithelial subsets

was performed on the Harmony batch-corrected PCA dimension reduction space. The

hierarchical tree was computed based on the Euclidean distance with Ward linkage. Dendrogram

was generated using the R function *plot.hc*.

Identification of malignant cells

Malignant cells were distinguished from non-malignant subsets based on information integrated

from multiple sources as described in our recent studies[15,41]: 1) Cluster distribution: due to the

high degree of inter-patient tumor heterogeneity, malignant cells often exhibit distinct cluster

distribution from that of normal epithelial cells. While non-malignant cells derived from

different patients often are clustered together by cell type, malignant cells from each patient's

tumor likely form a separate cluster. 2) Copy-number alterations (CNAs): We applied

inferCNV[17] (version 1.3.2) to infer large-scale CNAs in each individual cell with T cells as the

reference control to separate malignant cells from non-malignant cells. To quantify the cell level

copy number variations, CNV score were aggregated using a strategy similar to that described in

a previous study[17]. Briefly, we first computed arm-level CNV scores as the mean of the squares

of CNV values across each chromosomal arm. The arm-level CNV scores were further

aggregated across all chromosomal arms by taking the arithmetic mean value of the arm-level

scores using R function *mean*. 3) Marker gene expression: we also examined the expression of

lung epithelial lineage-specific genes and LUAD-related oncogenes in identified epithelial cell

clusters. 4) Cell level expression of $KRAS^{G12D}$ mutations. To extract the single cell somatic

variant for *KRAS* mutation in patients, the alignment records were extracted from the original

BAM file using the mutation location information. Unique mapping alignments (MAPQ=255)

labelled as either PCR duplication or secondary mapping were filtered out. The resulting somatic

variant carrying reads were evaluated using IGV[46] and the corresponding cell-specific barcodes

were then extracted and used for downstream analysis.

PCA analysis of malignant cells

Raw unique molecular identifier counts of identified malignant cells were log normalized and

used for PCA analysis using Seurat (*RunPCA* function). The PCA dimension reduction data was

extracted using *Embeddings* function. The top three most highly ranked PCs were exported for

visualization using JMP v15. The 3D scatterplot of PCA data was generated using the scatterplot

3D tool in JMP v15.

Determination of non-malignant cell types and states

The non-malignant cell types and states were determined based on unsupervised clustering

analysis following batch effects correction using Harmony[18]. Two rounds of clustering analysis

were performed on non-malignant cells to identify major cell types and cell transcriptional states

within the major cell types. The clustering and UMAP visualization of normal epithelial cells

(**Fig. 1b**) was performed using Seurat with the default parameters. Specifically, we used k.param

= 20 and resolution = 0.4 for the shared nearest-neighbor (SNN) graph construction and cluster

identification, respectively. UMAP visualization was performed with the default parameters

(min.dist = 0.3). For clustering analysis of airway epithelial cells and alveolar epithelial cells,

RunPCA function was used to determine the most contributing top PCs in each sub-population.

Same clustering approaches were used k.param = 20 and resolution = 0.4 for SNN graph

construction and clusters identification. UMAP plots were generated with min.dist = 0.3 using

RunUMAP function in Seurat. Density plots of alveolar intermediate cells (**Fig. 3h**) was

generated using *stat_densit_2d* function in R package ggplot2 (version: 3.3.5) with the first two

UMAP dimension reduction data as the input. For normal epithelial cells (fig 1.b), differentially

expressed genes (DEGs) of each cluster were identified using the *FindMarkers* function in

Seurat with the following parameters: FDR adjusted p < 0.05 and fold change > 1.2. We made

references to the expression of canonical epithelial marker genes of previously published

studies[15,47,48] to define the normal epithelial cell types and states. Bubble plots were generated for

selected DEGs and canonical markers to define AT1 (*AGER1*/*ETV5*/*PDPN*+), AT2

(*SFTPB*/*SFTPC*/*ETV5*+), bronchioalveolar (*SFTPC*/*SFTPB*+ and *SCGB1A1*/*SCGB3A1*+), AIC

(*AGER1*/*ETV5*/*PDPN*+ *and SFTPB*/*SFTPC*+), club and secretory

(*SCGB1A1*/*SCGB3A1*/*CYP2F1*+), basal (*KRT5*/*TP63*+), ciliated (*CAPS*/*PIFO*/*FOXJ1*+),

ionocyte (*ASCL3*/*FOXI1*+), neuroendocrine (*CALCA*/*ASCL1*+) and tuft

(*ASCL2*/*MGST2*/*PTGS1*+). We identified KACs as a subset of AICs by unsupervised clustering

of AICs and defined them based on marker genes identified in previous reports[30,31,49] including

significant up regulation of the following genes relative to other alveolar cells: *KRT8*, *CLDN4*,

PLAUR as well as *CDKN1A*/*2A*.

Scoring of curated gene signatures

We measured the expression levels of previously defined ITH meta-programs (MPs) by Gavish

et al.[21] which analyzed 1456 samples covering 24 cancer types and profiling 2,591,545 cells.

Among a total of 41 consensus ITH MPs identified, MPs with unassigned functional annotations

(Unassigned MPs 38-41; n = 4), neural/ hematopoietic lineage specific MPs (MPs 25-29, MPs

33-37; n = 10) as well as cell type specific MPs irrelevant to LUAD (MPs 22-24 secreted/cilia,

MP 32 skin-pigmentation; n = 4) were filtered out, resulting in 23 MPs which intimately

correlated with the hallmarks of cancer[50] for further analysis. The signature scores were

computed using the *AddModuleScore* function in R package *Seurat*, similarly as is described

previously[15,41]. The *KRAS* signature used in this study was derived by calculating the DEGs

between *KRAS*-mutant malignant cell enriched cluster and other malignant cells (FDR adjusted

p-value < 0.05, log fold-change > 1.2). The KACs signature used in this study (human and

mouse) was derived by calculating the DEGs between KACs and other alveolar cells (FDR

adjusted *p*-value < 0.05, log fold-change > 1.2). The KACs signature score and *KRAS* signature

score were calculated using the *AddModuleScore* function in Seurat.

Reconstruction of alveolar cell differentiation trajectories

Reconstruction of the differentiation trajectory of alveolar cells and malignant cells were

performed using Monocle 2[51] by inferring the pseudotemporal ordering of cells according to

their transcriptome similarity. Monocle 2 analysis of malignant cells from P14 was performed

with default parameters on the robustly expressed genes identified using the *detectGenes*

function with default parameters. Detected genes were further required to be expressed by at

least 50 cells. To show the expression dynamics of ITH meta-programs[21] as pseudotime

progresses, ITH meta-program scores were plotted along the pseudotime axis and the smoothed

lines was generated using the smoother tool in JMP Pro v15. CytoTRACE[20] was applied with

default parameters to infer the cellular differentiation states to complement the trajectory analysis

and robustly understand the cellular differentiation hierarchy. The *normalmixEM* function from

mixtools R package was used to determine the threshold of the CytoTRACE score in AICs

(**Extend Data Fig. 3c**) with k = 2. The final threshold calculated for dichotomizing the AICs into

high and low differentiation groups is 0.58.

Survival analysis

Survival analysis including overall survival (OS) status and survival time from TCGA LUAD

cohort was performed as is previously described[15]. *KRAS* mutation status in TCGA LUAD

samples was downloaded from cBioPortal (https://www.cbioportal.org, study ID:

luad_tcga_pan_can_atlas_2018). We used the log-rank test to calculate *P*-values between groups,

and the Kaplan–Meier method to generate survival curves. The statistical significance testing in

this analysis was two-sided. To control for multiple hypothesis testing, we applied the

Benjamini–Hochberg method to correct *P*-values and the FDR Q values were calculated where

applicable. Results were considered statistically significant at *P*-value or FDR Q value of < 0.05.

Bulk RNA extraction and WES

Total RNA was isolated from homogenized cryosections of human lung tissues and, when

available, from frozen PBMCs using Qiagen RNeasy mini kit (Qiagen) according to

manufacturer's recommendations. Qubit 4 Fluorometer (Thermo Fisher Scientific, Waltham,

MA) was used for measurement of RNA yield, and RNA quality was assessed by NanoDrop

1000 (Thermo Fisher Scientific) as well as the Agilent RNA 6000 Pico Kit (Agilent, Santa Clara,

CA) for computation of RNA integrity numbers. Whole exome sequencing (WES) was

performed to analyze recurrent driver mutations, using PBMCs, or most distant NL tissues when

blood draw was not consented, as germline control. WES data were processed and mapped to

human reference genome, somatic mutations were identified, annotated as previously

described[52,53] with further filtration steps. Briefly, only MuTect[54] calls marked as "KEEP" were

selected and taken into the next step. Mutations with a low variant allelic fraction (VAF<0.02) or

had a low alt allele read coverage (< 4) were removed. After that, common variants reported by

ExAc (the Exome Aggregation Consortium, http://exac.broadinstitute.org), Phase-3 1000

Genome Project (http://phase3browser.1000genomes.org/Homo_sapiens/Info/Index), or the

NHLBI GO Exome Sequencing Project (ESP6500) (http://evs.gs.washington.edu/EVS/) with the

minor allele frequency greater than 0.5% were further removed. The intronic mutations,

mutations at 3' or 5' UTR or UTR flanking regions, silent mutations were also removed. The

mutation load was calculated as the number of nonsynonymous somatic mutations (nonsense,

missense, splicing, stop gain, and stop loss substitutions and frameshift insertions and deletions)

called in a sample.

Statistical analyses

In addition to the algorithms and statistical analyses described above, all other basic statistical

analysis was performed in the R statistical environment (version 4.0.0). All statistical tests

performed in this study were two-sided. The Kruskal–Wallis H test was used to compare

variables of interests across three or more groups. Wilcoxon Rank-Sum test was used for paired

comparisons among matched samples from the same patients. Wilcoxon Rank-Sum test was also

used to compare other continuous variables such as gene expression levels, signature scores

between groups. The Spearman's correlation coefficient was calculated to assess the association

between two continuous variables (e.g., the cellular proportions and gene signature scores).

Fisher's exact test was used to identify enriched categories between two categorical variables.

For the TCGA dataset, the clinical data were downloaded from the PanCanAtlas study[20].

Benjamin-Hochberg method was used to control for multiple hypothesis testing. All statistical

significance testing in this study was two-sided and results were considered statistically

significant at *P*-values or FDR *q*-values < 0.05. When a *P*-value reported by R was smaller than

2.2e-16, it was reported as "P < 2.2×10^{-16}".

Digital spatial imaging of human tissues

We used standard immunofluorescence with the following antibodies: Claudin 4 (Clone 3E2C1,

AF594, LSBio, catalog number LS-C354893 concentration 0.5 µg/ml and Keratin 8 (Clone

EP1628Y, AF647, Abcam, catalog number ab192468, concentration 0.25 µg/ml). Optimization

of immunofluorescence biomarkers was previously performed with different dilutions using

colorectal carcinoma and lung adenocarcinoma tissue. Immunofluorescence staining was

performed with standard protocol for Morphology Biomarkers only using the GeoMx Digital

Spatial profiling protocol. Slides were scanned at 20X using the GeoMx Digital Spatial profiling

platform. Following completion of the scanning, multiplex immunofluorescence image slides

were visualized adjusting channel thresholds for each fluorophore. Biomarker expression was

assessed in the following histological areas: Adenocarcinoma cells, adjacent reactive lung tissue

and distant non-reactive lung tissue.

Animal housing and tobacco carcinogen exposure experiments

Animal experiments were conducted according to Institutional Animal Care and Use

Committee–approved protocols. Mice were maintained in a pathogen-free animal

facility. *Gprc5a*$^{-/-}$ mice were generated as previously described[24,55]. Sex- and age-

matched *Gprc5a*$^{-/-}$ mice were divided into starting groups of 12 mice per exposure (NNK or

saline control) and timepoint (end-of-exposure or 7 months post-exposure, n = 48 mice). 8-week

old mice were intraperitoneally injected with 50 mg/kg of body weight nicotine-specific

nitrosamine ketone (NNK, Toronto Research Chemicals) or vehicle 0.9% saline (control), three

times per week for 8 weeks. At the end-of-exposure or at 7 months post-exposure, lungs were

harvested for derivation of live single cells for scRNA-seq (4 mice/group) or were collected in

OCT for immunofluorescence (IF) analysis.

$Sftpc^{CreER/+}$; $Rosa^{Sun1GFP/+}$ were generously provided by Dr. Harold Chapman (University of

California, San Francisco) and were crossed to $Gprc5a^{-/-}$ mice to generate $Sftpc^{CreER/+}$;

$Rosa^{Sun1GFP/+}$; $Gprc5a^{-/-}$ mice for AT2-derived organoid experiments. $Sftpc^{CreER/+}$; $Rosa^{Sun1GFP/+}$;

$Gprc5a^{-/-}$ mice were treated with 50 mg/kg NNK or control saline (IP) (n = 6 each), three times

per week for 8 weeks. At week 6 of treatment (two weeks before the end of exposure), mice from

both groups received 250 µg (IP) Tamoxifen dissolved in corn oil for four consecutive days.

Lungs were harvested at the end of saline or NNK exposure and digested to derive live single

cells for organoid experiments.

Chemicals and reagents

Tobacco-specific carcinogen (nicotine-specific nitrosamine ketone; NNK) with a purity of

99.96% by HPLC was purchased from Toronto Research Chemicals (North York, ON, Canada).

Tamoxifen, as well as Hematoxylin and Eosin (H&E) staining reagents, were purchased from

Sigma Aldrich (St. Louis, MO).

Single-cell isolation from mouse lung tissues

Mouse lungs were collected at end-of-exposure or 7 months post exposure to NNK or saline and

freshly digested for derivation of viable single-cells. Mouse lungs were immediately collected in

6 cm dish containing ice-cold Leibovitz on ice. Extra-pulmonary airways were removed, and

lungs were transferred to a new dish on ice and minced into <1mm pieces with curved scissors.

Using wide-orifice low retention tips, tissue pieces were collected in freshly prepared ice-cold

enzyme solution (1.35 ml 1x Leibovitz + 150 µl of 20 mg/ml Collagenase Type I + 150 µl of 20

mg/ml Elastase + 40 µl 20 mg/ml DNaseI) and transferred to 5 ml LoBind Eppendorf tube.

Tissues were digested and RBCs were removed as described above for human tissues. Murine

single-cell were immediately stained with SYTOX blue, CD45-PE/Cy7 (BioLegend 103114),

Icam2-A647 (Life Technologies A15452), Epcam-BV421 (BioLegend 118225), and Ecad-A488

(eBioscience 53-3249-80). Viable (SYTOX Blue negative) epithelial (CD45-, Icam2-, Epcam+

and/or Ecad+) cells were sorted using FACS Aria flow cytometer and collected into PBS + 2%

FBS on ice. Cells were washed again to eliminate ambient RNA, and a sample was taken for

counting by Trypan Blue exclusion (T8154, Sigma Aldrich). Up to 10,000 cells per sample were

loaded on 10X Chromium microfluidic chips, followed by generation of single-cell libraries in

the manner described above for cells derived from human samples.

Histopathological and IF analysis of mouse lung tissues

After sacrifice, lungs were inflated by injection with 50% optimal cutting temperature (OCT)

reagent in PBS, excised for histopathological assessment. Lung surface lesions were examined

by macroscopic observation and immediately cryopreserved on dry ice. H&E stained

cryosections were digitally scanned using the Aperio ScanScope Turbo slide scanner (Leica

Microsystems Inc., Buffalo Grove, IL) at 200X magnification, and visualized by ImageScope

software (Leica Microsystems, Inc.). Lung sections from all mice were analyzed for lung tumor

incidence according to previously reported criteria that include assessment of tumor

differentiation and invasiveness as well as nuclear crowding, size, and atypia[56]. Total lung area,

as well as area of each of the tumor mass and malignant tumor lesions were selected and

measured by two experienced pathologists (JF) using the Aperio Image Toolbox (Leica

Microsystems Inc.). Lung tumors were enumerated from each group of mice by histopathological

analysis of H&E stained slides and according to previously reported criteria for characterization

of murine lung lesions[56]. Tumor volume (mm^3) was calculated by multiplying the malignant

tumor area (mm^2) by the thickness of tissue sections on each slide. Tumor burden was then

defined as percentage of tumor volume from the entire lung volume. Cryofixed lung tissue

sections were obtained for IF analysis of Lamp3 (clone 391005, Synaptic Systems), Krt8

(TROMA-I clone from the University of Iowa DSHB), and Pdpn (clone 8.1.1, from the

University of Iowa DSHB). Images were acquired using an immunofluorescence microscope.

3D culture and analysis of AT2-derived organoids

Sftpc^CreER/+; Rosa^Sun1GFP/+; Gprc5a^-/- treated with NNK or saline were sacrificed at end-of-

exposure. Lungs were harvested, dissociated into single cells, and live GFP+ single cells were

collected by flow cytometry using a FACS Aria II instrument as previously described[57]. GFP^+

AT2 cells from NNK- or saline-treated groups were immediately washed and re-suspended at a

concentration of 5,000 cells/50 µl of 3D media (F12 medium supplemented with

insulin/transferrin/selenium, 10% FBS, penicillin/streptomycin, L-glutamine). GFP+ cells were

mixed at a 1:1 ratio with 50,000 mouse endothelial cells (harvested from mouse lungs by CD31

selection and expanded *in vitro* as previously described [58]), and resuspended in 50 µl of

Geltrex™ reduced growth factor basement membrane matrix (Gibco, A1413301). 100 µl of 1:1

GFP^+:endothelial cell mixture was plated on transwell inserts with 0.4 µm pores and allowed to

solidify for 30 mins in a humidified CO_2 incubator. Each well was then supplemented with 3D

media containing ROCK inhibitor (Millipore, Y-27632) and recombinant mouse FGF-10 (R&D

Systems, 6224-FG), and plates were incubated at 37°C in a humidified CO_2 incubator. Wells

were replenished with 3D media every other day. At day 21 post-plating, 3D organoids were

harvested from the basement membrane matrix using Gentle Cell Dissociation Reagent

(StemCell Technologies, 100-0485), fixed with 4% paraformaldehyde, permeabilized, blocked,

and stained overnight at 4°C with a mixture of primary antibodies for immunofluorescence

analysis: guinea pig anti-lysosomal associated membrane protein 3 (LAMP3, 1:1000, Synaptic

Systems, 391005), chicken anti-green fluorescent protein (GFP, 1:5000, Abcam, AB13970), rat

anti-keratin8 (Krt8, 1:100, Developmental Studies Hybridoma Bank, TROMA-I). The next day,

organoids were washed and stained with fluorophore-conjugated secondary antibodies overnight

at 4°C while being protected from light. DAPI nuclear stain was added to the secondary antibody

mix for 30 minutes, after which organoids were washed, collected in Aqua-Poly/Mount

(Polysciences, 18606-20) and transferred to slides. Images of organoids were captured using an

Andor Revolution XDi WD Spinning Disk Confocal microscope and analyzed using Imaris

software.

1022

1023   **Data Availability**

1024   Sequencing data for P1-P5 were previously generated and deposited in the European Genome–

1025   phenome Archive (EGA) under the accession number EGAS00001005021[15]. All other

1026   sequencing data generated in this study (human and mouse) will be deposited in EGA and

1027   publicly available upon manuscript acceptance.

1028

1029

Code Availability

1030 **Code Availability**

1031 Detailed codes for all analyses will be shared via GitHub (https://github.com/) and publicly

1032 available very soon.

1033

1034

1035 **Methods references**

1036

40 Slyper, M. *et al.* A single-cell and single-nucleus RNA-Seq toolbox for fresh and frozen

human tumors. *Nature Medicine* **26**, 792-802, doi:10.1038/s41591-020-0844-1 (2020).

41 Wang, R. *et al.* Single-cell dissection of intratumoral heterogeneity and lineage diversity

in metastatic gastric adenocarcinoma. *Nature Medicine* **27**, 141-151, doi:10.1038/s41591-

020-1125-8 (2021).

42 McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in

Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-

337.e324, doi:10.1016/j.cels.2019.03.003 (2019).

43 Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nature*

biotechnology **36**, 411-420, doi:10.1038/nbt.4096 (2018).

44 Büttner, M., Miao, Z., Wolf, F. A., Teichmann, S. A. & Theis, F. J. A test metric for

assessing single-cell RNA-seq batch correction. *Nature methods* **16**, 43-49,

doi:10.1038/s41592-018-0254-1 (2019).

45 McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and

projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

46 Robinson, J. T., Thorvaldsdóttir, H., Wenger, A. M., Zehir, A. & Mesirov, J. P. Variant

Review with the Integrative Genomics Viewer. *Cancer Res* **77**, e31-e34,

doi:10.1158/0008-5472.Can-17-0337 (2017).

47 Sjöstedt, E. *et al.* An atlas of the protein-coding genes in the human, pig, and mouse

brain. *Science* **367**, doi:10.1126/science.aay5947 (2020).

48 Travaglini, K. J. *et al.* A molecular cell atlas of the human lung from single-cell RNA

sequencing. *Nature* **587**, 619-625, doi:10.1038/s41586-020-2922-4 (2020).

49 Maroni, G. *et al.* Identification of a targetable KRAS-mutant epithelial population in non-

small cell lung cancer. *Commun Biol* **4**, 370, doi:10.1038/s42003-021-01897-6 (2021).

50 Hanahan, D. & Weinberg, R. A. Hallmarks of cancer: the next generation. *Cell* **144**, 646-

674, doi:10.1016/j.cell.2011.02.013 (2011).

51 Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.

Nature **566**, 496-502, doi:10.1038/s41586-019-0969-x (2019).

52 Wang, R. *et al.* Multiplex profiling of peritoneal metastases from gastric adenocarcinoma

identified novel targets and molecular subtypes that predict treatment response. *Gut* **69**,

18-31, doi:10.1136/gutjnl-2018-318070 (2020).

53 Hao, D. *et al.* Integrated genomic profiling and modelling for risk stratification in patients

with advanced oesophagogastric adenocarcinoma. *Gut* **70**, 2055-2065,

doi:10.1136/gutjnl-2020-322707 (2021).

54   Cibulskis, K. *et al.* Sensitive detection of somatic point mutations in impure and

heterogeneous cancer samples. *Nature biotechnology* **31**, 213-219, doi:10.1038/nbt.2514

(2013).

55   Tao, Q. *et al.* Identification of the retinoic acid-inducible Gprc5a as a new lung tumor

suppressor gene. *J Natl Cancer Inst* **99**, 1668-1682, doi:10.1093/jnci/djm208 (2007).

56   Nikitin, A. Y. *et al.* Classification of proliferative pulmonary lesions of the mouse:

recommendations of the mouse models of human cancers consortium. *Cancer Res* **64**,

2307-2316 (2004).

57   Barkauskas, C. E. *et al.* Type 2 alveolar cells are stem cells in adult lung. *J Clin Invest*

123, 3025-3036, doi:10.1172/jci68782 (2013).

58   Lee, J. H. *et al.* Lung stem cell differentiation in mice directed by endothelial cells via a

BMP4-NFATc1-thrombospondin-1 axis. *Cell* **156**, 440-455,

doi:10.1016/j.cell.2013.12.039 (2014).

1085

1086

Acknowledgements

We appreciate all the patients who generously provided their samples for this research. We thank

Dr. Harold Chapman for kindly providing *Sftpc$^{CreER/+}$; Rosa$^{Sun1GFP/+}$* mice. This work was

supported in part by funding from Johnson and Johnson (to H.K.), the National Cancer Institute

(NCI) grants R01CA205608 (to HK), 1U2CCA233238 (to H.K., S.M.D., and A.E.S.), University

of Texas SPORE in Lung Cancer P50CA070907 and the P50 core grant CA016672, as well as

the Cancer Prevention and Research Institute of Texas grants RP150079 (to H.K.), RP220101 (to

H.K.), and RP160668 (to I.I.W.). The sequencing core facility at MD Anderson Cancer Center is

supported by the National Institutes of Health grant 1S10OD024977.

1096

1097

Author information

G.H., A.S., W.T., W.L., A.G.S, S.H.D., Z.R., X.C, E.D., L.I.B, E.R.P., S.J.M., J.C., J.F., P.S.,

I.I.W., L.W., and H.K. performed data curation and analysis. G.H., A.S., W.L., A.G.S, S.H.D.,

E.D., J.F., L.S., L.W., and H.K. developed methodology. G.H., A.S., W.L., A.G.S, S.D.H., L.S.,

L.W., and H.K. performed data validation. G.H., A.S., X.C, E.D., S.J.M.,J.C., S.M.D., P.S., I.I.W.,

C.S.S., A.E.S., L.W., and H.K. performed data visualization and interpretation. M.V.N., T.C.,

B.S., I.I.W., C.S.S., A.E.S., L.W., and H.K. provided resources for the work. C.S.S., A.E.S.,

L.W., and H.K. were involved in administration of the project. G.H., A.S., L.W. and H.K. wrote

the original draft. C.S.S., A.E.S., L.W., and H.K. supervised the study. L.W. and H.K. conceived

the study. All authors reviewed and/or edited the final version.

1108

1109

Ethics declarations

Competing interests

CCS and AES are employees of Johnson and Johnson. HK reports research funding from

Johnson and Johnson. MVN receives research funding to institution from Mirati, Novartis,

Checkmate, Alaunos/Ziopharm, AstraZeneca, Pfizer, Genentech, and consultant/advisory board

fees from: Mirati, Merck/MSD, Genentech. TC reports speaker fees/honoraria from The Society

for Immunotherapy of Cancer, Bristol Myers Squibb, Roche, Medscape, and PeerView; advisory

role/consulting fees from AstraZeneca, Bristol Myers Squibb, EMD Serono, Merck & Co.,

Genentech, and Arrowhead Pharmaceuticals; and institutional research funding from

AstraZeneca, Bristol Myers Squibb, and EMD Serono. SJM reports funding from Arrowhead

Pharma and Boehringer Ingelheim outside the scopes of submitted work. BS reports speaker fees

from AstraZeneca and PeerView, advisory fees from AstraZeneca, and consultation fees from

Medscape. IIW reports grants and personal fees from Genentech/Roche, grants and personal fees

from Bayer, grants and personal fees from Bristol-Myers Squibb, grants and personal fees from

AstraZeneca, grants and personal fees from Pfizer, grants and personal fees from HTG

Molecular, personal fees from Asuragen, grants and personal fees from Merck, grants and

personal fees from GlaxoSmithKline, grants and personal fees from Guardant Health, personal

fees from Flame, grants and personal fees from Novartis, grants and personal fees from Sanofi,

personal fees from Daiichi Sankyo, grants and personal fees from Amgen, personal fees from

Oncocyte, personal fees from MSD, personal fees from Platform Health, grants from Adaptive,

grants from Adaptimmune, grants from EMD Serono, grants from Takeda, grants from Karus,

grants from Johnson and Johnson, grants from 4D, from Iovance, from Akoya, outside the

submitted work.

1133

1134

1135   **Additional information**

1136   Supplementary Information is available for this paper.

1137

1138

1139   **Extended data figure**

1140

Extended Data Fig. 1. Summary metrics for data quality, identification of malignant cells and screening of driver mutations in LUAD samples. a, Number of cells retained for scRNA-seq in each sample. Y-axis is log-scaled. **b,** Fraction of mitochondrial genes (left) and the number of detected genes per cell (right). Box, median ± interquartile range; whiskers, 1.5× interquartile range; center line: median. **c,** Schematic plot illustrating the approach used to identify malignant cells in LUAD samples based on clustering, CNV, and expressed $KRAS^{G12D}$ mutations (Expr mutation). **d,** Recurrent driver mutations identified by whole exome sequencing. **e,** UMAP plot of 17,064 malignant cells colored by smoking status.

Extended Data Fig. 2. Characterization of inter- and intra-tumor heterogeneity of LUAD malignant cells. a, Harmony-corrected UMAP plot of malignant cells colored by cluster ID (left) and cluster distribution by sample (right). **b**, Enrichment of hierarchical clusters from Fig. 2d by patient. **c,** expression of 4 representative MP signatures among the 5 hierarchical clusters defined in Fig. 2d. Box, median ± interquartile range; whiskers, 1.5× interquartile range; center line: median. **d,** UMAP plots showing P14 malignant cells colored by expression of 3 MP signatures. **e**, Unsupervised clustering analysis of P14 malignant cells based on inferred CNV profiles (left). UMAP of P14 malignant cells (middle) and inferred trajectory (top right) colored by CNV clusters, as well as $KRAS^{G12D}$ mutation expression status along the developmental trajectory (bottom right). **f**, Signature score of alveolar lineage meta-program across the CNV clusters shown in panel e. ***: $p < 0.001$. P-value was calculated using Mann Whitney U test. **g,** Harmony-corrected UMAP plot of malignant cells colored by $KRAS$ signature score (left). Correlation between MP30 and $KRAS$ signature scores in malignant cells of KM-LUADs (right). P-values were calculated with Spearman correlation test. R denotes the Spearman correlation

coefficient. **h**, Validation of *KRAS* signature score in TCGA LUAD cohort. Left: Kaplan-Meier

plot showing the survival probability differences between samples with high level of *KRAS*

signature, those with low level of *KRAS* signature, and those with *KRAS*^{G12D} mutation. *KRAS* sig.

high: samples within top quartile of *KRAS* signature score. *KRAS* sig. low: samples below the

third quartile of *KRAS* signature score. mo: months. Right: heatmap showing the MP signature

score of TCGA LUAD samples. *: $p < 0.05$. *P*-value was calculated with Chi-square test.

1170

Extended Data Fig. 3. Phenotypic diversity and states of normal epithelial cells. a,

Composition of normal epithelial lineages among the spatial regions at sample level. BA:

bronchioalveolar. NE: neuroendocrine. Prolif: proliferative. **b**, Fractional changes of AT1, basal,

ciliated, and club and secretory cells along the continuum of the spatial samples. Box, median ±

interquartile range; whiskers, 1.5× interquartile range; center line: median. **c**, Distribution and

composition of AICs with low (left) or high (right) CytoTRACE score. **d**, Differentially

expressed genes between KACs and other AICs. **e,** Microphotographs of P10 (e) and P15 (f)

LUADs (1-5) and paired uninvolved NL tissue (6-10). Top panels: Hematoxylin & Eosin

staining showing LUAD tumor area (T) and tumor adjacent normal tissue (TAN; 1, top left),

while 6 shows normal-appearing lung parenchyma. Immunofluorescence staining with KRT8 (2

and 7, red), CLDN4 (3 and 8, yellow), PanCK (4 and 9, green), and composite image of all three

antibodies (5 and 10). Blue nuclear staining was done using Syto13. Magnification 20x. Bar 200

μm. **f,** AT1 signature expression across AT1, AT2, KACs and other AICs. Box definitions are

the same as in panel **b**. **g,** Expression of Alveolar lineage signature in TCGA LUAD samples and

matched NL tissues. ***: $p < 0.001$. *P*-value was calculated using Mann-Whitney U test. Box

definitions are the same as in panel **b**. **h,** *KRT8* signature expression in malignant cells of KM-

versus KRAS-wildtype LUADs. ***: *p*-value < 0.001. *P*-value was calculated using Mann-

Whitney U test. Box definitions are the same as in panel **b**.

1189

Extended Data Fig. 4. Identification of KACs in tobacco carcinogenesis mouse model of

KM-LUAD and the involvement of KACs in transition of AT2 to malignant cells during

AT2-to-AT1 differentiation. a-b, Expression levels of *Cd24a* and *Gkn2* in alveolar cells.

Enlarged UMAPs and violin plots of alveolar cells colored by the expression level of *Cd24a* (a)

and *Gkn2* (b). Box, median ± interquartile range; whiskers, 1.5× interquartile range; center line:

median. **c**, Proportions of normal epithelial cell lineages and malignant cells in each sample. **d**,

Fractional changes of malignant cells, AICs, AT2 and AT1 cells between EOE and 7 months

post NNK treatment. whiskers, 1.5× interquartile range; center dot: median. **e**,

Immunofluorescence analysis of Krt8 and Lamp3 in GFP-labelled AT2-derived mouse lung

organoids following end of exposure to saline or NNK. Scale bar: 10 μm. **f**, KAC signature

expression in samples of TCGA LUAD cohort (left) or preneoplasia cohort (right). AAH:

atypical adenomatous hyperplasia. *P*-values were calculated using Mann-Whitney U test. Box,

median ± interquartile range; whiskers, 1.5× interquartile range; center line: median.

1203

1204

1205   **Supplementary data table legends**

1206

1207   **Supplementary Data Table 1.** Sequencing data quality of all samples included in this study.

1208   Details of patient samples including sex, smoking status and sample location, as well as QC

1209   metrics of single-cell RNA-sequencing libraries in this study.

1210

1211 **Supplementary Data Table 2.** Fraction of epithelial cell lineages in each sample. Fraction of

1212 alveolar and airway lineages among total normal epithelial cells and across tumor and normal

1213 (tumor-adjacent, -intermediate, and -distant) samples of each patient.

1214

1215 **Supplementary Data Table 3.** Metaprogram scores across malignant cells (n = 17064). Scores

1216 of the 41 MPs from Gavish et al, CNV as well as differentiation, across individual malignant

1217 cells. Table also shows smoking and driver mutation status for each patient.

1218

1219 **Supplementary Data Table 4.** Significantly upregulated genes in KACs. List of top 50 most

1220 upregulated genes in human KACs. Average log fold change and adjusted p – values are

1221 indicated for each gene.

Fig. 1 | Transcriptional landscape of lung epithelial cells in early-stage LUAD patients. a, schematic overview of the experimental design and the analysis workflow. State: cellular transcriptional state. Composition: composition of cell subsets. Program: transcriptional programs in malignant cells. **b**, UMAP plots of 229,038 normal epithelial cells from 63 samples. Each dot represents a single cell colored by major cell lineages (left), airway sub lineages (top) and alveolar sub lineages (bottom). AIC: alveolar intermediate cell. **c**, Sub-lineage hierarchy within airways cells and alveolar cells (left) and expression (proportions and average expression levels) of selected marker genes for 10 normal epithelial cell types as defined in panel b (right) and malignant cells. **d**, Unsupervised clustering of 17,064 malignant cells colored by cluster ID. The zoom-in view (top left) showing malignant cells colored by $KRAS^{G12D}$ mutation status identified using scRNA-seq data. **e**, UMAP plots of malignant cells as shown in **d**, but colored by patient ID (left), inferred CNV score (middle), and driver mutations identified in each tumor sample using whole-exome sequencing (right).

Fig. 2. Deep single-cell analysis of malignant cells reveals unique signatures, states, and malignant programs. a, Principal component analysis (PCA) plot of malignant cells colored by driver mutations identified in each tumor sample using whole-exome sequencing. **b,** Harmony-corrected UMAP plots of malignant cells colored by patient ID (left, which were assigned to color palette groups based on driver mutation status), CNV scores (top middle), smoking status (top right). Comparison of CNV scores between malignant cells from samples carrying different driver mutations (bottom left) or between smokers and never smokers (bottom right). ***: FDR adjusted p-value < 0.001. Boxes: 25th to 75th percentile, whiskers: 1.5× interquartile range, center line: median. **c,** Landscape of malignant cell differentiation status (top left). Comparison of differentiation states (inferred by CytoTRACE) between malignant cells from samples with different driver mutations (bottom left). Per sample distribution of cell CytoTRACE score right). TMB: tumor mutation burden. Boxes definitions are the same as in panel **b**. **d,** Unsupervised clustering of malignant cells based on expression of 23 previously defined consensus intratumor heterogeneity meta-programs (MPs). **e,** Enrichment of clusters (C1-C5) from panel **d** in samples with different driver mutations. Distribution of CNV scores (top right) and CytoTRACE scores (bottom right) across the 5 clusters defined in panel **d**. Boxes definitions are the same as in panel **b**. **f,** Distribution of signature scores of 4 representative MPS: across clusters defined in panel **d**. **g,** Profiling of intratumoral heterogeneity in malignant cells from P14. UMAP plots show P14 malignant cells colored by (left to right) $KRAS^{G12D}$ mutation status, $KRAS$ signature expression, and cell differentiation status (CytoTRACE), respectively. Developmental trajectory of P14 malignant cells colored by inferred pseudotime, differentiation status, and the presence of $KRAS^{G12D}$ mutations. **h,** $KRAS$ signature expression in TCGA LUAD cohort. Enrichment of significantly correlated meta-programs with $KRAS$ signature in TCGA LUAD cohort are shown. Samples harboring somatic $KRAS^{G12D}$ mutations are annotated on the top. **i,** Survival probability differences between samples carrying somatic $KRAS^{G12D}$ mutation, and those with high or low level of $KRAS$ signature expression. $KRAS$ sig. high: samples showing high $KRAS$ signature expression (the top quartile) . $KRAS$ sig. low: samples with low $KRAS$ signature expression (below the third quartile). $KRAS^{G12D}$: samples with $KRAS^{G12D}$ mutation.

Figure 3

Fig. 3. | Identification and characterization of *KRT8+* alveolar intermediate cells that are implicated in human LUAD development. a, Composition of normal epithelial lineages across spatial regions as defined in **Fig. 1a**. Dis: distant normal. Int: intermediate normal. Adj: adjacent normal. AIC: alveolar intermediate cells. AT1: alveolar type 1. AT2: alveolar type 2. BA: bronchioalveolar. NE: neuroendocrine. **b**, Changes in cellular fractional of AT2 cells (left) and AICs (right) across the spatial samples. *P*-values were calculated with Kruskal-Wallis test. median ± interquartile range; whiskers, 1.5× interquartile range; center line: median. **c**, Pseudotime trajectory analysis of alveolar and malignant cells identified in this study. **d**, Distribution of CytoTRACE scores in AICs, AT1 and AT2 cells (top). Distribution of pseudotime scores in malignant cells of *EGFR*- and *KRAS*-mutant tumors. ***: *p*-value < 0.001. P-value was calculated with Mann-Whitney U test. Boxes definitions are the same as in panel **b**. **e**, Sub-clustering analysis of AICs (left). KACs: *KRT8+* alveolar intermediate cells. Expression levels and frequencies of representative marker genes of KACs (right). **f**, CytoTRACE score in KACs versus other AICs. Boxes definitions are the same as in panel **b**. **g**, Proportion of KACs among non-malignant epithelial cells across the spatial samples. P-value was calculated using Kruskal-Wallis test. Boxes definitions are the same as in panel **b**. **h**, Differences in cell density between LUAD (left) and normal lung tissues (middle). NL: normal lung. Barplot (right) shows fraction of alveolar cell subsets colored by sample type. ***: FDR adjusted *p*-value < 0.001. *P*-values were calculated using Fisher's exact tests. **i**, Microphotographs of P14 LUAD (1-5) and paired uninvolved NL tissue (6-10). Top panels: Hematoxylin & Eosin staining showing LUAD tumor area (T) and tumor adjacent normal tissue (TAN) displaying reactive hyperplasia of AT2 cells and presence of intra-alveolar macrophages (1, top left), while 6 shows TAN. Immunofluorescence staining with KRT8 (2 and 7, red), CLDN4 (3 and 8, yellow), PanCK (4 and 9, green), and composite image of all three antibodies (5 and 10). Blue nuclear staining was done using Syto13. Magnification 20x. Bar 200 *μm*. **j**, Enrichment of KRAS signature score (top left) and the alveolar lineage signature score (bottom left) across different subsets of alveolar cells. Correlation analysis between KAC signature scores and *KRAS* signature scores (top right) or alveolar lineage signature scores (bottom right). *P*-values were calculated with Spearman correlation test. R denotes the Spearman correlation coefficients. Boxes definitions are the same as in panel **b**. **k**, Enrichment of KAC signature score across samples of TCGA LUAD (top) and premalignancy (bottom) cohorts. AAH: atypical adenomatous hyperplasia. *P*-values were calculated using Wilcoxon signed-rank test.Boxes definitions are the same as in panel **b**. **l**, Fraction of *KRAS*-mutant samples among TCGA samples with high or low KAC signature score (left). MUT: *KRAS* mutant. WT: *KRAS* wild type. Low: LUAD samples with signature scores lower than the group median value. High: LUAD samples with signature scores higher than the group median value. Differences in overall survival probability between samples with high versus low KAC signature scores (right). Mo: months. **m**, Correlation between the expression levels of *LCN2* and KAC signature scores in the PROSPECT cohort (left). Overall survival probability differences between samples with high versus low KAC signature scores in PROSPECT cohort (right). Group median value of KAC signature scores was used dichotomize the high and low groups.

Fig. 4 | *Krt8+* alveolar intermediate cells evolve early and prior to tumor onset in vivo during tobacco-associated KM-LUAD pathogenesis. a, Schematic view of the experimental design. NNK: nicotine-derived nitrosamine ketone. **b**, UMAP view of mouse epithelial cell subsets. **c**, Proportions and average expression levels of selected marker genes for mouse normal epithelial cell lineages and malignant cells as defined in panel **a**. NE: neuroendocrine. KACs: *Krt8+* alveolar cells. **d**, UMAP distribution of alveolar and malignant cells colored by CNV score (top left), *Kras^{G12D}* mutation status (top right), and expression level of oncogenic markers *Kng2* and *Meg3*. **e**, Distribution of CNV scores (left) among alveolar and malignant cells and fraction of *Kras^{G12D}* mutant cells in KACs, malignant, AT1 and AT2 subsets (right). Absolute numbers of *Kras^{G12D}* mutant cells are indicated on top of each bar. **f**, Enlarged UMAPs of alveolar and malignant cells colored by cell subsets (left), *Kras^{G12D}* mutation status (middle) and CNV level (right), both at end-of-exposure to NNK (top row) and at 7 months post-NNK exposure (bottom row). EOE: end of exposure. Mo: months. **g**, Fraction of malignant cells (left) and KACs (right) across treatment groups and time points. *: FDR adjusted *p*-value < 0.05. P-values were calculated using Mann-Whitney U test. Box, median ± interquartile range; whiskers, 1.5× interquartile range; center line: median. **h**, Immunofluorescence of Krt8, Lamp3 and Pdpn level in mouse lung tissues. NL: normal lung. Scale bars: 10 μm.

Fig. 5 | *Krt8*+ alveolar cells are implicated in transition of alveolar cells to Kras-mutant malignant cells. a, Developmental trajectory of alveolar and malignant cells colored by inferred pseudotime (top left), cell differentiation status (top middle), and cell type (top right). Distribution of inferred pseudotime (bottom left) and CytoTRACE (bottom middle) across the indicated cell subsets. CytoTRACE score distribution in KACs at the two time points. EOE: end of exposure (bottom right). 7 mo: 7 months post exposure to NNK.
Box, median ± interquartile range; whiskers, 1.5× interquartile range; center line: median. **b,** Normalized expression level of differentially expressed genes in AT1, KACs, malignant cells and AT2 cells. **c,** Expression levels of representative KAC marker genes across 4 indicated cell subsets. Box, median ± interquartile range; whiskers, 1.5× interquartile range; center line: median.. **d,** Significantly enriched functional pathways of marker genes of KACs. **e,** Enrichment of KAC signature score, defined using marker genes of KACs, across the four cell subsets. ***: FDR q-value < 0.001; *: FDR q-value < 0.05. P-values were calculated using Mann-Whitney U test. Boxes definitions are the same as in panel b. **f,** Fraction of genes belonging to human KAC signature among mouse KAC genes versus in genome background. *P*-value was calculated using Fisher's exact test. **g,** Proportions and average expression levels of mouse orthologs of selected human KAC marker genes in murine KACs and other alveolar cells. **h,** Proposed model for alveolar plasticity dynamics in lung cancer whereby a subset of the AICs in the intermediate AT2-to-AT1 differentiation state can evolve into KACs and later, lead to LUAD development.